



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

1993, 4th Annual JWP Conference

May 8th, 9:30 AM - 4:30 PM

Parental Investment in the House Wren (*Troglodytes aedon*)

Doug Drexler

Illinois Wesleyan University

John Cavitt

Illinois State University

R. Given Harper, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Drexler, Doug; Cavitt, John; and Harper, Faculty Advisor, R. Given, "Parental Investment in the House Wren (*Troglodytes aedon*)" (1993). *John Wesley Powell Student Research Conference*. 41.

<http://digitalcommons.iwu.edu/jwprc/1993/posters/41>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

PARENTAL INVESTMENT IN THE
HOUSE WREN (*Troglodytes aedon*)

Doug Drexler, Dept. of Biology, IWU,
John Cavitt (Illinois State University), and R. Given Harper*

The parental investment hypothesis of resource utilization (Trivers 1972) states that a parent will maximize his or her lifetime reproductive fitness by using the most energy and time efficient method available in the process of reproduction. According to the parental investment hypothesis, if the parents and offspring are subjected to food shortages, the feeding activity of both parents will be necessary to maximize their reproductive success. In contrast, in a polygynous avian species such as the House Wren (*Troglodytes aedon*), if food is abundant during the nestling stage, the male should increase his reproductive success by seeking out extra-pair copulations or acquiring another mate.

The parental investment hypothesis was tested in House Wrens by simultaneously manipulating the brood size (the number of offspring in a nest) and the food availability to simulate conditions of food shortage and surplus. House Wren broods were manipulated to one of five treatments by adding or subtracting nestlings from the brood, and by the addition of a supplemental food source. Nestwatches were conducted on the experimental nests to determine adult feeding behavior and to determine the extent to which supplemental food was utilized.