



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2009, 20th Annual JWP Conference

Apr 18th, 11:00 AM - 12:00 PM

Observation of a Low-Lying Neutron Unbound State in ^{25}F

Alison Smith, '10

Illinois Wesleyan University

Nathan Frank, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Alison Smith, '10 and Nathan Frank, Faculty Advisor, "Observation of a Low-Lying Neutron Unbound State in ^{25}F " (April 18, 2009). *John Wesley Powell Student Research Conference*. Paper 2.
<http://digitalcommons.iwu.edu/jwprc/2009/oralpres7/2>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation O7.2

OBSERVATION OF A LOW-LYING NEUTRON UNBOUND STATE IN ^{25}F

Alison Smith and Nathan Frank*
Physics Department, Illinois Wesleyan University

A neutron unbound state from ^{25}F has been observed for the first time. Prior experimental data on neutron-rich fluorine isotopes indicate differences for bound excited states as compared to nuclear shell model predictions. The ^{25}F isotopes were produced by one-proton removal from an 86 MeV/u ^{26}Ne beam on a Beryllium target at the fast-fragmentation radioactive beam facility of the National Superconducting Cyclotron Laboratory at Michigan State University. The subsequent decay of the ^{25}F isotopes resulted in ^{24}F and neutrons which were detected in coincidence. The charged particles were bent away from the neutrons with a superconducting magnet and analyzed in a suite of charged particle detectors. The neutrons were detected in the Modular Neutron Array (MoNA). The properties of the charged particles and neutrons were used to reconstruct a decay energy spectrum for ^{25}F which was compared to simulations. Preliminary results indicating a resonant state at approximately 30 keV will be discussed.

Work supported by National Science Foundation Grants PHY-0606007 and PHY-055445