



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2002, 13th Annual JWP Conference

Apr 21st, 1:15 PM - 2:15 PM

Synthesis of Hexamolybdate Complexes with Difunctional Amines

David H. Aggen, '05

Illinois Wesleyan University

Rebecca Roesner, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

David H. Aggen, '05 and Rebecca Roesner, Faculty Advisor, "Synthesis of Hexamolybdate Complexes with Difunctional Amines" (April 21, 2002). *John Wesley Powell Student Research Conference*. Paper 25.
<http://digitalcommons.iwu.edu/jwprc/2002/posters3/25>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P2

SYNTHESIS OF HEXAMOLYBDATE COMPLEXES WITH DIFUNCTIONAL AMINES

David H. Aggen and Rebecca Roesner
 Department of Chemistry, Illinois Wesleyan University

Polyoxometalates, highly symmetric transition metal oxide clusters, are known to exhibit anti-viral activity. Modification of polyoxometalate compounds, through the addition of pendant organic groups, may enable the pharmaceutical targeting of diseased cells, which express specific biological macromolecules. Our research involves the synthesis of a naphthalene-hexamolybdate adduct from two equivalents of the hexamolybdate ion ($\text{Mo}_6\text{O}_{18}^{2-}$) and 1 equivalent of 1,5-diaminonaphthalene. Dicyclohexylcarbodiimide was utilized to facilitate the reaction, producing dicyclohexylurea as a byproduct.¹ This reaction was repeated several times, using different conditions, and the products were characterized by ^1H NMR spectroscopy. Downfield shifts of the aromatic, naphthalene resonances in these spectra suggest that the desired molybdenum-nitrogen bonds are forming. Yield and purity have been variable. Attempts to crystallize polyoxometalate compounds of this kind have produced spiny crystals, unsuitable for x-ray diffraction. We hope to explore other methods of purification and crystallization as our research continues.

