



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2000, 11th Annual JWP Conference

Apr 15th, 11:00 AM - 12:30 PM

A Survey of Organochlorine Pesticide Contamination in a Costa Rican Conservation Area

Mark Wieland

Illinois Wesleyan University

R. Given Harper, Faculty Advisor

Illinois Wesleyan University

Jeffrey A. Frick, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Mark Wieland; R. Given Harper, Faculty Advisor; and Jeffrey A. Frick, Faculty Advisor, "A Survey of Organochlorine Pesticide Contamination in a Costa Rican Conservation Area" (April 15, 2000). *John Wesley Powell Student Research Conference*. Paper 1.

<http://digitalcommons.iwu.edu/jwprc/2000/oralpres3/1>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation 3.5

A SURVEY OF ORGANOCHLORINE PESTICIDE CONTAMINATION IN A COSTA RICAN CONSERVATION AREA

Mark Wieland and R. Given Harper* and Jeffrey A. Frick*
Departments of Biology and Chemistry, Illinois Wesleyan University

Amphibians, turtles, rodents, and birds collected from a tropical conservation area in northwestern Costa Rica, where pesticides have not been directly applied, were analyzed for organochlorine (OC) pesticide contamination. Six of thirty-nine amphibians (three of eight species), three of six turtles (two species), one of eight rodents (one species), and nine of twenty-five birds (four species) contained OCs ranging from 2.77 ng/g to 277.70 ng/g in individual organisms. The most frequently detected compound (in thirteen organisms) was p,p'-DDE. Heptachlor, delta-BHC, dieldrin, endosulfan II, and p,p'-DDD were found in four or more organisms, while eight other OCs were found in one, two, or three organisms. The average body mass of contaminated amphibians was 156.40 g, compared to 56.89 g for uncontaminated amphibians, suggesting that increased body mass is indicative of greater susceptibility to contamination. The presence of OCs in taxa from the conservation area indicates the likelihood of long-distance transport of pesticides through the atmosphere. These contaminants may affect interactions between organisms in the tropical conservation area.