



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2006, 17th Annual JWP Conference

Apr 8th, 2:00 PM - 3:35 PM

Fisher Esterification of a Rhodium Substituted Keggin-Type Polytungstophosphate

Jeffrey W. Donelan

Illinois Wesleyan University

Rebecca Roesner, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Jeffrey W. Donelan and Rebecca Roesner, Faculty Advisor, "Fisher Esterification of a Rhodium Substituted Keggin-Type Polytungstophosphate" (April 8, 2006). *John Wesley Powell Student Research Conference*. Paper 32.
<http://digitalcommons.iwu.edu/jwprc/2006/posters2/32>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P18

**FISHER ESTERIFICATION OF A RHODIUM SUBSTITUTED KEGGIN-TYPE
POLYTUNGSTOPHOSPHATE**

Jeffrey W. Donelan and Rebecca Roesner*
Chemistry Department, Illinois Wesleyan University

Polyoxometalate (POM) chemistry is an emerging field with many applications in catalysis, electrochemistry, and medicine. Keggin-type POMs have the formula $[XM_{12}O_{40}]^n$, where M is an early transition metal and X can be almost any element. In recent years, chemists have become interested in attaching organic ligands to POMs for the purpose of creating immobilized polyoxoanion catalysts.¹ In our present work, a rhodium substituted Keggin-type POM with a pendent carboxylic functional group ($[PW_{11}O_{39}RhCH_2COOH]^{5-}$) was synthesized by previously reported methods.^{1,2} This POM was then converted to its methyl and ethyl ester derivatives through Fisher esterification with the appropriate alcohols. The ester products were then characterized by NMR and IR spectroscopies. A further effort is being made to immobilize the carboxylic acid functionalized polytungstophosphate through Fisher esterification with alcohol functionalized resin beads.

1. Wei, X. Dickman, M.H.; Pope, M.T. *J. Am. Chem. Soc.* **1998**, *120*, 10254-10255.
2. Unpublished results. Binder, J., Roesner, R. Illinois Wesleyan University.