



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2006, 17th Annual JWP Conference

Apr 8th, 9:00 AM - 10:00 AM

The Analysis of Amino Acid Substitutions at Positions ASP22 of FNR

Justin Ernat

Illinois Wesleyan University

Aaron Bailey

Illinois Wesleyan University

Laura Moore, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Justin Ernat; Aaron Bailey; and Laura Moore, Faculty Advisor, "The Analysis of Amino Acid Substitutions at Positions ASP22 of FNR" (April 8, 2006). *John Wesley Powell Student Research Conference*. Paper 1.
<http://digitalcommons.iwu.edu/jwprc/2006/posters/1>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P3

**THE ANALYSIS OF AMINO ACID SUBSTITUTIONS AT
POSITIONS ASP22 OF FNR**

Justin Ernat and Aaron Bailey and Laura Moore*
Chemistry Department, Illinois Wesleyan University

The FNR protein is an oxygen-sensing transcription factor in the facultative anaerobe, *Escherichia coli*. In wild-type FNR, a [4Fe-4S] cluster is incorporated into the protein. This results in a protein that dimerizes and can bind to specific sites in the DNA. In the presence of oxygen, this cluster degrades causing a conformational change that renders the protein inactive. The Fe-S cluster is ligated by cysteine residues at positions 20, 23, 29, and 122. Our work focuses on the effects on protein activity of amino acid substitutions at position Asp22. We studied the aerobic and anaerobic activity of the mutations Asp22Ala, Asp22Ser, Asp22Trp, Asp22Asn and Asp22Gly using Beta-galactosidase assays. Of these mutations Asp22Gly has been previously reported to be aerobically active, and it was suggested that this substitution stabilized the cluster to oxygen. Currently, under the low protein concentrations of our Beta-galactosidase assays, none substitutions have shown aerobic activity. We have also isolated Asp22Ala and Asp22Gly FNR protein. The Asp22Ala FNR protein has been characterized using absorption spectroscopy, which shows that the [4Fe-4S] cluster is not stable to the presence of oxygen. Further research will include isolation of other Asp22 FNR proteins as well as performing Beta-galactosidase assays with a higher concentration of protein.