



## Illinois Wesleyan University Digital Commons @ IWU

---

John Wesley Powell Student Research  
Conference

1998, 9th Annual JWP Conference

---

Apr 18th, 9:00 AM - 10:30 AM

# Synthesis of a Fully Conjugated Linear Diamine as a Potential Rotaxane Component

Matthew T. Lewellen

*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

---

Matthew T. Lewellen and Rebecca Roesner, Faculty Advisor, "Synthesis of a Fully Conjugated Linear Diamine as a Potential Rotaxane Component" (April 18, 1998). *John Wesley Powell Student Research Conference*. Paper 13.  
<http://digitalcommons.iwu.edu/jwprc/1998/posters/13>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation 13

SYNTHESIS OF A FULLY CONJUGATED LINEAR DIAMINE AS A POTENTIAL ROTAXANE COMPONENT

Matthew T. Lewellen and Rebecca A. Roesner\*  
 Department of Chemistry, Illinois Wesleyan University

Steps toward the preparation of a fully conjugated linear diamine are described. This diamine will be used as a linear component in the preparation of larger supramolecular species called rotaxanes. Rotaxanes are composed of a ring shaped molecule threaded onto a linear molecule with bulky stoppering groups used to prevent dethreading of the ring (Figure 1). The target rotaxane will be composed of two hexamolybdate ions, a beta-cyclodextrin molecule, and a linear diamine. In the first step of the diamine synthesis, *p*-aminotrimethylsilylethynylbenzene was prepared from 4-iodoaniline and trimethylsilyl acetylene according to Figure 2. The trimethylsilyl protecting group allows substitution to occur at only one C-H terminus of the acetylene molecule. In a subsequent step (Figure 3) the trimethylsilyl protecting group was removed from the *p*-aminotrimethylsilylethynylbenzene by addition of aqueous potassium hydroxide. In a third step, two equivalents of the newly formed *p*-aminoethynylbenzene will be coupled together to form 4,4'-bis(amino)diphenylbutadiyne (Figure 4).




Figure 1




Figure 2




Figure 3




Figure 4