



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2002, 13th Annual JWP Conference

Apr 21st, 1:15 PM - 2:15 PM

Complementation of *bchE* *R. Capsulatus* from *chlE* from the Cyanobacteria *Synechocystis* sp. 6803

Kristin Tracy, '03

Illinois Wesleyan University

David Bollivar, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Tracy, '03, Kristin and Bollivar, Faculty Advisor, David, "Complementation of *bchE* *R. Capsulatus* from *chlE* from the Cyanobacteria *Synechocystis* sp. 6803" (2002). *John Wesley Powell Student Research Conference*. 3.
<http://digitalcommons.iwu.edu/jwprc/2002/posters3/3>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P48

COMPLEMENTATION OF *BCH^E* IN *R. CAPSULATUS* WITH *CHL^E* FROM THE
CYANOBACTERIA *SYNECHOCYSTIS SP. 6803*

Kristin Tracy and David Bollivar*
Department of Biology, Illinois Wesleyan University

The bacterium *Rhodobacter capsulatus* has the ability to live both aerobically and anaerobically, and it uses photosynthesis to survive anaerobic conditions. One gene essential for photosynthesis in this organism is *bchE*, which codes for an enzyme in the pathway responsible for making bacteriochlorophyll. The cyanobacterium *Synechocystis sp. PCC 6803* has the related gene *chlE*, which codes for an enzyme involved in the chlorophyll biosynthesis pathway. In this experiment, the functional relationship of these two genes was tested through complementation. A plasmid was constructed that contained the *Synechocystis chlE* gene attached to an S-tag sequence, and the plasmid was then transformed into a strain of *E. coli* (S17-1/_ pir). This new strain of *E. coli* containing the *chlE* plasmid was then mated with a strain of *R. capsulatus* that had been engineered to lack a functional *bchE* gene and thus could not grow anaerobically. The transfer of the *chlE* plasmid and its ability to complement the non-functional *bchE* gene was demonstrated by the ability of the daughter cells that resulted from the mating (the ChlEexp strain) to grow anaerobically, as well as the presence of an S-tagged band of the appropriate size on a Western blot of cell extracts. As of yet, enzymatic assays have shown no activity in cell extracts, but from the data collected so far we can see that the cyanobacterial chlorophyll gene *chlE* has the ability to replace the function of the bacteriochlorophyll gene *bchE* in *R. capsulatus*.