



## Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research  
Conference

2006, 17th Annual JWP Conference

Apr 8th, 2:35 PM - 3:35 PM

# In Vivo Demonstration of Protein-Protein Interactions between Herpes Simplex Virus Type-1 Proteins UL20 and GK during Viral Egress from Infected Cells

Rachael Liesman

*Illinois Wesleyan University*

Elizabeth Balser, Faculty Advisor

*Illinois Wesleyan University*

Konstantin Kousoulas, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Rachael Liesman; Elizabeth Balser, Faculty Advisor; and Konstantin Kousoulas, Faculty Advisor, "In Vivo Demonstration of Protein-Protein Interactions between Herpes Simplex Virus Type-1 Proteins UL20 and GK during Viral Egress from Infected Cells" (April 8, 2006). *John Wesley Powell Student Research Conference*. Paper 19.  
<http://digitalcommons.iwu.edu/jwprc/2006/posters2/19>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P44

**IN VIVO DEMONSTRATION OF PROTEIN-PROTEIN INTERACTIONS  
BETWEEN HERPES SIMPLEX VIRUS TYPE-1 PROTEINS UL20 AND GK  
DURING VIRAL EGRESS FROM INFECTED CELLS**

Rachael Liesman and Elizabeth Balsler\* and Konstantin Kousoulas\*  
Biology Department, Illinois Wesleyan University

Recent evidence suggests a functional relationship between Herpes Simplex virus type-1 membrane proteins UL20 and gK. UL20 and gK likely interact to permit viral exodus from an infected cell, a process necessary for further viral infection of surrounding cells. We sought to demonstrate UL20 and gK interactions using a split-ubiquitin yeast two-hybrid system. Contrasting with traditional yeast two-hybrid experiments, the split-ubiquitin yeast two-hybrid system allows us to investigate interactions between integral membrane proteins such as gK and UL20. Confocal microscopy was also used to determine the intracellular localization of UL20 and gK. UL20 has been successfully fused into the bait vector and shown, through interactions with a control prey vector, to be properly inserted into the membrane. UL20 and gK have been shown to co-localize to the trans-golgi network (TGN) by confocal microscopy.

Continued study and increased understanding of viral patterns within infected cells can lead to better treatment and prevention of viral infections, as well as the use of HSV-1 as a medicinal vector to treat genetically-based disease.