



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2001, 12th Annual JWP Conference

Apr 21st, 11:00 AM - 12:30 PM

The Zoomin' HOTs: Dynamic Holographic Optical Tweezing

M. Bradley Taylor

Illinois Wesleyan University

Alexander J. Laurie

Illinois Wesleyan University

Gabriel C. Spalding, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Taylor, M. Bradley; Laurie, Alexander J.; and Spalding, Faculty Advisor, Gabriel C., "The Zoomin' HOTs: Dynamic Holographic Optical Tweezing" (2001). *John Wesley Powell Student Research Conference*. 4.
<http://digitalcommons.iwu.edu/jwprc/2001/oralpres2/4>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation 2.4

THE ZOOMIN' HOTs: DYNAMIC HOLOGRAPHIC OPTICAL TWEEZING

M. Bradley Taylor and Alexander J. Laurie and Gabriel C. Spalding*

Department of Physics, Illinois Wesleyan University

A new experimental technique, Holographic Optical Tweezing (H.O.T.), has been developed through a collaboration between our group at IWU and researchers at the University of Chicago.¹ Our addition of diffractive optics to earlier techniques for optical trapping significantly enhances the ability to control the assembly of biological cells or microscopic dielectric particles. In this new work, we describe original designs for optical systems using static holograms that nevertheless allow dynamic control over the number of traps in an assembly, their configuration/disorder (via spatial filtering), and even their spacings (via carefully designed zoom optics), as well as the strength of the traps (via the laser intensity). Although there are many possible applications for this new technology, as physicists our first interest is in exploring collective effects in optical binding, and in creating model systems that afford us many tunable parameters with which to explore many-body interactions. We are also interested in the onset of symmetry-breaking behavior (e.g., avalanches and the possibility of self-organized criticality, and the microscopic mechanisms underlying phase transitions in two and three dimensions).

¹Eric R. Dufresne, Gabriel C. Spalding, Matthew T. Dearing, Steven A. Sheets, David G. Grier, *Review of Scientific Instruments* 72, 1810 (2001).