



## Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research  
Conference

2008, 19th Annual JWP Conference

Apr 12th, 2:35 PM - 3:35 PM

# Synthesis of 30-Membered Azamacrocycles as Hosts for Polyoxoanion Guests

Melissa Myers

*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Melissa Myers and Rebecca Roesner, Faculty Advisor, "Synthesis of 30-Membered Azamacrocycles as Hosts for Polyoxoanion Guests" (April 12, 2008). *John Wesley Powell Student Research Conference*. Paper 18.  
<http://digitalcommons.iwu.edu/jwprc/2008/posters2/18>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

Poster Presentation P36

SYNTHESIS OF 30-MEMBERED AZAMACROCYCLES AS HOSTS FOR  
POLYOXOANION GUESTS

Melissa Myers and Rebecca Roesner\*  
Chemistry Department, Illinois Wesleyan University

The long term goal of this project is to create a capsule-shaped host molecule through the linkage of two macrocyclic compounds. This capsule is to be built around a polyoxometallate, a large, anionic cluster. Since most polyoxometallates decompose under basic condition, it may be possible to break down the polyoxoanion scaffold after the capsule is assembled. Size compatibility of the polyoxometallate and the macrocycles is essential, as these components must preferentially complex each other. Previous work on this project utilized fourteen membered macrocycles, which proved somewhat small to effectively complex with the polyoxometallate. Current work involves synthesis of much larger 30-membered macrocycles with the hope that they are more likely to provide a proper fit. In addition to steric factors, charge also plays an important role in determining whether the macrocycle and polyoxoanion will form an inclusion complex. Two different 30-membered rings are being investigated currently, one with four amine groups and the other with six. In acidic solution, the amine groups should form positive ammonium ions ideal for binding to the negatively charged polyoxometallate.

