



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2016, 27th Annual JWP Conference

Apr 16th, 9:00 AM - 10:00 AM

A Transposon Mutagenesis Screen for Heterocyst Production in a $\Delta HETP$ Strain of the Cyanobacterium *Anabaena*

Rachel Ende

Illinois Wesleyan University

Loralyn Cozy, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

 Part of the [Biology Commons](#), and the [Education Commons](#)

Ende, Rachel and Cozy, Faculty Advisor, Loralyn, "A Transposon Mutagenesis Screen for Heterocyst Production in a $\Delta HETP$ Strain of the Cyanobacterium *Anabaena*" (2016). *John Wesley Powell Student Research Conference*. 8. <http://digitalcommons.iwu.edu/jwprc/2016/posters/8>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Poster Presentation P17

**A TRANSPOSON MUTAGENESIS SCREEN FOR HETEROCYST
PRODUCTION IN A $\Delta HETP$ STRAIN OF THE
CYANOBACTERIUM *ANABAENA***

Rachel Ende and Loralyn Cozy*
Biology Department, Illinois Wesleyan University

The ability for organisms to genetically control cellular differentiation is a fundamental question of development. The multicellular cyanobacterium *Anabaena* maintains the ability for vegetative cells to differentiate into specialized heterocysts capable of nitrogen fixation. The genetic network responsible for initiating and regulating this differentiation is not fully understood. To identify genes within this differentiation network, a heterocysts deficient mutant of *Anabaena* was subjected to transposon mutagenesis. A transposon, a piece of DNA capable of random insertion into the genome was introduced into this heterocysts deficient strain. Following the introduction of the transposon, the cells were grown on nitrogen-limited media containing antibiotic to select for cells in which the transposon had inserted into the genome and restored heterocyst function. These colonies were then analyzed with light microscopy to characterize the amount and pattern of heterocysts development and inverse PCR was performed to identify the specific location of transposon insertion within the genome. The genes identified through this screen will be the subject of future research.