

**Efficacy of intermittent cooling strategies during tennis-specific treadmill exercise in hot, humid conditions
[abstract only]**

DEBNEY, Matthew, RUDDOCK, Alan <<http://orcid.org/0000-0002-7001-9845>>, O'HAGAN, Ciara and PURVIS, Alison <<http://orcid.org/0000-0002-3581-4990>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/32844/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

DEBNEY, Matthew, RUDDOCK, Alan, O'HAGAN, Ciara and PURVIS, Alison (2023). Efficacy of intermittent cooling strategies during tennis-specific treadmill exercise in hot, humid conditions [abstract only]. *Journal of Sports Sciences*, p. 26.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Efficacy of intermittent cooling strategies during tennis-specific treadmill exercise in hot, humid conditions

DR MATTHEW J. DEBNEY¹, DR ALAN RUDDOCK¹, DR CIARA J. O'HAGAN², DR ALISON J. PURVIS¹

¹Sheffield Hallam University, Sheffield, United Kingdom; ²Institute of Technology, Carlow, Ireland

Environmental conditions $\geq 30^{\circ}\text{C}$ wet-bulb globe temperature (WBGT) increases thermoregulatory and physiological demands for tennis players increasing the risk of heat related medical callouts (Smith et al. 2018, *Journal of Science and Medicine in Sport*, 21(5), 467-472). The aim of this study was to assess the efficacy of cooling strategies in alleviating thermal strain during simulated tennis match-play in a hot environment. Following institutional ethics approval, nine healthy, male amateur tennis players (mean \pm SD: age 26 ± 5 years, stature 180 ± 45 cm, body mass 77.7 ± 6.9 kg; $\dot{V}\text{O}_2\text{max}$ 52 ± 6 ml \cdot kg $^{-1}$ \cdot min $^{-1}$) completed the tennis-specific treadmill protocol (TSTP) (Debney et al. 2018, *Journal of Sport Sciences*, 36:sup1, 1-94) five times in hot, humid conditions ($36.3 \pm 0.7^{\circ}\text{C}$, $51.3 \pm 1.5\%$, $30.8 \pm 0.4^{\circ}\text{C}$ WBGT): cold water (CW); ice slurry (SLU); ice towel (TWL); ice towel and slurry (TWL+SLU); and a control (CTL). After every odd game, participants ingested 2 g \cdot kg $^{-1}$ thermoneutral water in CTL and TWL ($34.8 \pm 0.4^{\circ}\text{C}$); cold water ($10.7 \pm 2.5^{\circ}\text{C}$); or ice slurry (-0.5°C) in SLU and TWL+SLU. When appropriate, the ice towel was worn until 30 s before the next game beginning. After games 3, 5, 7 and 9 in all 3 sets of the TSTP, rectal temperature (T_c), mean skin temperature (T_{sk}), and thermal sensation (TS) were assessed; perceptual responses were assessed again after the rest. A three-way ANOVA (condition \times time (set) \times time (game)), with Bonferroni post-hoc analysis was used to assess statistically significant differences ($P < 0.05$) between conditions. Hedges g was used to assess effect size. There were no statistically significant differences between conditions until set 3, when thereafter, there was a lower T_c in SLU ($P = 0.009$, $g = -0.83$) and TWL+SLU ($P = 0.014$, $g = -1.07$) compared to CTL. T_c ($P = 0.034$, $g = -0.66$) and T_{sk} ($P = 0.007$, $g = -1.10$) were lower in TWL+SLU than CW. TS was lower after rest periods in all experimental conditions ($P < 0.05$) compared to CTL. Importantly, participants felt cooler before the rest period in TWL+SLU ($P = 0.002$, $g = -0.80$) compared to CTL. Combined cooling (TWL + SLU) was the most efficacious strategy, attenuating both physiological and perceptual thermal strain. If only one strategy can be implemented, internal cooling using SLU is likely more effective than external cooling at attenuating thermal strain during tennis-specific exercise in hot humid conditions.