



# Uncertainties in, and interpretation of, carbon flux estimates using the eddy covariance technique

H. W. Loescher,<sup>1</sup> B. E. Law,<sup>1</sup> L. Mahrt,<sup>2</sup> D. Y. Hollinger,<sup>3</sup> J. Campbell,<sup>1</sup> and S. C. Wofsy<sup>4</sup>

Received 29 November 2005; revised 18 April 2006; accepted 15 June 2006; published 14 November 2006.

[1] Whole ecosystem carbon dioxide (CO<sub>2</sub>) exchange estimated with the eddy covariance (EC) technique has been central to studies on the responses of terrestrial ecosystems to disturbance and intra-annual and interannual variations in climate, but challenges exist in understanding and reducing the uncertainty in estimates of net ecosystem exchange (NEE) of CO<sub>2</sub>. We review the potential uncertainties associated with the eddy covariance technique, including systematic errors from insensitivity to high-frequency turbulence, random errors from inadequate sample size associated with averaging period, vertical and horizontal advection issues, and selection criteria for removing periods of inadequate mixing from further analyses. We also discuss benefits and caveats of using independent measurements to evaluate EC-derived NEE, such as comparisons of EC-derived annual NEE and allometric net ecosystem production estimates (NEP) and interpretation of nighttime NEE with scaled chamber-based estimates of ecosystem respiration.

**Citation:** Loescher, H. W., B. E. Law, L. Mahrt, D. Y. Hollinger, J. Campbell, and S. C. Wofsy (2006), Uncertainties in, and interpretation of, carbon flux estimates using the eddy covariance technique, *J. Geophys. Res.*, *111*, D21S90, doi:10.1029/2005JD006932.

## 1. Introduction

[2] The use of eddy covariance (EC) has improved our understanding of temporally and spatially integrated net ecosystem exchange rates (NEE) of CO<sub>2</sub> between ecosystems and the atmosphere, identifying key biotic and abiotic processes that control these rates [Loescher *et al.*, 2003; Bowling *et al.*, 2001; Katul *et al.*, 1998; Lee, 1998; Goulden *et al.*, 1996; Grace *et al.*, 1996; Hollinger *et al.*, 1994] and improving ecosystem process models [e.g., Thornton *et al.*, 2002; Law *et al.*, 2002; Williams *et al.*, 2001; Aber *et al.*, 1996]. As the EC technique matures, efforts are focused on reducing uncertainty in NEE estimates.

[3] The EC technique was pioneered over grass and croplands with long fetch and short roughness lengths [Kaimal and Wyngaard, 1990; Verma *et al.*, 1989; Lemon, 1960; Monteith and Szeicz, 1960]. Researchers have since applied this approach over structurally complex ecosystems in nonideal terrain, introducing new challenges in the interpretation of results and reduction of uncertainties. The EC technique is a direct, nondestructive micrometeorological approach derived through the simplification of the conservation equation [Baldocchi, 2003; Baldocchi and

Meyers, 1998; Baldocchi *et al.*, 2000; Shen and Leclerc, 1997]. EC is used to estimate NEE through the addition of above-canopy turbulent exchange and the change in CO<sub>2</sub> storage in the canopy air space (i.e., the temporal change in carbon concentration integrated from ground level to the point of measured turbulent exchange, term I, equation (1a)):

$$NEE_c = \underbrace{\int_0^Z \frac{\partial \bar{c}}{\partial t} dz}_I + \underbrace{\int_0^Z \frac{\partial \overline{w'c'}}{\partial z} dz}_II + \underbrace{\int_0^Z \frac{\partial \overline{u'c'}}{\partial x} dz}_III + \underbrace{\int_0^Z \frac{\partial \overline{v'c'}}{\partial y} dz}_IV + \underbrace{\int_0^Z \frac{\partial \overline{u\bar{c}}}{\partial x} dz}_V + \underbrace{\int_0^Z \frac{\partial \overline{v\bar{c}}}{\partial y} dz}_{VI} + \underbrace{\int_0^Z \frac{\partial \overline{w\bar{c}}}{\partial z} dz}_{VII} \quad (1a)$$

$$NEE_c = \underbrace{\int_0^Z \frac{\partial \bar{c}}{\partial t} dz}_I + \overline{w'c'}(z=Z) = \underbrace{\int_0^Z \frac{\partial \bar{c}}{\partial t} dz}_I + \underbrace{\overline{w'c'}}_{II} \quad (1b)$$

where  $c$  is the scalar quantity such as CO<sub>2</sub>,  $u$ ,  $v$ , and  $w$  are horizontal, cross-wind, and vertical windspeeds (m s<sup>-1</sup>), respectively,  $x$ ,  $y$ , and  $z$  are Cartesian coordinates,  $t$  is time, and  $Z$  is the measurement height (m), the overbars indicate a time average and prime denotes turbulent fluctuations (i.e., deviations from a mean quantity). Term I in equation (1a) represents the time rate-of-change of  $c$  in the vertical column (i.e., storage), and is considered to be equal to zero over a 24-hour period, but can be significant over shorter time intervals. Terms II–IV represent the turbulent flux

<sup>1</sup>Department of Forest Science, College of Forestry, Oregon State University, Corvallis, Oregon, USA.

<sup>2</sup>College of Oceanic and Atmospheric Science, Oregon State University, Oregon, USA.

<sup>3</sup>U.S. Forest Service, U.S. Department of Agriculture, Durham, New Hampshire, USA.

<sup>4</sup>Division of Engineering and Applied Science, Harvard University, Cambridge, Massachusetts, USA.

divergence. Terms V–VII represent advection through the layer between the surface and sensor. The partial derivatives usually are estimated from data at points in the domain, and thus are replaced here by finite differences. Assuming that the measurement height is sufficient and the surface characteristics are horizontally homogeneous, terms III and IV are often thought to be 0 and ignored. Terms IV–VII are inherently difficult to measure well, but thought to be small and thus not often estimated. Hence only terms I and II remain. Term II in equation (1b) is commonly referred to as the eddy covariance (additional assumptions in its estimation are discussed later). *NEE* is typically estimated over a 30-min period from high-frequency measurements, and integrated further to estimate daily, season and annual fluxes. The relative importance in the sources of error differs across timescales, spatial extent and site conditions. For example, for *NEE* at minute-to-hour timescales (compare with photosynthetic uptake and respiratory efflux), systematic and random errors associated with instrumentation precision, calibration and placement, and turbulent transport become significant. In scaling *NEE* from day-to-season (months), random errors are reduced, but other errors are added because of filtering and filling gaps in data. Consequently, the sources of error in applying EC to understand rapid biological response to meteorological variables are quite different from those errors associated with determining the environmental controls on phenology, annual carbon exchange, effects from disturbance, or whether mature forested ecosystems continue to accumulate carbon. Of course the relative contribution of all errors becomes large in either example (i.e., minute-to-hour and day-to-season) when the mean flux is close to 0.

[4] Some of these errors or uncertainties stem from flows not fully accounted for within and below the sensor field, which can result in potential violation of the assumptions. Other uncertainties result from the stochastic nature of turbulence. A key stratagem of the long-term flux network, AmeriFlux, is to assure accurate estimates within and among flux sites for synthesis activities and regional analysis. The AmeriFlux science plan has outlined rigorous quality control protocols for within site measurements (see <http://public.ornl.gov/ameriflux/About/scif.cfm>) to avoid large systematic biases so that subtle spatial and temporal trends may be discerned. Quality assurance across the network is assessed through direct comparison of software routines and instrumentation. Using an independent raw data file developed by the Euroflux and AmeriFlux networks (“gold files” for closed- and open-path infrared gas analyzers can be found at <http://public.ornl.gov/ameriflux/standards-gold.shtml>), researchers can process flux data sets through their own software routine and check estimates against a standard processed file. Site-to-site differences among instrument configurations are estimated by an independent portable flux measurement system that visits each site. Consistency and rigor in sample design, analysis, diagnostics, and data quality checks help to ensure data are comparable across sites.

[5] In this paper, we review the current progress in estimating ecosystem level carbon exchange using the eddy covariance approach across different temporal scales, and suggest future research directions that can be enhanced by the EC technique. Explicit discussion of instrumentation

errors and calibrations are not covered here, and are given by *Foken and Oncley* [1995], *Massman and Lee* [2002], and *Foken et al.* [2004], and related discussions on surface energy balance closure are given by *Aubinet et al.* [2000], *Mahrt* [1998], *Twine et al.* [2000], and *Foken et al.* [2004].

## 2. Challenges in Measurements, Analyses, and Interpretation of EC Fluxes


### 2.1. Flux Measurement

[6] Estimates of the total CO<sub>2</sub> flux from a vegetated canopy should be made in the surface layer, above the roughness sublayer, the depth of which changes with stability [*Raupach*, 1991; *Raupach and Thom*, 1981]. In the surface layer, sometimes referred to as the inertial layer, Monin-Obukhov similarity theory applies in homogenous stationary atmospheric conditions [e.g., *Kaimal and Finnigan*, 1994]. The level close to the surface where the winds asymptotically become zero by extrapolating Monin-Obukhov similarity theory downward is  $d + z_0$ , where  $d$  is the zero-plane displacement and  $z_0$  is the characteristic roughness length [see *Panofsky and Dutton*, 1984]. Combined,  $d$  and  $z_0$  determine the bottom boundary of the surface layer and  $d$  can vary with changes in stability [*Loescher et al.*, 2003]. Consequently,  $d$  and  $z_0$  are not easily estimated for forest canopies and are not formally defined for below-canopy environments. Estimates of  $z_0$  range from 0.01 m for grass [*Hansen*, 1993] to >2.4 m over tropical forests [*Loescher et al.*, 2003].

[7] The use of eddy covariance relies on conditions which induce turbulence in the surface layer (i.e., shear stress, surface heating), and sometimes these conditions are weak or absent. Nighttime conditions can be particularly problematic as surface heating is absent and radiational cooling produces stable thermal stratification of the air column that acts to suppress shear induced turbulence. The measurements above the canopy may become partially decoupled from the surface and the computed fluxes become sensitive to the method of calculation. Researchers have, by necessity, found that empirical relationships between the friction velocity ( $u^*$ ) and scalar fluxes can determine conditions that can assist in the estimation of robust turbulent exchanges in 30-min averages. The  $u^*$  is normally defined as [cf. *Weber*, 1999],

$$u^* = \left[ (\overline{w'u'})^2 + (\overline{w'v'})^2 \right]^{0.25} \quad (2)$$

[8] Examining the sensitivity in annual integrals to different  $u^*$  thresholds was introduced by *Goulden et al.* [1996] and is now a common method to determine “well-mixed” conditions [cf. *Aubinet et al.*, 2000; *Gu et al.*, 2005]. The impact of accepting data on the basis of different  $u^*$  thresholds is illustrated in annual integrals from Howland Forest, an AmeriFlux site, where the annual sums do not change above a  $u^*$  threshold of  $\sim 0.25 \text{ m s}^{-1}$  (Figure 1a). At other sites, a  $u^*$  threshold however, may not always be as evident. Using 30-min-averaged data for example, the nighttime EC flux continued to increase with increasing  $u^*$  values from a structurally simple and topographically flat cocoa plantation (Figure 1b), suggesting inadequate mixing to fully estimate the respiratory losses using 30-min-averaged turbulent data.



**Figure 1.** The relationship between  $u^*$  threshold and (a) annual integrals for 1999 and 2000 from a coniferous forest at Howland, Maine ( $45^{\circ}12'14''\text{N}$ ,  $68^{\circ}44'25''\text{W}$ ), and (b) nighttime 30-min data from a 13-year-old, even-aged cocoa plantation, Sulawesi, Indonesia ( $1^{\circ}8'27''\text{S}$ ,  $120^{\circ}3'34''\text{E}$ ). Indonesian data were collected from 4.3 m above a 10.3 m canopy, from February 2002 to April 2003, and the site was flat for  $\sim 6$  km in any direction from the tower. Cocoa data were bin averaged across  $u^*$  intervals of  $0.1 \text{ m s}^{-1}$ , and error bars are  $\pm 1$  SD. For additional site and data description for Figure 1a, see *Hollinger et al.* [2004], and for Figure 1b, see *Falk* [2004].

[9] Presently, 80% of flux sites surveyed report the use of a  $u^*$  threshold criteria for acceptance of nocturnal data with values ranging between  $0.0\text{--}0.6 \text{ m s}^{-1}$  (Table 1). The number of nocturnal 30-min periods excluded from further analysis by a  $u^*$  threshold vary from site-to-site. An extreme example is found in the tropics where *Miller et al.* [2004] report that as much as  $\sim 80\%$  of nighttime data are removed depending on the value of the  $u^*$  threshold. The choice of

the threshold value in this case resulted in a  $\sim 5 \text{ t C ha}^{-1} \text{ y}^{-1}$  range in annual *NEE*. The flux community has developed no standardized method to determine  $u^*$  thresholds, although a statistical approach was developed by *Gu et al.* [2005]. Use and applicability of a  $u^*$  threshold to determine carbon flux remains an active area of research, but the general concept of restricting analysis to periods with sufficient atmospheric mixing is sound.

**Table 1.** List of Studies That Discuss the Relationship Between  $u^*$  and Annual NEE<sup>a</sup>

Study	Location	Vegetation type	Year	Annual NEE	$u^*$ Threshold	
<i>Anthoni et al.</i> [1999] <sup>b</sup>	Oregon	ponderosa pine	1996	-3.2	none	
			1997	-2.7	none	
<i>Anthoni et al.</i> [2002] <sup>c</sup>	Oregon	ponderosa pine	2000		$0.1 < u^* < 0.55$	
<i>Barford et al.</i> [2001]	Massachusetts	maple/oak	1993–2000	-2.0	0.2	
<i>Berbigier et al.</i> [2001] <sup>d</sup>	France	pine	1997	-4.3	0.4	
<i>Black et al.</i> [1996] <sup>e</sup>	Saskatchewan	aspen	1994	-1.6	0.15	
			1996	-0.8		
			1997	-1.2		
			1998	-2.9		
<i>Clark et al.</i> [1999]	Florida	slash pine	1996	-7.4	0.2	
			1997	-6.1		
		cypress	1996	-0.8		
			1997	-0.4		
<i>Goulden et al.</i> [1997]	Canada	black spruce	1995	0.7	0.2	
			1996	0.2		
			1997	-0.1		
<i>Goulden et al.</i> [1996]	Massachusetts	maple/oak	1991	-2.8	0.2	
			1992	-2.2		
			1993	-1.4		
			1994	-2.1		
			1995	-2.7		
<i>Grace et al.</i> [1996]	Brazil	tropical humid	1994	-1.0	none	
	France	beech	1996	-2.2	0.00	
<i>Granier et al.</i> [2000] <sup>f</sup>			1997	-2.6	0.00	
<i>Hollinger et al.</i> [1999] <sup>g</sup>	Maine	mixed conifer	1996	-2.1	0.15	
			<i>Hollinger et al.</i> [2004]	1996		0.25, 0.1–0.5
				1997		0.25, 0.1–0.5
				1998		0.25, 0.1–0.5
<i>Jarvis et al.</i> [1997]	Canada	black spruce	1994	-2.0	0.4	
<i>Lee et al.</i> [1999]	Ontario	maple	1996	-0.8	0.15	
			1997	-2.7		
			1998	-2.0		
			1998	-2.0		
<i>Lindroth et al.</i> [1998]	Sweden	pine	1995	0.9	0.4	
			1996	-0.05		
			1997	0.8		
<i>Lindroth et al.</i> [1998]	Sweden		1997	-1.9	0.4	
<i>Loeschler et al.</i> [2003] <sup>h</sup>	Costa Rica	tropical wet	1998	0.05	0.45	
			1999	-1.53		
			2000	-5.97		
			1995	-5.9		
			1995	-0.7		
<i>Malhi et al.</i> [1998]	Brazil	tropical humid	1995		none	
<i>Malhi et al.</i> [1999]	Saskatchewan	spruce	1995		yes	
<i>Meyers</i> [2001]	Oklahoma	grassland	1997	0.41		
<i>Oechel et al.</i> [2000]	Alaska	tundra		0.4		
<i>Paw U et al.</i> [2000]	Washington	mixed conifer	1998	-1.7		
<i>Pilegaard et al.</i> [2001]	Denmark	beech	1997	-1.7	0.25	
			1998	-1.2		
			1998	-2.4		
<i>Schmid et al.</i> [2000] <sup>i</sup>	Indiana	deciduous	1998	-2.4	0.2	
<i>Schmid et al.</i> [2003]	Michigan	deciduous	1999	-1.7	0.35	
			2000	-1.60	0.35	
			2001	-0.8	0.35	
			1997	-2.67	0.2	
			1998	-4.5	0.2	
<i>Suyker et al.</i> [2003]	Oklahoma	prairie	1997			
<i>Valentini et al.</i> [2000]	Italy	spruce	1998	-4.5	0.2	
		oak	1997	-6.6		
		spruce?	1998	-4.5	0.2	
		beech	1994	-4.7	0.17	
		mixed deciduous	1997	-6.6	0.2	
		Germany	spruce	1996	-3.3	0.3
		Germany	spruce	1997	-0.8	0.2
		Germany	spruce	1997	-0.8	
		Germany	conifer	1996	-3.1	0.4
				1997	-4.9	0.4
		Netherlands	pine	1997	-2.1	0.25
		Iceland	poplar	1997	-1.0	0.2
		UK	spruce	1997	-2.1	0.2
				1998	-6.7	0.2
		Finland	pine	1997	-2.3	0.3
				1998	-2.6	
				1999	-1.9	
	Belgium	mixed	1997	-4.3	0.4	
<i>Valentini et al.</i> [2000] <sup>j</sup>			1997	-1.6	0.0	
<i>Vourlitis and Oechel</i> [1997]	Alaska	wet tundra	1995	-0.156	none	
		moist tundra	1995	-0.276	none	
<i>Vourlitis and Oechel</i> [1999] <sup>k</sup>	Alaska	tussock tundra	1994	-0.395	none	

**Table 1.** (continued)

Study	Location	Vegetation type	Year	Annual NEE	$u^*$ Threshold
<i>Wilson et al.</i> [2001]	Tennessee	maple/oak	1995	-0.552	none
			1994	-5.25	
			1995	-6.1	
			1996	-6.0	
			1997	-6.5	
			1998	-6.6	
			1999	-7.4	
<i>Yamamoto et al.</i> [1999]	Japan	deciduous	1994	-1.2	
			1995	-0.7	
			1996	-1.4	
			1997	-1.5	
			1998	-1.4	

<sup>a</sup>NEE is reported in units of  $\text{t C ha}^{-1} \text{y}^{-1}$ , and  $u^*$  is reported in  $\text{m s}^{-1}$ .

<sup>b</sup>Included a vertical advection term following *Lee et al.* [1999] and scaled chamber measurements to estimate  $R_c$ .

<sup>c</sup>Seasonal fluxes only.

<sup>d</sup>Used data from nights when  $u^* > 0.4$  for at least 2.5 hours and had storage flux for 1 month.

<sup>e</sup>Used sigma u criterion to develop a temperature response, then modeled  $R_c$ .

<sup>f</sup>No correlation found between NEE and  $u^*$ .

<sup>g</sup>Assigned uncertainty of 25%, some of the integrated flux (DOY 145–265); the data was filled with a functional relationship using  $u^* > 0.4$ .

<sup>h</sup>Fixed nighttime  $R_c$  based on data with  $u^* > 0.45$ .

<sup>i</sup>The  $u^*$  correction reduced sink by  $\sim 50\%$ .

<sup>j</sup>No relationship between  $u^*$  and NEE.

<sup>k</sup>Estimated sink for growing season only.

### 2.1.1. Systematic Errors

[10] There are different sources of systematic errors in EC measurements. Systematic errors occur as a result of instrument response time, pathlength averaging, and physical separation between sonic anemometers and the measurement system for  $\text{CO}_2$  or other trace gases [*Massman and Lee*, 2002]. In addition, selective systematic errors can result from both the exclusion of high-frequency turbulence due to low-pass filtering (used to eliminate noise) and the exclusion of low-frequency flows due to the length or method of averaging time [*Mann and Lenschow*, 1994; *Foken and Wichura*, 1996; *Vickers and Mahrt*, 1997]. For example, the transport of carbon dioxide due to slowly moving boundary layer eddies (i.e., conditions with weak winds and significant surface heating), may be excluded by traditional averaging periods [*Sakai et al.*, 2001]. The overall effect is analogous to inadequate spatial averaging, as is the case with using too short of an averaging distance to define the turbulent fluctuations in aircraft observations [*Betts et al.*, 1990; *Desjardins et al.*, 1997]. Systematic errors occur within each of the averaging periods (i.e., 30 min) and are present in the EC term (equation (1b)) and scale in the same proportion with longer time integrals (i.e., 1 year, Table 2). Systematic errors associated with tilt corrections, flux divergence and advection between the surface and EC measurement level can exist because of inaccurate definition of the flux field, or when particular conditions arise causing one or more of terms III–VII (equation (1a)) to be not equal to zero, and these errors may not become apparent until data are examined across longer time intervals, from day-to-day to season-to-year.

### 2.1.2. Defining the Flux Field

[11] The partitioning between turbulent and nonturbulent motions is influenced by the coordinate rotation method used to align the sonic anemometer with respect to the surface [*Wilczak et al.*, 2001] and can lead to systematic errors. While investigators have been attempting to correct for possible sonic misalignment for at least 25 years, the methods for such corrections have been detailed in the

refereed literature only recently. One method, planar rotation [*Wilczak et al.*, 2001], was developed in the late 1970s by Steven Stage. For nighttime stable conditions, rotating the coordinate system for individual (e.g., 30 min) records can lead to erratic results [*Finnigan et al.*, 2003]. Applying new rotation angles for each 30-min record to eliminate the cross-wind Reynolds stresses cannot always be justified for such nocturnal conditions. Several alternative methods have been developed which compute rotation angles across longer or ecologically significant periods (month-season-annum). Some of these methods are classified by *Paw U et al.* [2000] and discussed in extensive detail by *Wilczak et al.* [2001]. Alternatively, *Lee* [1998] rotated the coordinate system for the entire data sample, independently for each wind direction group. This approach tends to better capture the directional influence of slopes on turbulence, and the directionally dependent flow distortions associated with the instruments, mounting brackets and so forth. This planar fit method however, is sensitive to remounting of the sonic anemometer, which will result in the necessary calculation of new rotation angles.

[12] *Mahrt et al.* [2000] compared different coordinate rotation methods for (carefully aligned) sonic anemometers, and found that differences were insignificant, except for the estimation of momentum flux under weak wind conditions ( $< 1 \text{ m s}^{-1}$ ) or for scalar fluxes under very weak winds ( $< \sim 0.25 \text{ m s}^{-1}$ ). Such conditions would typically be eliminated by the  $u^*$  threshold discussed previously. They recommended a slightly modified version of the rotation scheme used by *Lee* [1998]. For misaligned sonic anemometers, more serious errors of  $\sim 14\%$  and  $3\%$  per degree tilt were found for momentum and scalar fluxes [*Moncrieff et al.*, 1996].

[13] Instrument-induced flow distortions may be larger below canopy because the vertical velocity is not small compared to the total horizontal velocity, i.e., large attack angles. Careful examination of coordinate rotation methodologies is needed on below-canopy fluxes and should also take deviation in flow by trunks and branches into account, particularly for sites located on challenging topography.

**Table 2.** Different Sources of Error in 30 min and Annual NEE Estimates<sup>a</sup>

Source of Error	Systematic Error, 30 min	Random Error, 30 min	Time-Integrated Error for Annum Estimates	Notes, Annual NEE t C ha <sup>-1</sup> y <sup>-1</sup>	Reference
Instrument noise	2%	11%	2%	...	<i>Kruijt et al.</i> [2004]
	~0%	10–20%	~0%	...	<i>Weseley and Hart</i> [1985]
	...	15–20%	...	due to large eddies sampled at 495 m	<i>Berger et al.</i> [2001]
Rotation scheme and averaging operator	10–25%	...	10–25%	...	<i>Kruijt et al.</i> [2004]
	14%	...	...	...	<i>Moncrieff et al.</i> [1996]
	~0–20%	...	5%	uniform systematic error	<i>Goulden et al.</i> [1996]
calibration	~0–20%		0–6%	due to signal and span drift	<i>Kruijt et al.</i> [2004]
Frequency loss	8.4%			...	<i>Kruijt et al.</i> [2004]
	<5%			for daytime <i>NEE</i> estimates	<i>Berger et al.</i> [2001]
	<12%			for nighttime <i>NEE</i> estimates	“ ”
	0–9%			cospectral similarity	<i>Goldstein et al.</i> [2000]
	6–16%			“ ”	<i>Loescher et al.</i> [2003]
Advection			(–) 10–15%	during June and July at WLEF, WI	<i>Yi et al.</i> [2000]
			(–) 5–20%	Carson, WA (1.9, 1998–1999)	M. Falk et al. (manuscript in preparation)
Total uncertainty			(–) 40%	vertical advection term only	<i>Baldocchi et al.</i> [2000]
			12.5%	Jaru, Brazil site (~5.0, 1999)	<i>Kruijt et al.</i> [2004]
			32%	Manaus, Brazil site (~8.4, 2000)	“ ”
			14%	Petersham, MA site	<i>Goulden et al.</i> [1996]
			21%	<sup>b</sup> Oak Ridge, TN site (2.1, 1994)	<i>Wilson and Baldocchi</i> [2001]
			54%	<sup>c</sup> Metolius, OR site (3.2, 1996)	<i>Anthoni et al.</i> [1999]
			13%	<sup>d</sup> “ ”	“ ”
			<27%	<sup>e</sup> Tapajos, Brazil site (2.7, 1997)	<i>Saleska et al.</i> [2003]
Gap filling			119%	<sup>c,f</sup> WLEF, WI (0.16, 1997)	<i>Davis et al.</i> [2006]
			264–527%	<sup>f,g</sup> “ ”	“ ”
			3–20%	...	<i>Kruijt et al.</i> [2004]
		3.5–20%	8.3–11%	for daytime <i>NEE</i> estimates	<i>Falge et al.</i> [2001]
		1.5–4%	7–52%	for nighttime <i>NEE</i> estimates	“ ”

<sup>a</sup>The errors attributable to gap filling were not available for estimates of total uncertainty (except as noted).

<sup>b</sup>Constrained by soil respiration estimates.

<sup>c</sup>Error of ~1/2 due to errors in the respiration estimate.

<sup>d</sup>Error due to gap filling.

<sup>e</sup>Random uncertainty only.

<sup>f</sup>High % due to relatively low estimate of NEE.

<sup>g</sup>Systematic errors only.

### 2.1.3. Random Errors


[14] Sources of random error differ in flux estimates made within an averaging period (e.g., 30 min) and those associated with adding averaging periods together to achieve estimates across longer time integrals, that is, random flux errors generally decrease with increasing record length. The random flux error that *Moncrieff et al.* [1996] refers to is the variability among averaging periods, which includes both the random error that is added from averaging period to averaging period and the variability due to synoptic changes in ambient conditions. Diurnal variations were filtered out of their calculation. Sources of random error within an averaging period result primarily from an inadequate sample size [*Hollinger and Richardson*, 2005; *Kruijt et al.*, 2004; *Finkelstein and Sims*, 2001; *Vickers and Mahrt*, 1997; *Mann and Lenschow*, 1994; *Lumley and Panofsky*, 1964]. This does not mean to imply however, that sampling more frequently (e.g., from 10 to 50 Hz) would necessarily reduce the effects of random error. Needed instead, is the ability to sample fast enough to resolve all transport scales, but also long enough to adequately sample the infrequent large turbulent motions that transport flux. The Allen variance can be used to assess the relative contribution of instrument random and systematic error under static conditions and can assist in determining an appropriate averaging time to reduce random error [*Allen*, 1966; *Barnes and Allen*,

1990] [cf. *Bowling et al.*, 2003; *Loescher et al.*, 2005] (Figure 2). This is strictly true only if the random error is normally distributed which is apparently not the case for (dynamic) flux data [*Hollinger and Richardson*, 2005; *Richardson et al.*, 2006]. In the case of a closed path sensor, longer averaging times used to reduce random error can introduce systematic error (e.g., sensor drift), which makes frequent calibrations and instrument stability (i.e., pressure control and temperature regulation) increasingly important to reduce uncertainty. With regard to instrumentation noise, the random error generally becomes small with averaging times >15 min while the effects from systematic error scale among different time integrals.

[15] Unfortunately, increasing the averaging period to capture a large enough sample size also has the potential to increase the overall contribution of nonstationarity of the ambient environment, which in turn, can bias the computed fluxes [*Lenschow et al.*, 1994]. Techniques for assessing nonstationarity in geophysical turbulence are given by *Gluhovsky and Agee* [1994], *Foken and Wichura* [1996], and *Mahrt* [1998].

[16] To reduce random errors, two averaging times are used to compute the turbulent flux, such that,

$$s' = s - \langle s \rangle \quad (3)$$



**Figure 2.** Allan variance calculated for sampled  $\text{CO}_2$  from a closed-path infra-red sensor (model Li-7000, Li-Cor Inc., Lincoln, Nebraska). The Li-7000 was placed in a temperature-controlled chamber ( $31.00 \pm 0.02^\circ\text{C}$ ), and after an equilibration period ( $\sim 12$  hours), 20 Hz data were collected for  $\sim 3$  hours. Flow of  $\text{N}_2$  was regulated through both reference and sample cells. Dashed line indicates a 30-min averaging period.

where  $s$  is the measured scalar quantity, and the angle brackets refer to the averaging operator that defines the turbulent fluctuations (noted by the prime). The corresponding perturbations ideally include all of the turbulent motions but exclude influences from mesoscale motions. The vertical motion at lower frequencies cannot be adequately measured by sonic anemometers, although it is often thought that their overall contribution to the total surface flux is small and can hence be ignored. Inadvertent inclusion of mesoscale motions however, can lead to large random flux error [Vickers and Mahrt, 2006a; Sun et al., 1998]. Such contamination most likely occurs with stable conditions in the presence of gravity waves, meandering motions, nonstationary drainage flows and other nameless mesoscale motions of unknown origin [see references in Mahrt et al., 2001b].

[17] The vertical flux is then computed as  $\langle w's' \rangle$  where the angle brackets represent the averaging time on a longer timescale to reduce the random flux error (i.e., averaging the products of perturbation). The longer averaging time (30 min to 1-year scale) is sometimes referred to as the “flux averaging time” and leads to the  $NEE$  estimates. Both the overbar and angle brackets must be chosen as simple unweighted averages, particularly to satisfy some averaging schemes [e.g., Reynolds, 1901]. Sometimes the averaging period in equation (3) and that associated with the overbar (compare with equation (1b)) are the same, in which case, random flux errors may be large. For example, the largest turbulent motion included in equation (3) would be of the same timescale as the flux averaging time, and therefore only one sample of this motion would be captured. Such

errors can contribute to large variability in process-oriented studies that use 30-min averages, but are less important when averaged fluxes are summed together for longer time periods, as in the calculation of seasonal and annual  $NEE$  estimates, or when inferring relationships such as light response to daytime  $NEE$  with a large number of 30-min average data sets.

[18] At times, the distinction between turbulent and nonturbulent motions is difficult to detect and may overlap in scale. The inspection of integrated cospectra (ogives), however, can assist in determining the temporal scale for transport of carbon [Friehe et al., 1991; Oncley et al., 1990]. On the basis of examination of a large number of multiresolution cospectra, under a variety of conditions, Vickers and Mahrt [2006a] demonstrate that the ideal averaging time for defining the turbulent fluctuations (equation (3)) decreases with increasing stability. However, they also show that averaging times increase with height above the canopy because of increases in eddy size with height. Hence the use of variable averaging times seeks to minimize the influence of random errors in very stable conditions due to inadvertent capture of mesoscale motions and to minimize systematic flux loss in unstable weak wind conditions due to inadvertent omission of slowly moving large eddies. Use of variable averaging times, however, may be impractical for routine calculations, and the applicability of this approach to below-canopy flux calculations is not known. In many cases however, it is necessary to characterize the absolute or relative magnitude of random flux measurement errors. Maximum likelihood and Kalman filter approaches, for example, require such estimates.

## 2.2. Flux Divergence Below the Measurement Level

[19] Environmental conditions that contribute toward flux divergence (i.e., terms II–IV  $\neq 0$ , equation (1a)) often also affect storage flux (i.e., term I  $\neq 0$ , equation (1a)) and advective flows (i.e., terms V–VII  $\neq 0$ , equation (1a)), which can be confounded and difficult to resolve. Under very stable conditions with semiopen canopies, nocturnal radiative cooling at the ground surface can cause strong atmospheric stratification below the canopy, suppressing turbulent mixing [e.g., *Mahrt et al.*, 2000, 2001a], despite the fact that the openness of some canopies would presumably encourage greater mixing between the below-canopy environment and the atmosphere. With strong below-canopy stratification, below-canopy flows may become decoupled from the above-canopy flow, particularly with formation of below-canopy drainage flows (discussed further below). Under these conditions the vertical CO<sub>2</sub> flux decreases rapidly with height above the source area, sometimes in the lowest few meters above the surface [*Soler et al.*, 2002; *Wilson and Meyers*, 2001; *Lee et al.*, 1995]. As a result, the magnitude of the turbulent flux may be much smaller at the above-canopy EC measurement level than the total flux [*Staebler and Fitzjarrald*, 2004]. When the below-canopy environment is (partially) decoupled, CO<sub>2</sub> concentrations increase and can be advected horizontally and vented at heterogeneous locations downwind [*Sun et al.*, 1998] or during the early morning transition periods [*Grace et al.*, 1995; *Lee and Black*, 1993; *Yang et al.*, 1999].

[20] With very stable conditions (smaller eddy size and weak mixing), the influence from local heterogeneities and microscale turbulent structure becomes increasingly important in quantifying the flux. Deploying sonic anemometers closer to surface can reduce the influence of vertical flux divergence, but in doing so, increases the flux loss due to pathlength averaging and instrument separation, and reduces the representativeness of the flux measurement (i.e., smaller footprint [*Schmid*, 1997; *Foken et al.*, 2004]). Therefore the effects of inadequate representation of the source area can become large when the understory, litter, and soil fluxes are significantly heterogeneous in space. Increasing the EC measurement height increases the representativeness of the measurements, but may also increase uncertainties due to vertical flux divergence.

## 2.3. Estimation of Advection

### 2.3.1. Vertical Advection

[21] There have been several recent studies examining the effects of localized advective flows on tower-based estimates [e.g., *Staebler and Fitzjarrald*, 2004]. Vertical advection at night may be significant even though the mean vertical windspeed near the surface is very small (i.e.,  $\sim 0 \text{ m s}^{-1}$ ). Vertical advection has sometimes accounted for the largest fraction of the variation in nocturnal 30-min *NEE* estimates [*Lee*, 1998], and can greatly alter annual *NEE* estimates [*Baldocchi et al.*, 2000; *Lee*, 1998]. *Finnigan* [1999], *Paw U et al.* [2000] and *Baldocchi et al.* [2000] further argue that if a vertical advection term is included in the conservation equation, then the horizontal advection terms should also be included. Complications further arise in some forests when advection terms during the night differ between above- and below-canopy environments. *Vickers and Mahrt* [2006b] have concluded that the mean vertical

motion cannot be estimated from sonic anemometers and that the mean vertical motion must be estimated from mass continuity.

### 2.3.2. Horizontal Advection

[22] Respired CO<sub>2</sub> may be advected horizontally below the EC measurement level, leading to underestimation of ecosystem respiration ( $R_e$ ). Horizontal advection can be partitioned into separate types of flows: (1) synoptic scale occurring over a deep atmospheric layer, often deeper than the boundary layer, (2) propagating transient mesoscale flows, and (3) localized stationary flows associated with surface heterogeneity, sometimes occupying only the lower part of the boundary layer [*Lee*, 1998, and references therein]. The direct effect of synoptic advection is generally small within the canopy environment, except for the passage of frontal systems. A wide variety of physical processes contribute to mesoscale motions.

[23] Winds over complex topography can modify the mean wind shear and generate turbulence and waves. The latter can lead to local advection and breakdown of the nocturnal boundary layer. Recently, such investigations have examined the effects of small amplitude waves at intermediate periods of 1–5 min (D. Anderson, personal communication, 2002) and stronger rotor motions at longer periods of about 20-min [*Turnipseed et al.*, 2003] on CO<sub>2</sub> budgets.

[24] Cold air drainage is generated by a cooling air mass over sloped terrain [i.e., *Gedayloo et al.*, 1980; *Doran and Zhong*, 1994; *Doran*, 1991]. Its development is opposed by the Bernoulli pressure gradient, vertical stress divergence and drag forces [*Staebler and Fitzjarrald*, 2005; *Lee*, 1998; *Mahrt*, 1992]. Drag forces are often associated with the spatial distribution of aboveground vegetation [*Staebler and Fitzjarrald*, 2005], but may include other small-scale topographical features such as boulders, ledges and dunes. With clear skies and weak mesoscale flow, nocturnal drainage flows can develop on extremely small angled slopes [*Caughey et al.*, 1979; *Lee and Hu*, 2002], and sometimes develop entirely below canopy [*Lee*, 1998; *Miller et al.*, 1983; *Staebler and Fitzjarrald*, 2004]. Examination of drainage flows can be particularly difficult because they can occur in a thin stratified layer, particularly in the early evening [*Manins*, 1992; *Soler et al.*, 2002], sometimes becoming nonstationary or pulsating [*Doran and Horst*, 1981]. Additional advective flows can develop from local circulations generated over contrasting landscape types, e.g., nocturnal updrafts over warm surfaces such as lakes [*Sun et al.*, 1998], and daytime updrafts and large eddies over a forest initiated from adjoining clearcut areas [*Loescher et al.*, 2006; *Leclerc et al.*, 2003].

[25] The magnitude and sometimes even the sign of the advection can be questioned for all but the simplest topography. For example, air temperature along a slope does not vary monotonically and the temperature gradient can reverse in sign because of variations in soil heat flux, enhanced mixing related to changes in roughness, and increased shear-induced mixing associated with drainage flows. Increases in the vertical mixing of heat imply increased mixing of CO<sub>2</sub>. A reversal of horizontal gradients can occur at midslope where a thermal belt occurs because of the downward mixing of warmer air at midslope, in concert with colder air trapped in the valley [*Yoshino*, 1975;



Geiger, 1961]. These spatial variations in flow regimes not only limit the utility of flux measurements from all but the most ideal tower sites but also prevent precise estimates of horizontal CO<sub>2</sub> gradients without a detailed measurement system. *Staebler and Fitzjarrald* [2004] present an approach on the basis of spatial coherence calculations to determine the spatial resolution needed to adequately estimate horizontal gradients. They investigated horizontal advection associated with surface horizontal heterogeneities at the Harvard Forest and Camp Borden AmeriFlux sites. They found that obstructions (tree stems) can explain below-canopy directional flow patterns, and also found that nighttime thermal stratification could produce drainage flows ~92% of the time, and that buoyancy exerted more control on these drainage flows than stress divergence and pressure gradients ~58% of the time. While these findings are not universally transferable to all sites, *Staebler and Fitzjarrald* [2004] outline a methodology to determine the magnitude of controls on below-canopy advection.

[26] Exact contributions of horizontal advection to *NEE* are difficult to estimate and cannot be determined by the current EC technique (i.e., single-point, tower measurement). Instead, the current micrometeorological methods used to estimate vertical and horizontal components of advection [cf. *Lee*, 1998; *Paw U et al.*, 2000; *Finnigan*, 1999] may be more appropriate in assessing when advection becomes significant [*Lee and Hu*, 2002; *Gudiksen et al.*, 1992], determining bounds on *NEE* [*Baldocchi et al.*, 2000], and estimating appropriate  $u^*$  thresholds that may remove much of the data for conditions when advection or flux divergence is prevalent [*Gu et al.*, 2005]. Nascent, site-specific estimates of advection range from 5–40% across seasonal to annual intervals (Table 2).

[27] Applying a classical fluid dynamic approach of estimating budgets in a control volume may be an alternative solution to the CO<sub>2</sub> advection problem [*Sun et al.*, 1998; *Finnigan et al.*, 2003]. Developing this approach is difficult however, because it requires detailed estimates of storage, horizontal and vertical advection by the mean flow, and the horizontal flux divergence. Aircraft observations [*Betts et al.*, 1990, 1992] can estimate vertical transport through the “top of the box”, provided that spatial heterogeneity is not too great. Since aircraft EC measurements are relatively expensive, they are limited to a small fraction of the desired observational period and are most meaningful when used in concert with tower measurements [*Vickers and Mahrt*, 2003].

[28] Other approaches include the use of a tracer in conjunction with independent scalar flux measurements [*Martens et al.*, 2003], or Lagrangian and analytical model efforts [*Leclerc et al.*, 2003]. *Martens et al.* [2003] used a tracer (i.e., Radon-222 which occurs naturally and is biologically inactive) to provide an independent estimate of advection and to examine the use of a  $u^*$  threshold on  $R_e$ . They found  $u^*$  corrected EC estimates agree well with radon-derived  $R_e$ . Their result supports the use of a  $u^*$  threshold when estimating  $R_e$  from EC, and suggests drainage flows export carbon when the canopy is not well mixed. However, comparisons between advection rates derived from a tracer or traditional micrometeorological methods can be confounded by use of inappropriate parameters, such as auto-

correlated turbulent diffusion coefficients or mismatched characterization of above- and below-canopy environments.

## 2.4. Gap Filling

[29] Other errors in annual estimates of *NEE* may arise through the filling of the unavoidable gaps in the EC data sets so that integrated fluxes may be calculated daily to annually, and in estimating daytime respiration or gross ecosystem exchange. *Falge et al.* [2001] found that errors did not differ much among three methods for gap filling: mean diurnal variation, nonlinear regressions, and look-up tables based on meteorological and seasonal conditions. Each of these three methods provided a good approximation of the original integrated values when artificial gaps were created, even when large percentages of data were missing. They also reported that (1) the mean diurnal variations were best simulated when estimated for 7-d windows for nighttime data and 14-d windows for daytime data and (2) look-up tables and nonlinear regressions provided higher levels of accuracy across seasons-to-year intervals and among large temperature ranges. The authors did not however, recommend a particular method because much of the variation among methodologies was confounded by differences in data preparation and in identifying nocturnal periods with insufficient turbulence for valid EC measurements. In the simulations of *Falge et al.* [2001], errors attributable to gap filling were  $<0.50 \text{ Mt C ha}^{-1} \text{ y}^{-1}$ , even with ~65% of the annual data missing.

[30] The construction and use of physiologically based models are central to the analysis of EC data [*Hollinger and Richardson*, 2005]. Models of varying complexity are also used to test our understanding of ecosystem processes. The recent availability of EC data has led to the increased use of data-based modeling (also called inverse modeling) where model parameters are estimated directly from the data. In order to do this correctly, and to estimate the uncertainty in the model parameters, we need to know the uncertainty (random error) in the underlying EC measurements, and have a methodology that can explicitly express the nonlinearity between EC derived fluxes and abiotic parameters.

[31] *Hollinger and Richardson* [2005] examined the characteristics of random error in EC measurements with a unique two-tower system, and found that flux data do not conform to least squares assumptions of error term variance homogeneity and normality. Using a data binning approach, they confirmed that errors in flux data are heteroscedastic and not normally distributed at a variety of flux sites ranging from crop fields to large towers [*Richardson et al.*, 2006]. Flux data errors appear to be better represented by a double exponential probability density function (PDF) than a Gaussian PDF.

[32] There are potentially serious consequences from using ordinary least squares (OLS) fitting techniques with flux data, including incorrect determination of model parameter values and bias in annual *NEE* sums [*Hollinger and Richardson*, 2005; *Richardson and Hollinger*, 2005]. Instead of OLS methods, approaches such as maximum likelihood provide unbiased estimates of model parameters, and when coupled with, for example, Monte Carlo methods, also provide confidence intervals for parameter estimates. Several authors have applied these or related approaches (e.g., neural networks) to EC data [*Papale and Valentini*,

2004, 2003; Schulz *et al.*, 2001; van Wijk and Bouten, 2002]. Richardson and Hollinger [2005] showed that OLS fitting methods resulted in models that overestimated nocturnal respiration (and hence underestimated *NEE*) relative to the maximum likelihood approach.

[33] Kalman filter techniques are another approach for estimating model parameter values and filling gaps in a data record, and have recently been applied to EC data [Jarvis *et al.*, 2004; Williams *et al.*, 2005; Gove and Hollinger, 2006]. The Kalman filter is a recursive algorithm for estimating the state of a process in a way that minimizes the error [Kalman, 1960]. It has a structure that is ideal for time series where the data structure is autocorrelated, which make it an attractive candidate for estimating missing flux data. The equations for a Kalman filter consist of two parts; time update or “predictor” equations, and measurement update or “control” equations [Welch and Bishop, 2001]. The time update projects the current state (surface flux) and covariance estimates forward in time, and a measurement update adjusts the projected estimate on the basis of an actual measurement made at that time. Because the original Kalman filter was developed to estimate the state of a system on the basis of linear stochastic difference equations, and many interesting processes are nonlinear (including ecosystem atmosphere exchanges), a variety of approaches have been developed for the nonlinear problem. Williams *et al.* [2005] use the ensemble Kalman filter to assimilate EC and other ecosystem data into an ecosystem C balance model [Evensen, 2003] while Gove and Hollinger [2006] employ the unscented Kalman filter [Julier and Uhlmann, 1997] to estimate *NEE* model parameters and fill gaps in an EC record.

### 3. Measurements of Respiration Components for Diagnosing EC Data and Understanding Sources of Respired CO<sub>2</sub>

[34] Ecosystem respiration ( $R_c$ ) can be estimated with EC during the night, but may vary in temporal resolution, from 30 min to 12 hours, because longer averaging periods may be needed to assess the flux divergence, advection and storage terms (equation (1a)). Independent measurements of respiration from foliage, wood and soil have also been made with chambers, scaled to a site, and used to examine contributions of different processes to ecosystem productivity [Harmon *et al.*, 2004; Bolstad *et al.*, 2004; Granier *et al.*, 2000; Law *et al.*, 1999a; Lavigne *et al.*, 1997; Goulden *et al.*, 1996]. Attention is being focused on the spatial and temporal representativeness of soil chamber measurements because soil CO<sub>2</sub> efflux accounts for ~50 to 70% of  $R_c$  [Davidson *et al.*, 2006; Law *et al.*, 1999a; Goulden *et al.*, 1996] and because of large spatial variability in soil properties, litterfall, coarse woody debris, and microbial activity. In this section we discuss the uncertainty in chamber measurements over soil, which also apply toward other chamber-based measurements.

[35] In an attempt to estimate both the temporal and spatial variability in soil gas exchange, sampling designs often include the combination of automated chambers (closed dynamic [e.g., Lavigne *et al.*, 1997; Norman *et al.*, 1997; Goulden and Crill, 1997]) to assess temporal changes in soil fluxes (e.g., hourly timescale), and manual chambers (closed static, e.g., Li-6400-09, Li-Cor Inc.)


measured at many locations weekly to seasonally to assess spatial variation for scaling the automated chamber measurements to the site or approximate footprint of the tower.

[36] Sampling errors can also originate from not fully accounting for the spatial heterogeneity in soil CO<sub>2</sub> flux [Drewitt *et al.*, 2002; Law *et al.*, 2001a; Janssens *et al.*, 2001; Rayment and Jarvis, 2000; Law *et al.*, 1999a; Fang *et al.*, 1998; Lavigne *et al.*, 1997]. It is essential to characterize the spatial variability and adjust the sampling design appropriately for scaling to site and to ultimately compare with EC data. However, this can be quite a challenge if the chamber and tower source areas are mismatched.

[37] Capturing the spatial heterogeneity and achieving high accuracy in soil flux measurements are not necessarily the same thing. A biased chamber design can measure the spatial heterogeneity while still being inaccurate. Accuracy in chamber estimates can only be established by measuring known fluxes under defined conditions. This has been done very infrequently [cf. Davidson *et al.*, 2002; Butnor and Johnsen, 2004; Fang and Moncrieff, 1998]. For example, Butnor *et al.* [2005] found that dynamic closed chambers underestimated fluxes through porous media such that air-filled porosity of soil alters the effective chamber volume. If systematic uncertainties are uniform among chambers, the coefficient of variation ( $CV = \sigma/\mu$  where  $\sigma$  is the standard deviation and  $\mu$  is the mean) can be used to determine the number of chambers needed to adequately represent the spatial variability. The CV can be used to estimate the precision in the *measured* (population) mean (see Figures 3a–3b). Reported values of CVs range 0.15–0.55 (Table 3). These CVs are relatively high and they place limits on the precision we can expect from a reasonable number of chambers. Furthermore, CVs may also change with seasonal climatic conditions, for example, it is possible for CVs to increase to 0.8 during spring snow melt with increased and spatially variable microbial activity (for Metolius, J. Irvine, personal communication, 2005), or also increase with the onset of wet season rains. As a general rule, increasing the number of chambers will enhance the precision in the *measured* mean.

[38] Estimates from both chamber types, custom built and manufactured alike, are subject to systematic and sampling errors that can be large (Table 3). Estimating the systematic uncertainties can determine how accurate the *measured* mean is to a *true* mean from the soil surface. Sources of this type of error include the size and dynamic volume of the chamber, differences between the external and internal chamber environments such as; wind, light [Dore *et al.*, 2003], pressure [Lund *et al.*, 1999; Massman *et al.*, 1997], and temperature can affect measurements, but once identified, can be remedied [Dore *et al.*, 2003; Davidson *et al.*, 2002; Fang and Moncrieff, 1998]. There are efforts underway to calculate soil fluxes on the basis of a dynamic soil volume to address changes in effective soil volume with moisture, particularly in porous soils (L. Xu, personal communication, 2005).


[39] Most importantly, automated chambers assess the temporal variability in soil respiration and a large number of spatially distributed, manually sampled dynamic closed chambers can assess the spatial variability. If all the chambers are in the representative tower source area, these two methodologies can be used in conjunction to estimate soil



**Figure 3a.** Mean number of chambers needed to be within a percentage around the mean across different  $\alpha$ -values from two different contrasting soils types. Estimates are from the (top) Metolius, Oregon, intermediate ponderosa pine stand, measured with 32 manually operated, open-dynamic chambers (Li-6400-09, Li-Cor Inc.) across five sampling periods within one season (DOY 219, 226, 234, 249, and 283 in 2002). (bottom) Relationship between the number of chambers needed and precision (with an  $\alpha = 0.1$ ) of the measured soil flux as a function of the coefficient of variations (CV) for the same data set. CV's were adjusted by changing the standard deviation as a proportion of the mean flux. Mean flux ranged 3.37–1.71 and 4.49–3.05  $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  from Metolius and La Selva sites, respectively.

respiration at comparable scales to those measured by EC. Protocols for measurement, sampling strategies and how soil respiration estimates can be used in conjunction with EC are discussed by *Ryan and Law* [2005].

[40] Deploying EC systems below forest canopies along with chamber measurements has increased our understanding of functional and physical relationships governing *NEE*. Below-canopy flux measurements have provided useful



**Figure 3b.** Mean number of chambers needed to be within a percentage around the mean across different  $\alpha$ -values from two different contrasting soils types. Estimates are from (top) La Selva, Costa Rica (old growth wet tropical forest), measured with 24 closed dynamic chambers across 12 sampling periods within one (dry) season (DOY 365 in 1998, and DOY 14, 26, 35, 49, 63, 83, 98, 112, 133, 141, and 155 in 1999). (bottom) As for Figure 3a.

temporal information for understanding seasonal differences in diel patterns and turbulent structure, while chamber-based measurements can characterize the partitioning of respiration from the different sources, including soil, foliage, and bole [Law *et al.*, 1999b]. Taken together, Law *et al.* [2001b] determined the vertical distribution of respiration components and provided an estimate of the below-canopy  $\text{CO}_2$  source area. The information can be used to evaluate component processes in models, particularly belowground processes that are poorly understood. Data

from chamber measurements have also been used to fill gaps in EC data sets, particularly when a (large) representative portion of the ecosystem is included in the measurements [Dore *et al.*, 2003; Bubier *et al.*, 2002; Oechel *et al.*, 1998; Drake *et al.*, 1996]. However, this is extremely labor intensive when foliage and wood respiration measurements need to be sampled frequently, in addition to automated measurements of soil respiration, so it is not a feasible approach to regularly replace nighttime respiration estimates from EC measurements.

**Table 3.** Reported Sources of Error and Coefficient of Variation (As an Indication of How Accurately Spatial Heterogeneity is Estimated) Among Different Chamber Designs and Environments<sup>a</sup>

Source of Uncertainty	Percent Error of Estimate	Environment	Reference
<i>Chamber Design</i>			
CS, CD, OD	(-) 7–45%	black spruce/jack pine, boreal	<i>Norman et al.</i> [1997] <sup>b</sup>
CD, SL	(-) 1–46%	Scots pine, subboreal	<i>Janssens et al.</i> [2000] <sup>b</sup>
CD	< (-) 3%	beech/Douglas fir, temperate	<i>Longdoz et al.</i> [2000] <sup>c</sup>
CD	(-) 15%	laboratory	<i>Nay et al.</i> [1994] <sup>d</sup>
CD	(-) 8–14%	ponderosa pine, temperate	<i>Irvine and Law</i> [2002]
CD	(-) 20–70%	grassland, temperate	<i>Lund et al.</i> [1999] <sup>e</sup>
SL	25%, -75%	laboratory	<i>Nay et al.</i> [1994], <i>Nay</i> [1994] <sup>d,f</sup>
OD	(-) 0–10%	black spruce/jack pine, boreal	<i>Rayment</i> [2000], <i>Rayment and Jarvis</i> [1997] <sup>b,g</sup>
OD	~200%	slash pine, subtropical	<i>Fang and Moncrieff</i> [1998] <sup>h</sup>
OD	0–15%	eastern United States, mixed, temperate	<i>Davidson et al.</i> [2002] <sup>i</sup>
<i>Coefficient of variation</i>			
CD	~0.30	old growth, wet tropical	(H. Loescher and P. Crill, personal communication, 2004)
CD	~0.35	ponderosa pine, temperate	(H. Loescher and J. Irvine, personal communication, 2004)
CD	0.15–0.20	ponderosa pine, temperate	<i>Irvine and Law</i> [2002]
CD	0.16–0.45	aspen, boreal	<i>Russell and Voroney</i> [1998]
OD	0.48	old growth, wet tropical	(H. Loescher and P. Crill, personal communication, 2004)
OD	0.45	ponderosa pine, temperate	(H. Loescher and J. Irvine, personal communication, 2004)
OD	0.55	slash pine, subtropical	<i>Fang and Moncrieff</i> [1998] <sup>g</sup>
OD	0.30	eastern US, mixed, temperate	<i>Davidson et al.</i> [2002]
OD	~0.30	ponderosa pine, temperate	<i>Xu et al.</i> [2001]

<sup>a</sup>CS = closed static, i.e., syringe samples, CD = closed dynamic, e.g., automated chambers, OD = open dynamic, i.e., scrubs headspace below ambient, SL = soda lime.

<sup>b</sup>Includes the potential of an altered diffusion gradient and the spatial heterogeneity on the natural environment.

<sup>c</sup>Systematic leakage for CO<sub>2</sub>.

<sup>d</sup>Measurements made over a uniform synthetic medium.

<sup>e</sup>Chamber overpressurization by 0.5 Pa.

<sup>f</sup>SL tended to underestimate high fluxes (~4.9 μmol m<sup>-2</sup> s<sup>-1</sup>) and overestimated low fluxes (<1.5 μmol m<sup>-2</sup> s<sup>-1</sup>).

<sup>g</sup>Remedied the large effects due to pressure differentials within open systems.

<sup>h</sup>Chamber pressure < atmospheric pressure by ~0.6 Kpa.

<sup>i</sup>Lack of equalizing pressure to ambient.

[41] In tall forests, the structural complexity often does not allow well-mixed conditions to prevail through the canopy profile even under high  $u^*$  values [*Saleska et al.*, 2003; *Loescher et al.*, 2003; *Miller et al.*, 2004], which increase the uncertainty of  $R_e$  estimates. When nocturnal mixing of the canopy airspace was minimal ( $u^* < 0.25 \text{ m s}^{-1}$ ), some studies found chamber-based estimates of total ecosystem respiration were correlated with the change in storage in the canopy air space, which can dominate  $NEE$  estimates during calm wind conditions [*Law et al.*, 1999a; *Lavigne et al.*, 1997]. However, when  $u^* \sim 0.25\text{--}0.3 \text{ m s}^{-1}$ ,  $NEE$  was significantly less than chamber estimates and subcanopy flux measurements [*Law et al.*, 1999b], suggesting that under weak wind conditions, flows with low-CO<sub>2</sub> concentration developed below—and compromised measurements from the above-canopy EC system.

[42] Even under conditions of sustained turbulence, chamber and EC estimates of nocturnal respiration do not always agree [*Chambers et al.*, 2004; *Dore et al.*, 2003; *Drewitt et al.*, 2002; *Law et al.*, 1999b; *Bolstad et al.*, 2004; *Cook et al.*, 2004; *Lavigne et al.*, 1997; *Goulden et al.*, 1996; *Davidson et al.*, 2006]. Because uncertainties in these techniques are dominated by different factors (spatial variability for chambers and intermittent turbulence for EC), it is often difficult to interpret differences between chamber and EC-based estimates of ecosystem respiration. However,

such comparisons may still be used to partition the relative importance of component to the ecosystem flux, determine the process level controls on these fluxes, and constrain parameters used to model and scale estimates of carbon flux [*Law et al.*, 2006].

#### 4. Comparisons Between EC and Allometric Estimates of Net Ecosystem Production

[43] Determining net ecosystem production ( $NEP$ ) by assessing the change in ecosystem carbon mass over time can provide an independent check of integrated EC measurements, and can partition  $NEE$  into the component fluxes.

[44] A mass balance approach for estimating  $NEP$  from biological measurements in forests separates gross photosynthate into three carbon allocation pathways [*Law et al.*, 2003; *Campbell et al.*, 2004; *Randerson et al.*, 2002]:

$$NEP_{\text{foliage}} = NPP_{\text{foliage}} - \text{Litterfall} \quad (4a)$$

$$NEP_{\text{wood}} = NPP_{\text{wood}} - RH_{\text{woody debris}} \quad (4b)$$

$$NEP_{\text{root}} = \Delta\text{coarse root} + \Delta\text{fine root} \quad (4c)$$

$$NEP_{\text{total}} = NEP_{\text{foliage}} + NEP_{\text{wood}} + NEP_{\text{root}} + \Delta\text{soil C} \quad (4d)$$

Where  $NPP_{\text{foliage}}$  is net primary production of tree, shrub, and herb foliage and litterfall is assumed to approximate the heterotrophic decomposition of dead leaves,  $NPP_{\text{wood}}$  is the net primary production of the bole, branches, and bark of trees and shrubs,  $RH_{\text{woody debris}}$  is the heterotrophic respiration from fine and coarse woody debris, and  $\Delta_{\text{coarse root}}$  and  $\Delta_{\text{fine root}}$  are the changes in each of the root carbon pools over time, respectively. Total net ecosystem production ( $NEP_{\text{total}}$ ) can be then calculated by the addition of equations (4a)–(4c) with  $\Delta_{\text{soil C}}$  accounting for any accumulation or depletion of carbon in the mineral soil (e.g., through leaching or erosion). The  $NEP$  of foliage, wood, and root are typically determined by measuring annual growth increment of these tissues, while the change in soil C is typically assessed by the difference between carbon content at two points in time (e.g., five years apart), or by chronosequences and the use of paired plots, trading space for time.

[45] In an alternative mass balance approach,  $NEP$  can be approximated in ecosystems as;

$$NEP_{\text{total}} = NPP_{\text{total}} - HR_{\text{detritus}} - HR_{\text{soil}} \quad (5)$$

where  $NPP_{\text{total}}$  is the total net primary production,  $HR_{\text{detritus}}$  is heterotrophic respiration from all detritus, and  $HR_{\text{soil}}$  is heterotrophic respiration from the soil. Estimating  $NEP$  in this manner requires an estimate of annual soil respiration (typically modeled from periodic measurements) and the fraction of which is heterotrophic metabolism. Separating autotrophic and heterotrophic sources of soil respiration can, however, be difficult. In some cases, where vegetation has transitioned between  $C_3$  and  $C_4$  cover, stable isotopic analysis of respired  $CO_2$  can be used to partition rhizosphere respiration and decomposition [Rochette et al., 1999]. Alternatively, isotopic tracers can be added [Pataki et al., 2003]. In other cases, separation between autotrophic and heterotrophic respiration can be made using trenched plots [cf. Epron et al., 1999] and girdling [cf. Hogberg et al., 2001]. More commonly, the heterotrophic contribution to total soil respiration is estimated by measuring the  $CO_2$  evolved from freshly excised roots [Law et al., 2001b]. There are often other “missing” fluxes such as dissolved organic carbon [Neff and Asner, 2001], herbivory [Clark et al., 2001], and harvest [Scott et al., 2004] that need to be estimated to determine if these components are significant at the measured timescales.

[46] While  $NEP$  derived from mass balance approaches has the potential to serve as a cross check of  $NEE$ , its utility as a validation tool is limited by both the mismatch in temporal scale between the two metrics and the various inherent measurement errors. For the most part,  $NEP$  is integrated over a one-to-five year interval rendering it insensitive to the seasonal changes in carbon flux resolved by EC. While chronosequences (typically spanning decades) are adequate to detect long-term changes in soil C, they are insensitive to assess the short-term changes (e.g., <1 year), which may be contained in  $NEE$  measurements. Alternatively, some studies assume that soil C is in steady state at interdecadal timescales, implying the allocation patterns are also at steady state at shorter timescales (e.g., interannually). The extent that these assumptions are met however, is not clear. Ironically, it is the incongruence

between  $NEE$  and  $NEP$  that may best address the assumptions of short-term steady stasis in soil C.

[47] To properly assess EC estimates of  $NEE$  with mass balance estimates of  $NEP$ , it is important to recognize the various sources of error in  $NEP$ . We recognize three categories:

[48] 1. Measurement error arising from instrument or human error. This error can be random or systematic but is typically small. Examples include the error in measuring tree height or shrub cover.

[49] 2. Sampling error arising from the use of limited sample points to represent a contiguous area or time interval. This error can be large but is typically random. Examples include scaling soil C from a subsample of cores to a hectare plot or by using monthly soil respiration measurements to compute an annual flux.

[50] 3. Parameterization error arising from the use of scaling parameters derived from other studies. These systematic errors can be large or small. Examples include the use of non-site-specific allometric biomass equations or decomposition constants.

[51] Generally speaking, the measurement error of values used to compute biomass such as tree diameter and tree height is low with an estimated CV of <0.02 from a structurally complex tropical forest (D.B. Clark, personal communication, 2003), inferring the CV (precision) is likely closer to 0 in structurally less complex ecosystems. This precision can also be enhanced when care is taken to standardize sampling protocols used from plot-to-plot and year-to-year (e.g., reducing human measurement error, measuring the circumference at the same location on the bole).

[52] Sampling error, on the other hand, can be quite large. Measuring radial bole increment on only one fifth of the trees in a hectare plot and inferring the growth of the remaining boles from diameter measurements imposed a CV of 0.20 of the stand-wide bole increment (i.e., error 2 above [Campbell et al., 2004]). Alternatively, Keller et al. [2001] used a resampling approach (i.e., Monte Carlo) to simulate the uncertainty associated with scaling based on the measured spatial variability in DBH measurements, and found that 15 0.5 ha plots were needed for estimates to be within 20% of the measured mean (and 95% CI, i.e., error 2 above).

[53] Parameterization error arises largely because of the inability to estimate all necessary parameters for a complete budget. The most common is the use of non-site-specific wood decomposition constants and non-site-specific estimates of the heterotrophic contribution to soil respiration [Hanson et al., 2000]. Perhaps more problematic is the use of non-site-specific allometric equations that predict biomass from variables such as height and stem diameter. The log-log nature of most allometric equations mean that even subtle differences in these equations can lead to large systematic errors in biomass estimates [Gower et al., 1996]. Differences among published allometric equations led to a 200% range in biomass estimates for a tropical moist forest in Thailand [Clark et al., 2001] and an 83% range in biomass estimates for a conifer forests in the Pacific NW, USA [Law et al., 2006]. This approach however, assumes that each parameter estimate is normally distributed, which is often not met, and may exaggerate uncertainty in  $NEP$  estimates. For example, Keller et al.

[2001] report the uncertainty in the allometric equation used to estimate the biomass from tropical trees with DBH > 35 cm resulted in estimates within ~20% of the mean (i.e., error 3 above).

[54] Despite a general awareness of the uncertainties associated with mass balance estimates of *NEP*, quantifying these uncertainties in a consistent manner among sites remains a challenge for most researchers. Most studies rely upon their own expert knowledge of a particular ecosystem and/or technique to assign liberal estimates of uncertainty to each component of *NEP* [Harmon *et al.*, 2004; Campbell *et al.*, 2004]. Aggregating the component errors can be done as the root sum square of the component errors or through more sophisticated stochastic resampling procedures (e.g., Monte Carlo), which can account for probability distributions for each specific component variable as well as the covariance in error among component variables [Harmon *et al.*, 2004; Campbell *et al.*, 2004]. Furthermore, second-order uncertainties can be estimated by comparing alternative approaches to estimating *NEP* such as the method defined in equations (4a)–(4d) to those defined in equation (5) [Law *et al.*, 2001a].

[55] For annual integrals of *NEE*, uncertainties can be estimated for gap filling techniques [Falge *et al.*, 2001, 2002],  $u^*$  filtering [e.g., Miller *et al.*, 2004], and bounds established by advection estimates. No matter what the approach, the magnitude of random uncertainty scales with the flux (compare with multisite comparison [Richardson *et al.*, 2006]).

[56] Reducing known systematic errors (by, for example, minimizing pressure differentials between ambient and chamber environments, ogive analyses, and rigorous testing of  $u^*$  filtering, for EC) and maintaining high-quality assurance in data acquisition (e.g., using traceable calibrations, appropriate and frequent maintenance of sensors, established and tested protocols for both allometric and EC methods) is essential in estimating *NEP* and *NEE*. If the magnitude of the summed errors is small (e.g., <50% of the total estimate), then absolute comparisons between EC and allometric estimates may be appropriate, but if they are large (e.g., >1.5x the mean estimate), comparisons are likely to be more qualitative. As a result, single year comparisons of *NEE* and *NEP* may not be reliable, but rather comparisons over multiple years may help to place bounds on expected values and increase our confidence that apparent trends in partitioned fluxes are real.

[57] Comparisons of *NEE* and *NEP* have been used (1) to directly assess long- and short-term climatic effects of the changes in C stores, (2) to test functional relationships used to scale carbon flux from plot-to-region [e.g., Vourlitis *et al.*, 2000; Oechel *et al.*, 1998], and (3) in determining appropriate conditions to apply  $u^*$  thresholds on EC measurements. For example, Curtis *et al.* [2002] reported annual *NEE* and *NEP* from five temperate deciduous forests to be within the range of expected estimates, but found that it took at least five years for the cumulative estimates of *NEE* and *NEP* to converge. Biometric estimates of decomposition of coarse woody debris may have to be averaged over several years for robust comparisons of annual estimates of net uptake. Hence these comparisons are not expected to be good in systems with slow turnover rates.

[58] Saleska *et al.* [2003] and Miller *et al.* [2004] estimated uncertainty in both *NEE* and allometric measures of C flux on the basis of a rigorous empirical determination of  $u^*$  thresholds and confidence intervals, respectively (cf. from an old growth tropical humid forest). Saleska *et al.* [2003] constructed a C budget that approximates *NEP* but lacked litterfall and respiration estimates (except for a rough estimate of coarse woody detritus respiration). Miller *et al.* [2004] used only live aboveground biomass estimates which included trees with a diameter >55 cm, lianas, vines, and hemi-epiphytes. Both of these studies estimated annual *NEE* that were within error bounds established by the allometric estimates, but the allometric-based errors were large,  $\pm 1.6$  and  $2.0 \text{ t C ha}^{-1} \text{ y}^{-1}$  compared to  $\pm 0.5$  and  $0.4 \text{ t C ha}^{-1} \text{ y}^{-1}$  from *NEE* estimates (Saleska *et al.* [2003] and Miller *et al.* [2004], respectively). Including measured and scaled respiration budgets into these estimates would likely add many additional sources of error. This highlights problems associated with the interpretation and utility of incomplete *NEP* budgets for comparison purposes. However, for robust comparisons, all *NEP* components should be estimated (either measured or modeled), particularly belowground processes, and uncertainty estimates provided for both the *NEP* components and *NEE* estimates alike.

## 5. Conclusions

[59] A goal of all flux sites should be to identify and if possible reduce systematic and random errors, thereby increasing the accuracy and precision in *NEE* estimates. Some errors associated with low-frequency flux loss can be addressed through a more rigorous use of ogive analyses (integrated cospectra). At a minimum, ogives can be assessed over time spans that capture the effects of local and mesoscale motions to better determine the appropriate averaging times. However, additional flux loss may occur under specific climate conditions that would not be seen by ogive analyses. Exploration of some as yet undefined similarity functions may be able to determine specific conditions under which low-frequency transport of carbon occurs.

[60] Comparing *NEP* to *NEE* helps validate both estimates and enhances our understanding of carbon sequestration. Such comparisons are also useful for determining the underlying controls on productivity within a particular ecosystem. *NEP* estimates allow the partitioning of carbon stores and fluxes at coarse timescales (annual to multi-annual). In contrast, *NEE* estimates integrate over larger spatial areas and can determine environmental controls on fluxes across fine-to-coarse timescales (half-hourly to multi-annual). Even though chamber measurements have additional utility in partitioning EC *NEE*, the spatial and temporal variability in these measurements are a large source of uncertainty. The number of chambers needed to reduce this uncertainty (i.e., 95% confident to be within 10% of a measured mean soil flux (Figure 3a, top)) can be impractical and unrealistic for many ecosystems. New approaches are needed to reduce the measured variability in respiratory losses.

[61] All sites are subject to potential losses of  $\text{CO}_2$  via advection. Many ecosystems that are important to our understanding of terrestrial carbon budgets are in challenging topography. For this reason, we need to be aware of

conditions that lead to advection on flat and hilly terrain alike. Improvements to current approaches are necessary to increase the precision of within-site measurements and across-site comparisons, and to increase our ability to predict regional carbon flux. Possible approaches may include some combination of designs that can better define and quantify the types of advective flows that exert the greatest influence on carbon flux [e.g., *Staebler*, 2003] coupled with *NEP* estimates to constrain the fraction of advection not accounted for meteorologically, or constraining productivity by other surrogate measures that can utilize multiple constraints derived from functional relationships of *NEP* or *NEE*. In all cases, more effort is needed to assess errors in the multiple approaches to estimating whole ecosystem fluxes and contributing processes.

[62] While it would be nice to suggest a uniform criteria for measurement accuracy and precision, (e.g., to provide comparisons with *NEE*, components *A*, *B* and *C* of *NEP* should be measured with accuracy of *D* while components *X*, *Y* and *Z* may be safely approximated) this is not possible since such guidance is inherently site specific. The sources and characteristics of measurement uncertainty are contingent on the measurement technique used and the physical and biological system under investigation. We support current calls [cf. *Randerson et al.*, 2002; *Chapin et al.*, 2006] for consistency in *NEP* estimation among sites.

[63] The largest challenge in reducing the uncertainties in EC measurements today, is addressing our inability to properly quantify the advection terms. We support exploring new techniques and technologies that can resolve the nighttime CO<sub>2</sub> gradients at the relevant temporal and spatial scales to account for these otherwise missed flows.

[64] **Acknowledgments.** This research was supported by the Office of Science (BER), U.S. Department of Energy, grants DE-FG02-03ER63624 and DE-FG03-01ER63278. Authors wish to thank E. Veldkamp, P. Crill, M. Ryan, M. Goulden, L. Schwendenmann, S. Miller, U. Falk, and J. Irvine for data access, P. Curtis, T. Ocheltree, W. Massman, and the anonymous reviewers for their thoughtful comments.

## References

- Aber, J. D., P. B. Reich, and M. L. Goulden (1996), Extrapolating leaf CO<sub>2</sub> exchange to the canopy: A generalized model of forest photosynthesis compared with measurements by eddy correlation, *Oecologia*, *106*, 257–265.
- Allen, D. W. (1966), Statistics of atomic frequency standards, *Proc. IEEE*, *54*, 221–231.
- Anthoni, P. M., B. E. Law, and M. H. Unsworth (1999), Carbon and water vapor exchange of an open-canopied ponderosa pine ecosystem, *Agric. For. Meteorol.*, *95*, 151–168.
- Anthoni, P. M., M. H. Unsworth, B. E. Law, J. Irvine, D. D. Baldocchi, S. Van Tuyl, and D. Moore (2002), Seasonal differences in carbon and water vapor exchange in young and old-growth Ponderosa pine ecosystems, *Agric. For. Meteorol.*, *111*, 203–222.
- Aubinet, M., et al. (2000), Estimates of the annual net carbon and water exchange of forests: The EUROFLUX methodology, *Adv. Ecol. Res.*, *30*, 113–175.
- Baldocchi, D. D. (2003), Assessing the eddy-covariance technique for evaluating the carbon dioxide exchange rates of ecosystems: Past, present and future, *Global Change Biol.*, *9*, 479–492.
- Baldocchi, D. D., and T. Meyers (1998), On using eco-physiological, micrometeorological and biogeochemical theory to evaluate carbon dioxide, water vapor and trace gas fluxes over vegetation: A perspective, *Agric. For. Meteorol.*, *90*, 1–25.
- Baldocchi, D. D., J. Finnigan, K. Wilson, K. T. Paw U, and E. Falge (2000), On measuring net ecosystem exchange over tall vegetation on complex terrain, *Boundary Layer Meteorol.*, *96*, 257–291.
- Barford, C. C., S. C. Wofsy, M. L. Goulden, J. W. Munger, E. H. Pyle, S. P. Urbanski, L. Hutya, S. R. Saleska, D. Fitzjarrald, and K. Moore (2001), Factors controlling long- and short-term sequestration of atmospheric CO<sub>2</sub> in a mid-latitude forest, *Science*, *294*, 1688–1691.
- Barnes, J. A., and D. W. Allen (1990), Variances based on data with dead time between measurements, *Tech. Note 1318*, Natl. Inst. of Standards and Technol., Gaithersburg, Md.
- Berbigier, P., J. M. Bonnefond, and P. Mellmann (2001), CO<sub>2</sub> and water vapour fluxes for 2 years above Euroflux forest site, *Agric. For. Meteorol.*, *108*, 183–197.
- Berger, B. W., K. J. Davis, C. Yi, P. S. Bakwin, and C. Zhao (2001), Long-term carbon dioxide fluxes from a very tall tower in a northern forest: Long measurement methodology, *J. Atmos. Oceanic Technol.*, *18*, 529–542.
- Betts, A. K., R. L. Desjardins, J. I. MacPherson, and R. D. Kelly (1990), Boundary-layer heat and moisture budgets from FIFE, *Boundary Layer Meteorol.*, *50*, 109–138.
- Betts, A. K., R. L. Desjardins, and J. I. MacPherson (1992), Budget analysis of the boundary-layer grid flights during FIFE 1987, *J. Geophys. Res.*, *97*, 18,533–18,546.
- Black, T. A., et al. (1996), Annual cycles of water vapour and carbon dioxide fluxes in and above a boreal aspen forest, *Global Change Biol.*, *2*, 219–229.
- Bolstad, P. V., K. J. Davis, J. Martin, B. D. Cook, and W. Wang (2004), Component and whole-system respiration fluxes in northern deciduous forests, *Tree Physiol.*, *24*, 493–504.
- Bowling, D. R., P. P. Tans, and R. K. Monson (2001), Partitioning net ecosystem carbon exchange with isotopic fluxes of CO<sub>2</sub>, *Global Change Biol.*, *7*, 127–145.
- Bowling, D. R., S. D. Sargent, B. D. Tanner, and J. R. Ehleringer (2003), Tunable diode laser absorption spectroscopy for stable isotope studies of ecosystem-atmosphere CO<sub>2</sub> exchange, *Agric. For. Meteorol.*, *118*, 1–19.
- Bubier, J., P. Crill, and A. Mosedale (2002), Net ecosystem CO<sub>2</sub> exchange measured by autochambers during the snow-covered season at a temperate peatland, *Hydrol. Process.*, *16*, 3667–3682.
- Butnor, J. R., and K. H. Johnsen (2004), Calibrating soil respiration measures with a dynamic flux apparatus using artificial soil media of varying porosity, *Eur. J. Soil Sci.*, *55*, 639–647.
- Butnor, J. R., K. H. Johnsen, and C. A. Maier (2005), Soil properties differently influence estimates of soil CO<sub>2</sub> efflux from three chamber-based measurement systems, *Biogeochemistry*, *73*, 283–301.
- Campbell, J. L., O. J. Sun, and B. E. Law (2004), Disturbance and net ecosystem production across three climatically distinct forest landscapes, *Global Biogeochem. Cycles*, *18*, GB4017, doi:10.1029/2004GB002236.
- Caughey, S. J., J. C. Wyngaard, and J. C. Kaimal (1979), Turbulence in the evolving stable boundary layer, *J. Atmos. Sci.*, *36*, 1041–1052.
- Chambers, J. Q., E. S. Tribuzy, L. C. Toledo, B. F. Crispim, N. Higuchi, J. dos Santos, A. C. Araújo, B. Kruijt, A. D. Nobre, and S. E. Trumbore (2004), Respiration from a tropical forest ecosystem: Partitioning of sources and low carbon use efficiency, *Ecol. Appl.*, *14*, S72–S88.
- Chapin, F. S., III, et al. (2006), Reconciling carbon-cycle concepts, terminology, and methodology, *Ecosystems*, in press.
- Clark, D. A., S. Brown, D. W. Kicklighter, J. Q. Chambers, and J. R. Thomlinson (2001), Measuring net primary production in forests: Concepts and field methods, *Ecol. Appl.*, *11*, 356–370.
- Clark, K. L., H. L. Gholz, J. B. Moncrieff, F. Cropley, and H. W. Loescher (1999), Environmental controls over net exchanges of carbon dioxide from contrasting Florida ecosystems, *Ecol. Appl.*, *9*, 936–948.
- Cook, B. D., et al. (2004), Carbon exchange and venting anomalies in an upland deciduous forest in northern Wisconsin, USA, *Agric. For. Meteorol.*, *126*, 271–295.
- Curtis, P. S., P. J. Hanson, P. Bolstad, C. Barford, J. C. Randolph, H. P. Schmid, and K. Wilson (2002), Allometric and eddy-covariance based estimates of annual carbon storage in five eastern North American deciduous forests, *Agric. For. Meteorol.*, *113*, 3–19.
- Davidson, E. A., K. Savage, L. V. Verchot, and R. Navarro (2002), Minimizing artifacts and biases in chamber-based measurements of soil respiration, *Agric. For. Meteorol.*, *113*, 21–37.
- Davidson, E. A., A. D. Richardson, K. E. Savage, and D. Y. Hollinger (2006), A distinct seasonal pattern of the ratio of soil respiration to total ecosystem respiration in a spruce-dominated forest, *Global Change Biol.*, *12*, 230–239.
- Davis, K. J., P. S. Bakwin, C. Yi, B. W. Berger, C. Zhao, R. M. Teclaw, and J. G. Isebrands (2006), The annual cycles of CO<sub>2</sub> and H<sub>2</sub>O exchange over a northern mixed forest as observed from a very tall tower, *Ecol. Appl.*, in press.
- Desjardins, R. L., et al. (1997), Scaling up flux measurements for the boreal forest using aircraft-tower combination, *J. Geophys. Res.*, *102*, 29,125–29,133.
- Doran, J. C. (1991), The effects of ambient winds on valley drainage flows, *Boundary Layer Meteorol.*, *55*, 177–189.
- Doran, J. C., and T. W. Horst (1981), Velocity and temperature oscillations in drainage winds, *J. Appl. Meteorol.*, *20*, 361–364.



- Doran, J. C., and S. Zhong (1994), Regional drainage flows in the Pacific-Northwest, *Mon. Weather Rev.*, *122*, 1158–1167.
- Dore, S., G. J. Hymus, D. P. Johnson, C. R. Hinkle, R. Valentini, and B. G. Drake (2003), Cross validation of open-top chamber and eddy covariance measurements of ecosystem CO<sub>2</sub> exchange in a Florida scrub-oak ecosystem, *Global Change Biol.*, *9*, 84–95.
- Drake, B. G., M. S. Muehe, G. Peresta, M. A. Gonzalez-Meler, and R. Matamala (1996), Acclimation of photosynthesis, respiration and ecosystem carbon flux of a wetland on Chesapeake Bay, Maryland to elevated atmospheric CO<sub>2</sub> concentration, *Plant Soil*, *187*, 111–118.
- Drewitt, G. B., T. A. Black, Z. Nestic, E. R. Humphreys, E. M. Jork, R. Swanson, G. J. Ethier, T. Griffiths, and K. Morgenstern (2002), Measuring forest floor CO<sub>2</sub> fluxes in a Douglas-fir forest, *Agric. For. Meteorol.*, *101*, 299–317.
- Epron, D., L. Farque, E. Lucot, and P. M. Badot (1999), Soil CO<sub>2</sub> efflux in a beech forest: The contribution of root respiration, *Ann. For. Sci.*, *56*, 289–295.
- Evensen, G. (2003), The ensemble Kalman filter: Theoretical formulation and practical implementation, *Ocean Dyn.*, *53*, 343–367.
- Falge, E., et al. (2001), Gap filling strategies for defensible annual sums of net ecosystem exchange, *Agric. For. Meteorol.*, *107*, 43–69.
- Falge, E., et al. (2002), Seasonality of ecosystem respiration and gross primary production as derived from FLUXNET measurements, *Agric. For. Meteorol.*, *113*, 53–74.
- Falk, U. (2004), Turbulent fluxes of CO<sub>2</sub>, H<sub>2</sub>O and energy in the atmospheric boundary layer above tropical vegetation investigated by eddy-covariance measurements, Ph.D. dissertation, Inst. of Bioclimatol., Univ. Göttingen, Germany.
- Fang, C., and J. B. Moncrieff (1998), An open-top chamber for measuring soil respiration and the influence of pressure difference on CO<sub>2</sub> efflux measurement, *Functional Ecol.*, *12*, 319–325.
- Fang, C., J. B. Moncrieff, H. L. Gholz, and K. L. Clark (1998), Soil CO<sub>2</sub> efflux and its spatial variation in a Florida slash pine plantation, *Plant Soil*, *205*, 135–146.
- Finkelstein, P. L., and P. F. Sims (2001), Sampling error in eddy correlation flux measurements, *J. Geophys. Res.*, *106*, 3503–3509.
- Finnigan, J. J. (1999), A comment on the paper by Lee (1998): “On micrometeorological observations of surface-air exchange over tall vegetation”, *Agric. For. Meteorol.*, *97*, 55–64.
- Finnigan, J. J., R. Clement, Y. Malhi, R. Leuning, and H. A. Cleugh (2003), A re-evaluation of long-term flux measurement techniques, part I: Averaging and coordinate rotation, *Boundary Layer Meteorol.*, *107*, 1–48.
- Foken, T., and S. P. Oncley (1995), Results of the workshop “Instrumental and methodological problems of land surface flux measurements”, *Bull. Am. Meteorol. Soc.*, *76*, 1191–1193.
- Foken, T., and B. Wichura (1996), Tools for quality assessment of surface-based flux measurements, *Agric. For. Meteorol.*, *78*, 83–105.
- Foken, T., M. Gockede, M. Mauder, L. Mahrt, B. Amiro, and J. Munger (2004), Post-field data quality control, in *Handbook of Micrometeorology: A Guide for Surface Flux Measurements*, edited by X. Lee, pp. 181–207, Springer, New York.
- Friehe, C. A., W. J. Shaw, D. P. Rogers, K. L. Davidson, W. G. Large, S. A. Stage, G. H. Crescenti, S. J. S. Khalsa, G. K. Greenhut, and F. Li (1991), Air-sea fluxes and surface-layer turbulence around a sea surface temperature front, *J. Geophys. Res.*, *96*, 8593–8609.
- Gedayloo, T., W. E. Clements, S. Barr, and J. A. Archuleta (1980), Nocturnal drainage wind characteristics in two converging airsheds, *Bull. Am. Meteorol. Soc.*, *61*, 89–89.
- Geiger, R. (1961), *The Climate Near the Ground*, Harvard Univ. Press, 611 pp., Cambridge, Mass.
- Gluhovsky, A., and E. Agee (1994), A definitive approach to turbulence statistical studies in planetary boundary layers, *J. Atmos. Sci.*, *51*, 1682–1690.
- Goldstein, A. H., N. E. Hultman, J. M. Fracheboud, M. R. Bauer, J. A. Panek, M. Xu, Y. Qi, A. B. Guenther, and W. Baugh (2000), Effects of climate variability on the carbon dioxide, water and sensible heat fluxes above a ponderosa pine plantation in the Sierra Nevada (CA), *Agric. For. Meteorol.*, *101*, 113–129.
- Goulden, M. L., and P. M. Crill (1997), Automated measurements of CO<sub>2</sub> exchange at the moss surface of a black spruce forest, *Tree Physiol.*, *17*, 537–542.
- Goulden, M. L., J. W. Munger, S.-M. Fan, B. C. Daube, and S. C. Wofsy (1996), Measurements of carbon sequestration by long-term eddy covariance: Methods and a critical evaluation of accuracy, *Global Change Biol.*, *3*, 169–182.
- Goulden, M. L., B. C. Daube, S. M. Fan, D. J. Sutton, A. Bazzaz, J. W. Munger, and S. C. Wofsy (1997), Physiological responses of a black spruce forest to weather, *J. Geophys. Res.*, *102*, 28,987–28,996.
- Gove, J. H., and D. Y. Hollinger (2006), Application of a dual unscented Kalman filter for simultaneous state and parameter estimation in problems of surface-atmosphere exchange, *J. Geophys. Res.*, *111*, D08S07, doi:10.1029/2005JD006021.
- Gower, S. T., S. Pongracic, and J. J. Landsberg (1996), A global trend in belowground carbon allocation: Can we use the relationship at smaller scales?, *Ecol.*, *77*, 1750–1755.
- Grace, J., et al. (1995), Fluxes of carbon-dioxide and water-vapor over an undisturbed tropical forest in south-west Amazonia, *Global Change Biol.*, *1*, 1–12.
- Grace, J., Y. Malhi, J. Lloyd, J. McIntyre, A. C. Miranda, P. Meir, and H. S. Miranda (1996), The use of eddy covariance to infer the net carbon dioxide uptake of Brazilian rain forest, *Global Change Biol.*, *2*, 209–217.
- Granier, A., et al. (2000), The carbon balance of a young Beech forest, *Functional Ecol.*, *14*, 312–325.
- Gu, L., et al. (2005), Objective threshold determination for nighttime eddy flux filtering, *Agric. For. Meteorol.*, *128*, 179–197.
- Gudiksen, P. H., J. M. Leone, C. W. King, D. Ruffieux, and W. D. Neff (1992), Measurements and modeling of the effects of ambient meteorology on nocturnal drainage flows, *J. Appl. Meteorol.*, *9*, 1023–1032.
- Hansen, F. V. (1993), Surface roughness lengths, ARL technical report, U.S. Army, White Sands Missile Range, N. M.
- Hanson, P. J., N. T. Edwards, C. T. Garten Jr., and J. A. Andrews (2000), Separating root and soil microbial contributions to soil respiration: A review of methods and observations, *Biogeochemistry*, *48*, 115–146.
- Harmon, M. E., K. Bible, M. G. Ryan, D. C. Shaw, H. Chen, J. Klopatek, and X. Li (2004), Production, respiration and overall carbon balance in an old-growth *Pseudotsuga-Tsuga* forest ecosystem, *Ecosystems*, *7*, 498–512.
- Hogberg, P., A. Nordgren, N. Buchmann, A. F. S. Taylor, A. Ekblad, M. N. Hogberg, G. Nyberg, M. Ottoson-Lofvenius, and D. J. Read (2001), Large-scale forest gridding shows that current photosynthesis drives soil respiration, *Nature*, *411*, 789–792.
- Hollinger, D. Y., and A. D. Richardson (2005), Uncertainty in eddy covariance measurements and its application to physiological models, *Tree Physiol.*, *25*, 873–885.
- Hollinger, D. Y., F. M. Kelliher, J. N. Byers, J. E. Hunt, T. M. McSeveny, and P. L. Weir (1994), Carbon dioxide exchange between an undisturbed old-growth temperate forest and the atmosphere, *Ecology*, *75*, 134–150.
- Hollinger, D. Y., S. M. Goltz, E. A. Davidson, J. T. Lee, K. Tu, and H. T. Valentine (1999), Seasonal patterns and environmental control of carbon dioxide and water vapour exchange in an ecotonal boreal forest, *Global Change Biol.*, *5*, 891–902.
- Hollinger, D. Y., et al. (2004), Spatial and temporal variability in forest-atmosphere CO<sub>2</sub> exchange, *Global Change Biol.*, *10*, 1689–1706.
- Irvine, J., and B. E. Law (2002), Contrasting soil respiration in young and old-growth ponderosa pine forests, *Global Change Biol.*, *8*, 1183–1194.
- Janssens, I. A., A. S. Kowalski, B. Longdoz, and R. Ceulemans (2000), Assessing forest soil CO<sub>2</sub> efflux: An *in situ* comparison of four techniques, *Tree Physiol.*, *20*, 23–32.
- Janssens, I. A., A. S. Kowalski, and R. Ceulemans (2001), Forest floor CO<sub>2</sub> fluxes estimated by eddy covariance and chamber-based model, *Agric. For. Meteorol.*, *102*, 61–69.
- Jarvis, P. G., J. M. Massheder, S. E. Hale, J. B. Moncrieff, M. Rayment, and S. L. Scott (1997), Seasonal variation of carbon dioxide, water vapor, and energy exchanges of a boreal black spruce forest, *J. Geophys. Res.*, *102*, 28,953–28,966.
- Jarvis, A. J., V. J. Stauch, K. Schulz, and P. C. Young (2004), The seasonal temperature dependency of photosynthesis and respiration in two deciduous forests, *Global Change Biol.*, *10*, 939–950.
- Julier, S., and J. Uhlmann (1997), A new extension of the Kalman filter to nonlinear systems, *Proc. SPIE*, *3068*, 182–193.
- Kaimal, J. C., and J. S. Finnigan (1994), *Atmospheric Boundary-Layer Flows: Their Structure and Measurement*, 453 pp., Oxford Univ. Press, New York.
- Kaimal, J. C., and J. C. Wyngaard (1990), The Kansas and Minnesota experiments, *Boundary Layer Meteorol.*, *50*, 31–47.
- Kalman, R. E. (1960), A new approach to linear filtering and prediction problems, *Trans. ASME J.*, *82*, 35–45.
- Katul, G. G., C. D. Geron, C. I. Hsieh, B. Vidakovic, and A. B. Guenther (1998), Active turbulence and scalar transport near the forest-atmosphere interface, *J. Appl. Meteorol.*, *37*, 1533–1546.
- Keller, M., M. Palace, and G. Hurtt (2001), Biomass estimation in the Tapajos National Forest, Brazil—Examination of sampling and allometric uncertainties, *For. Ecol. Manage.*, *154*, 371–382.
- Kruijt, B., J. A. Elbers, C. von Randow, A. C. Araújo, P. J. Oliveira, A. Culf, A. O. Manzi, A. D. Nobre, P. Kabat, and E. J. Moors (2004), The robustness of eddy correlation fluxes for Amazon rain forest conditions, *Ecol. Appl.*, *14*, S101–S113.
- Lavigne, M. B., et al. (1997), Comparing nocturnal eddy covariance measurements to estimates of ecosystem respiration made by scaling chamber

- measurements at six coniferous boreal sites, *J. Geophys. Res.*, *102*, 28,977–28,985.
- Law, B. E., M. G. Ryan, and P. M. Anthoni (1999a), Seasonal and annual respiration of a ponderosa pine ecosystem, *Global Change Biol.*, *5*, 169–182.
- Law, B. E., D. D. Baldocchi, and P. M. Anthoni (1999b), Below-canopy and soil CO<sub>2</sub> fluxes in a ponderosa pine forest, *Agric. For. Meteorol.*, *94*, 171–188.
- Law, B. E., P. Thornton, J. Irvine, S. Van Tuyl, and P. Anthoni (2001a), Carbon storage and fluxes in ponderosa pine forests at different developmental stages, *Global Change Biol.*, *7*, 755–777.
- Law, B. E., F. M. Kelliher, D. D. Baldocchi, P. M. Anthoni, J. Irvine, D. Moore, and S. Van Tuyl (2001b), Spatial and temporal variation in respiration in a young ponderosa pine forest during a summer drought, *Agric. For. Meteorol.*, *110*, 27–43.
- Law, B. E., et al. (2002), Environmental controls over carbon dioxide and water vapor exchange of terrestrial vegetation, *Agric. For. Meteorol.*, *113*, 97–120.
- Law, B. E., O. Sun, J. L. Campbell, S. Van Tuyl, and P. Thornton (2003), Changes in carbon storage and fluxes in a chronosequence of ponderosa pine, *Global Change Biol.*, *9*, 510–524.
- Law, B. E., D. Turner, J. L. Campbell, M. Leftsky, M. Guzy, O. Sun, S. Van Tuyl, and W. Cohen (2006), Carbon fluxes across regions: Observational constraints at multiple scales, in *Scaling and Uncertainty Analysis in Ecology: Methods and Applications*, edited by J. Wu et al., pp. 167–190, Springer, New York.
- Leclerc, M. Y., A. Karipot, T. Prabha, G. Allwine, B. Lamb, and H. L. Gholz (2003), Impact of non-local advection on flux footprints over a tall forest canopy: A tracer flux experiment, *Agric. For. Meteorol.*, *115*, 19–30.
- Lee, I. Y., R. L. Coulter, H. M. Park, and J. H. Oh (1995), Numerical-simulation of nocturnal drainage flow properties in a rugged canyon, *Boundary Layer Meteorol.*, *72*, 305–321.
- Lee, X. (1998), On micrometeorological observations of surface-air exchange over tall vegetation, *Agric. For. Meteorol.*, *91*, 39–49.
- Lee, X. H., and T. A. Black (1993), Atmospheric-turbulence within and above a Douglas-fir stand: Statistical properties of the velocity-field, *Boundary Layer Meteorol.*, *64*, 149–174.
- Lee, X. H., and X. Z. Hu (2002), Forest-air fluxes of carbon, water and energy over non-flat terrain, *Boundary Layer Meteorol.*, *103*, 277–301.
- Lee, X., J. D. Fuentes, R. M. Staebler, and H. H. Neumann (1999), Long-term observation of the atmospheric exchange of CO<sub>2</sub> with a temperate deciduous forest in southern Ontario, Canada, *J. Geophys. Res.*, *104*, 15,975–15,984.
- Lemon, E. R. (1960), Photosynthesis under field conditions. II. An aerodynamic method for determining the turbulent carbon dioxide exchange between the atmosphere and a corn field, *J. Agron.*, *52*, 697–703.
- Lenschow, D. H., J. Mann, and L. Kristensen (1994), How long is long enough when measuring fluxes and other turbulence statistics?, *J. Atmos. Oceanic Technol.*, *11*, 661–673.
- Lindroth, A., A. Grelle, and A. S. Moren (1998), Long-term measurements of boreal forest carbon balance reveal large temperature sensitivity, *Global Change Biol.*, *4*, 443–450.
- Loescher, H. W., S. F. Oberbauer, H. L. Gholz, and D. B. Clark (2003), Environmental controls on net ecosystem-level carbon exchange and productivity in a Central American tropical wet forest, *Global Change Biol.*, *9*, 396–412.
- Loescher, H. W., et al. (2005), Comparison of temperature and wind statistics in contrasting environments among different sonic anemometer-thermometers, *Agric. For. Meteorol.*, *133*, 119–139.
- Loescher, H. W., G. Starr, T. A. Martin, M. Binford, and H. L. Gholz (2006), The effect of daytime circulations on eddy covariance carbon dioxide flux measurements over a *Pinus elliottii* canopy, *J. Appl. Meteorol.*, in press.
- Longdoz, B., M. Yernaux, and M. Aubinet (2000), Soil CO<sub>2</sub> efflux measurements in a mixed forest: Impact of chamber disturbances, spatial variability and seasonal evolution, *Global Change Biol.*, *6*, 907–917.
- Lumley, J. L., and H. A. Panofsky (1964), *The Structure of Atmospheric Turbulence*, 239 pp., Wiley-Interscience, Hoboken, N. J.
- Lund, C. P., W. J. Riley, L. L. Pierce, and C. B. Field (1999), The effects of chamber pressurization on soil-surface CO<sub>2</sub> flux and the implications for NEE measurements under elevated CO<sub>2</sub>, *Global Change Biol.*, *5*, 269–281.
- Mahrt, L. (1992), Momentum balance of gravity flows, *J. Atmos. Sci.*, *39*, 2701–2711.
- Mahrt, L. (1998), Flux sampling strategy for aircraft and tower observations, *J. Atmos. Oceanic Technol.*, *15*, 416–429.
- Mahrt, L., X. Lee, A. Black, H. Neumann, and R. M. Staebler (2000), Nocturnal mixing in a forest subcanopy, *Agric. For. Meteorol.*, *101*, 67–78.
- Mahrt, L., D. Vickers, R. Nakamura, J. Sun, S. Burns, D. Lenschow, and M. Soler (2001a), Shallow drainage flows, *Boundary Layer Meteorol.*, *101*, 243–260.
- Mahrt, L., E. Moore, D. Vickers, and N.-O. Jensen (2001b), Dependence of turbulent and mesoscale velocity variances on scale and stability, *J. Appl. Meteorol.*, *40*, 628–641.
- Malhi, Y., A. D. Nobre, J. Grace, B. Kruijff, M. G. P. Pereira, A. Culf, and S. Scott (1998), Carbon dioxide transfer over a central Amazonian rain forest, *J. Geophys. Res.*, *103*, 31,593–31,612.
- Malhi, Y., D. D. Baldocchi, and P. G. Jarvis (1999), The carbon balance of tropical, temperate and boreal forests, *Plant Cell Environ.*, *22*, 715–740.
- Manins, P. (1992), Vertical fluxes in katabatic flows, *Boundary Layer Meteorol.*, *60*, 169–178.
- Mann, J., and D. H. Lenschow (1994), Errors in airborne flux measurements, *J. Geophys. Res.*, *99*, 14,519–14,526.
- Martens, C. S., et al. (2003), Radon fluxes in tropical forest ecosystems of Brazilian Amazonia: Night-time CO<sub>2</sub> net ecosystem exchange derived from radon and eddy covariance, *Global Change Biol.*, *10*, 618–629.
- Massman, W. J., and X. Lee (2002), Eddy covariance flux corrections and uncertainties in long term studies of carbon and energy, *Agric. For. Meteorol.*, *113*, 121–144.
- Massman, W. J., R. A. Sommerfeld, A. R. Mosier, K. F. Zeller, T. J. Hehn, and S. G. Rochelle (1997), A model investigation of turbulence-driven pressure-pumping effects on the rate of diffusion of CO<sub>2</sub>, N<sub>2</sub>O, and CH<sub>4</sub> through layered snowpacks, *J. Geophys. Res.*, *102*, 18,851–18,863.
- Meyers, T. P. (2001), A comparison of summertime water and CO<sub>2</sub> fluxes over rangeland for well watered and drought conditions, *Agric. For. Meteorol.*, *106*, 205–214.
- Miller, D. R., J. D. Bergen, and G. Neuroth (1983), Cold air drainage in a narrow forested valley, *For. Sci.*, *29*, 357–370.
- Miller, S. D., M. L. Goulden, M. C. Menton, H. R. da Rocha, H. C. de Freitas, A. M. S. Figueira, and C. A. D. de Sousa (2004), Biometric and micrometeorological measurements of tropical forest carbon balance, *Ecol. Appl.*, *14*, S114–S126.
- Moncrieff, J. B., Y. Malhi, and R. Leuning (1996), The propagation or errors in long-term measurements of land-atmosphere fluxes of carbon and water, *Global Change Biol.*, *2*, 231–240.
- Monteith, J. L., and G. Szeicz (1960), The CO<sub>2</sub> flux over a field of sugar beets, *Q. J. R. Meteorol. Soc.*, *86*, 205–214.
- Nay, S. M. (1994), Evaluation methods for measuring soil CO<sub>2</sub>-efflux in artificial, controlled and natural ecosystems, M.S. thesis, 69 pp., Oregon State Univ., Corvallis.
- Nay, S. M., K. G. Mattson, and B. T. Borman (1994), Biases of chamber methods for measuring CO<sub>2</sub> efflux demonstrated with laboratory apparatus, *Ecology*, *75*, 2460–2463.
- Neff, J. C., and G. P. Asner (2001), Dissolved organic carbon in terrestrial ecosystems: Synthesis and a model, *Ecosystems*, *4*, 29–48.
- Norman, J. M., C. J. Kurharik, S. T. Gower, D. D. Baldocchi, P. M. Crill, M. Rayment, K. Savage, and R. G. Striegl (1997), A comparison of six methods for measuring soil-surface carbon dioxide fluxes, *J. Geophys. Res.*, *102*, 28,771–28,777.
- Oechel, W. C., G. L. Vourlitis, S. Brooks, T. L. Crawford, and E. Dumas (1998), Intercomparison among chamber, tower, and aircraft net CO<sub>2</sub> and energy fluxes measured during the Arctic System Science Land-Atmosphere-Ice Interactions (ARCSS-LAII) flux study, *J. Geophys. Res.*, *103*, 28,993–29,004.
- Oechel, W. C., G. Vourlitis, S. J. Hastings, R. C. Zulueta, L. Hinzman, and D. Kane (2000), Acclimation of ecosystem CO<sub>2</sub> exchange in the Alaskan Arctic in response to decadal climate warming, *Nature*, *406*, 978–981.
- Oncley, S. P., J. A. Businger, E. C. Itsweire, C. A. Friche, J. C. LaRue, and S. S. Chang (1990), Surface layer profiles and turbulence measurements over uniform land under near-neutral conditions, paper presented at the 9th Symposium on Boundary Layer and Turbulence, Am. Meteorol. Soc., Roskilde, Denmark, 30 April to 3 May.
- Panofsky, H. A., and J. A. Dutton (1984), *Atmospheric Turbulence*, 397 pp., John Wiley, Hoboken, N. J.
- Papale, D., and R. Valentini (2003), A new assessment of European forests carbon exchanges by eddy fluxes and artificial neural network spatialization, *Global Change Biol.*, *9*, 525–535.
- Papale, D., and R. Valentini (2004), Spatial and temporal assessment of biospheric carbon fluxes at continental scale by neural network optimization, in *Forest at the Land-Atmosphere Interface*, edited by M. Mencuccini et al., pp. 203–229, Cab Int., Wallingford, U. K.
- Pataki, D. E., et al. (2003), Tracing changes in ecosystem function under elevated carbon dioxide conditions, *BioScience*, *53*, 805–818.
- Paw U, K. T., D. Baldocchi, T. P. Meyers, and K. B. Wilson (2000), Correction of eddy-covariance measurements incorporating both advective effects and density fluxes, *Boundary Layer Meteorol.*, *97*, 487–511.
- Pilegaard, K., P. Hummelshøj, N.-O. Jensen, and Z. Chen (2001), Two years of continuous CO<sub>2</sub> eddy-flux measurements over a Danish beech forest, *Agric. For. Meteorol.*, *107*, 29–41.

- Randerson, J. T., F. S. Chapin, J. W. Harden, J. C. Neff, and M. E. Harmon (2002), Net ecosystem production: A comprehensive measure of net carbon accumulation by ecosystems, *Ecol. Appl.*, *12*, 937–947.
- Raupach, M. R. (1991), Vegetation-atmosphere interaction in homogeneous and heterogeneous terrain: Some implications of mixed-layer dynamics, *Vegetation*, *91*, 105–120.
- Raupach, M. R., and A. S. Thom (1981), Turbulence in and above plant canopies, *Annu. Rev. Fluid Mech.*, *13*, 97–129.
- Rayment, M. B. (2000), Closed chamber systems underestimate soil CO<sub>2</sub> efflux, *Eur. J. Soil Sci.*, *51*, 107–110.
- Rayment, M. B., and P. G. Jarvis (2000), Temporal and spatial variation of soil CO<sub>2</sub> efflux in a Canadian boreal forest, *Soil Biol. Biochem.*, *32*, 35–45.
- Reynolds, O. (1901), *Papers on Mechanical and Physical Subjects 1881–1900: Collected Work*, vol. 2, 733 pp., Cambridge Univ. Press, New York.
- Richardson, A. D., and D. Y. Hollinger (2005), Statistical modeling of ecosystem respiration using eddy covariance data: Maximum likelihood parameter estimation, and Monte Carlo simulation of model and parameter uncertainty, applied to three simple models, *Agric. For. Meteorol.*, *31*, 191–208.
- Richardson, A. D., et al. (2006), A multi-site analysis of random error in tower-based measurements of carbon and energy fluxes, *Agric. For. Meteorol.*, *136*, 1–18.
- Rochette, P., L. B. Flanagan, and E. G. Gregorich (1999), Separating soil respiration into plant and soil components using analysis of the natural abundance of carbon-13, *Soil Sci. Soc. Am. J.*, *63*, 1207–1213.
- Russell, C. A., and R. P. Voroney (1998), Carbon dioxide efflux from the floor of a boreal aspen forest. I. Relationship to environmental variables and estimates of C respired, *Can. J. Soil Sci.*, *78*, 301–310.
- Ryan, M. G., and B. E. Law (2005), Interpreting, measuring, and modeling soil respiration, *Biogeochemistry*, *73*, 3–27.
- Sakai, R., D. Fitzjarrald, and K. E. Moore (2001), Importance of low-frequency contributions to eddy fluxes observed over rough surfaces, *J. Appl. Meteorol.*, *40*, 2178–2192.
- Saleska, S. R., et al. (2003), Carbon fluxes in old-growth Amazonian rainforests: Seasonality and disturbance-induced net carbon loss, *Science*, *302*, 1554–1557.
- Schmid, H. P. (1997), Experimental design for flux measurements: Matching scales of observations and fluxes, *Agric. For. Meteorol.*, *87*, 179–200.
- Schmid, H. P., C. S. B. Grimmond, F. Cropley, B. Offerle, and H.-B. Su (2000), Measurements of CO<sub>2</sub> and energy fluxes over a mixed hardwood forest in the mid-western United States, *Agric. For. Meteorol.*, *103*, 357–374.
- Schmid, H. P., H.-B. Su, C. S. Vogel, and P. S. Curtis (2003), Ecosystem-atmosphere exchange of carbon dioxide over a mixed hardwood forest in northern lower Michigan, *J. Geophys. Res.*, *108*(D14), 4417, doi:10.1029/2002JD003011.
- Schulz, K., A. Jarvis, and K. Beven (2001), The predictive uncertainty of land surface fluxes in response to increasing ambient carbon dioxide, *J. Clim.*, *14*, 2551–2562.
- Scott, N. A., C. A. Rodrigues, H. Hughes, J. T. Lee, E. A. Davidson, D. B. Dail, P. Malerba, and D. Y. Hollinger (2004), Changes in carbon storage and net carbon exchange one year after an initial shelterwood harvest at Howland Forest, ME, *Environ. Manage.*, *33*, suppl. 1, S9–S22.
- Shen, S. H., and M. Y. Leclerc (1997), Modeling the turbulence structure in the canopy layer, *Agric. For. Meteorol.*, *87*, 3–25.
- Soler, M. R., C. Infante, P. Buenestado, and L. Mahrt (2002), Observations of nocturnal drainage flow, *Boundary Layer Meteorol.*, *105*, 253–273.
- Staebler, R. (2003), Forest subcanopy flows and micro-scale advection of carbon dioxide, Ph.D. dissertation, 193 pp., State Univ. of N. Y., Albany.
- Staebler, R., and D. R. Fitzjarrald (2004), Observing subcanopy CO<sub>2</sub> advection, *Agric. For. Meteorol.*, *122*, 139–156.
- Staebler, R., and D. R. Fitzjarrald (2005), Measuring canopy structure and the kinematics of subcanopy flows in two forests, *J. Appl. Meteorol.*, *44*, 1161–1179.
- Sun, J., R. Desjardins, L. Mahrt, and I. MacPherson (1998), Transport of carbon dioxide, water vapor and ozone by turbulence and local circulations, *J. Geophys. Res.*, *103*, 25,873–25,885.
- Suyker, A. E., S. B. Verma, and G. G. Burba (2003), Interannual variability in net CO<sub>2</sub> exchange of a native tallgrass prairie, *Global Change Biol.*, *9*, 255–265.
- Thornton, P. E., et al. (2002), Modeling and measuring the effects of disturbance history and climate on carbon and water budgets in evergreen needleleaf forests, *Agric. For. Meteorol.*, *113*, 185–222.
- Turnipseed, A. A., D. E. Anderson, P. D. Blanken, W. M. Baugh, and R. K. Monson (2003), Airflows and turbulent flux measurements in mountainous terrain. part I. Canopy and local effects, *Agric. For. Meteorol.*, *119*, 1–21.
- Twine, T. E., W. P. Kustas, J. M. Norman, D. R. Cook, P. R. Houser, T. P. Meyers, J. H. Prueger, P. J. Starks, and M. L. Wesely (2000), Correcting eddy-covariance flux underestimates over a grassland, *Agric. For. Meteorol.*, *103*, 279–300.
- Valentini, R., et al. (2000), Respiration as the main determinant of carbon balance in European forests, *Nature*, *404*, 861–865.
- van Wijk, M. T., and W. Bouten (2002), Simulating daily and half-hourly fluxes of forest carbon dioxide and water vapor exchange with a simple model of light and water use, *Ecosystems*, *5*, 597–610.
- Verma, S., J. Kim, and R. J. Clement (1989), Carbon dioxide, water vapor and sensible heat fluxes over a tall grass prairie, *Boundary Layer Meteorol.*, *46*, 53–67.
- Vickers, D., and L. Mahrt (1997), Quality control and flux sampling problems for tower and aircraft data, *J. Atmos. Oceanic Technol.*, *14*, 512–526.
- Vickers, D., and L. Mahrt (2003), The cospectral gap and turbulent flux calculations, *J. Atmos. Oceanic Technol.*, *20*, 660–672.
- Vickers, D., and L. Mahrt (2006a), A solution for flux contamination by mesoscale motions with very weak turbulence, *Boundary Layer Meteorol.*, *118*, 431–447.
- Vickers, D., and L. Mahrt (2006b), Contrasting mean vertical motion from tilt correction methods and mass continuity, *Agric. For. Meteorol.*, in press.
- Vourlitis, G. L., and W. C. Oechel (1997), Landscape-scale CO<sub>2</sub>, H<sub>2</sub>O vapour and energy flux of moist-wet coastal tundra ecosystems over two growing seasons, *J. Ecol.*, *85*, 575–590.
- Vourlitis, G. L., and W. C. Oechel (1999), Eddy covariance measurements of CO<sub>2</sub> and energy fluxes of an Alaskan tussock tundra ecosystem, *Ecology*, *80*, 686–701.
- Vourlitis, G. L., W. C. Oechel, A. Hope, D. Stow, B. Boynton, J. Verfaillie Jr., R. Zuluetta, and S. J. Hastings (2000), Physiological models for scaling plot measurements of CO<sub>2</sub> flux across an Arctic tundra landscape, *Ecol. Appl.*, *10*, 60–72.
- Weber, R. O. (1999), Remarks on the definition and estimation of friction velocity, *Boundary Layer Meteorol.*, *93*, 197–209.
- Welch, G., and G. Bishop (2001), *An Introduction to the Kalman Filter; Course 8*, Siggraph, Los Angeles, Calif. (available at [http://www.cs.unc.edu/~tracker/media/pdf/SIGGRAPH2001\\_CoursePack\\_08.pdf](http://www.cs.unc.edu/~tracker/media/pdf/SIGGRAPH2001_CoursePack_08.pdf))
- Wesely, M. L., and R. L. Hart (1985), Variability of short term eddy-correlation estimates of mass exchange, in *The Forest-Atmosphere Interaction*, edited by B. A. Hutchison and B. B. Hicks, pp. 591–612, Springer, New York.
- Wilczak, J. M., S. P. Oncley, and S. A. Stage (2001), Sonic anemometer tilt correction algorithms, *Boundary Layer Meteorol.*, *99*, 127–150.
- Williams, M., B. E. Law, P. M. Anthoni, and M. H. Unsworth (2001), Use of a simulation model and ecosystem flux data to examine carbon-water interactions in ponderosa pine, *Tree Physiol.*, *21*, 287–298.
- Williams, M., P. A. Schwarz, B. E. Law, J. Irvine, and M. R. Kurpius (2005), An improved analysis of forest carbon dynamics using data assimilation, *Global Change Biol.*, *11*, 89–105.
- Wilson, K. B., and D. D. Baldocchi (2001), Comparing independent estimates of carbon dioxide exchange over 5 years at a deciduous forest in the southeastern United States, *J. Geophys. Res.*, *106*, 34,167–34,178.
- Wilson, K. B., and T. P. Meyers (2001), The spatial variability of energy and carbon dioxide fluxes at the floor of a deciduous forest, *Boundary Layer Meteorol.*, *98*, 443–473.
- Xu, M., T. A. DeBiase, Y. Qi, A. Goldstein, and Z. Liu (2001), Ecosystem respiration in a young ponderosa pine plantation in the Sierra Nevada mountains, California, *Tree Physiol.*, *21*, 309–318.
- Yamamoto, S., S. Murayama, N. Saigusa, and H. Kondo (1999), Seasonal and inter-annual variation of CO<sub>2</sub> flux between a temperate forest and the atmosphere in Japan, *Tellus, Ser. B*, *51*, 402–413.
- Yang, P. C., T. A. Black, H. H. Neumann, M. D. Novak, and P. D. Blanken (1999), Spatial and temporal variability of CO<sub>2</sub> concentration and flux in a boreal aspen forest, *J. Geophys. Res.*, *104*, 27,653–27,661.
- Yi, C., K. J. Davis, P. S. Bakwin, B. W. Berger, and L. C. Marr (2000), Influence of advection on measurements of the net ecosystem-atmosphere exchange of CO<sub>2</sub> from a very tall tower, *J. Geophys. Res.*, *105*, 9991–9999.
- Yoshino, M. M. (1975), *Climate in a Small Area*, 549 pp., Tokyo Press, Tokyo.

J. Campbell, B. E. Law, and H. W. Loesch, Department of Forest Science 321 Richardson Hall, Oregon State University, Corvallis, OR 97331, USA. ([hank.loesch@oregonstate.edu](mailto:hank.loesch@oregonstate.edu))

D. Y. Hollinger, USDA Forest Service, Northern Research Station, 271 Mast Road, P.O. Box 640, Durham, NH 03824, USA. ([dhollinger@fs.fed.us](mailto:dhollinger@fs.fed.us))

L. Mahrt, College of Oceanic and Atmospheric Science, 104 Ocean Admin. Building, Oregon State University, Corvallis, OR 97331, USA. ([mahrt@coas.oregonstate.edu](mailto:mahrt@coas.oregonstate.edu))

S. C. Wofsy, Division of Engineering and Applied Science/Department of Earth and Planetary Science, Pierce Hall, Harvard University, 29 Oxford Street, Cambridge, MA 02138-2901, USA. ([wofsy@fas.harvard.edu](mailto:wofsy@fas.harvard.edu))