

QUANTIFYING LOSSES OF  
NITROGEN FROM  
LAND-APPLIED DAIRY MANURES

*BY*

*DAVID D. MYROLD,  
NANCY C. BAUMEISTER,  
AND  
JAMES A. MOORE*



*OREGON STATE UNIVERSITY  
CORVALLIS, OREGON*

**WRRI-115**

**NOVEMBER, 1992**

**QUANTIFYING LOSSES OF NITROGEN FROM  
LAND-APPLIED DAIRY MANURES**

by

David D. Myrold, Nancy C. Baumeister, and James A. Moore  
Departments of Crop and Soil Science and Bioresources Engineering  
Oregon State University

Final Technical Completion Report  
Project Number G1609-04

Submitted to  
United States Department of the Interior  
Geological Survey  
Reston, Virginia 22092

Project Sponsored by  
Oregon Water Resources Research Institute  
Oregon State University  
Corvallis, OR 97331

The activities on which this report is based were financed in part by the Department of the Interior, U.S. Geological Survey, through the Oregon Water Resources Research Institute. The contents of this publication do not necessarily reflect the views and policies of the Department of the Interior, nor does mention of trade names or commercial products constitute their endorsement by the United States Government.

## **FOREWORD**

The Oregon Water Resources Research Institute (WRRRI), located on the Oregon State University campus, serves the State of Oregon. The Institute fosters, encourages and facilitates water resources research and education involving all aspects of the quality and quantity of water available for beneficial use. The Institute administers and coordinates statewide and regional programs of multidisciplinary research in water and related land resources. The Institute provides a necessary communications and coordination link between the agencies of local, state and federal government, as well as the private sector, and the broad research community at universities in the state on matters of water-related research. The Institute also coordinates the interdisciplinary program of graduate education in water resources at Oregon State University.

It is Institute policy to make available the results of significant water-related research conducted in Oregon's universities and colleges. The Institute neither endorses nor rejects the findings of the authors of such research. It does recommend careful consideration of the accumulated facts by those concerned with the solution of water-related problems.

## TABLE OF CONTENTS

<b>LIST OF FIGURES</b> .....	iv
<b>LIST OF TABLES</b> .....	v
<b>ABSTRACT</b> .....	vi
<b>INTRODUCTION</b> .....	1
<b>MATERIALS AND METHODS</b> .....	2
<b>Soil and Site Description</b> .....	2
<b>Experimental Design</b> .....	2
<b>Manure Applications</b> .....	2
<b>Denitrification Measurements</b> .....	2
<b>Soil Core-Ion Exchange Resin Method</b> .....	3
<b>Modifications and Improvements</b> .....	4
<b>Measurement of Total Soil N</b> .....	5
<b>Statistical Analysis</b> .....	5
<b>RESULTS AND DISCUSSION</b> .....	6
<b>Seasonal Patterns in Denitrification Rates</b> .....	6
<b>Sources of variability for denitrification rates</b> .....	6
<b>Annual Denitrification Losses</b> .....	10
<b>Net N Mineralization</b> .....	11
<b>Plant N Uptake.</b> .....	11
<b>Total N method.</b> .....	12
<b>Core-IER Method.</b> .....	13
<b>Nitrogen Leaching Estimates</b> .....	16
<b>CONCLUSIONS</b> .....	18
<b>REFERENCES</b> .....	19
<b>APPENDIX</b> .....	42

## LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1a Denitrification and respiration rates in Waldo soil. Statistical significance is indicated by the following symbols: "+" for $0.05 < p \leq 0.10$ , "*" for $0.01 < p \leq 0.05$ , and ":" for $p \leq 0.01$ . Respiration rates are averaged over all treatments.	23
1b Denitrification and respiration rates in Amity soil. Statistical significance is indicated by the following symbols: "+" for $0.05 < p \leq 0.10$ , "*" for $0.01 < p \leq 0.05$ , and ":" for $p \leq 0.01$ . Respiration rates are averaged over all treatments.	24
1c Denitrification and respiration rates in Quillamook soil. Statistical significance is indicated by the following symbols: "+" for $0.05 < p \leq 0.10$ , "*" for $0.01 < p \leq 0.05$ , and ":" for $p \leq 0.01$ . Respiration rates are averaged over all treatments.	25
2a Soil $\text{NO}_3^-$ concentration in the Waldo soil. Means followed by the same letter are not significantly different (FPLSD, $\alpha = 0.05$ ). However, the control was significantly lower than the manure treatments at the June 1990 ( $p=0.09$ ) and October 1990 ( $p=0.08$ ) samplings.	26
2b Soil $\text{NO}_3^-$ concentration in the Amity soil. Means followed by the same letter are not significantly different (FPLSD, $\alpha = 0.05$ ).	27
2c Soil $\text{NO}_3^-$ concentration in the Quillamook soil. Means followed by the same letter are not significantly different (FPLSD, $\alpha = 0.05$ ).	28
3a Difference between bulk soil water content and water content of incubated soil for the Waldo soil.	29
3b Difference between bulk soil water content and water content of incubated soil for the Amity soil.	30
3c. Difference between bulk soil water content and water content of incubated soil for the Quillamook soil.	31
4 Net N mineralization rates and water content expressed as percent of water-holding capacity for individual cores (Waldo soil, fall/winter set, control treatment) $n=16$ .	32

## LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Selected characteristics of the three western Oregon soils used in this study.	33
2	Manure application dates.	34
3	Correlation coefficients (r) for denitrification. Denitrification and respiration rates were log transformed. Values in table were significant at $\alpha = 0.10$ .	35
4	Annual denitrification losses by manure treatment. Figures in parentheses is the percent of applied manure N lost, after subtraction of the control. Means followed by the same letter within a column are not significantly different (Tukey, $p=0.10$ ).	36
5	Measurement of net N mineralization rates in unfertilized controls by three different methods for the 1990 growing season (mean and 95% confidence interval).	37
6	Initial (March, 1990) and final (October, 1991) soil organic N.	38
7	Net N mineralization by the core-IER method.	39
8	$\text{NO}_3^-$ and $\text{NH}_4^+$ recovered from the resin bags as a percent of total net mineralized N.	40
9	Percent of mineralized N nitrified, averaged over all treatments.	41
A1	Nitrogen in harvested biomass. Mean $\pm$ standard deviation; $n=3$ for Amity and Quillamook, $n=4$ for Waldo. Data courtesy of J.A. Moore and M.J. Gamroth.	42

## ABSTRACT

Dairy manures are often applied to cropland for disposal and as a source of N fertilizer. Recent trends in the dairy industry toward larger herd sizes, and increases in land costs resulting in less land available to dairy farmers for disposal of manure result in the possibility of excessively high manure application rates. Nitrogen in excess of crop needs may leach as  $\text{NO}_3^-$  into groundwater or be lost by denitrification to the atmosphere. A high level of  $\text{NO}_3^-$  in groundwater is hazardous to human and animal health. Denitrification losses can also have negative environmental impacts, because  $\text{N}_2\text{O}$ , one of the denitrification products, is a radiatively active trace gas that contributes to global warming. Additionally,  $\text{N}_2\text{O}$  in the stratosphere is oxidized to  $\text{NO}$ , which catalyses the destruction of  $\text{O}_3$ .

A field study was conducted to measure two components of N cycling in manure fertilized pastures: denitrification losses and net N mineralization. Plots were established in three soil types representing a range of drainage classes. Plots received 0 to 450 kg manure-N  $\text{ha}^{-1} \text{y}^{-1}$  in five to seven split applications.

Denitrification, soil respiration, moisture content, and temperature were measured periodically for 15 months. Denitrification was measured by the  $\text{C}_2\text{H}_2$  inhibition method on intact soil cores. Nitrogen mineralization rates were measured by a soil core-ion exchange resin (core-IER) method. Three successive incubations were summed to produce an estimate for annual net N mineralization.

Annual denitrification losses were increased by manure applications and varied by soil type. Highest losses (108 kg N  $\text{ha}^{-1} \text{y}^{-1}$  at the highest manure rate) were in the well drained Quillamook soil. Lowest losses (33 kg N  $\text{ha}^{-1} \text{y}^{-1}$  at the highest manure rate) were in the poorly drained Waldo soil. Denitrification losses ranged from 5 to 16% of applied N. Denitrification was a significant component of the N budget in these manure fertilized pastures. Denitrification in the fall and early winter removed  $\text{NO}_3^-$  which would likely have leached from the soils. In the poorly drained Waldo soil, denitrification rates did not significantly increase beyond the lowest manure rate, and were probably limited by  $\text{NO}_3^-$ . Restricted nitrification or plant or microbial competition for  $\text{NO}_3^-$  kept  $\text{NO}_3^-$  concentrations below 3 mg  $\text{NO}_3^-$ -N  $\text{kg}^{-1}$  soil. In the

well drained Quillamook soil, denitrification rates increased linearly with each increment in manure rate.

Net N mineralization as measured by the soil core-IER method were several times higher than either of two alternate methods (plant N uptake and loss of soil organic N) for measuring net N mineralization, although the core-IER and plant uptake methods ranked the soils the same. It appears likely that the core-IER method positively biases net N mineralization through the effect of the exclusion of plant roots on soil water relations.



## INTRODUCTION

Animal manures are a valuable source of plant nutrients, in particular N. Manure can replace or supplement inorganic fertilizers in cropping systems, however effective use of the manure N by farmers is limited by several factors. In modern, specialized farming operations, production of manure is concentrated in dairy and feedlot operations. Hauling costs for manure are high relative to inorganic fertilizer sources of N. Additionally, the timing of N availability is critical to optimum crop yields. Nitrogen in manure exists in organic and inorganic forms, and the exact timing of N mineralization from organic forms is dependent on a number of factors and is difficult to predict.

For these reasons, most manure produced by dairy farmers is applied where it is produced, typically on land used for pasture, hay or silage crops. The high cost of farmland and economic pressure to increase dairy herd sizes, with concomitant increases in the volume of manure produced, has resulted in the possibility of excessively high rates of manure being applied to land.

Nitrogen applied in excess of plant uptake can become a hazard in the environment (Keeney, 1982). Nitrate not used by plants can leach into groundwater and localized high levels of  $\text{NO}_3^-$  have been found in groundwater in Oregon (Pettit and Thomas, 1986). Denitrification is an alternate fate for unused  $\text{NO}_3^-$ . In this microbial process,  $\text{NO}_3^-$  is converted to  $\text{N}_2$  gas and a small and variable portion of  $\text{N}_2\text{O}$  and  $\text{NO}$  (Firestone, 1982). Nitrous oxide as a radiatively active gas is directly involved in greenhouse warming.  $\text{NO}$  is directly involved in ozone destruction and indirectly in global warming.

This study is part of a larger effort to develop a N budget for grass pastures receiving dairy manure. The objectives of this portion of the study were: (1) to measure denitrification losses from dairy manure surface applied to grass pastures in three soil types and relate those losses to soil variables, (2) to measure net N mineralization in three ways using the core-IER method, plant uptake, and change in soil organic N, and (3) to estimate leaching losses.

## MATERIALS AND METHODS

### Soil and Site Description

Plots were established in the fall of 1989 on mixed perennial ryegrass (*Lolium perenne*) and orchardgrass (*Dactylis glomerata*) pastures in three soil types representing a range of drainage classes (Table 1). The Amity silt loam (fine, mixed, mesic Argiaquic Xeric Argialboll) and Waldo silty clay loam (fine, mixed, mesic Fluvaquentic Haplaquoll) are in the Willamette Valley of western Oregon (44.30° N, 123.20°W). The Willamette Valley has a Mediterranean climate of hot, dry summers and cool, moist winters. The Quillamook silt loam (medial, isomesic Alic Pachic Melandudand) is in Tillamook County, on the Pacific coast of Oregon (45.50°N, 123.90°W), in a maritime climate.

### Experimental Design

The experimental design was a randomized complete block. Manure treatments were completely randomized within each soil. Three replicate plots were used at the Amity and Quillamook sites. The Waldo site has two replicate plots for manure treatments and four replicates for the control (no manure) treatment.

### Manure Applications

Fresh dairy manure containing a mixture of urine and feces was used (1.6 g organic N kg<sup>-1</sup> manure, 1.1 g NH<sub>4</sub><sup>+</sup>-N kg<sup>-1</sup> manure) was applied at 0, 150, 300, and 450 kg manure-N ha<sup>-1</sup> y<sup>-1</sup> for two growing seasons. In order to maximize plant N uptake manure applications were split into five to seven parts (Table 2). After each application, regrowth of the plots was monitored. When the grass reached 20 to 25 cm, the plots were harvested, grass removed, and manure was applied again.

### Denitrification Measurements

At each sampling date, three soil cores were taken from randomly selected points in each plot and soil temperatures at 10 cm were recorded. Denitrification rates were measured on intact soil cores by the acetylene (C<sub>2</sub>H<sub>2</sub>) inhibition technique. With

minor variations, the protocol described by Tiedje et al. (1989) was followed. Soil cores were taken with a steel corer, which extracts a relatively intact soil core about 15 cm long and 2 cm in diameter, encased in a 20-cm-long polyacrylic tube. To facilitate diffusion of  $C_2H_2$  throughout the soil, the tube is slightly larger in diameter than the soil core. The tube was stoppered at both ends and 6 mL of air was added to the tube. The headspace was mixed with a 30-mL syringe. A 5.5-mL gas sample was put into a nominal 3-mL draw vacutainer (Becton-Dickson, Rutherford, NJ) for later analysis. Six mL of  $C_2H_2$ , producing a partial pressure of at least 100 kPa, was injected into the tube and the headspace was mixed again. Acetylene was made in the field by mixing water with calcium carbide in a sealed serum bottle. Cores were incubated at ambient soil temperature for 24 hours. At the end of the incubation, the tube headspace was mixed and a final 5.5-mL gas sample was withdrawn from the tube and put into a vacutainer. A sample of the soil was dried at 105°C to determine gravimetric water content.

Gas samples were analyzed by gas chromatography (Varian 3700 equipped with a  $^{63}Ni$  electron capture detector) using a Porapak Q column. The carrier gas was 95% argon and 5% methane at a flow rate of 40 mL  $min^{-1}$ . The column was maintained at 35°C. The electron capture detector was operated at 350°C.

Denitrification and respiration rates were calculated by multiplying the increase in concentration of  $N_2O$  and  $CO_2$  in the headspace over the incubation period by the total void volume of the tube. Dissolved  $N_2O$  was calculated from the Ostwald coefficient (Wilhelm et al., 1977). Total void volume in the tube was calculated by subtracting the volume occupied by water and the volume occupied by soil from the total volume of the tube. Annual losses were calculated from the area under the curve defined by rate measurements for each replicate plot.

#### **Soil Core-Ion Exchange Resin Method**

For each of three seasons (spring, summer, and fall/winter), a set of PVC tubes (5 cm in diameter and 50 cm long; 25 cm long for the fall/winter set) were installed at random locations in the manure treated plots described above. A metal cap was placed on the tubes and a sledgehammer was used to drive the sharpened tubes into

the ground. Compaction was minimal on most tubes. The tubes were removed from the ground, extracting a soil core containing about 1 kg soil. Approximately 2 cm of soil was removed from the bottom of the soil core so that a nylon mesh bag containing 10 g (dry weight) mixed bed ion exchange resin (Rexyn 300, Fisher Scientific) could be placed there. The resin bag was held in place by a piece of cheesecloth or fiberglass screening while the tube was reinserted into its hole.

Manure applications (after the first manure application) were made individually to the soil cores, at first with a modified syringe using diluted manure and later with frozen manure pellets, at or close to the same time as applications were made to the whole plots.

At the end of the incubation period, the soil cores and resin bags were removed from the ground. The resin bags were extracted in 100 mL of 2N KCl solution. Soil from the tube was mixed and a subsample was extracted in 2N KCl. Extracts were colorimetrically analyzed for  $\text{NH}_4^+$  by the salicylate/nitroprusside method and  $\text{NO}_3^-$  by Cd reduction and diazotization (Alpkem, Clackamas, OR).

At the beginning and end of each incubation period, bulk soil was sampled to the appropriate depth (50 or 25 cm) and the  $\text{NH}_4^+$  and  $\text{NO}_3^-$  content determined.

### **Modifications and Improvements**

The method was modified after the observation was made at the end of the first (spring) set that annual grasses had invaded some tubes and in others, captured ryegrass crowns were growing. In the second (summer) set, living crowns were uprooted and replaced in the tube. That was moderately effective at controlling grass regrowth, but not completely. In the third (fall/winter) set, Simazine, a residual herbicide, was used. This was effective; no living vegetation was observed in the tubes.

Additional problems were encountered with the fall/winter set. Since the spring and summer sets were installed into moderately moist soils, it was possible to drive them with a cap and sledgehammer to a depth of 50 cm. Some compaction was noted, but it was not judged to be a serious problem. However, the fall/winter set was installed at the end of summer, into hard, dry soils. The 50-cm-long tubes broke while

being installed. However, shorter tubes were installable, so 25-cm tubes were used for the fall/winter set.

### **Measurement of Total Soil N**

Total Kjeldahl N (TKN) was determined on soil samples taken in March, 1990, prior to beginning manure application and October, 1991, following two growing seasons of manure applications. Soil was sampled to a depth of 15 cm. Soils were digested for TKN as described by Bremner and Mulvaney (1982). Ammonium from the TKN digests was determined as above. Inorganic N present at the final sampling time was not separately determined and is therefore included in the estimate of organic N.

### **Statistical Analysis**

Treatment effects within a sampling date for each soil were determined on log transformed denitrification and respiration rates by standard analysis of variance (ANOVA) using SAS statistical software package (SAS Institute, Cary, NC) for a completely randomized design with three replicates and three subsamples (four or two replicates and four or six subsamples in the Waldo soil) and Tukey's HSD procedure to compare means at the  $p=0.10$  level. Significance of treatment effects for annual loss were determined by ANOVA using the annual loss calculated for each replicate plot. Statistical significance for soil  $\text{NO}_3^-$  concentrations was determined by F-protected least significant difference (FPLSD).

Annual net N mineralization rates were determined by summing the plot means for each of the three seasons together. Treatment effects were determined by standard analysis of variance (ANOVA) using SAS statistical package (SAS Institute, Cary, NC) for a randomized complete block design. Plant uptake data were analyzed by ANOVA. A paired t-test was used to detect significant changes in soil organic N. If the 95% confidence interval of the difference (final TKN - initial TKN) for each treatment did not include zero, then there was a significant net accumulation or loss of organic N. Analysis of variance on the net accumulation or depletion was also run and 95% confidence intervals were constructed from the ANOVA results.

## RESULTS AND DISCUSSION

### Seasonal Patterns in Denitrification Rates

The two Willamette Valley sites are on different soil series but are located within 500 m of each other. They had similar temporal patterns of denitrification (Fig. 1a and 1b). In the Amity and Waldo soils, 77 and 83% of the annual loss occurred in the four months from mid-October to mid-February. There were two peaks in denitrification rates: following the first fall rains and following a five-day freeze.

The Quillamook site is about 100 miles northwest of the Willamette Valley sites. Denitrification rates peaked a month later there and only one peak was observed (Fig. 1c), however at this site a four-month time period (late October to late February) accounted for a similar portion of the annual loss (73%).

### Sources of variability for denitrification rates

Because the factors that drive denitrification (denitrifier populations, soil C availability, restricted aeration, and availability of  $\text{NO}_3^-$ ) are strongly affected by climatic events such as rainfall, freeze/thaw cycles, etc., denitrification rates are highly variable in time and can change by an order of magnitude in a week (Ball and Ryden, 1984). Denitrification rates also respond rapidly to perturbations such as fertilization with  $\text{NO}_3^-$  (Jarvis et al., 1991), irrigation (Rolston et al., 1982) or manure applications (Thompson et al., 1987). Denitrification rates increased within a day following injection of fermentation residues (Rice et al., 1988).

In this study, rapid changes in denitrification rates followed two climatic events: the first fall rains and a winter freeze/thaw cycle. For example, at the Willamette valley sites, rates increased 10 to 100 times (depending on manure treatment) in the two weeks following the first significant rainfall of the fall. During this time the respiration rate doubled (Fig. 1a and 1b). Wetting of dry soil (Patten et al., 1980) has been shown to stimulate denitrification rates, most likely by increasing the availability of soil C to denitrifying microorganisms. In both Willamette Valley sites, there was an approximately three-fold increase in denitrification following a freeze/thaw cycle. Freeze/thaw cycles have been shown to stimulate denitrification

rates (Edwards and Kilham, 1986). Thompson (1989) suggests that high denitrification rates following thawing of frozen soil may be due to enhanced breakdown rates of organic matter. In the Amity soil, respiration rates increased following the thaw, but in the Waldo soil they decreased. Decreased heterotrophic respiration in the Waldo may be attributed to near saturated conditions.

Treatment effects on denitrification rates were significant only when rates were high. For the Waldo soil, this was the period from late October to early January. For the Quillamook, the June sampling date, and the four sampling dates from late October to early December, had significant treatment effects. In the Amity soil, spatial variability was higher, probably because of damage to the pasture by gophers. This obscured separation of manure rate effects, but the manured plots, considered as a group, had significantly ( $p \leq 0.10$ ) higher denitrification rates than the control plots on five dates (Fig. 1b).

Manure had no effect on soil respiration rates. The average respiration rates were 18 for the Amity, 32 for the Waldo, and 28 mg CO<sub>2</sub>-C g<sup>-1</sup> soil d<sup>-1</sup> for the Quillamook soil.

Soil temperature at 10-cm depth varied between a low of 6°C in December and a high of 16°C in June in the Willamette Valley sites. The mean soil temperature over the active denitrification period was 9°C. The Quillamook soil was slightly warmer, reaching a low of 10°C in the winter and a high of 20°C in July. Soil temperature was not included in the correlation analysis because the data set is incomplete and might yield biased results since not all sampling dates would be represented. Myrold (1988) found a weak negative correlation between denitrification rate and soil temperature. This is counter-intuitive, because one would expect microbial activity to be higher in warmer soils. This relationship is probably a result of a negative correlation between soil temperature and soil water content and a positive relationship between microbial activity and soil water content. In the Mediterranean climate of western Oregon, most rainfall occurs in the fall, winter, and spring, when soils are cooler.

Soil inorganic N content was measured four times (Fig 2a, 2b, and 2c). In the Willamette Valley sites, soil NO<sub>3</sub><sup>-</sup> was always less than 4 mg NO<sub>3</sub><sup>-</sup>-N kg<sup>-1</sup> soil. In the

Quillamook soil,  $\text{NO}_3^-$  reached a high of  $29 \text{ mg NO}_3^- \text{-N kg}^{-1}$  soil in the fall and manure treatment effects on soil  $\text{NO}_3^-$  concentration were significant ( $p \leq 0.05$ ). These  $\text{NO}_3^-$  concentrations are below the  $5 \text{ mg NO}_3^- \text{-N kg}^{-1}$  soil which Ryden (1986) suggests as a minimum requirement for significant denitrification to occur. However it must be noted that these values represent only bulk soil concentrations and that microsite concentrations could be much higher.

Spatial variability of denitrification rates was high. The distribution was positively skewed and resembled a log-normal distribution. The coefficient of variation (CV) was generally between 100 and 200% (before log transformation). The CV is a normalized measure of variability defined as the sample standard deviation divided by the sample mean (Snedecor and Cochran, 1980). This variability was comparable to the variability commonly encountered in studies of denitrification. A highly skewed distribution can be produced under a multiplicative model, where the effect (denitrification rate) is the product, rather than the sum, of several other variables. A stochastic, multiplicative model using as variables respiration rates and denitrification enzyme activity was able to successfully model measured denitrification rates (Parkin and Robinson, 1989).

Although denitrification requires  $\text{NO}_3^-$ , low  $\text{O}_2$  concentration, and denitrifying organisms in addition to a source of available C, many investigators have concluded that the availability of C is the primary factor driving denitrification rates (Groffman and Tiedje, 1991). One rationale behind this is the observation that very high rates of  $\text{O}_2$  consumption can create anaerobic conditions even where the diffusion of  $\text{O}_2$  is not impeded, and even in fully aerobic atmospheres. Parkin (1987), dissected several soil cores in an attempt to localize the denitrifying activity. In the most extreme case an 80 mg leaf fragment was identified which had a specific denitrification rate (under fully aerobic conditions) more than 20,000 times the median denitrification rate of the bulk soil. He calculated that a water film on the decaying leaf, combined with a very high rate of oxygen consumption, could create an anaerobic condition at the leaf surface. In light of these findings, factors that increase the spatial variability of organic C might be expected to increase the variability of denitrification rates. Goulding and Webster (1989) did find that applications of farm yard manure as



contrasted with inorganic fertilizers increased both the magnitude and the variability of denitrification losses. Thompson (1989) found that with N held constant, denitrification decreased five times when the applied cattle slurry was diluted by one-half.

In other investigations into the source of variability in denitrification, Parkin et al. (1987) measured denitrification rates and phase I denitrification enzyme assays (PDA) on a large number of soil samples. Coefficients of variation for the denitrification rate measurements varied between 200 and 300. In this assay, a C source and  $\text{NO}_3^-$  are supplied in excess and only the distribution and activity of denitrifying organisms is measured. Even with two sources of variability removed, the CV was still around 40%.

In light of the complexity of the factors driving denitrification, it is no surprise that attempts to correlate measured variables with denitrification rate have met with only limited success. Correlation analysis by Burton and Beauchamp (1985) of 13 variables on three dates against denitrification rates detected only five significant relationships and no variable explained more than 35% of the denitrification rate variability. Myrold (1988) correlated denitrification with three variables (soil water content, respiration, and temperature). No single variable explained more than 29% of the variability, but a multiple regression including all variables was able to explain 43% of the variability. Elliott et al. (1991) measured the denitrification rates of bulk soil and earthworm castings, along with respiration rate, moisture content, and inorganic N content, under a range of fertilizer and drainage regimes. For bulk soil, no correlations were significant. However for the earthworm castings,  $\text{NH}_4^+$ , moisture content and  $\text{NO}_3^-$  explained 25, 25, and 36% of the variability, respectively.

Parsons et al. (1991), measured six variables and incorporated them into a multiple regression with denitrification rates as the dependent variable. By using mean values by sampling date they were able to improve the  $r^2$  of the multiple regression from 0.27 to 0.74. In this experiment, using mean values by date ( $n=13$ ) improved correlations (Table 3). However, more variables were significantly correlated when spatial variability was incorporated by using individual data points ( $n \approx 400$ ). Soil water content was the best predictor of denitrification in the Willamette

valley soils, while respiration was best in the Quillamook soil. For each soil, correlations were further strengthened by restricting the dates used to the periods of most active denitrification (Table 3). This suggests that the measured variables are not sufficient to predict which soil conditions will produce high or low denitrification rates. They are better able to predict relative denitrification rate when soils are actively denitrifying.

### Annual Denitrification Losses

Manure applications increased denitrification losses in all soils (Table 4) however kinetics of the response to increasing manure rate varied. In the Quillamook soil, annual denitrification losses increased linearly with increasing manure rates but in the Willamette Valley sites, denitrification response was hyperbolic, approaching saturation at 300 kg manure-N ha<sup>-1</sup> y<sup>-1</sup>. These kinetics suggest that some factor related to the availability of one of the critical components (C, NO<sub>3</sub><sup>-</sup>, or anaerobiosis) is limiting the denitrification rate in the Willamette Valley soils.

Because these three soils differ in both drainage class and climate, separating the contribution of climate and drainage to denitrification losses is not possible. However, some observations can be made. Past comparisons of the effects of soil drainage on denitrification losses have given contradictory results. Installation of mole drains in a poorly drained clay soil receiving NO<sub>3</sub><sup>-</sup> fertilizer decreased denitrification losses by almost one-half (Colbourn and Harper, 1987). But a well-drained loam receiving cattle slurry had losses five times greater than a poorly drained silty clay loam (Thompson and Pain, 1989). These differences are probably because of the necessity for mineralization and nitrification of organic N before the N is available for denitrification. In a study directly comparing NO<sub>3</sub><sup>-</sup> fertilizer with cattle slurry, imperfectly drained soils receiving NO<sub>3</sub><sup>-</sup> directly had much greater denitrification losses than soils receiving the same amount of N as cattle slurry (Egginton and Smith, 1986). Where an organic N source is involved, either the mineralization or the nitrification step may be rate limiting. The restricted aeration of the poorly drained Waldo soil, although providing the anaerobic volume necessary to denitrification, could also slow both mineralization of applied manure and nitrification of NH<sub>4</sub><sup>+</sup>.

There is indirect evidence for slower mineralization of manure N in the Waldo soil. Plant N uptake was highly correlated with denitrification loss in the Waldo soil ( $R^2=0.89$ ), but not in any other soil. This suggests that both denitrification and plant N uptake may have been N-limited in the Waldo soil because well-established grass swards are effective N scavengers. Also, the  $\text{NO}_3^-$  concentration of the Waldo soil was always the lowest of the three soils (Fig. 2a, 2b, and 2c).

Although there was a tendency towards increasing plant N uptake with manure treatments, manure treatment effects were not significant in the first year although there were significant differences in yield between soils ( $p \leq 0.001$ ). For annual denitrification losses, however, both soil and manure effects were significant. This implies that manure applications affected denitrification rates in some way other than through N availability, probably through the stimulating effect of the available C in manure on the soil microbial populations.

### **Net N Mineralization**

Net N mineralization from soil organic matter (SOM) was estimated by three different methods: the soil core-IER method described above, plant N uptake in unfertilized control plots, and direct measurement of the soil organic N (SON) pool. The methods generally agreed poorly (Table 5). However, the plant uptake and core-IER methods ranked the soils the same. In all soils, the core-IER estimates were higher than any other method.

**Plant N Uptake.** Plant N uptake in unfertilized plots is often used to measure net N mineralization from soil (Legg and Meisinger, 1982). The amount of N mineralized varies considerably. Perennial ryegrass monocultures removed an average of  $46 \text{ kg N ha}^{-1} \text{ y}^{-1}$  over the first three years after establishment (Harkess and Frame, 1986). Rangeley and Newbould (1986) estimated soil derived N as  $91 \text{ kg N ha}^{-1} \text{ y}^{-1}$  in the second year of perennial ryegrass. Plant uptake in annual and perennial grasslands in a Mediterranean climate was about  $100 \text{ kg N ha}^{-1} \text{ y}^{-1}$  in annual and perennial grasslands (Joffre, 1990). Hatch et al. (1991) measured net N mineralization by plant uptake of  $263 \text{ kg N ha}^{-1}$  method in unfertilized perennial grass swards.

Estimating N mineralization by plant uptake has the advantage of sampling a much larger volume of soil than incubation methods so consequently spatial variability is lower. However, there are disadvantages as well. Leaching or gaseous losses will not be included, and an increase in belowground (root) biomass N will not be measured. Whitehead et al. (1990) has shown that grass swards can continue to accrete belowground biomass at least to 15 years in age. Also, if plant uptake is limited by a factor other than N, it will not accurately reflect N mineralization.

Among the three soils studied (Waldo, Amity, and Quillamook), there were almost four-fold differences in plant N uptake in the unfertilized control plots (Table 5). The effects of manure treatment were not significant in the first year within a soil, however the soils were significantly different from each other ( $p \leq 0.0001$ ). In the second year of treatment, N yields from the control plots dropped, whereas N yields from the manured plots increased. Second year N yields from the control plots were significantly lower ( $p \leq 0.01$ ) than the manured plots in the Waldo and Quillamook, but not the Amity. Although the difference in N yield between the average of the manured plots and control plots was similar for all soils (about 50 kg N), variability in the Amity soil was higher and treatment effects were not significant (Appendix). The increased variability was probably due to gopher damage to the Amity plots. The soil by treatment interaction was not significant in either year, indicating that manure had a similar effect on N yield within each soil. The error associated with plant uptake was much smaller than the error associated with either the core-IER or the SON method, providing more sensitive detection of treatment effects.

**Total N method.** Direct measurement of the SON pool by Kjeldahl digestion of soil has been used to measure net N mineralization. However, because of the large size of the total N pool relative to the active, or mineralizable, portion small errors in measurement produce a large uncertainty in the estimated net loss or gain of soil N. Net N mineralization in the control plots was 53, 94, and 25 kg N ha<sup>-1</sup> y<sup>-1</sup> for the Waldo, Amity, and Quillamook soils (Table 5). These figures are surprisingly close to the N mineralization rates calculated by plant uptake for the Amity and Waldo soils, although the associated error was much higher.

These N mineralization rates are equivalent to 1.3, 2.1, and 0.3 % of the initial SON content (Table 6). Again, the Waldo and Amity turnover rates are similar to other estimates for the rate of turnover of organic N in grassland soils. Whitehead (1984) (as discussed in Whitehead, 1986) in a greenhouse study of 21 UK soils, found that from 1.5 to 4% of the total soil N was mineralized over a year. Jenkinson (1990) suggests 2.8 and 2.9% per year as typical for turnover of organic N in pasture and grassland (again in the UK). Differences in the absolute amount of N mineralized and the portion of the total N pool mineralized have been related to the management history of the site, particularly additions of organic materials and tillage (Johnston et al., 1989):

**Core-IER Method.** Incubation methods involving varying degrees of soil disturbance have been used to measure or predict net N mineralization in agricultural soils (Stanford, 1982). The core-IER estimates for net N mineralization were much higher than either of the other methods, particularly for the Quillamook soil, and it appears likely that there is a positive bias in the method. Additionally, although a trend towards higher net N mineralization in the manure treatments can be seen (Table 7), because of the high spatial variability, the differences were not significant, except in the Waldo soil, where net N mineralization in the control plot was lower than the manured plots ( $p \leq 0.05$ ).

Where both plant uptake and an incubation method have been used and can be compared, agreement has varied from good to poor. The reason for the variability of agreement of incubation methods with plant uptake methods is presumably the effect of incubation conditions and amount and C/N ratio of mineralizable organic material in the soil cores on the opposing processes of N mineralization and immobilization. If both occur at the same rate in the incubated soil as in the bulk soil, or if both are equally enhanced, the method will give an accurate estimate of net N mineralization.

Good agreement between incubation and plant uptake methods has been achieved by some investigators (Joffre, 1990; Hatch, 1991), but not by others. Rees (1989) measured net mineralization rates in an incubation method less than 11% of plant soil-derived N uptake. He suggested that microbial immobilization of N was responsible for the underestimate; increases in microbial biomass in the incubated soil

accounted for 56% of the deficit. Raison et al. (1987) suggested that decomposing excised plant roots can enhance immobilization of N, causing underestimation of net N mineralization rates, although they did not find any evidence of that bias in their experiment.

Overestimation of net N mineralization may result if either the gross N mineralization rate is enhanced or gross N immobilization is decreased. A mechanism for reducing gross immobilization has been suggested by Hart and Firestone (1988), who measured net N mineralization by buried bag and core-IER methods in young and old growth forests. The two methods agreed well for the young forest. However in the old growth forest net N mineralization by the core-IER method was twice that of the buried bag method. The authors suggest that the  $\text{NO}_3^-$  adsorbed on the ion exchange resins may be protected against re-immobilization, thus reducing gross immobilization. Although this may have happened some extent, this would not explain the overestimation of net N mineralization in this experiment since on an annual basis, most (83, 76, and 73% for the Waldo, Amity, and Quillamook respectively) of the mineralized N was recovered from the soil, not the resin bags (Table 8). Only in the fall/winter set was any substantial amount of N leached to the resin bags.

The enhanced moisture status of the soil cores relative to the bulk soil may be a more important source of bias. Because the core-IER method depends on excluding plant roots to prevent plant N uptake, a side effect is that there are no transpirational losses. Soil water content in the cores was always higher than in the bulk soil (Fig. 3a, 3b, and 3c), and tended to increase over the incubation period, even while the bulk soil was drying from evapo-transpirational losses. The more favorable water content of the tubes relative to the bulk soil could well enhance microbial activity and the gross N mineralization rate in the soil cores.

Net N mineralization is most rapid at a water content close to field capacity, and falls off sharply as the soil approaches saturation (Myers et al., 1982). Linn and Doran (1984) confirmed the many previous reports that maximum microbial activity occurs at approximately 60% of a soils water holding capacity (WHC). The soil cores varied in water content within a season, even in the fall/winter set where complete

control of grass regrowth was achieved. For the fall/winter set for the Waldo soil, soil water content varied between 0.31 and 0.41 g water g<sup>-1</sup> soil (68 to 92% WHC) and net N mineralization and soil water content (Fig 4) were positively correlated ( $r=0.54$ ,  $p\leq 0.0001$ ). Site-specific data are not available, but by extrapolation from Soil Conservation Service data (Huddleston, 1982) for a similar soil, field capacity in the Waldo should be around 0.36 g water g<sup>-1</sup> soil (79% WHC). A similar, but weaker, pattern was seen in the Quillamook soil ( $r=0.36$ ,  $p=0.05$ ). Estimation of the Quillamook field capacity is a bit more speculative, since the bulk density of the pasture site is considerably lower than the published value (Huddleston, 1982) for this soil. Field capacity, based on extrapolation from published values, is probably between 0.80 and 1.00 g water g<sup>-1</sup> soil, while soil water contents were between 0.60 and 0.85 g H<sub>2</sub>O g<sup>-1</sup> soil.

There was a positive correlation between soil water content on net N mineralization in other seasons as well. Unfortunately, control of grasses was not as good, so it is possible that unmeasured plant N uptake occurred.

There is a large pool of easily decomposable organic materials in established swards, which increases with the age of the sward, and with utilization by grazing rather than cutting. Whitehead (1990) measured the N content of stubble, litter, root, and soil macro-organic matter fractions of 8 and 15-year-old perennial ryegrass swards. The fractions contained, on average, 68, 12, 249, and 240 kg N ha<sup>-1</sup> for a total of 569 kg N ha<sup>-1</sup>. He suggested that 364 kg N ha<sup>-1</sup>, or 64 % of this N is labile or readily mineralizable under disturbance such as tillage. The core-IER method does not involve a disturbance such as mixing, which would mimic tillage. However, no roots or intact plant parts were observed in the cores after the three or six-month incubation. It should also be noted that, whatever the size of the potentially mineralizable organic matter fractions in these soils, this material is given three opportunities to contribute to the annual mineralization rate: once for each of three incubation periods.

Although the cores were wetter than the bulk soil, they did not seem to be wet enough to inhibit nitrification (Table 9). Net nitrification was about 74% over all soils and seasons. Least nitrification occurred in the Waldo soil in the fall/winter set. This is consistent with the seasonal high water table which occurs in this soil.

The Quillamook soil was anomalous in several ways. First, net N mineralization by either the core-IER method or by plant N uptake was three to four times greater than in the Amity or Waldo soils. The total N content was almost twice that of the Willamette Valley sites. Second, the spatial variability of almost every measured parameter was higher. These differences were to some extent expected. The Quillamook soil is a deep, well drained productive andisol in a coastal, maritime climate ideal for grass production. The Quillamook site, because of its close proximity to a heifer barn, has been heavily grazed and manured. Pasture utilization by grazing causes uneven distribution of N which would be expected to increase spatial variability. Manure applications leave an easily decomposable pool of organic N. In contrast, the Amity and Waldo sites have historically not been intensively managed. Estimated past N fertilization rates are 40 to 50 kg N ha<sup>-1</sup> y<sup>-1</sup> for the Waldo and 70 to 80 kg N ha<sup>-1</sup> y<sup>-1</sup> for the Amity (M.J. Gamroth, personal communication). They have been infrequently grazed. The different field histories of the sites would have effects on N availability lasting at least several years into the current treatment regime. Thus the treatment effects on N availability reflect past management as well as current management of the soil.

Plant uptake can be considered a minimum estimate for net N mineralization since there will always be some denitrification and leaching losses. Denitrification losses were 10, 20, and 10 kg N ha<sup>-1</sup> y<sup>-1</sup> in unfertilized control plots for the Waldo, Amity and Quillamook soils and most of the loss occurred in the winter months (Table 4). Leaching losses, measured by porous cup lysimeters at 50 cm depth, were about 7 kg N ha<sup>-1</sup> y<sup>-1</sup> (Moore et al., 1992).

### **Nitrogen Leaching Estimates**

The core-IER method was intended to produce an independent estimate of leaching losses by placing an IER bag below the rooting zone and measuring the portion of mineralized N which moved below the rooting zone. In the spring and summer sets very little N was leached to the resin bag; even though a large amounts of N was mineralized, it accumulated in the soil. The lack of accumulation on the resin bags cannot be interpreted as indicating that leaching of NO<sub>3</sub><sup>-</sup> did not occur,



because the amount of N mineralized was far in excess of what the pasture grasses took up in that period (Appendix).

However, it may be reasonable to accept the fall/winter period as a less biased measurement. The soil cores were only slightly wetter than the bulk soil, and the measured N mineralization rates were much lower. The 6-month period (October through March) accounted for 12 and 18% of annual net N mineralization in the Waldo and Amity soils, and 22% in the Quillamook soil (Table 6), or 32, 54 and 224 kg N ha<sup>-1</sup> y<sup>-1</sup>, respectively. If those mineralization rates represent the upper end of net N mineralization, then plant uptake in that period can account for at least 50% of mineralized N for the Waldo and Amity soils, but only about 13% for the Quillamook. Leaching losses in the 0 to 20 kg N ha<sup>-1</sup> range would be predicted for the Waldo and Amity soils. These compare well with the losses of about 7 kg N ha<sup>-1</sup> measured by Moore et al. (1992) using lysimeters and would probably be considered acceptable losses from the standpoint of groundwater protection.

The status of the Quillamook soil with respect to N mineralization rates and potential leaching losses is somewhat more difficult to discern. It seems likely that the core-IER method does overestimate net N mineralization, however the exact extent of the overestimation is not known. We could put an upper limit to the bias by accepting plant N uptake as accurately representing net N mineralization. In that case, the core-IER estimate for annual net N mineralization is five times greater. If we then adjust the fall/winter net N mineralization by that ratio, we would suggest that plant uptake can account for most of that N, leaving a lower limit of 7 kg N ha<sup>-1</sup> to be leached from the soil. If we use an a lower enhancement factor of three (from the Amity soil), then we would calculate leaching losses of 36 kg N ha<sup>-1</sup>. If we do not adjust the figure at all, then we estimate that 196 kg N ha<sup>-1</sup> would be leached. Perhaps coincidentally, of the 224 kg N ha<sup>-1</sup> mineralized in the cores during that period, 190 kg N ha<sup>-1</sup> was recovered from the resin bag (at 20 cm depth) and 34 kg N ha<sup>-1</sup> was recovered from the soil while plant uptake for that period was 28 kg N ha<sup>-1</sup>. Estimates for leaching losses in the Quillamook soil range therefore from 7 to 196 kg N ha<sup>-1</sup>, which is inclusive of the 6 to 28 kg N ha<sup>-1</sup> annual losses obtained with porous cup lysimeters (Moore et al., 1992).

## CONCLUSIONS

Denitrification can be a significant route for N loss from fertilized pastures. From 5 to 16% of the applied N was lost through denitrification. In the Willamette Valley soils, percent loss decreased with increasing fertilization rate, whereas in the Quillamook soil, the percent lost to denitrification increased with increasing manure rate. This indicates that denitrification in less well drained soils under Mediterranean climates cannot be relied upon to remove more than 20 to 30 kg N ha<sup>-1</sup> y<sup>-1</sup>, even under high loading rates. The capacity of the Quillamook soil to remove N by denitrification was much higher. The denitrification response was linear even up to the highest rate applied (586 kg manure-N ha<sup>-1</sup> y<sup>-1</sup>). It is likely that the more moderate climate, increased rainfall of the Pacific coast and good drainage of the Quillamook soil act in concert increase the rate of N cycling, so these results should not be extrapolated to the poorly drained soils also found along the Pacific coast.

Plant uptake is probably superior to the core-IER incubation method for measurement of net N mineralization, at least in unfertilized well-established swards, where extensive root systems allow relatively little leaching. At high rates of fertilizer, plant uptake will not reflect net N mineralization as well and other methods must be brought into play. More replication in the core-IER method could reduce error and improve detection of treatment effects, and the method could be treated at least as an index of mineralization. More frequent sampling would minimize differences between bulk soil water content and incubated soil water content, and also further reduce variability.

Leaching losses measured by the accumulation of inorganic N on the resin bags of the N mineralization cores ranged from 0 to 20 kg N ha<sup>-1</sup>, which agrees well with independent measurements. At the Quillamook site, however, the inorganic N trapped on the resin bags greatly exceeded lysimeter estimates, probably because the core-IER method enhances net N mineralization. As a general rule, leaching losses of N were about half those of denitrification.

## REFERENCES

- Ball, F.R., and J.C. Ryden. 1984. Nitrogen relationships in intensively managed temperate grasslands. *Plant Soil* 76:23-33.
- Bremner, J.M., and C.S. Mulvaney. 1982. Nitrogen-total. p. 595-624. *In* A.L. Page et al. (ed.) *Methods of Soil Analysis. Part 2. Agron. Monogr. 9.* Amer. Soc. Agron., Madison, WI.
- Burton, D.L., and E.G. Beauchamp. 1986. Denitrification rate relationships with soil parameters in the field. *Commun. Soil Sci. Plant Anal.* 16:539-549.
- Colbourn, P., and I.W. Harper. 1987. Denitrification in drained and undrained arable clay soil. *J. Soil Sci.* 38:531-539.
- Edwards, A.C., and K. Killham. 1986. The effect of freeze/thaw on gaseous nitrogen loss from upland soils. *Soil Use Manage.* 2:86-91.
- Egginton, G.W., and K.A. Smith. 1986. Losses of nitrogen by denitrification from grassland soil fertilized with cattle slurry and calcium nitrate. *J. Soil Sci.* 37:69-80.
- Elliott, P.W., D. Knight, and J.M. Anderson. 1991. Variables controlling denitrification from earthworm casts and soil in permanent pastures. *Biol. Fertil. Soils* 11:24-29.
- Firestone, M.K. 1982. Biological denitrification. p. 289-318. *In* F.J. Stevenson (ed.) *Nitrogen in Agricultural Soils. Agron. Monogr. 22.* Amer. Soc. Agron., Madison, WI.
- Goulding, K.W.T., and C.P. Webster. 1989. Denitrification losses of nitrogen from arable soils as affected by old and new organic matter from leys and farmyard manure. p. 225-234. *In* J.A. Hansen and K. Hendriksen (ed.) *Nitrogen in Organic Wastes Applied to Soils.* Academic Press, San Diego, CA.
- Groffman, P.M., and J.M. Tiedje. 1991. Relationships between denitrification, CO<sub>2</sub> production and air-filled porosity in soils of different texture and drainage. *Soil Biol. Biochem.* 23: 299-302.
- Harkess, R.D., and J. Frame. 1986. Efficient use of fertilizer nitrogen on grass swards: effects of timing, cutting management and secondary grasses. p. 29-38. *In* H.G. van der Meer, J.C. Ryden, and G.C. Ennik (ed.) *Nitrogen Fluxes in Intensive Grassland Systems.* Martinus Nijhoff Publishers, Dordrecht, The Netherlands.

- Hart, S.J., and M.K. Firestone. 1988. Evaluation of three in situ soil nitrogen availability assays. *Can. J. For. Res.* 19:185-191.
- Hatch, D.J., S.C. Jarvis, and S.E Reynolds. 1991. An assessment of the contribution of net mineralization to N cycling in grass swards using a field incubation method. *Plant Soil* 138:23-32.
- Huddleston, J.H. 1982. Soils of Oregon: Summaries of Physical and Chemical Data. Spec. Rep. 662. Oregon State Extension Service, Corvallis, OR.
- Jarvis, S.C., D. Barraclough, J. Williams, and A.J. Rook. 1991. Patterns of denitrification loss from grazed grassland: effects of N fertilizer inputs at different sites. *Plant Soil* 131:77-88.
- Jenkinson, D.S. 1990. Turnover of organic C and N in the soil. *Phil. Trans. R. Soc. London.* B329:361-368.
- Joffre, R. 1990. Plant and soil nitrogen dynamics in mediterranean grasslands: a comparison of annual and perennial grasses. *Oecologia* 85:142-149.
- Johnston, A.E., S.P. McGrath, P.R. Poulton, and P.W. Lane. 1989. Accumulation and loss of nitrogen from manure, sludge and compost: Long-term experiments at Rothamsted and Woburn. p. 126-139. *In* J.A. Hansen and K. Henriksen (ed.) *Nitrogen in Organic Wastes Applied to Soils.* Academic Press, San Diego, CA.
- Keeney, D.R. 1982. Nitrogen management for maximum efficiency and minimum pollution. p. 605-641. *In* F.J Stevenson (ed.) *Nitrogen in Agricultural Soils.* Agron. Monogr. 22. Amer. Soc. Agron., Madison, WI.
- Legg J.O., and J.J. Meisinger. 1982. Soil nitrogen budgets. p. 503-566. *In* *Nitrogen in Agricultural Soils.* Agron. Monogr 22. Amer. Soc. Agron., Madison, WI.
- Linn, D.M., and J.W. Doran. 1984. Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and non-tilled soils. *Soil Sci. Soc. Am. J.* 48:1267-1272.
- Moore, J.M, M.J. Gamroth, S. Skarda, S. Kelly, N. Baumeister, and D. Myrold. 1992. Pathways of Nitrogen Movement from Land Spread Manure. Final Report to Soil Conservation Service. 17 pp.
- Myers, R.J.K., C.A. Campbell, and K.L. Weier. 1982. Quantitative relationship between net nitrogen mineralization and moisture content of soils. *Can. J. Soil Sci.* 62:111-124.
- Myrold, D.D. 1988. Denitrification in ryegrass and winter wheat cropping systems of western Oregon. *Soil Sci. Soc. Am. J.* 52:412-416.

- Parkin, T.B. 1987. Soil microsites as a source of denitrification variability. *Soil Sci. Soc. Am. J.* 51:1194-1199.
- Parkin, T.B., and J.A. Robinson. 1989. Stochastic models of soil denitrification. *Appl. Environ. Microbiol.* 55:72-77.
- Parkin, T.B., J.L. Starr, and J.J. Meisinger. 1987. Influence of sample size on measurement of soil denitrification. *Soil Sci. Soc. Am. J.* 51:1492-1501.
- Parsons, L.L. R.E. Murray, and M.S. Smith. 1991. Soil denitrification dynamics: spacial and temporal variations of enzyme activity populations and nitrogen gas loss. *Soil Sci. Soc. Am. J.* 55:90-95.
- Patten D.K., J.M. Bremner, and A.M. Blackmer. 1980. Effects of drying and air-dry storage of soils on their capacity for denitrification of nitrate. *Soil Sci. Soc. Am. J.* 44:67-70.
- Pettit, G., and T.K. Thomas. 1986. Oregon Groundwater Quality. p. 423-428. *In* National Water Summary 1986--Groundwater Quality: State Summaries. U.S. Geological Survey, Portland, OR.
- Pratt, P.F., S. Davis, and R.G. Sharpless. 1976. A four-year field trial with animal manures I. Nitrogen balances and yields. *Hilgardia* 44:99-112.
- Raison, R.J., M.J. Connell, and P.K. Khanna. 1987. Methodology for studying fluxes of soil mineral-N in situ. *Soil Biol. Biochem.* 19:521-530.
- Rangeley, A., and P. Newbould. 1986. The response to nitrogen fertilizer from a cut perennial ryegrass (*Lolium perenne* L.) pasture in the scottish uplands relative to efficiency of fertilizer use and provision of herbage to animals. p. 19-28. *In* H.G. van der Meer, J.C. Ryden and G.C. Ennik (ed.) Nitrogen Fluxes in Intensive Grassland Systems. Martinus Nijhoff Publishers, Dordrecht, The Netherlands.
- Rees, R.M. 1989. Measurement of nitrogen mineralization in soil incubations. p. 11-24. *In* J.A. Hansen and K. Hendriksen (ed.) Nitrogen in Organic Wastes Applied to Soils. Academic Press, San Diego, CA.
- Rice, C.W., P.E. Sierzega, J.M. Tiedje, and L.W. Jacobs. 1988. Simulated denitrification in the microenvironment of a biodegradable organic waste injected into soil. *Soil Sci. Soc. Am. J.* 52:102-108.
- Rolston, D.E., A.N. Sharpley, D.W. Toy, and F.E. Broadbent. 1982. Field measurement of denitrification: III. Rates during irrigation cycles. *Soil Sci. Soc. Am. J.* 46:289-296.

- Ryden, J.C. 1986. Gaseous losses of nitrogen from grassland. p. 59-73. *In* H.G. van der Meer, J.C. Ryden, and G.C. Ennik (ed.) *Nitrogen Fluxes in Intensive Grassland Systems*. Martinus Nijhoff Publishers, Dordrecht, The Netherlands.
- Snedecor, G.W., and W.G. Cochran. 1980. The mean and standard deviation. p. 26-38. *In* *Statistical Method, Seventh Edition*. Iowa State University Press, Ames, IA.
- Stanford, G. 1982. Assessment of soil nitrogen availability. p. 651-688. *In* *Nitrogen in Agricultural Soils*. Agron. Monogr. 22. Amer. Soc. Agron., Madison, WI.
- Thompson, R.B. 1989. Denitrification in slurry treated soil: occurrence at low temperatures, relationship with soil nitrate and reduction by nitrification inhibitors. *Soil Biol. Biochem.* 21:875-882.
- Thompson, R.B., and B.F. Pain. 1989. Denitrification from cattle slurry applied to grassland. p. 247-260. *In* J.A. Hansen and K. Hendriksen (ed.) *Nitrogen in Organic Wastes Applied to Soils*. Academic Press, San Diego, CA.
- Thompson, R.B., J.C. Ryden, and D.R. Lockyer. 1987. Fate of nitrogen in cattle slurry following surface application or injection to grassland. *J. Soil Sci.* 38:689-700.
- Tiedje, J.M., S. Simkins, and P.M. Groffman. 1989. Perspectives on measurement of denitrification in the field including recommended protocols for acetylene based methods. p. 217-240. *In* M. Clarholm and L. Bergstrom (ed.) *Ecology of Arable Land*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Whitehead, D.C. 1986. Sources and transformations of organic nitrogen in intensively managed grassland soils. p. 47-58. *In* H.G. van der Meer, J.C. Ryden, and G.C. Ennik (ed.) *Nitrogen Fluxes in Intensive Grassland Systems*. Martinus Nijhoff Publishers, Dordrecht, The Netherlands.
- Whitehead, D.C., A.W Bristow, and D.R. Lockyer. 1990. Organic matter and nitrogen in the unharvested fraction of grass swards in relation to the potential for nitrate leaching after ploughing. *Plant Soil* 123:39-49.
- Wilhelm, I., R. Battino, and R.J. Wilcock. 1977. Low-pressure solubility of gases in liquid water. *Chem. Rev.* 77:219-262.




Figure 1a. Denitrification and respiration rates in the Waldo soil. Statistical significance is indicated by the following symbols: "+" for  $0.05 < p \leq 0.10$ , "\*" for  $0.01 < p \leq 0.05$ , and ":" for  $p \leq 0.01$ . Respiration rates are averaged over all treatments.




Figure 1b. Denitrification and respiration rates in Amity soil. Statistical significance is indicated by the following symbols: "+" for  $0.05 < p \leq 0.10$ , "\*" for  $0.01 < p \leq 0.05$ , and ":" for  $p \leq 0.01$ . Respiration rates are averaged over all treatments.






Figure 1c. Denitrification and respiration rates in Quillamook soil. Statistical significance is indicated by the following symbols: "+" for  $0.05 < p \leq 0.10$ , "\*" for  $0.01 < p \leq 0.05$ , and ":" for  $p \leq 0.01$ . Respiration rates are averaged over all treatments.




Figure 2a. Soil NO<sub>3</sub><sup>-</sup> concentration in the Waldo soil. Means followed by the same letter are not significantly different (FPLSD,  $\alpha = 0.05$ ). However, the control was significantly lower than the manure treatments at the June 1990 ( $p=0.09$ ) and October 1990 ( $p=0.08$ ) samplings.




Figure 2b. Soil NO<sub>3</sub><sup>-</sup> concentration in the Amity soil. Means followed by the same letter are not significantly different (FPLSD,  $\alpha = 0.05$ ).




Figure 2c. Soil NO<sub>3</sub><sup>-</sup> concentration in the Quillamook soil. Means followed by the same letter are not significantly different (FPLSD,  $\alpha = 0.05$ ).




Figure 3a. Difference between bulk soil water content and water content of incubated soil for the Waldo soil.




Figure 3b. Difference between bulk soil water content and water content of incubated soil for the Amity soil.




Figure 3c. Difference between bulk soil water content and water content of incubated soil for the Quillamook soil.




Figure 4. Net N mineralization rates and water content expressed as percent of water-holding capacity for individual cores (Waldo soil, fall/winter set, control treatment). n=16.



Table 1. Selected characteristics of the three western Oregon soils used in this study.

Soil	MAP†	Drainage class	Organic C	Total N	Bulk density	pH‡
	mm		—g kg <sup>-1</sup> —		Mg m <sup>-3</sup>	
Amity	1070	somewhat poor	33.9	2.5	1.2	5.9
Waldo	1070	poor	32.9	2.3	1.2	6.1
Quillamook	2350	well	90.0	7.0	0.7	5.5

† Mean Annual Precipitation

‡ 2:1 H<sub>2</sub>O:soil

Table 2. Manure application dates.

	Amity	Waldo	Quillamook
	date		
1990	10/15/89	10/15/89	03/22/90
	04/02/90	04/02/90	04/26/90
	05/01/90	05/01/90	05/24/90
	06/01/90	06/01/90	06/21/90
	07/05/90	08/02/90	07/30/90
	11/01/90	11/01/90	09/14/90
			11/20/90
Annual total† (kg N ha <sup>-1</sup> )	150	150	175
1991	04/22/91	04/22/91	03/22/91
	05/28/91	05/28/91	05/01/91
	07/10/91	07/10/91	06/12/91
	08/20/91	08/20/91	07/15/91
	10/17/91	10/17/91	08/12/91
			10/09/91

† Amount applied at the 150 kg N ha<sup>-1</sup> rate; amounts for the 300 and 450 kg N ha<sup>-1</sup> treatments are proportional to these actual rates.

Table 3. Correlation coefficients (r) for denitrification. Denitrification and respiration rates were log transformed. Values in table were significant at  $\alpha = 0.10$ .

		Soil		
Variable		Amity	Waldo	Quillamook
All data	Soil water content	.56	.35	.19
(n=400)	Respiration	.09	.18	.41
Mean (by date)	Soil water content	.53	.67	
(n=13)	Respiration			.70
Mean (active period†)	Soil water content	.75	.77	
	Respiration			.89

† Amity (n=6), Waldo (n=7), Quillamook (n=8).

Table 4. Annual denitrification losses by manure treatment. Figures in parentheses is the percent of applied manure N lost, after subtraction of the control. Means followed by the same letter within a column are not significantly different (Tukey,  $p=0.10$ ).

Treatment	Soil		
	Waldo†	Amity‡	Quillamook‡
	kg N ha <sup>-1</sup>		
control	11a§	21a	12a
150 kg N	24ab (8%)	36a (10%)	38ab (13%)
300 kg N	31b (6%)	44a (7%)	58b (12%)
450 kg N	33b (5%)	47a (5%)	108c (16%)

† n=4 for control treatment, n=2 for manure treatments

‡ n=3

Table 5. Measurement of net N mineralization rates in unfertilized controls by three different methods for the 1990 growing season (mean and 95% confidence interval).

Soil	Method		
	core-IER	plant N uptake†	SON‡
	kg N ha <sup>-1</sup> y <sup>-1</sup>		
Amity	338 ± 111	118 ± 18	94 ± 102
Waldo	237 ± 88	58 ± 18	53 ± 139
Quillamook	1213 ± 373	196 ± 16	25 ± 284

† Data courtesy of J.A. Moore and M.J. Gamroth (Appendix).

‡ Half of the net loss over two growing seasons. The error was also halved to provide an estimated confidence interval.

Table 6. Initial (March, 1990) and final (October, 1991) soil organic N.

Soil	Treatment	Total Kjeldahl N		
		Initial	Final	net change †
		kg N ha <sup>-1</sup>		
Amity	Control	4,615	4,327	-288 ± 203
	150 kg N	4,485	4,442	-43 ± 203
	300 kg N	4,342	4,530	188 ± 203
	450 kg N	4,718	4,740	22 ± 203
Waldo	Control	4,172	4,066	-105 ± 278
	150 kg N	4,208	4,434	226 ± 393
	300 kg N	4,090	4,236	147 ± 393
	450 kg N	3,912	3,956	44 ± 393
Quillamook	Control	7,117	7,068	-49 ± 568
	150 kg N	7,310	7,454	144 ± 568
	300 kg N	7,582	7,597	16 ± 568
	450 kg N	7,417	7,537	324 ± 695
All soils	Control	-	-	-143 ± 174
	150 kg N	-	-	94 ± 195
	300 kg N	-	-	113 ± 195
	450 kg N	-	-	114 ± 209

† Mean ± 95% confidence interval.

Table 7. Net N Mineralization by the core-IER method. Data for seasonal periods are means and annual estimates are means  $\pm$  90 % confidence intervals.

Treatment	Waldo			
	Spring	Summer	Fall/winter	Annual§
	kg N ha <sup>-1</sup>			
Control§	110	142	32	264 $\pm$ 95
150 kg N†	73	173	50	296 $\pm$ 134
300 kg N†	87	296	42	425 $\pm$ 134
450 kg N†	141	265	63	469 $\pm$ 134
	Amity			
	Spring	Summer	Fall/winter	Annual
	kg N ha <sup>-1</sup>			
Control†	194	91	54	338 $\pm$ 111
150 kg N‡	87	115	64	252 $\pm$ 91
300 kg N‡	185	96	62	352 $\pm$ 91
450 kg N‡	200	150	64	414 $\pm$ 91
	Quillamook			
	Spring	Summer	Fall/winter	Annual
	kg N ha <sup>-1</sup>			
Control‡	370	454	224	1048 $\pm$ 360
150 kg N‡	310	316	257	923 $\pm$ 360
300 kg N‡	246	558	211	1012 $\pm$ 360
450 kg N‡	397	738	256	1391 $\pm$ 360

† n=2

‡ n=3

§ n=4

Table 8.  $\text{NO}_3^-$  and  $\text{NH}_4^+$  collected on the resin bags as a percent of total net mineralized N.

Season	Soil		
	Waldo	Amity	Quillamook
Spring	5	11	20
Summer	14	23	4
Fall/winter	63	73	85



Table 9. Percent of mineralized N nitrified, averaged over all treatments.

Season	Soil		
	Waldo	Amity	Quillamook
	%		
Spring	87	75	71
Summer	70	62	81
Fall/winter	58	71	88

## APPENDIX

Table A1. Nitrogen in harvested biomass. Mean  $\pm$  standard deviation; n=3 for Amity and Quillamook, n=4 for Waldo. Data courtesy of J.A. Moore and M.J. Gamroth.

Soil	Treatment	Year	
		1990	1991
— kgN ha <sup>-1</sup> —			
Amity	Control	123 $\pm$ 8	148 $\pm$ 13
	150 kg N	116 $\pm$ 4	176 $\pm$ 36
	300 kg N	139 $\pm$ 15	181 $\pm$ 19
	450 kg N	141 $\pm$ 13	220 $\pm$ 76
Waldo	Control	58 $\pm$ 15	53 $\pm$ 17
	150 kg N	71 $\pm$ 18	86 $\pm$ 11
	300 kg N	75 $\pm$ 10	104 $\pm$ 8
	450 kg N	70 $\pm$ 9	129 $\pm$ 20
Quillamook	Control	196 $\pm$ 11	176 $\pm$ 11
	150 kg N	216 $\pm$ 12	206 $\pm$ 11
	300 kg N	209 $\pm$ 16	246 $\pm$ 16
	450 kg N	202 $\pm$ 4	239 $\pm$ 18