



Contribution to Special Issue: 'Towards a Broader Perspective on Ocean Acidification Research' Original Article

Interpretation and design of ocean acidification experiments in upwelling systems in the context of carbonate chemistry co-variation with temperature and oxygen

Jonathan C. P. Reum^{1*†}, Simone R. Alin², Chris J. Harvey¹, Nina Bednaršek², Wiley Evans^{2,3}, Richard A. Feely², Burke Hales⁴, Noelle Lucey⁵, Jeremy T. Mathis^{2,3}, Paul McElhany¹, Jan Newton⁶, and Christopher L. Sabine²

¹Conservation Biology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration, 2725 Montlake Blvd. E, Seattle, WA 98112, USA

²Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, 7600 Sand Point Way NE, Seattle, WA 98115, USA

³Ocean Acidification Research Center, School of Fisheries and Ocean Sciences, University of Alaska Fairbanks, 245 O'Neill Bldg, Fairbanks, AK 99775, USA

⁴College of Earth, Ocean and Atmospheric Sciences, Oregon State University, Corvallis, OR 97331, USA

⁵Marine Environment and Sustainable Development Unit ENEA, PO Box 224, La Spezia, Italy

⁶Applied Physics Laboratory, University of Washington, Box 355640, Seattle, WA 98105, USA

*Corresponding author: tel: +1 206 860 3204; fax: +1 206 860 3217; e-mail: jonathan.reum@noaa.gov

†Present address: Washington Sea Grant, University of Washington, Box 355640, Seattle, WA, USA

Reum, J. C. P., Alin, S. R., Harvey, C. J., Bednaršek, N., Evans, W., Feely, R. A., Hales, B., Lucey, N., Mathis, J. T., McElhany, P., Newton, J., and Sabine, C. L. Interpretation and design of ocean acidification experiments in upwelling systems in the context of carbonate chemistry co-variation with temperature and oxygen. – ICES Journal of Marine Science, 73: 582–595.

Received 22 July 2014; revised 24 November 2014; accepted 26 November 2014; advance access publication 7 January 2015.

Coastal upwelling regimes are some of the most productive ecosystems in the ocean but are also among the most vulnerable to ocean acidification (OA) due to naturally high background concentrations of CO₂. Yet our ability to predict how these ecosystems will respond to additional CO₂ resulting from anthropogenic emissions is poor. To help address this uncertainty, researchers perform manipulative experiments where biological responses are evaluated across different CO₂ partial pressure (*p*CO₂) levels. In upwelling systems, however, contemporary carbonate chemistry variability remains only partly characterized and patterns of co-variation with other biologically important variables such as temperature and oxygen are just beginning to be explored in the context of OA experimental design. If co-variation among variables is prevalent, researchers risk performing OA experiments with control conditions that are not experienced by the focal species, potentially diminishing the ecological relevance of the experiment. Here, we synthesized a large carbonate chemistry dataset that consists of carbonate chemistry, temperature, and oxygen measurements from multiple moorings and ship-based sampling campaigns from the California Current Ecosystem (CCE), and includes fjord and tidal estuaries and open coastal waters. We evaluated patterns of *p*CO₂ variability and highlight important co-variation between *p*CO₂, temperature, and oxygen. We subsequently compared environmental *p*CO₂ – temperature measurements with conditions maintained in OA experiments that used organisms from the CCE. By drawing such comparisons, researchers can gain insight into the ecological relevance of previously published OA experiments, but also identify species or life history stages that may already be influenced by contemporary carbonate chemistry conditions. We illustrate the implications co-variation among environmental variables can have for the interpretation of OA experimental results and suggest an approach for designing experiments with *p*CO₂ levels that better reflect OA hypotheses while simultaneously recognizing natural co-variation with other biologically relevant variables.

Keywords: California Current, climate change, hypoxia, multistressor experiment, pH.

Introduction

Coastal upwelling systems located along the eastern boundary of ocean basins are some of the most productive ecosystems in the ocean but are also among the most vulnerable to OA (Feely *et al.*, 2008; Gruber *et al.*, 2012). The net transport of deep, nutrient-rich waters to the sunlit surface by upwelling-favourable winds promotes high rates of primary production which in turn supports productive foodwebs and major fisheries (Fréon *et al.*, 2009). However, subsurface and newly upwelled waters naturally exhibit low O₂ and high CO₂ concentrations due to the remineralization of organic material exported from surface layers. Consequently, they have a reduced capacity to buffer against changes in carbonate chemistry resulting from ocean uptake of anthropogenic CO₂ relative to open-ocean surface waters (Feely *et al.*, 2008; Fassbender *et al.*, 2011; Harris *et al.*, 2013). In eastern Pacific systems such as the California Current, the CO₂ burden and O₂ drawdown due to respiration are high because the source waters transported to upwelling centres along the coast have been isolated from the surface for a few decades (Feely *et al.*, 2008; Hauri *et al.*, 2009). In the California Current, anthropogenic CO₂ has already lowered pH by ~0.1, causing the depth of undersaturation with respect to aragonite to shoal and expanding the spatial extent of undersaturated surface waters (Feely *et al.*, 2008; Gruber *et al.*, 2012; Harris *et al.*, 2013). In only a few more decades, models suggest that the depth of undersaturation may shoal into the upper 75 m of the water column in some regions year-round (Gruber *et al.*, 2012; Hauri *et al.*, 2013). Given the economic, ecological, and biogeochemical importance of eastern boundary upwelling regions, understanding how species that compose these ecosystems will respond to OA has emerged as a high research priority (Fabry *et al.*, 2008; Gruber, 2011; Doney *et al.*, 2012).

To evaluate the sensitivity of species to OA, researchers commonly rely on manipulative experiments where organisms are exposed to different carbonate chemistry conditions. Typically, experiments include “control” conditions that attempt to simulate contemporary or preindustrial CO₂ concentrations and “acidified” treatments that correspond to potential future CO₂ uptake by the oceans. For studies focused on organisms from low productivity, open-ocean surface waters researchers can rely on IPCC scenarios of atmospheric pCO₂ concentrations to identify potential carbonate chemistry treatments because assumptions of air–sea pCO₂ equilibrium are often nearly met (Barry *et al.*, 2010; Orr, 2011). However, in upwelling systems, CO₂ levels are more variable relative to open ocean waters due to the outcropping of high-CO₂ subsurface waters and high rates of primary production and respiration that strongly modulate seawater carbonate chemistry (Hales *et al.*, 2005; Feely *et al.*, 2008; Borges and Abril, 2011; Fassbender *et al.*, 2011). Consequently, OA experiments that use organisms from these habitats and that rely on IPCC future atmospheric CO₂ scenarios to devise control and acidified seawater treatments may inadequately replicate contemporary carbonate chemistry or include treatments that fail to reflect realistic future OA hypotheses (Barry *et al.*, 2010; Andersson and Mackenzie, 2012; McElhany and Busch, 2012).

Recognition of the importance of including environmentally relevant pCO₂ levels in OA experiments has grown considerably, and has led to the use of seawater chemistry monitoring programmes to inform treatment design in several recent OA studies (Hofmann *et al.*, 2014). Less appreciated from an experimental perspective, however, is the possibility that carbonate chemistry

conditions may also naturally co-vary with other biologically relevant variables including temperature and O₂ over multiple spatial and temporal scales (Reum *et al.*, 2014). This may have important implications for the design and interpretation of ocean acidification (OA) experiments because of the potential for non-additive interactions between carbonate chemistry, temperature, and O₂ on organismal performance and ecological interactions (Wernberg *et al.*, 2012, Harvey *et al.*, 2013; Koch *et al.*, 2013; Kroeker *et al.*, 2013). For many organisms, aerobic capacity and metabolic scope (the amount of energy that can be allocated to activities beyond those required for basic existence) may be influenced strongly by temperature, and adversely impacted by reductions in ambient oxygen availability or increases in CO₂ concentrations (Pörtner, 2010, 2012).

Given the combined effects that temperature and environmental O₂ and pCO₂ have on organismal physiology and experimental evidence indicating non-additive interactions on response variables related to fitness and ecosystem function (Pörtner and Farrell, 2008; Pörtner, 2010; Harvey *et al.*, 2013; Kroeker *et al.*, 2013), knowledge of their co-variability is essential for designing OA experiments that adequately characterize biological performance under contemporary relative to future acidified conditions. Yet for workers focused on laboratory OA experiments, these relationships are rarely incorporated into experimental designs. This poses important potential drawbacks. Foremost, if carbonate chemistry strongly covaries with temperature or O₂, researchers risk running experiments with control-water characteristics that are atypical of the habitat to which a focal organism/life stage may have acclimated or adapted to. Recently, pCO₂ was shown to range widely (~200–2500 µatm) and co-vary with temperature and O₂ in a fjord located in the northeast Pacific (Reum *et al.*, 2014). In that system, the strength and direction of the relationships changed with season, but also between subregions that differed in terms of vertical mixing. Consequently, within a given season and region, organisms occurring in low pCO₂ waters experience temperatures and O₂ levels that differed from those experienced by organisms that occupy high pCO₂ waters, and this has direct implications for how ecologically relevant OA experiments should be designed (Reum *et al.*, 2014).

Here, we expand on the topic and evaluate the potential importance of this issue for upwelling systems. To do so, we have assembled a large dataset of carbonate chemistry measurements from a variety of habitats throughout the California Current Ecosystem (CCE), a major eastern boundary upwelling system that supports highly productive foodwebs. Along the coast, equatorward winds during spring and summer drive surface flow offshore in an Ekman layer, leading to the upwelling of cold, salty, O₂-poor, and nutrient- and CO₂-rich subsurface water. In fall and winter, the wind direction reverses resulting in downwelling and shoreward advection of oceanic waters which are relatively warm, fresh, O₂-saturated, nutrient-deplete, and near air–sea CO₂ equilibrium (Evans *et al.*, 2011; Harris *et al.*, 2013). The dataset presented here consists of pCO₂ measurements from multiple moorings and ship-based sampling campaigns that collectively span 14° of latitude, and includes data from estuary and open coastal water habitats. Although the extent of data is constrained in either time or space for any single sampling campaign, the synthesis of many datasets offers an overview of the potential range of pCO₂, temperature, and O₂ conditions experienced by organisms from this region. Subsequently, we compared environmental pCO₂ and temperature measurements to conditions in OA experiments performed on organisms obtained from populations that reside within the CCE. In doing so, we place these

experiments in a larger environmental context and draw on observations that may help guide OA researchers in considering experimental designs that are more appropriate for organisms that occur in the CCE and similar coastal systems.

Material and methods

Carbonate chemistry data

Our analysis focused on coastal waters within the domain of the northern and central CCE (Figure 1). The region extends from northern Vancouver Island, British Columbia (50°N) to Point Conception, California (34°N), and is typified by distinct wind-driven upwelling (May–October) and downwelling (November–April) oceanographic seasons (Checkley and Barth, 2009). The region includes large expanses of open coastal habitat and many inlets and estuaries that range considerably in terms of surface area, depth, substrate, and connectivity with the open ocean (Hickey and Banas, 2003). To obtain an overview of the potential range of temperatures and carbonate chemistry conditions, we retrieved all publicly available records of measurements sampled within our region of interest from the Carbon Dioxide Information Analysis Center (CDIAC) and Surface Ocean CO₂ Atlas (SOCAT, Bakker et al., 2013). The dataset includes carbonate chemistry measurements from discrete water samples collected at depth, shipboard measurements of surface water *p*CO₂, and mooring *p*CO₂ time-series of surface waters off the coast of California and Washington (Supplementary Table S1). In addition to CDIAC and SOCAT data, we also included unpublished carbonate chemistry data obtained from the 2011 and 2012 NOAA Ocean Acidification Programme West Coast Cruise

survey, which sampled waters along the continental shelf of western North America; mooring *p*CO₂ time-series of surface waters from an open coast location (Newport, Oregon), and a shallow tidal estuary (Netarts Bay, Oregon); and published carbonate chemistry data from late summer and fall surveys in a large fjord estuary complex (Strait of Juan de Fuca and Puget Sound, Washington; Supplementary Table S1). Further details of the survey, sampling dates, sampling method, and habitats from which data were obtained are provided in Supplementary Table S1. Although the dataset does not include all habitat types that occur within the CCE, we believe these data are sufficiently representative enough to offer an overview of key patterns of carbonate chemistry, temperature, and O₂ co-variation. For the purposes of the present study, we included all measurements extending up to 200 km from the coast.

For our analysis, we focused on *p*CO₂ rather than other parameters of the carbonate system because *p*CO₂ is directly changed by anthropogenic CO₂ emissions and is the most commonly used treatment variable in species-exposure OA experiments. Further, direct measurements of *p*CO₂ were more widely available than other parameters in the CDIAC and SOCAT databases, and when not available were readily estimated from other measured parameters of the carbonate system (see Supplementary Table S1 for details on estimation methods). We recognize that organisms may potentially be more sensitive to other variables of the carbonate system such as pH or to changes in the calcium carbonate saturation state of seawater (Barton et al., 2012; Waldbusser and Salisbury, 2013) and that waters with similar *p*CO₂ values may differ with respect to aragonite or calcite saturation states if total alkalinities differ. However, *p*CO₂ is routinely reported in OA studies and is the parameter that allowed us to best standardize comparisons among experiments and between experiments and environmental carbonate chemistry measurements. Further, the focus on *p*CO₂ prevented the use of some carbonate chemistry datasets such as those including information only on pH (e.g. Hofmann et al., 2011). We limited our analysis to *p*CO₂ measurements taken from the top 50 m of the water column because the species and life history stages for which OA experiments have been performed from the northern and central CCE typically occur in waters within this depth range, although individuals of some populations may occur for periods below this depth.

We evaluated separate relationships between *p*CO₂ and temperature and *p*CO₂ and O₂ within summer upwelling and winter downwelling periods. Co-variation patterns for *p*CO₂ and temperature were examined for data pooled from shipboard underway and discrete sample measurements collected along the West Coast continental shelf. In addition, we also examined co-variation patterns for Puget Sound and the Strait of Juan de Fuca discrete sample measurements (hereafter referred to as “Puget Sound”). A subset of these data were examined in an earlier study (Reum et al., 2014), but we include them here for completeness and to facilitate comparisons among habitats represented within the CCE. We chose to present co-variation patterns for Puget Sound separately from the West Coast because Puget Sound exhibits slow exchange with open coastal waters and high rates of primary productivity and respiration (Feely et al., 2010). The resulting *p*CO₂–temperature relationships therefore likely differ considerably from patterns observed in shelf waters. For the time-series data, we estimated *p*CO₂–temperature relationships for upwelling and downwelling seasons, but examined co-variation patterns separately for each mooring to evaluate site-level differences.

To evaluate *p*CO₂ and O₂ relationships and facilitate comparisons across survey types and seasons, we converted all O₂ concentration




Figure 1. Map of coordinates where (a) environmental carbonate chemistry data were obtained from the northern and central CCE from moorings and ship-based underway and discrete samples, (b) the locations where organisms (or their broodstock) included in published OA experiments were collected. In (a), samples coded as Puget Sound also include measurements from the adjoining Strait of Juan de Fuca. For additional details on environmental carbonate chemistry datasets, see Supplementary Table S1.

measurements to micromole per kilogramme. As with the $p\text{CO}_2$ and temperature data, we examined co-variation within data pooled from shipboard West Coast measurements, Puget Sound, and each mooring and by season. To improve assumptions of normality in the residual error structure for the $p\text{CO}_2$ –temperature and $p\text{CO}_2$ – O_2 relationships, we \log_{10} -transformed $p\text{CO}_2$ values before estimating linear relationships using least-squares regression. For presentation purposes, the linear relationships were back-transformed to the original $p\text{CO}_2$ scale. Our goal was to estimate overall mean relationships within seasons and survey types to facilitate visual inspection and to evaluate co-variation patterns in an exploratory manner. We therefore did not test for significant differences between seasons or survey types. All linear relationships were fitted using the R version 2.11 statistical software package (R Development Core Team, 2011).

OA experimental studies

We searched the published literature for OA experiments that included organisms obtained directly from habitats within the northern and central CCE or that originated from broodstock collected from the region. To do so, we searched Google Scholar, ISI databases (Web of Science, Current Contents), and references included in recently published reviews and meta-analyses that address OA (Dupont *et al.*, 2010; Wernberg *et al.*, 2012; Harvey *et al.*, 2013; Kroeker *et al.*, 2013; Wittmann and Pörtner, 2013). We used the search terms “ocean acidification”, “carbon dioxide”, “experiment”, and “manipulation”. We included all studies published through 15 October 2013 that were found under these search criteria.

From each study, we retrieved information on the collection site of the organism (or their broodstock) and the temperatures and $p\text{CO}_2$ levels at which the experiment was performed. For nearly all studies, experimental O_2 concentrations were not reported nor could saturation conditions be safely assumed. We therefore focused our comparison of environmental and experimental conditions to $p\text{CO}_2$ and temperature. Following the authors’ interpretation of the results, we recorded whether the dependent biological variables differed significantly in a positive or negative direction or showed no significant difference relative to the control $p\text{CO}_2$ level specified by the author. If authors did not explicitly designate a control $p\text{CO}_2$ level in their study, we considered treatments with $p\text{CO}_2$ levels closest to present-day air $p\text{CO}_2$ levels ($\sim 400 \mu\text{atm}$) as the control to facilitate comparisons across studies. When more than one response variable was tested in an experiment, we coded the net outcome of the experiment at a given treatment level based on the result of the variable most sensitive to $p\text{CO}_2$.

From the outset, we recognized that the species and life history stages for which published OA experimental data were available held distributions that spanned large sections of the northern and central CCE or were planktonic with high dispersal potential. Further, the exact dispersal patterns of many of these species are not well understood. Although larvae of some species may be functionally limited to a subset of the CCE system (e.g. some species may occur primarily in bays or estuaries), “spillage” into adjoining habitats and waters through advective processes is also probable. We therefore used the full dataset of field $p\text{CO}_2$ and temperature measurements to demarcate the potential $p\text{CO}_2$ –temperature space organisms may encounter. We acknowledge that the actual $p\text{CO}_2$ levels and temperatures experienced by organisms will differ for populations across locations, seasons, and due to possible interannual variation in upwelling and climate forcing. Our main intention, however, was to draw comparisons between the conditions maintained in OA experiments and the full range of environmental

$p\text{CO}_2$ and temperature values based on empirical observations. In the absence of detailed information on the fine-scale distribution and movement patterns of most species and life history stages, and environmental $p\text{CO}_2$ –temperature measurements of matching resolution, comparisons at finer spatio-temporal scales were not possible.

Results

$p\text{CO}_2$, temperature, and O_2 co-variation

Temperature and $p\text{CO}_2$ values in the upper 50 m of the CCE spanned 6 – 19°C and 100 – $1500 \mu\text{atm}$, respectively, based on the pooled shipboard underway and discrete water sample measurements along the open coast. $p\text{CO}_2$ values tended to increase with decreasing water temperature during summer upwelling months (Figure 2a). In Puget Sound, a similar but steeper relationship was apparent relative to the open coastal waters, reflecting CO_2 -enriched waters (Figure 2a). Along the open coast, the range of $p\text{CO}_2$ values was also wider at cool relative to warm temperatures. For instance, $p\text{CO}_2$ values at 9°C ranged from 320 to $1400 \mu\text{atm}$, while at 16°C , the range extended from 130 to $420 \mu\text{atm}$. In Puget Sound, the $p\text{CO}_2$ range was also wider at cooler temperatures (Figure 2a).

Although fewer measurements were available during winter in general (Supplementary Table S1), the data showed weak co-variation between $p\text{CO}_2$ and temperature in open coastal waters and the overall range of $p\text{CO}_2$ values and temperatures narrowed relative to summer (Figure 2b). This is due to the relative absence of the cold, high- $p\text{CO}_2$ upwelled source waters. In contrast, winter $p\text{CO}_2$ and temperature values in Puget Sound positively covaried, and the range of $p\text{CO}_2$ values and temperatures also narrowed relative to summer (Figure 2b).

Time-series data collected from moored platforms in open coastal waters from four locations also indicated co-variation between $p\text{CO}_2$ and temperature (Figure 3a–c). During summer upwelling months, $p\text{CO}_2$ again generally declined with increasing temperature (Figure 3a–c). The range of $p\text{CO}_2$ levels among stations, however, differed substantially. Time-series data collected at moorings in shelf waters off Washington and near Point Conception, CA. (Figure 3a and b) ranged in $p\text{CO}_2$ from 200 to 600 and 300 to $600 \mu\text{atm}$, respectively, while off the coast of Oregon the range spanned 200 – $1100 \mu\text{atm}$ (Figure 3c). The differences among stations reflect considerable spatial variation in the supply of newly upwelled waters.

Time-series data collected from Netarts Bay, a shallow tidal estuary in northern Oregon, indicated summer range in $p\text{CO}_2$ values was also considerable, spanning 300 – $800 \mu\text{atm}$ and reflected high rates of primary production and respiration (Figure 3d). The overall relationship between $p\text{CO}_2$ and temperature, however, was also negative. In winter, co-variation between $p\text{CO}_2$ and temperature at all mooring stations was weaker relative to summer but variable across stations (Figure 3d). In addition, the range of temperatures and $p\text{CO}_2$ values were generally lower in winter relative to summer in two of the five stations (Figure 3c and d).

Measurements of O_2 were available for a subset of $p\text{CO}_2$ records included in the complete environmental $p\text{CO}_2$ and temperature dataset (Supplementary Table S1). For all survey types, mooring time-series, and seasons, $p\text{CO}_2$ levels decreased with increasing O_2 concentration (Figure 4). For most moorings on the shelf, O_2 ranged from 230 to $350 \mu\text{mol kg}^{-1}$ (both seasons combined). Summer O_2 levels at Newport Oregon, however, ranged from 120 to $350 \mu\text{mol kg}^{-1}$ (Figure 4). Summer O_2 measurements from discrete samples from




Figure 2. (a) Relationship between $p\text{CO}_2$ and temperature in the top 50 m of the water column in the northern and central CCE during summer upwelling season which approximately spans May through October and (b) winter (November through April) when downwelling-favourable winds predominate. All non-time-series data are displayed. Measurements of $p\text{CO}_2$ from cool waters in Puget Sound are elevated relative to other regions sampled in the CCE. Regression lines are overlaid to aid evaluation of patterns. Lines labelled P and C denote relationships for Puget Sound and open coastal locations, respectively. For reference, approximate present-day $p\text{CO}_2$ levels ($\sim 390 \mu\text{atm}$) are indicated by the dashed horizontal line.

Puget Sound and the West Coast ranged more widely, from 40 to $400 \mu\text{mol kg}^{-1}$, where concentrations below $\sim 60 \mu\text{mol kg}^{-1}$ reflect hypoxic conditions (Figure 4). The general negative relationship between $p\text{CO}_2$ and O_2 corresponds to the well-understood effects of aerobic respiration and photosynthesis in marine ecosystems. When aerobic respiration dominates, CO_2 is remineralized and O_2 levels are drawn down, while the reverse occurs when photosynthesis dominates.

OA studies

In total, 26 experiments from 22 published OA studies included organisms (or broodstock) from the northern and central CCE (Figure 1b) and provided sufficient information on treatment water conditions to compare with the CCE $p\text{CO}_2$ –temperature space defined by combining all environmental datasets (Table 1). In terms of taxonomic diversity, 50, 35, and 15% of experiments evaluated responses in echinoderms, molluscs, and teleosts, respectively. However, 35% of all experiments focused on responses in a single species, the echinoderm *Strongylocentrotus purpuratus* (Table 1). Of the experiments, 81% examined gamete, egg, or larval performance (survival, growth, calcification rates), while the remaining studies evaluated performance metrics (growth, development, or calcification rates) in juvenile or adult life history stages (Table 1). One study measured genetic diversity in echinoderm larvae after exposure to different $p\text{CO}_2$ treatments to evaluate evolutionary potential.

To facilitate visual inspection and comparison of experimental $p\text{CO}_2$ and temperature treatments with the $p\text{CO}_2$ –temperature space defined by the complete set of environmental measurements in our dataset, we examined echinoderm and non-echinoderm studies separately (Figure 5). Overall, OA experiments included $p\text{CO}_2$ treatment levels that extended from 200 to $4000 \mu\text{atm}$; 42% of experiments included a single elevated $p\text{CO}_2$ treatment in addition to a control at a given temperature, 38% considered two

different elevated $p\text{CO}_2$ treatments, and 19% included three or more elevated treatments (Figure 5). In total, 72% of experiments included at least one control and one elevated $p\text{CO}_2$ treatment that occurred within the observed range of $p\text{CO}_2$ and temperature values in the CCE. Of those, 63% (10 out of 16) observed negative biological effects, 18% observed positive effects, and 18% observed no effect relative to control $p\text{CO}_2$ levels. Only two studies included $p\text{CO}_2$ treatments below approximate present-day atmospheric levels ($\sim 400 \mu\text{atm}$; Figure 5).

Compared with the $p\text{CO}_2$ –temperature space defined by the complete set of environmental measurements in our dataset, five experiments were performed at temperatures that matched or exceeded the warmest observed values ($\sim 19^\circ\text{C}$; Figure 5). These included three experiments on the early life history stages of the native Olympia oyster, an experiment on sand dollar larvae, and an experiment on the non-native Pacific oyster which is routinely reared at $\sim 20^\circ\text{C}$ to optimize survival under commercial hatchery conditions. One experiment included a 2.1°C treatment; though this temperature was meant to simulate cool conditions in Alaskan waters, the source stock was collected near Puget Sound (Table 1).

A review of each study indicated that Intergovernmental Panel on Climate Change (IPCC) estimates of future global surface ocean mean $p\text{CO}_2$ levels were used as the sole rationale for selecting OA treatments in 45% of studies, while 31% cited a combination of regional modelling studies, local field measurements, and IPCC estimates to support their choice of experimental $p\text{CO}_2$ treatment levels. Of the remaining studies, 13% provided no rationale for their choice of $p\text{CO}_2$ treatment levels, one based the high $p\text{CO}_2$ treatment level on observations of contemporary upwelling conditions, and one noted natural high carbonate chemistry variability in coastal upwelling systems which necessitated the need to test biological responses to a wide range of $p\text{CO}_2$ levels. In terms of temperature, 80% of studies did not provide a rationale for their choice of




Figure 3. Relationship between $p\text{CO}_2$ and temperature of surface waters measured during summer (light grey open circle symbols) and winter (dark grey) at open coastal moorings in (a) Washington, (b) California, and (c) Oregon, and (d) a tidal estuary in Oregon. Regression lines are overlaid and labelled S and W to indicate summer or winter, respectively. Dashed line convex hulls demarcate data ranges where seasons overlap. For reference, approximate present-day $p\text{CO}_2$ levels ($\sim 390 \mu\text{atm}$) are indicated by the dashed horizontal line.

experimental levels. The remaining studies cited similarity to local field conditions as a rationale for selecting temperatures. Of the two studies that performed experiments that included crossed temperature and $p\text{CO}_2$ treatments meant to correspond to conditions in the CCE, the warmer treatments were chosen in reference to IPCC global average temperature increase projections.

Discussion

Single-species response experiments have offered important initial evidence that some species and life history stages may be adversely impacted by OA (Kroeker *et al.*, 2013; Wittmann and Pörtner, 2013), but there is a pressing need for the development of experiments that include more appropriate reference treatments that reflect $p\text{CO}_2$ levels species have acclimated or adapted to and acidified treatments that more closely reflect natural heterogeneity in carbonate chemistry (Barry *et al.*, 2010; Andersson and Mackenzie, 2012; McElhany and Busch, 2012; Waldbusser and Salisbury, 2013). Our analysis shows that an additional source of

concern is natural co-variation between carbonate chemistry and other biologically relevant variables including temperature and O_2 . These findings have direct consequences for designing experiments that aim to include control treatments that are similar to natural water conditions and for selecting elevated $p\text{CO}_2$ treatment levels that more closely correspond to OA hypotheses. Further, by placing the findings of published OA experiments into a larger environmental context, we gain information on the ecological relevance of experimental water conditions and insight into the potential sensitivity of some species and life stages to carbonate chemistry conditions that already occur in the CCE.

Although researchers increasingly recognize the importance of basing experimental $p\text{CO}_2$ levels on *in situ* carbonate chemistry observations (Yu *et al.*, 2011; Evans *et al.*, 2013; Hofmann *et al.*, 2014), the implications of natural co-variation with temperature or O_2 to experimental design and inference are only beginning to be explored. Given experimental evidence and theoretical expectations of interactive or synergistic effects between $p\text{CO}_2$, temperature, and O_2 on organisms




Figure 4. Relationships between $p\text{CO}_2$ and O_2 in the top 50 m of the water column during summer upwelling (May through October) and winter downwelling seasons (November through April). Regression lines are overlaid for summer and winter; lines labelled M, P, and C correspond to mooring, Puget Sound, and open coast datasets, respectively. For reference, approximate present-day $p\text{CO}_2$ levels ($\sim 390 \mu\text{atm}$) are indicated by the dashed horizontal line, and the hypoxia threshold ($60 \mu\text{mol kg}^{-1}$) is indicated with a dashed vertical line.

(Pörtner, 2010, 2012; Harvey et al., 2013), we suggest that failure to account for natural co-variation among these variables in habitats from the CCE may lead to results with diminished relevance for making predictions. To illustrate this point, we present an example multistressor experimental scheme typical of published OA experiments in which temperature (three levels at 8, 12, and 16°C) is crossed with two $p\text{CO}_2$ levels that correspond to approximate global surface ocean present-day (400 μatm) and future (800 μatm) conditions (Figure 6a). Under the conventional method, treatments are fully orthogonal which permits estimates of the effect sizes of the individual predictor variables and of their interaction on the response variable. The method holds merit as a tool for comparing the relative influence that each predictor has on the response variable, is widely applied in multistressor experiments, and facilitates the development of mechanistic models. However, if a goal of a study is to evaluate the potential sensitivity of organisms to future OA as is often the case, the design may be inadequate, given natural $p\text{CO}_2$ –temperature co-variation within different habitats and water masses. For example, assuming an organism of interest occurs in shelf waters off Oregon during summer upwelling months (e.g. a pelagic larval invertebrate), the assumption that 800 μatm corresponds to a future OA prediction across all temperatures is not accurate. At the Newport, Oregon, mooring $p\text{CO}_2$ levels of 800 μatm already occur at 8°C under present-day conditions and “control” 400 μatm waters do not (Figure 6a). At temperatures above 13°C, the mean $p\text{CO}_2$ values approach air–sea equilibrium conditions. We do not doubt that simple crossed experimental designs will provide information on the interactive effects of $p\text{CO}_2$ and temperature, but we do question the efficacy of the design for testing OA hypotheses on future ecological response to OA, given naturally occurring $p\text{CO}_2$ –temperature relationships and the wide range of both variables in the CCE.

In light of potential co-variation between carbonate chemistry and other important environmental variables, how should

researchers select $p\text{CO}_2$ treatments that correspond to OA hypotheses? As a starting point, we recommend that OA experimental designs include multiple controls that reflect the span of $p\text{CO}_2$ levels and temperatures likely to be experienced by the organism under study (Figure 5b). To design $p\text{CO}_2$ treatments that represent future OA scenarios in productive coastal systems, we suggest that researchers focus on changes in the anthropogenic contribution to *in situ* dissolved inorganic carbon, DIC (e.g. Feely et al., 2008, 2010; Barry et al., 2010; Melzner et al., 2012; Shaw et al., 2013). At the Newport, Oregon, mooring, newly upwelled waters exhibit $p\text{CO}_2$ values that are elevated relative to air–sea equilibrium conditions due to the remineralization of organic material before surfacing (Evans et al., 2011). However, after surfacing CO_2 concentrations can be drawn down rapidly by photosynthesis (Hales et al., 2005; Hales et al., 2006; Evans et al., 2011), often at rates that typically far exceed CO_2 equilibration times across the air–sea interface (e.g. van Geen et al., 2000; Fassbender et al., 2011). Consequently, the anthropogenic CO_2 burden of upwelled waters is primarily acquired when they were last in contact with the atmosphere and before DIC changes due to biological processes post-surfacing. In our example, $p\text{CO}_2$ treatments reflecting future OA hypotheses could be obtained by increasing *in situ* DIC concentrations by an increment (ΔDIC) expected under a given CO_2 emissions scenario. The future DIC estimate ($\Delta\text{DIC} + \text{in situ DIC}$), along with a second parameter from the carbonate system, could then be used to recalculate the carbonate system to estimate treatment $p\text{CO}_2$ levels.

Because the Newport, Oregon, time-series consists only of $p\text{CO}_2$, we first needed to estimate *in situ* DIC. To do so, we estimated total alkalinity (TA) from salinity measurements using a linear model parameterized with data from the CCE (Gray et al., 2011). The relationship has relatively low residual error (approximately $\pm 20 \mu\text{mol kg}^{-1}$) and was shown previously to adequately predict TA for the purposes of estimating the carbonate system when using $p\text{CO}_2$ as the second parameter (Harris et al., 2013). We used the estimated TA and *in situ* $p\text{CO}_2$, salinity, and temperature measurements to solve the carbonate system and calculate *in situ* DIC. To estimate ΔDIC , we solved the carbonate system based on estimates of TA, and *in situ* temperature, and salinity, but assuming seawater equilibrium with average atmospheric $p\text{CO}_2$ levels during the period of the moored observations ($\sim 390 \mu\text{atm}$; Harris et al., 2013) and those predicted under an emissions scenario for the year 2100 (788 μatm ; Intergovernmental Panel on Climate Change IS92a “continually increasing” emissions scenario). The present-day air–sea equilibrium DIC estimate was subtracted from the future equilibrium estimate to obtain ΔDIC . Although the same present day and future $p\text{CO}_2$ levels were used to calculate ΔDIC for all samples, ΔDIC values differ among samples because estimates of DIC at air–sea $p\text{CO}_2$ equilibrium vary based on the TA, temperature, and salinity of the sample. Across all water samples, ΔDIC spanned ~ 75 – $105 \mu\text{mol kg}^{-1}$ depending on the TA and temperature of samples. We then obtained future $p\text{CO}_2$ estimates by solving the carbonate system using the estimated TA and *in situ* DIC + ΔDIC values (Figure 6b). Versions of the method have been described previously (e.g. Barry et al., 2010) and can be used to estimate preindustrial carbonate chemistry conditions (e.g. Feely et al., 2008; Harris et al., 2013).

Under this approach, and assuming the same number of treatments is used as depicted in Figure 6a, the effects of temperature and $p\text{CO}_2$ can no longer be separated because orthogonality in the design is lost (Figure 6b). However, a more realistic set of control treatments are included that offer a firmer basis for drawing

Table 1. Brief summary of experimental outcomes of reviewed OA studies that focus on organisms (or their broodstock) collected from the CCE.

Study number	Reference	Collection site	Species (common name)	Life stage	Duration (days, hours)	T (°C)	Control pCO ₂ (µatm)	Treatment pCO ₂ (µatm)	Outcome relative to control
1	Gooding et al. (2009)	Jaricho Beach, BC	<i>Pisaster ochraceus</i> (purple sea star)	Juvenile	70 d	12, 15	380	780	Increased growth rate, reduced calcified mass; feeding and growth increased with temperature
2	Reuter et al. (2011)	Barkley Sound, BC	<i>Strongylocentrotus franciscanus</i> (red sea urchin)	Sperm/eggs	1 h	10.2	400	800, 1800	Decreased range of sperm concentrations over which high fertilization success was likely
3	Nienhuis et al. (2010)	Barkley Sound, BC	<i>Nucella lamellose</i> (Frieded dogwinkle)	Adults	6 d	9.0	380	780, 1585	Enhanced shell dissolution
4	Sunday et al. (2011)	Barkley Sound, BC	<i>Strongylocentrotus franciscanus</i> (red sea urchin)	Larvae	1 d	12.0	400	1000	Reduced larval size, large variation among families
4	Sunday et al. (2011)	Barkley Sound, BC	<i>Mytilus trossulus</i> (bay mussel)	Larvae	1 d	12.0	400	1000	Reduced larval size, small variation among families
5	Crim et al. (2011)	Barkley Sound, BC	<i>Haliotis kamtschatkana</i> (northern abalone)	Larvae	8 d	12.0	400	800, 1800	Reduced survival, increased abnormalities in shell structure, reduced size in normal-shelled larvae
6	Chan et al. (2011)	Orcas Island, WA	<i>Dendraster excentricus</i> (Pacific sand dollar)	Larvae	10 d	20.0	380	1000	Reduced body size and stomachs; no effect on swimming speed
7	Timmins-Schiffman et al. (2012)	San Juan Island, WA	<i>Crassostrea gigas</i> (Pacific oyster)	Larvae	3 d	20.0	468	847, 1065	Increased calcification at day 1, but smaller average size on day 3
8	O'Donnell et al. (2013)	San Juan Island, WA	<i>Mytilus trossulus</i> (bay mussel)	Adults	20 d	10.4	432	299, 575, 736, 980, 1057, 1180, 1322, 1498	Weaker, less flexible byssal threads at treatments above 1200 µatm
9	Hurst et al. (2012)	Port Townsend, WA	<i>Theragra chalcogramma</i> (walleye pollock)	Yearlings	42 d	8.8	414	478, 815, 1805	No effect on growth but increase in otolith deposition rate, biological implication is unclear
9	Hurst et al. (2012)	Port Townsend, WA	<i>Theragra chalcogramma</i> (walleye pollock)	Sub-yearlings warm	196 d	8.3	596	828, 1285, 2894	Increase growth at pCO ₂ higher than 900 µatm; no change in condition factor
9	Hurst et al. (2012)	Port Townsend, WA	<i>Theragra chalcogramma</i> (walleye pollock)	Sub-yearlings cool	196 d	2.4	386	225, 643, 1543	No difference in growth or condition factor across treatments
10	Hurst et al. (2013)	Port Townsend, WA	<i>Theragra chalcogramma</i> (walleye pollock)	Eggs/larvae	35 d	8.1	442	296, 871, 1844	No difference in egg hatch rate, size, or survival, but longer time till hatching; authors think elevated pCO ₂ treatments have minor effect
11	Evans et al. (2013)	Fogarty Creek, OR	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	96 h	12.8	435	813, 1255	Transcriptome up-regulation of genes related to calcification at intermediate pCO ₂ , no change at high level
12	Pespeni et al. (2013)	OR to CA., various locations	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	17 d	14.1	428	897	Reduction in larval body length, change in allele frequency, no change in timing of settlement or competence to metamorphose

Continued

Table 1. Continued

Study number	Reference	Collection site	Species (common name)	Life stage	Duration (days, hours)	T (°C)	Control pCO ₂ (µatm)	Treatment pCO ₂ (µatm)	Outcome relative to control
13	LaVigne et al. (2012)	OR to CA, various locations	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	50 d	14.1	490	1001	No change Sr/Mg composition in spines; though a difference was observed for a Santa Barbara population of urchins
14	Kelly et al. (2013)	Northern and southern CA, various locations	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	5 d	13.0	424	1210	Reduced size in larvae from dams and sires taken from northern and southern locations; no difference in metabolism
15	Place and Smith (2012)	Bodega Bay, CA	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Embryos	7 d	15.1	378	1486, 4110	No disruption to cell cycle in fertilized eggs
16	Hettinger et al. (2012)	Tomales Bay, CA	<i>Ostreola lurida</i> (Olympia oyster)	Larvae/ juveniles	45 d	20.0	739	933, 1355	Reduced larval and juvenile growth
17	Hettinger et al. (2013a)	Tomales Bay, CA	<i>Ostreola lurida</i> (Olympia oyster)	Larvae	22 d	19.4	520	1075	Reduced number of settlers; no difference in shell size or larval dry weight
18	Hettinger et al. (2013b)	Tomales Bay, CA	<i>Ostreola lurida</i> (Olympia oyster)	Larvae/ juveniles	127 d	20.0	485	1060	Reduced larval survival and growth; reduced growth in juveniles outplanted to an estuary
19	Gaylord et al. (2011)	Tomales Bay, CA	<i>Mytilus californianus</i> (California mussel)	Larvae	8 d	15.4	380	540, 970	Thinner, weaker shells
20	Padilla-Gamiño et al. (2013)	Santa Barbara, CA	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	75 h	13, 18	400	1100	Reduced body size and respiration rate; difference in transcriptome observed
21	Matson et al. (2012)	Santa Barbara, CA	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	6 d	15.6	365	1038, 1444	Reduced arm length; no difference in utilization of energy lipid reserves and protein content remained unchanged
22	Yu et al. (2011)	Santa Barbara, CA	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	6 d	15.6	372	1057, 1469	Slight reduction in size
22	Yu et al. (2011)	Santa Barbara, CA	<i>Strongylocentrotus purpuratus</i> (purple sea urchin)	Larvae	6 d	15.6	432	1441	Slight reduction in size




Figure 5. Environmental $p\text{CO}_2$ and temperature measurements (top 50 m) from the northern and central CCE and conditions maintained in OA experiments performed on organisms from the region. Dark grey circles correspond to environmental measurements from Puget Sound, Washington; light grey circles correspond to environmental measurements from all other regions. $p\text{CO}_2$ treatment levels included in an individual experiment at a given temperature are connected by solid black lines. Numbers denote study code (see Table 1). A convex hull (solid grey line) demarcating the extent of all environmental $p\text{CO}_2$ and temperature measurements is depicted to aid visual comparisons. For reference, approximate present-day atmospheric $p\text{CO}_2$ levels ($\sim 390 \mu\text{atm}$) are indicated by the dashed horizontal line.




Figure 6. Schematic of potential experimental approaches to evaluate OA effects, given co-variation between $p\text{CO}_2$ and temperature. To illustrate the benefits and drawback of each approach, *in situ* $p\text{CO}_2$ and temperature measurements from the N10 mooring near Newport, Oregon, during summer upwelling season (2008) are depicted (grey, filled squares; bars indicate standard deviation). (a) A conventional temperature (three levels: 8, 12, and 16°C) by $p\text{CO}_2$ experimental design in which control $p\text{CO}_2$ values are based on approximate present-day global average surface ocean $p\text{CO}_2$ levels and the acidified treatments corresponding to IPCC emissions scenario IS92a projections for year 2100 (390 and 788 μatm ; open square and circle symbols, respectively). Arrows indicate statistical comparisons permitted by the design. (b) Experimental design informed by *in situ* $p\text{CO}_2$ and temperature measurements. Under this design, three controls are included to account for natural co-variation in temperature and $p\text{CO}_2$. Treatment levels that more closely correspond to an OA hypothesis were obtained by specifying an increase in DIC attributed to anthropogenic CO_2 emissions (see Discussion for details). The future DIC estimate and estimates of TA were used to recalculate the carbonate system to obtain target $p\text{CO}_2$ treatment levels. We calculated $p\text{CO}_2$ using the R library “seacarb” (Lavigne and Gattuso, 2010) with dissociation constants from Lueker *et al.* (2000).

inferences about future OA impacts at a given temperature. The experimental design could be improved further by using O_2 concentrations that currently occur at the three different $p\text{CO}_2$ -temperature controls.

The method we use to estimate future $p\text{CO}_2$ requires several important assumptions. First, the approach implicitly assumes that TA, salinity, and temperature will remain unchanged and that future difference in DIC between the observed *in situ* values and

those calculated assuming present-day air–sea equilibrium will remain the same. Further, to estimate ΔDIC , we assumed that all water properties measured at the time of sampling are the same as when the water mass last approached air–sea equilibrium. If waters were cooler at that time, this would result in a slight overestimate of ΔDIC of $\sim 1.6 \mu\text{mol kg}^{-1}$ for each degree Celsius. Following earlier studies, we also assumed that waters upwelled at the mooring location possessed an anthropogenic CO_2 burden that approximated present-day atmospheric $p\text{CO}_2$ conditions (Harris et al., 2013). In other locations within the CCE, such assumptions may not be justified because subsurface waters upwelled on to the shelf may last have had contact with the surface decades prior and therefore would contain less anthropogenic CO_2 (Feely et al., 2008; Harris et al., 2013).

The OA $p\text{CO}_2$ –temperature relationship depicted in Figure 6b corresponds to one simple hypothesis for how present-day carbonate chemistry conditions might change with a simple augmentation of CO_2 . However, marine organisms will be subjected to multiple stressors in the future, including warmer temperatures, lower ambient O_2 concentrations, cultural eutrophication, and pollution (Boyd, 2011; Gruber, 2011; Doney et al., 2012). Treatments corresponding to an OA + warming hypothesis could be created by adding to *in situ* measurements of both temperature and DIC and recalculating the carbonate system to obtain appropriate estimates of $p\text{CO}_2$ (Melzner et al., 2012). If a hypoxia + OA treatment is sought, any assumed reduction in O_2 owing to aerobic respiration necessarily corresponds to an increase in CO_2 (Melzner et al., 2012; Sunda and Cai, 2012). The corresponding increase in DIC beyond that attributed to OA could be estimated based on the molar ratio of O_2 consumed to CO_2 released in the respiratory consumption of organic matter (Sunda and Cai, 2012). We caution, however, that the appropriateness of these simple methods for estimating treatment levels should be thoughtfully considered in the light of the physical and biological attributes of the system under study. Such simplifications do not take into account the indirect and cascading impacts that changes in individual properties such as temperature will have on ecosystem metabolism and thus the distribution and concentration of O_2 and CO_2 (e.g. Keeling et al., 2010; Gruber, 2011; Doney et al., 2012), nor would they reflect potential changes in large-scale circulation patterns or productivity regimes which might fundamentally alter relationships between $p\text{CO}_2$ and other variables (e.g. Rykaczewski and Dunne, 2010). Despite these uncertainties, developing experimental designs that include controls that reflect present-day $p\text{CO}_2$ –temperature– O_2 relationships in upwelling systems should become the cornerstone of experiments that aim to quantify the potential response of organism to future predicted changes in their environment. The development of experimental systems that permit simultaneous control over $p\text{CO}_2$, temperature, and O_2 conditions remains a technical challenge, but a growing number of OA research facilities are acquiring the capacity to do so (Bockmon et al., 2013).

Our compilation of environmental carbonate chemistry data is meant to offer an initial overview of the ranges of $p\text{CO}_2$, temperature, and O_2 in the CCE and the extent to which their relationships vary between seasons and regions. We examined patterns of co-variation between $p\text{CO}_2$ and temperature at coarse seasonal time-scales, but note that co-variation patterns may differ depending on temporal scale. On diel time-scales, solar heating and photosynthesis and respiration may drive strong cyclical patterns in temperature, $p\text{CO}_2$, and O_2 (Barton et al., 2012; Frieder et al., 2012; Waldbusser and Salisbury, 2013), while abrupt events such as

storms or the advection of a different water mass into a region can result in rapid change in water characteristics that may persist for several days (Frieder et al., 2012). Over interannual time-scales, large-scale climate phenomena like *El Niño*/Southern Oscillation can influence upwelling patterns, coastal productivity, carbonate chemistry (Chavez et al., 1999; Friederich et al., 2002), and thus potentially relationships among $p\text{CO}_2$ and temperature and O_2 . To date, studies evaluating patterns of co-variation between $p\text{CO}_2$ and other biologically relevant variables over a range of temporal scales are sparse, but the topic is an area of research we are currently exploring. Incorporating both temporal dynamics and multiple stressors in experimental systems is technically challenging, but may be necessary for ecologically relevant predictions.

Conclusions

OA is expected to have far-reaching impacts on the structure and function of marine ecosystems by altering biogeochemical processes and the productivity and distribution of species (Fabry et al., 2008; Doney et al., 2009, 2012; Mora et al., 2013; Waldbusser and Salisbury, 2013). Our ability to predict the response of complex ecological systems to OA, however, remains limited and is highly constrained by major uncertainties in the response of species to both direct (e.g. physiology, neurological impairment) and indirect (e.g. trophic interactions) processes that may be vulnerable to OA (Fabry et al., 2008; Hofmann et al., 2010). While authors have noted the need for OA researchers to use $p\text{CO}_2$ levels that correspond to ambient conditions a study species or life history stage is likely to experience (Andersson and Mackenzie, 2012; McElhany and Busch, 2012), patterns of co-variation with temperature and O_2 have yet to be incorporated into OA experimental designs (Reum et al., 2014). As demonstrated here, this issue should be of concern to researchers in upwelling systems and other coastal environments where water conditions are highly dynamic over a range of spatial and temporal scales and where co-variation between $p\text{CO}_2$, temperature, and O_2 are generally expected. Because inferences on the potential response of organisms to future conditions are necessarily premised on the notion that experimental controls reflect present-day conditions, we strongly recommend researchers consider how $p\text{CO}_2$ naturally varies with other biologically important variables in their experimental designs. Importantly, the studies we reviewed for the CCE indicate that several species may be sensitive to carbonate chemistry conditions that already occur, and suggest that present-day variability in carbonate chemistry may be more important to contemporary ecological patterns than previously thought. With the continued collection of high-quality carbonate chemistry measurements and their archival on freely accessible databases, analyses like the one we present here for the CCE may yield further insight into the relevance of carbonate chemistry variability to contemporary ecological processes as well as guide OA experimental design in other marine systems.

Supplementary data

Supplementary material is available at the ICESJMS online version of the manuscript.

Acknowledgements

The authors wish to thank many research technical staff in the laboratories that contributed data used in this manuscript, particularly the scientific and engineering staff in NOAA's Pacific Marine Environmental Laboratory (PMEL) carbon, CTD- O_2 , and engineering groups; UW's Applied Physics Laboratory (UW/APL) and

School of Oceanography; OSU's College of Earth, Ocean, and Atmospheric Sciences Ocean Ecology and Biogeochemistry division; and the Scripps Institution of Oceanography Ocean Time Series Group for the tremendous effort involved in making these datasets available. In addition, we thank the officers and crew on all spatial surveys, underway observing ships, and mooring deployments for their able assistance during field operations. Funding for JCPR was provided through a National Research Council Fellowship. Funding for NOAA/PMEL contributions to the mooring time-series, underway $p\text{CO}_2$, and spatial survey data used in these analyses were provided by NOAA's Climate Programme Office's Global Carbon Cycle Programme (grants GC05-288 and GC10-102), NOAA's Ocean Acidification Programme, and NOAA/PMEL. The Chá bã mooring off La Push, Washington, was also supported by the Murdock Charitable Trust, NANOOS (Northwest Association of Networked Ocean Observing Systems, a Regional Association of NOAA's US Integrated Ocean Observing System Programme), UW/APL, UW College of the Environment, and UW Provost's Office. The PRISM cruises were supported by the State of Washington through UW Oceanography and APL. The NH10 mooring is funded by NOAA through NANOOS, and by NSF through CMOP (Science and Technology Center for Coastal Margin Observation and Prediction OCE-0424602). Surface ocean $p\text{CO}_2$ data collected on the NH10 mooring were supported by NSF Chemical Oceanography OCE-0752576 awarded to BH and P.G. Strutton. I. Kaplan, T. Klinger, and three anonymous reviewers provided valuable comments on early versions of the manuscript. This is PMEL contribution number 4100.

References

- Andersson, A. J., and Mackenzie, F. T. 2012. Revisiting four scientific debates in ocean acidification research. *Biogeosciences*, 9: 893–905.
- Barry, J. P., Tyrrell, T., Hansson, L., Plattner, G., and Gattuso, J. 2010. Atmospheric CO_2 targets for ocean acidification perturbation experiments. *In* Guide to Best Practices for Ocean Acidification Research and Data Reporting, pp. 53–66. Ed. by U. Riebesell, V. J. Fabry, L. Hansson, and J.-P. Gattuso. Publications Office of the European Union, Luxembourg.
- Bakker, D. C. E., Pfeil, B., Smith, K., Hankin, S., Olsen, A., Alin, S. R., Cosca, C., *et al.* 2013. An update to the Surface Ocean CO_2 Atlas (SOCAT version 2). *Earth System Science Data*, 6: 1–22, 6, doi:10.5194/essd-6-69-2014, 2014
- Barton, A., Hales, B., Waldbusser, G. G., Langdon, C., and Feely, R. A. 2012. The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: implications for near-term ocean acidification effects. *Limnology and Oceanography*, 57: 698–710.
- Bockmon, E., Frieder, C., Navarro, M., White-Kershek, L., and Dickson, A. 2013. Controlled experimental aquarium system for multi-stressor investigation: carbonate chemistry, oxygen saturation, and temperature. *Biogeosciences*, 10: 5967–5975.
- Borges, A. V., and Abril, G. 2011. Carbon dioxide and methane dynamics in estuaries. *In* Treatise on Estuarine and Coastal Science, 5, pp. 119–161. Ed. by D. McClusky, and E. Wolanski. Academic Press, Waltham.
- Boyd, P. W. 2011. Beyond ocean acidification. *Nature Geoscience*, 4: 273–274.
- Chan, K. Y. K., Grünbaum, D., and O'Donnell, M. J. 2011. Effects of ocean-acidification-induced morphological changes on larval swimming and feeding. *The Journal of Experimental Biology*, 214: 3857–3867.
- Chavez, F. P., Strutton, P. G., Friederich, C. E., Feely, R. A., Feldman, G. C., Foley, D. C., and McPhaden, M. J. 1999. Biological and chemical response of the equatorial Pacific Ocean to the 1997–98 El Niño. *Science*, 286: 2126–2131.
- Checkley, D. M., and Barth, J. A. 2009. Patterns and processes in the California Current System. *Progress in Oceanography*, 83: 49–64.
- Crim, R. N., Sunday, J. M., and Harley, C. D. 2011. Elevated seawater CO_2 concentrations impair larval development and reduce larval survival in endangered northern abalone (*Haliotis kamtschatkana*). *Journal of Experimental Marine Biology and Ecology*, 400: 272–277.
- Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A. 2009. Ocean acidification: the other CO_2 problem. *Annual Review of Marine Science*, 1: 169–192.
- Doney, S. C., Ruckelshaus, M., Emmett Duffy, J., Barry, J. P., Chan, F., English, C. A., Galindo, H. M., *et al.* 2012. Climate change impacts on marine ecosystems. *Annual Review of Marine Science*, 4: 11–37.
- Dupont, S., Ortega-Martínez, O., and Thorndyke, M. 2010. Impact of near-future ocean acidification on echinoderms. *Ecotoxicology*, 19: 449–462.
- Evans, T. G., Chan, F., Menge, B. A., and Hofmann, G. E. 2013. Transcriptomic responses to ocean acidification in larval sea urchins from a naturally variable pH environment. *Molecular Ecology*, 22: 1609–1625.
- Evans, W., Hales, B., and Strutton, P. G. 2011. Seasonal cycle of surface ocean $p\text{CO}_2$ on the Oregon shelf. *Journal of Geophysical Research*, 116: C05012.
- Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C. 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES Journal of Marine Science*, 65: 414–432.
- Fassbender, A. J., Sabine, C. L., Feely, R. A., Langdon, C., and Mordy, C. W. 2011. Inorganic carbon dynamics during northern California coastal upwelling. *Continental Shelf Research*, 31: 1180–1192.
- Feely, R. A., Alin, S. R., Newton, J., Sabine, C. L., Warner, M., Devol, A., Krembs, C., *et al.* 2010. The combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized estuary. *Estuarine Coastal and Shelf Science*, 88: 442–449.
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B. 2008. Evidence for upwelling of corrosive “acidified” water onto the continental shelf. *Science*, 320: 1490–1492.
- Fréon, P., Barange, M., and Aristegui, J. 2009. Eastern boundary upwelling ecosystems: integrative and comparative approaches. *Progress in Oceanography*, 83: 1–14.
- Frieder, C. A., Nam, S. H., Martz, T. R., and Levin, L. A. 2012. High temporal and spatial variability of dissolved oxygen and pH in a near-shore California kelp forest. *Biogeosciences*, 9: 3917–3930.
- Friederich, G., Walz, P., Burczynski, M., and Chavez, F. 2002. Inorganic carbon in the central California upwelling system during the 1997–1999 El Niño–La Niña event. *Progress in Oceanography*, 54: 185–203.
- Gaylord, B., Hill, T. M., Sanford, E., Lenz, E. A., Jacobs, L. A., Sato, K. N., Russell, A. D., *et al.* 2011. Functional impacts of ocean acidification in an ecologically critical foundation species. *The Journal of Experimental Biology*, 214: 2586–2594.
- Gooding, R. A., Harley, C. D., and Tang, E. 2009. Elevated water temperature and carbon dioxide concentration increase the growth of a keystone echinoderm. *Proceedings of the National Academy of Sciences of the United States of America*, 106: 9316–9321.
- Gray, S. E., DeGrandpre, M. D., Moore, T. S., Martz, T. R., Friederich, G. E., and Johnson, K. S. 2011. Applications of in situ pH measurements for inorganic carbon calculations. *Marine Chemistry*, 125: 82–90.
- Gruber, N. 2011. Warming up, turning sour, losing breath: ocean biogeochemistry under global change. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369: 1980–1996.
- Gruber, N., Hauri, C., Lachkar, Z., Loher, D., Frölicher, T. L., and Plattner, G.-K. 2012. Rapid progression of ocean acidification in the California Current System. *Science*, 337: 220–223.

- Hales, B., Karp-Boss, L., Perlin, A., and Wheeler, P. A. 2006. Oxygen production and carbon sequestration in an upwelling coastal margin. *Global Biogeochemical Cycles*, 20: GB002517.
- Hales, B., Takahashi, T., and Bandstra, L. 2005. Atmospheric CO₂ uptake by a coastal upwelling system. *Global Biogeochemical Cycles*, 19: GB1009.
- Harris, K. E., DeGrandpre, M. D., and Hales, B. 2013. Aragonite saturation state dynamics in a coastal upwelling zone. *Geophysical Research Letters*, 40: 2720–2725.
- Harvey, B. P., Gwynn-Jones, D., and Moore, P. J. 2013. Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming. *Ecology and Evolution*, 3: 1016–1030.
- Hauri, C., Gruber, N., Plattner, G.-K., Alin, S., Feely, R. A., Hales, B., and Wheeler, P. A. 2009. Ocean acidification in the California current system. *Oceanography*, 22: 60–77.
- Hauri, C., Gruber, N., Vogt, M., Doney, S. C., Feely, R. A., Lachkar, Z., Leinweber, A., et al. 2013. Spatiotemporal variability and long-term trends of ocean acidification in the California Current System. *Biogeosciences*, 10: 193–216.
- Hettinger, A., Sanford, E., Hill, T. M., Hofsfelt, J. D., Russell, A. D., and Gaylord, B. 2013a. The influence of food supply on the response of Olympia oyster larvae to ocean acidification. *Biogeosciences*, 10: 6629–6638.
- Hettinger, A., Sanford, E., Hill, T. M., Lenz, E. A., Russell, A. D., and Gaylord, B. 2013b. Larval carry-over effects from ocean acidification persist in the natural environment. *Global Change Biology*, 19: 3317–3326.
- Hettinger, A., Sanford, E., Hill, T. M., Russell, A. D., Sato, K. N., Hoey, J., Forsch, M., et al. 2012. Persistent carry-over effects of planktonic exposure to ocean acidification in the Olympia oyster. *Ecology*, 93: 2758–2768.
- Hickey, B. M., and Banas, N. S. 2003. Oceanography of the US Pacific Northwest Coastal Ocean and estuaries with application to coastal ecology. *Estuaries*, 26: 1010–1031.
- Hofmann, G. E., Barry, J. P., Edmunds, P. J., Gates, R. D., Hutchins, D. A., Klinger, T., and Sewell, M. A. 2010. The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective. *Annual Review of Ecology, Evolution, and Systematics*, 41: 127–147.
- Hofmann, G. E., Evans, T. G., Kelly, M. W., Padilla-Gamiño, J. L., Blanchette, C. A., Washburn, L., Chan, F., et al. 2014. Exploring local adaptation and the ocean acidification seascape—studies in the California Current Large Marine Ecosystem. *Biogeosciences*, 11: 1053–1064.
- Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F., Paytan, A., et al. 2011. High-frequency dynamics of ocean pH: a multi-ecosystem comparison. *PLoS ONE*, 6: e28983.
- Hurst, T. P., Fernandez, E. R., and Mathis, J. T. 2013. Effects of ocean acidification on hatch size and larval growth of walleye pollock (*Theragra chalcogramma*). *ICES Journal of Marine Science*, 70: 812–822.
- Hurst, T. P., Fernandez, E. R., Mathis, J. T., Miller, J. A., Stinson, C. M., and Ahgeak, E. F. 2012. Resiliency of juvenile walleye pollock to projected levels of ocean acidification. *Aquatic Biology*, 17: 247–259.
- Keeling, R. F., Körtzinger, A., and Gruber, N. 2010. Ocean deoxygenation in a warming world. *Annual Review of Marine Science*, 2: 199–229.
- Kelly, M. W., Padilla-Gamiño, J. L., and Hofmann, G. E. 2013. Natural variation and the capacity to adapt to ocean acidification in the keystone sea urchin *Strongylocentrotus purpuratus*. *Global Change Biology*, 19: 2536–2546.
- Koch, M., Bowes, G., Ross, C., and Zhang, X. H. 2013. Climate change and ocean acidification effects on seagrasses and marine macroalgae. *Global Change Biology*, 19: 103–132.
- Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M., et al. 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biology*, 19: 1884–1896.
- LaVigne, M., Hill, T. M., Sanford, E., Gaylord, B., Russell, A. D., Lenz, E. A., and Hofsfelt, J. D. 2012. The elemental composition of purple sea urchin (*Strongylocentrotus purpuratus*) calcite and potential effects of pCO₂ during early life stages. *Biogeosciences*, 10: 3465–3477.
- Lavigne, H., and Gattuso, J. 2010. Seacarb: seawater carbonate chemistry with R. R package version 2.4.3.
- Lueker, T. J., Dickson, A. G., and Keeling, C. D. 2000. Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry*, 70: 105–119.
- Matson, P. G., Pauline, C. Y., Sewell, M. A., and Hofmann, G. E. 2012. Development under elevated pCO₂ conditions does not affect lipid utilization and protein content in early life-history stages of the purple sea urchin, *Strongylocentrotus purpuratus*. *The Biological Bulletin*, 223: 312–327.
- McElhany, P., and Busch, D. S. 2012. Appropriate pCO₂ treatments in ocean acidification experiments. *Marine Biology*, 160: 1807–1812.
- Melzner, F., Thomsen, J., Koeve, W., Oschlies, A., Gutowska, M. A., Bange, H. W., Hansen, H. P., et al. 2012. Future ocean acidification will be amplified by hypoxia in coastal habitats. *Marine Biology*, 160: 1875–1888.
- Mora, C., Wei, C.-L., Rollo, A., Amaro, T., Baco, A. R., Billett, D., Bopp, L., et al. 2013. Biotic and Human vulnerability to projected changes in ocean biogeochemistry over the 21st century. *PLoS Biology*, 11: e1001682.
- Nienhuis, S., Palmer, A. R., and Harley, C. D. G. 2010. Elevated CO₂ affects shell dissolution rate but not calcification rate in a marine snail. *Proceedings of the Royal Society B: Biological Sciences*, 277: 2553–2558.
- Orr, J. 2011. Recent and future changes in ocean carbonate chemistry. *In* Ocean Acidification. Ed. by J. Gattuso, and L. Hansson. Oxford University Press, New York, NY.
- O'Donnell, M. J., George, M. N., and Carrington, E. 2013. Mussel byssus attachment weakened by ocean acidification. *Nature Climate Change*, 3: 587–590.
- Padilla-Gamiño, J. L., Kelly, M. W., Evans, T. G., and Hofmann, G. E. 2013. Temperature and CO₂ additively regulate physiology, morphology and genomic responses of larval sea urchins, *Strongylocentrotus purpuratus*. *Proceedings of the Royal Society B: Biological Sciences*, 280: 20130155.
- Pespeni, M. H., Sanford, E., Gaylord, B., Tessa, M. H., Hofsfelt, J. D., Jaris, H. K., LaVigne, M., et al. 2013. Evolutionary change during experimental ocean acidification. *Proceedings of the National Academy of Sciences of the United States of America*, 110: 6937–6942.
- Place, S. P., and Smith, B. W. 2012. Effects of seawater acidification on cell cycle control mechanisms in *Strongylocentrotus purpuratus* embryos. *PLoS ONE*, 7: e34068.
- Pörtner, H. O. 2010. Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. *The Journal of Experimental Biology*, 213: 881–893.
- Pörtner, H. O. 2012. Integrating climate-related stressor effects on marine organisms: unifying principles linking molecule to ecosystem-level changes. *Marine Ecology Progress Series*, 470: 273–290.
- Pörtner, H. O., and Farrell, A. P. 2008. Physiology and climate change. *Science*, 322: 690–692.
- R Development Core Team. 2011. R: a Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Reum, J. C., Alin, S. R., Feely, R. A., Newton, J., Warner, M., and McElhany, P. 2014. Seasonal carbonate chemistry covariation with temperature, oxygen, and salinity in a fjord estuary: implications

- for the design of ocean acidification experiments. *PLoS ONE*, 9: e89619.
- Reuter, K. E., Lotterhos, K. E., Crim, R. N., Thompson, C. A., and Harley, C. D. 2011. Elevated $p\text{CO}_2$ increases sperm limitation and risk of polyspermy in the red sea urchin *Strongylocentrotus franciscanus*. *Global Change Biology*, 17: 163–171.
- Rykaczewski, R. R., and Dunne, J. P. 2010. Enhanced nutrient supply to the California Current Ecosystem with global warming and increased stratification in an earth system model. *Geophysical Research Letters*, 37: GL045019.
- Shaw, E. C., McNeil, B. I., Tilbrook, B., Matear, R., and Bates, M. L. 2013. Anthropogenic changes to seawater buffer capacity combined with natural reef metabolism induce extreme future coral reef CO_2 conditions. *Global Change Biology*, 19: 1632–1641.
- Sunda, W. G., and Cai, W. J. 2012. Eutrophication induced CO_2 -acidification of subsurface coastal waters: interactive effects of temperature, salinity, and atmospheric $p\text{CO}_2$. *Environmental Science and Technology*, 46: 10651–10659.
- Sunday, J. M., Crim, R. N., Harley, C. D. G., and Hart, M. W. 2011. Quantifying rates of evolutionary adaptation in response to ocean acidification. *PLoS ONE*, 6: e22881. doi:10.1371/journal.pone.0022881.
- Timmins-Schiffman, E., O'donnell, M. J., Friedman, C. S., and Roberts, S. B. 2012. Elevated $p\text{CO}_2$ causes developmental delay in early larval Pacific oysters, *Crassostrea gigas*. *Marine Biology*, 160: 1973–1982.
- van Geen, A., Takesue, R. K., Goddard, J., Takahashi, T., Barth, J. A., and Smith, R. L. 2000. Carbon and nutrient dynamics during coastal upwelling off Cape Blanco, Oregon. *Deep Sea Research*, 47: 975–1002.
- Waldbusser, G. G., and Salisbury, J. E. 2013. Ocean acidification in the coastal zone from an organism's perspective: multiple system parameters, frequency domains, and habitats. *Annual Review of Marine Science*, 6: 221–247.
- Wernberg, T., Smale, D. A., and Thomsen, M. S. 2012. A decade of climate change experiments on marine organisms: procedures, patterns and problems. *Global Change Biology*, 18: 1491–1498.
- Wittmann, A. C., and Pörtner, H-O. 2013. Sensitivities of extant animal taxa to ocean acidification. *Nature Climate Change*, 3: 995–1001.
- Yu, P. C., Matson, P. G., Martz, T. R., and Hofmann, G. E. 2011. The ocean acidification seascape and its relationship to the performance of calcifying marine invertebrates: Laboratory experiments on the development of urchin larvae framed by environmentally-relevant $p\text{CO}_2/\text{pH}$. *Journal of Experimental Marine Biology and Ecology*, 400: 288–295.

Handling editor: Howard Browman