

Motor skills, calibrated severity & autism

The relationship of motor skills and adaptive behavior skills in young children with autism spectrum disorders

Corresponding Author:

Megan MacDonald, PhD
Oregon State University
School of Biological & Population Health Sciences
202 Women's Building
Corvallis, OR
97331
541-737-3273
megan.macdonald@oregonstate.edu

Catherine Lord, PhD
Weill Cornell Medical College
New York Presbyterian Hospital
Center for Autism and the Developing Brain
21 Bloomingdale Rd.
White Plains, NY
10605
914-997-5848
cal2028@med.cornell.edu

Dale Ulrich, PhD
University of Michigan
School of Kinesiology
1402 Washington Heights
Ann Arbor, MI
734-615-1904
ulrichd@umich.edu

Author Note: Dr. Lord receives royalties from Western Psychological Service for the sale of diagnostic instruments which she has co-authored. Dr. Lord donates all royalties from these projects and clinics to not-for-profit autism charities, specifically Have Dreams which is located in Chicago, Illinois. Have Dreams offers wraparound services for children and adults with autism. Drs. MacDonald & Ulrich have no disclosures.

Support for this project was provided in part from funding awarded to Dr. Lord from the Simons Foundation, First Words and the following grants: NICHD U19 HD35482-01. The Neurobiology and Genetic of Autism. 06/01/97-05/31/07 (Lord). NIMH RO1 MH081873-01A1. Longitudinal Studies of Autism Spectrum Disorders: 2 to 23. 09/01/08-05/31/13 (Lord). Blue Cross Blue Shield Foundation of Michigan Grant

Motor skills, calibrated severity & autism

number 1687.SAP (MacDonald).

Motor skills, calibrated severity & autism

Abstract

Objective: To determine the relationship of motor skills and the core behaviors of young children with autism, social affective skills and repetitive behaviors, as indicated through the calibrated autism severity scores.

Design: The univariate GLM tested the relationship of gross and fine motor skills measured by the gross motor scale and the fine motor scale of the MSEL with autism symptomology as measured by calibrated autism severity scores.

Setting: Majority of the data collected took place in an autism clinic.

Participants: A cohort of 159 young children with ASD (n=110), PDD-NOS (n=26) and non-ASD (developmental delay, n=23) between the ages of 12- 33 months were recruited from early intervention studies and clinical referrals. Children with non-ASD (developmental delay) were included in this study to provide a range of scores indicated through calibrated autism severity.

Interventions: Not applicable.

Main Outcome Measures: The primary outcome measures in this study were calibrated autism severity scores.

Results: Fine motor skills and gross motor skills significantly predicted calibrated autism severity ($p < 0.01$). Children with weaker motor skills displayed higher levels of calibrated autism severity.

Conclusions: The fine and gross motor skills are significantly related to autism symptomology. There is more to focus on and new avenues to explore in the realm of discovering how to implement early intervention and rehabilitation for young children with autism and motor skills need to be a part of the discussion.

Motor skills, calibrated severity & autism

Key words: Autism, motor skills, young children, calibrated severity

Corresponding Author:

Megan MacDonald, PhD
Oregon State University
School of Biological & Population Health Sciences
202 Women's Building
Corvallis, OR
97331
541-737-3273
megan.macdonald@oregonstate.edu

1
2
3
4
5
6
7
8
9
10
11
12
13

Abstract

Motor skill deficits are pervasive and persistent in young children with autism spectrum disorder (ASD). Yet the focus of intervention is on core impairments, which are part of the diagnostic criteria for ASD, deficits in social communication skills. The purpose of this study is to determine whether a relationship exists between the motor skills of young children with ASD (14- 49 months) and adaptive behavior skills. Two hundred and thirty three children with ASD between the ages of 14- 49 months participated in this study. A multiple regression analysis tested the relationship of motor skills on the adaptive behavior composite, daily living, adaptive social and adaptive communicative skills holding constant age, non-verbal problem solving, and clinical ASD diagnosis. Fine motor skills significantly predicted all adaptive behavior skills ($p < 0.01$). Gross motor skills were predictive of daily living skills ($p < 0.05$). Children with weaker motor skills displayed greater deficits in adaptive behavior skills.

Motor skills, adaptive behavior skills & ASD

1 Autism spectrum disorder (ASD) is a pervasive developmental disorder characterized by
2 deficits in social communicative skills and repetitive or restricted interests (APA, 1994, 2000,
3 2013). ASD affects 1 in 88 individuals (CDC, 2012) and a recent national survey indicated that
4 ASD affects 1 in 50 school-aged children (Blumberg et al., 2013). In addition to the hallmark
5 characteristics of ASD, social communicative deficits, children also display motor skill deficits
6 (Landa & Garrett-Mayer, 2006; Lloyd, MacDonald, & Lord, 2013; Provost, Heimerl, & Lopez,
7 2007; Provost, Lopez, & Heimerl, 2007; Teitelbaum, Teitelbaum, Nye, Fryman, & Maurer,
8 1998; Yirmiya & Ozonoff, 2007). In fact, in some of the original clinical descriptions of what is
9 now called ASD, Asperger attached considerable weight to motor clumsiness (Frith, 1991).

10 Motor skill deficits appear early in life for many young children with ASD, and become
11 prominent around 14- 24 months of age (Chawarska et al., 2007). Delayed infant motor
12 milestones have been reported and these late milestones (especially walking) have acted as initial
13 developmental concerns to parents, who later received an ASD diagnosis for their child
14 (Chawarska et al., 2007; Flanagan, Landa, Bhat, & Bauman, 2012; Landa & Garrett-Mayer,
15 2006; Lloyd et al., 2013; Teitelbaum, et al., 1998).

16 For the most part the use of motor skill performance in an ASD diagnosis is embedded in
17 gestures, stereotypies and imitation (Lord et al., 2000; Luyster et al., 2009). However it has been
18 suggested that one of the earliest detectable signs and a cardinal feature of the disability may
19 reside in motor skill ability early in life (Flanagan et al., 2012; Fournier, Hass, Naik, Lodha, &
20 Cauraugh, 2010; Sutera, Pandey, Esser, & Rosenthal, 2007; Teittelbaum, Teittelbaum, Nye,
21 Fryman, & Maurer, 1998). When a group of children at high-risk for ASD (children who had a
22 sibling diagnosed with ASD), were studied prospectively three groups emerged for comparison,
23 children with language delay, ASD and typical development (Landa & Garrett-Mayer, 2006).

Motor skills, adaptive behavior skills & ASD

1 Deficits in fine motor skills were evident in children with ASD at 6 months of age and
2 significantly worsened fine and gross motor skills were evident at 14 and 24 months of age
3 (Landa & Garrett-Mayer, 2006). This prospective study suggested early motor skill deficits as a
4 potential diagnostic distinction between children with ASD and children with other
5 developmental delays (language delay) (Landa & Garrett-Mayer, 2006). These findings
6 corroborated previous studies suggesting motor skill delays early in development may act as
7 some of the first signs for developmental concern (Teittelbaum et al., 1998). Through video
8 analysis Teitelbaum et al. (1998) found oral motor deficits as well as delays in motor milestones
9 such as lying, righting, sitting, crawling and walking were present in young children with ASD
10 long before a diagnosis was conclusive.

11 In a large cross-sectional study of young children with ASD (N= 172, aged 14- 36
12 months) fine and gross motor skill deficits became significantly worse within a short
13 chronological timeframe (6- 18 months) (Lloyd et al., 2013). The same study confirmed cross-
14 sectional findings through a longitudinal analysis of children who had two motor skill
15 assessments approximately one-year apart. When the same children were studied over time, they
16 displayed significantly worse motor skill deficits as they aged (Lloyd et al., 2013).

17 Other motor deficits areas have been found in gait and postural control. Children with
18 ASD (aged 4- 6 years) demonstrated a short step length and irregular body oscillations during
19 locomotion consistent with a less stable and more variable posture compared to a control group
20 without a disability (Vernazza-Martin et al., 2005). Young children with ASD (less than 2 years)
21 displayed similar gait and postural control deficits (Esposito & Venuti, 2008). At 20 months of
22 age children with ASD showed deficits in gait parameters which included performing abnormal
23 heel-to-toe patterns, odd arm posturing and generally higher frequencies of anomalies in

Motor skills, adaptive behavior skills & ASD

1 movement including waddle walking (Esposito & Venuti, 2008). Postural sway in school-aged
2 children with ASD was significantly greater than typically developed controls in mediolateral,
3 anteroposterior and normalized sway (Fournier et al., 2010; Memari et al., 2013). Similarly,
4 postural sway in the mediolateral direction was greater in children with autism during dynamic
5 movements such as walking (Fournier, Kimberg, et al., 2010).

6 Consistent with other motor skill deficits at a young age, motor planning deficits were
7 evident when children with ASD were unable translate motor intention into a global motor action
8 (mean age 7.6 years), but rather treated each motor task involved in an overall action as an
9 independent task (Fabbri-Destro, Cattaneo, Boria, & Rizzolatti, 2009). Vernazza-Martin et al.
10 (2005) found that young children with ASD (4- 6 years) had difficulty defining the goal of the
11 motor action. Even when the task was adapted in a highly motivating fashion, children appeared
12 to understand the instruction, marked in their purposeful action of moving toward the object, but
13 the final motor goal could not be completed, indicating shortfalls in motor planning (Vernazza-
14 Martin, et al., 2005). These children were able to perform simple motor tasks but unable to chain
15 multiple motor tasks together into a more complex motor action.

16 Descriptive studies have clearly demonstrated that motor skill deficits exist in children
17 with ASD at a young age and have even gone so far as to suggest early motor skill deficits as a
18 preliminary diagnostic marker and a cardinal characteristic of the disability (Flanagan et al.,
19 2012; Fournier, Hass, et al., 2010; Matson, Mahan, Fodstad, Hess, & Neal, 2010; Teittelbaum et
20 al., 1998). These deficits in young children with autism range in nature and across tasks
21 (Berkeley, Zittel, Pitney, & Nichols, 2001; Rinehart et al., 2006; Staples & Reid, 2009; Sutera et
22 al., 2007; Teittelbaum et al., 1998; Vernazza-Martin et al., 2005). Even though evidence suggests
23 motor deficits are prominent and pervasive across time and age very little has been discussed in

Motor skills, adaptive behavior skills & ASD

1 terms of how motor skill deficits interact with the core-characteristics of autism, deficits in social
2 communication skills (Berkeley, et al., 2001; Fournier, Hass et al., 2010; Landa & Garrett-
3 Mayer, 2006; Staples & Reid, 2009). It is well known that deficits in social communicative
4 skills are core diagnostic characteristics of ASD. For young children this includes skills such as
5 imitation, joint attention, general social interactions with others, as well as functional and
6 symbolic play (Kasari et al., 2005).

7 The importance of improving social communicative skills in children with autism early in
8 life has been clearly indicated (Dawson et al., 2010; Helt et al., 2008; Kasari, et al., 2005; Kasari,
9 Gulsrud, Wong, Kwon, & Locke, 2010; Kasari, Paparella, Freeman, & Jahromi, 2008; Kelley,
10 Naigles, & Fein, 2010; Wong & Kwan, 2010). The purpose of this study is to test the
11 relationship of fine and gross motor skills on the behavior composite, daily living, adaptive
12 social and communicative skills of children with autism. It is hypothesized that children with
13 better motor skills will have better adaptive behavior skills.

14 **Method**

15 **Participants**

16 The Institutional Review Board approved all methods and procedures for this study.
17 Young children with ASD, pervasive developmental disorder- not otherwise specified (PDD-
18 NOS) and non-ASD (developmental delay) between the ages of 12- 60 months were recruited
19 from early intervention studies and clinical referrals to university-based clinics. Generally,
20 children were recruited through autism support groups, study flyers and clinical referrals from
21 pediatricians and teachers were informed of the study during clinical visits.

1 All participants (N=233) in this study were children between the ages of 14- 49 months
2 (mean age= 30.36 months) with a confirmatory diagnosis of ASD, PDD-NOS or non-ASD (non-
3 ASD developmentally delayed children) based on the DSM-IV(see Table 1).

4 **Measurements**

5 **Child diagnostic measures.** All participants were administered the Autism Diagnostic
6 Observation Schedule (ADOS) (Lord, et al., 2000; Lord, Rutter, DiLavore, & Risi, 1999) or its
7 precursor, the Pre-Linguistic Autism Diagnostic Observation Schedule (DiLavore, Lord, &
8 Rutter, 1995), in order to acquire diagnostic information through direct observation of the
9 children by a trained clinician. The participants at one university-based center were administered
10 the Toddler module of the ADOS (Luyster, et al., 2009). Diagnosis was determined by
11 standardized algorithms established for the Autism Diagnostic Observation Schedule (Gotham,
12 Pickles, & Lord, 2009; Lord, et al., 2000). A clinical psychologist or a trainee who completed
13 research training and met standard requirements for research reliability administered the ADOS
14 (Lord, et al., 1999).

15 Revised algorithms for the ADOS modules 1, 2 and 3 have been published with stronger
16 specificity and sensitivity (Gotham, Risi, Pickles, & Lord, 2007). Standardized scores of
17 calibrated severity are available using raw scores from the revised algorithms of the ADOS
18 (Gotham, et al., 2009). Calibrated autism severity scores have been indicated optimal for
19 comparisons of assessments across time (and age), and to identify different trajectories of autism
20 severity independent of verbal IQ (Gotham, et al., 2009).

21 Each member of the research team, at all centers, established inter-rater reliability
22 exceeding 90% exact agreement ($\kappa > 0.70$) for all items on the ADI-R and 80% exact
23 agreement ($\kappa > 0.60$) on codes for the PL-ADOS, ADOS, and Toddler module for three

Motor skills, adaptive behavior skills & ASD

1 consecutive administrations before the study began. Reliability was maintained over time
2 through consensus coding of approximately every sixth administration with a second rater who
3 was blind to referral status.

4 **Developmental level measurement.** The Mullen Scales of Early Learning (MSEL)
5 (Mullen, 1995) was used to assess cognitive development. This test of development provides
6 reliable and valid information for children from birth to 68 months of age. The subscales on the
7 MSEL are organized into 5 domains: gross motor, fine motor, visual reception (nonverbal
8 problem solving), receptive language, and expressive language. An early learning composite
9 score is derived from the fine motor, visual reception, receptive language and expressive
10 language scales.

11 **Adaptive skills.** The Vineland Adaptive Behavior Scales, 2nd Ed. (VABS) (Sparrow,
12 Cicchetti, & Balla, 2005) were used to assess adaptive skills. The VABS is a standardized
13 parent-report measure of everyday adaptive functioning and yields domain scores in the areas of
14 communication, daily living skills, social skills, and motor development (fine motor and gross
15 motor). A standardized behavior composite score is derived from all domains. This assessment
16 was administered by interview or over the phone, to the parent or primary caregiver of the
17 participant.

18 **Motor skill measurement.** The gross motor and fine motor scales of the MSEL were
19 used to assess motor skills (Mullen, 1995). A gross motor and fine motor age equivalent score
20 was used for analysis as standardized T-scores are not provided for children beyond 33 months
21 of age on the gross motor scale, even though most children with ASD did not meet the ceiling
22 requirements for this scale.

Motor skills, adaptive behavior skills & ASD

1 were conducted for each subscale of interest, this included the standard daily living, adaptive
2 social behavior and adaptive communicative behavior. For each dependent variable the multiple
3 regression analysis included non-verbal problem solving, ethnicity and calibrated autism severity
4 scores. Age was not included as a predictor as age and non-verbal problem solving were
5 collinear. There were no interactions.

6 **Fine motor skills and adaptive behavior composite, daily living, adaptive social**
7 **behavior & adaptive communicative behavior.** A multiple regression analysis tested the
8 relationship of fine motor skills, visual receptive organization (non-verbal problem solving),
9 calibrated autism severity and ethnicity on the adaptive behavior composite, daily living skills,
10 adaptive social behavior and adaptive communicative behavior of young children with autism.
11 For each dependent variable the multiple regression analysis included fine motor skills, non-
12 verbal problem solving, ethnicity and calibrated autism severity. There were no interactions.
13 Fine motor skills and calibrated autism severity were significant predictors of adapted behavior
14 composite ($p < .001$), daily living skills ($p < .001$), adaptive social skills ($p < .05$) and adaptive
15 communicative skills ($p < .001$) (Table 2).

16 **Gross motor skills and adaptive behavior composite, daily living, adaptive social**
17 **behavior & adaptive communicative behavior.** A multiple regression analysis tested the
18 relationship of gross motor skills, visual receptive organization (non-verbal problem solving),
19 calibrated autism severity and ethnicity on the adaptive behavior composite, daily living skills,
20 adaptive social behavior and adaptive communicative behavior of young children with autism.
21 There were no interactions.

Motor skills, adaptive behavior skills & ASD

1 significantly improves behavior in the social communicative domain as well as other aspects of
2 ASD symptomology (Dawson, et al., 2010; Kasari, et al., 2010; Wong & Kwan, 2010). These
3 improvements have included IQ, language, play skills, adaptive behavior and ASD diagnosis (ie.
4 moving from ASD, to PDD-NOS) (Dawson, et al., 2010; Kasari, Freeman, & Paparella, 2006;
5 Kasari, et al., 2008). Although there is widespread agreement on the necessity of early
6 intervention, there is less consistent agreement on early intervention content (Kasari, et al.,
7 2005). A report from the National Research Council (NRC; 2001) indicated that motor skills
8 should be emphasized in specialized early intervention for young children with ASD. Yet, more
9 than a decade later, motor skill content does not appear to be a priority in early intervention
10 models.

11 In a group of school-aged children with ASD fundamental motor skills were
12 significantly related to social communicative skills (MacDonald, et al., 2013). The authors
13 concluded that motor skills needed to be included in the discussion of content for social skills-
14 based programs for school-aged children. It has been previously acknowledged that a potential
15 relationship exists between the motor domain and the social communicative domains, however a
16 direct relationship between these domains has not been planned in young children with ASD
17 (Landa & Garrett-Mayer, 2006; Ozonoff, et al., 2008; Sauter, et al., 2007). In the present study a
18 direct relationship was found between fine motor skills and adaptive social and adaptive
19 communication skills of young children with ASD, while holding other important variables
20 constant including calibrated autism severity and non-verbal problem solving. Motor skills may
21 be a missing component of current early intervention programs and an additive piece to further
22 improve motor skills and possibly social communicative skills in young children with ASD
23 (MacDonald, et al., 2013; Rosenbaum, 2005). In this study gross motor skills were less

Motor skills, adaptive behavior skills & ASD

1 predictive of the adaptive social and adaptive communicative skills of these young children.
2 However, they were predicted of daily living skills. Although the MSEL is a standardized
3 measure of motor skills, it is typically used for developmental assessments, a more sensitive
4 measure of motor skills, such as the Peabody Motor Development Scales- 2nd Edition (Folio &
5 Fewell, 2000), may have been more informative. For an extensive review of motor skill
6 assessments for children with ASD please see (Staples, MacDonald, & Zimmer, 2012).

7 With the best content for early intervention under investigation, and with early
8 intervention on the forefront of ASD research, the importance of motor skills should also be
9 considered (Dawson, et al., 2010; Kasari, et al., 2005; Matson & Kozlowski, 2011; Sauter, et al.,
10 2007). The Denver model, an early intervention for young children with ASD, might be the
11 closest early intervention that successfully intertwines motor skills, social communicative skills
12 and social success; however how proficient motor skills impacts social success has not been
13 measured (Rogers, 2000). In fact, many early intervention models imply the use of games or
14 activities to facilitate social learning and some of the activities require relatively proficient motor
15 skills for success. For instance games like “chase”, songs that require actions, climbing activities
16 (like sliding or playing on equipment), as well as activities that include coloring or drawing are
17 often important components of early intervention programs. Arguably, these activities may be
18 more successful if the participants have more proficient motor skills. Unmistakably, the
19 emphasis of early intervention programs is social communicative, but it is possible that these two
20 domains, motor and social communicative, are already working together. In short, teaching age-
21 appropriate motor skills might add to a practitioner’s repertoire of intervention strategies aimed
22 at improving social success (Lloyd, et al., 2013; Sauter, et al., 2007; Swiezy, 2008).

1 The prevalence and pervasiveness of motor skill deficits in young children with ASD has
2 been clearly indicated across tasks (Chawarska, et al., 2007; Matson, et al., 2010; Ozonoff, et al.,
3 2008; Teittelbaum, et al., 1998; Vernazza-Martin, et al., 2005). Even at a young age, the motor
4 skill deficits of children with ASD widen as the children become older (age range 14- 36
5 months) (Lloyd, et al., 2013). Teaching motor skills to children with ASD, through adapted
6 physical activity, has been successful (MacDonald et al., 2012). How to best implement motor
7 skill improvements for young children with ASA and how to combine early intervention to
8 include motor skill development is an area of future research. However, programs such as those
9 implemented in pre-school adapted physical education settings and which include the use of
10 adapted equipment and adapted instruction with a focus on motor skill improvement would be a
11 good starting point (Goodway & Branta, 2003). Children with ASD have unique characteristics,
12 which will be important to consider in motor skill-based programming. Given the relationship
13 between motor skills and adaptive social communicative skills found in this study, including
14 motor skill development in early intervention might be a significant additive piece for young
15 children diagnosed with ASD in addition to more traditional early intervention programs.

16 **Future directions**

17 The necessity of early intervention for young children with ASD has clearly been
18 indicated. However, a more concerning feature is agreement on program content (Kasari, et al.,
19 2005). Retrospective comparisons of early intervention programs are difficult given the use of
20 various outcome measures (Lord et al., 2005; Matson & Sipes, 2010). Clearly, motor skills
21 deficits have been indicated, and this study shows direct relationships between fine and gross
22 motor skills and adaptive behavioral skills in young children with ASD. A critical next step
23 includes better understanding how motor skills interact with other types of social

Motor skills, adaptive behavior skills & ASD

1 communicative skills. Given the motor skill deficits in this group of children, group-based
2 programs to improve these skills are imperative and a consideration for future studies. If motor
3 skill interventions are successful, how motor skill interventions improve social communicative
4 success may also provide important information.

5 **Limitations**

6 Limitations to this study include the retrospective nature of the work and secondary data
7 analysis. Age equivalent motor scores were used in order to standardize scores across age (as
8 standard scores were only available up to 33 months). More sensitive motor skill assessments
9 need to be implemented to better understand this relationship, such as the Peabody Motor
10 Developmental Scales- 2nd Edition (Folio & Fewell, 2000; Staples, et al., 2012). Another
11 limitation to this study included the use of motor skills in the adaptive behavior composite score
12 derived from the Vineland Adaptive Behavior Scales. Although the specific motor scales of the
13 Vineland were not assessed, motor scales are embedded in the adaptive behavior composite
14 score. Another limitation to this study included the administration of the Vineland Adaptive
15 Behavior Scales. It appears that the Vineland was implemented using two different methods,
16 face-to-face interview and over-the-phone administration. Although similar, the reliability of
17 these methods has not been tested, to our knowledge.

18 **Conclusion**

19 The findings of this study acknowledge that a relationship exists between motor skills
20 and the core characteristics of ASD, deficits in the social and communicative domain. Motor
21 skills need to be considered and included in early intervention programming.

22

1 **References**

2 American Psychological Association (APA). (1994). *Diagnostic and statistical manual of mental*
3 *disorders* (4th ed.). Washington, DC: Author.

4 APA. (2000). *Diagnostic and statistical manual of mental disorders* (4th, text rev. ed.).
5 Washington, DC: Author.

6 APA. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC:
7 Author.

8 Berkeley, S. L., Zittel, L. L., Pitney, L. V., & Nichols, S. E. (2001). Locomotor and object control
9 skills of children diagnosed with autism. *Adapted Physical Activity Quarterly*, 18(4), 405-
10 416.

11 Blumberg, S., Bramlett, M., Kogan, M., Schieve, L., Jones, J., & Lu, M. (2013). Changes in
12 prevalence of parent-reported autism spectrum disorder in school-aged U.S children:
13 2007 to 2011- 2012. *National Health Statistics Report*, 65.

14 Center for Disease Control & Prevention (CDC). (2012).

15 Chawarska, K., Paul, R., Klin, A., Hannigen, S., Dichtel, L., & Volkmar, F. (2007). Parental
16 recognition of developmental problems in toddlers with autism spectrum disorders.
17 *Journal of Autism and Developmental Disorders*, 37(1), 62-72.

18 Dawson, G., Rogers, S., Munson, J., Smith, M., Winter, J., Greenson, J., . . . Varley, J. (2010).
19 Randomized, controlled trial of an intervention for infants with autism: The Early Start
20 Denver Model. *Pediatrics*, 125(1), 17-23.

21 DiLavore, P. C., Lord, C., & Rutter, M. (1995). Pre-Linguistic Autism Diagnostic Observation
22 Schedule (PLADOS). *Journal of Autism and Developmental Disorders*, 25(4), 355-379.

Motor skills, adaptive behavior skills & ASD

- 1 Esposito, G., & Venuti, P. (2008). Analysis of toddlers' gait after six months on independent
2 walking to identify autism: A preliminary study. *Perceptual and Motor Skills, 106*, 259-
3 269.
- 4 Fabbri-Destro, M., Cattaneo, L., Boria, S., & Rizzolatti, G. (2009). Planning action in autism.
5 *Experimental Brain Research, 192*(3), 521-525.
- 6 Flanagan, J., Landa, R., Bhat, A., & Bauman, M. (2012). Head lag in infants at risk for autism: a
7 preliminary study. *The American Journal of Occupational Therapy, 66*(5), 577-585. doi:
8 10.5014/ajot.2012.004192
- 9 Folio, M., & Fewell, R. (2000). *Peabody Developmental Motor Scales* (2nd ed.). San Antonio, TX:
10 Pearson.
- 11 Fournier, K., Hass, C., Naik, S., Lodha, N., & Cauraugh, J. (2010). Motor coordination in autism
12 spectrum disorders: A synthesis and meta-analysis. *Journal of Autism and Developmental*
13 *Disorders, 40*(10), 1227-1240. doi: 10.1007/s10803-010-0981-3
- 14 Fournier, K., Kimberg, C., Radonovich, K., Tillman, M., Chow, J., Lewis, M., . . . Hass, C. (2010).
15 Increased static and dynamic postural control in children with autism spectrum disorder.
16 *Gait Posture, 32*(1), 6-9. doi: 10.1016/j.gaitpost.2010.02.007
- 17 Frith, U. (1991). *Autism and Asperger syndrome*. Cambridge: Cambridge University Press.
- 18 Goodway, J. & Branta, C. (2003). Influence of a motor skill intervention on fundamental motor
19 skill development of disadvantaged preschool children. *Research Quarterly for Exercise*
20 *and Sport, 74*(1), 36-46.

Motor skills, adaptive behavior skills & ASD

- 1 Gotham, K., Pickles, A., & Lord, C. (2009). Standardizing ADOS scores for a measure of severity
2 in autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 39(5),
3 693-705. doi: 10.1007/s10803-008-0674-3
- 4 Gotham, K., Risi, S., Pickles, A., & Lord, C. (2007). The Autism Diagnostic Observation Schedule:
5 Revised algorithms for improved validity. *Journal of Autism and Development Disorders*,
6 37(4), 613- 627. doi: 10.1007/s10803-006-0280-1
- 7 Helt, M., Kelley, E., Kinsbourne, M., Pandey, J., Boorstein, H., Herbert, M., & Fein, D. (2008). Can
8 children with autism recover? If so, how? *Neuropsychological Review*, 18, 339-366.
- 9 Kasari, C., Freeman, S., & Paparella, T. (2006). Joint attention and symbolic play in young
10 children with autism: a randomized controlled intervention study. *Journal of Child*
11 *Psychology and Psychiatry*, 47(6), 611-620.
- 12 Kasari, C., Freeman, S., Paperella, T., Wong, C., Kwon, S., & Gultrud, A. (2005). Early intervention
13 of core deficits in autism. *Clinical Neuropsychiatry*, 2(6), 380-388.
- 14 Kasari, C., Gulsrud, A., Wong, C., Kwon, S., & Locke, J. (2010). Randomized controlled caregiver
15 mediated joint engagement intervention for toddlers with autism. *Journal of Autism and*
16 *Development Disorders*, 40(9), 1045-1056. doi: 10.1007/s10803-010-0955-5
- 17 Kasari, C., Paparella, T., Freeman, S., & Jahromi, L. (2008). Language outcome in autism:
18 Randomized comparison of joint attention and play interventions. *Journal of consulting*
19 *and clinical psychology*, 75(1), 125-137.
- 20 Kelley, E., Naigles, L., & Fein, D. (2010). An in-depth examination of optimal outcome children
21 with a history of autism spectrum disorders. *Research in Autism Spectrum Disorders*, 4,
22 526-538.

Motor skills, adaptive behavior skills & ASD

- 1 Landa, R., & Garrett-Mayer, E. (2006). Development in infants with autism spectrum disorders:
2 A prospective study. *Journal of Child Psychology and Psychiatry*, 47(6), 629-638. doi:
3 10.1111/j.1469-7610.2006.01531.x
- 4 Lloyd, M., MacDonald, M., & Lord, C. (2013). Motor skills of toddlers with autism spectrum
5 disorders. *Autism*, 17(2), 133-146. doi: 10.1177/1362361311402230
- 6 Lord, C., Risi, S., Lambrecht, L., Cook, E., Leventhal, B., DiLavore, P., . . . Rutter, M. (2000). The
7 Autism Diagnostic Observation Schedule- Generic: A standard measure of social and
8 communication deficits associated with the spectrum of autism. *Journal of Autism and*
9 *Developmental Disorders*, 30(3), 205-223.
- 10 Lord, C., Rutter, M., DiLavore, P. C., & Risi, S. (1999). *Autism Diagnostic Observation Schedule*
11 Los Angeles: Western Psychological Services.
- 12 Lord, C., Wagner, A., Rogers, S., Szatmari, P., Aman, M., Charman, T., . . . Yoder, P. (2005).
13 Challenges in evaluating psychosocial interventions for autistic spectrum disorders.
14 *Journal of Autism and Developmental Disorders*, 35(6), 695-708. doi: 10.1007/s10803-
15 005-0017-6
- 16 Luyster, R., Gotham, K., Guthrie, W., Coffing, M., Petrak, R., Pierce, K., . . . Lord, C. (2009). The
17 Autism Diagnostic Observation Schedule-toddler module: a new module of a
18 standardized diagnostic measure for autism spectrum disorders. *Journal of Autism &*
19 *Developmental Disorders*, 39(9), 1305-1320. doi: 10.1007/s10803-009-0746-z
- 20 MacDonald, M., Esposito, P., Hauck, J., Jeong, I., Hornyak, J., Argento, A., & Ulrich, D. (2012).
21 Successful bicycle training for youth with Down syndrome and autism spectrum

Motor skills, adaptive behavior skills & ASD

- 1 disorders. *Focus on Autism & Other Developmental Disabilities*. doi:
2 10.1177/1088357611428333
- 3 MacDonald, M., Jaszewski, C., Esposito, P., & Ulrich, D. (2011). The effect of learning to ride a
4 two-wheel bicycle on the social development of children with autism spectrum disorder:
5 a qualitative study. *Palestra, 25*(4), 37-42.
- 6 MacDonald, M., Lord, C., & Ulrich, D. (2013). The relationship of motor skills and social
7 communicative skills in school-aged children with autism spectrum disorder. *Adapted*
8 *Physical Activity Quarterly*.
- 9 Matson, J., & Kozlowski, A. M. (2011). The increasing prevalence of autism spectrum disorders.
10 *Research in Autism Spectrum Disorders, 5*, 418-425. doi:
11 <http://dx.doi.org/10.1016/j.rasd.2010.06.004>
- 12 Matson, J., Mahan, S., Fodstad, J. C., Hess, J. A., & Neal, D. (2010). Motor skills abilities in
13 toddlers with autistic disorder, pervasive developmental disorder-not otherwise
14 specified, and atypical development *Research in Autism Spectrum Disorders, 4*, 444-449.
- 15 Matson, J., & Sipes, M. (2010). Methods of early diagnosis and tracking for autism and Pervasive
16 Developmental Disorder Not Othewise Specified (PDDNOS). *Journal of Developmental*
17 *and Physical Disabilities, 22*, 343-358. doi: 10.1007/s10882-009-9184-2
- 18 Memari, A., Ghanouni, P., Gharibzadeh, S., Eghlidi, J., Ziaee, V., & Moshayedi, P. (2013). Postural
19 sway patterns in children with autism spectrum disorder compared with typically
20 developing children. *Research in autism spectrum disorders, 7*, 325-332. doi:
21 10.1016/j.rasd.2012.09.010

Motor skills, adaptive behavior skills & ASD

- 1 Mullen, E. (1995). *Mullen scales of early learning (AGS ed.)*. Circle Pines: MN: American
2 Guidance Services.
- 3 National Research Council (NRC). (2001).
- 4 Ozonoff, S., Young, G. S., Goldring, S., Greiss-Hess, L., Herrera, A. M., Steele, J., . . . Rogers, S. J.
5 (2008). Gross motor development, movement abnormalities, and early identification of
6 autism. *Journal of Autism and Developmental Disorders*, *38*, 644-656.
- 7 Provost, B., Heimerl, S., & Lopez, B. R. (2007). Levels of gross and fine motor development in
8 young children with autism spectrum disorder. *Physical & Occupational Therapy in*
9 *Pediatrics*, *27*(3).
- 10 Provost, B., Lopez, B. R., & Heimerl, S. (2007). A comparison in motor delays in young children:
11 Autism spectrum disorder, developmental delay, and developmental concerns. *Journal*
12 *of Autism and Developmental Disorders*, *37*, 321-328.
- 13 Rinehart, N., Tonge, B., Bradshaw, J., Iansek, R., Enticott, P., & Johnson, K. (2006). Movement
14 related potential in high-functioning autism and Asperger's disorder. *Developmental*
15 *medicine and child neurology*, *48*, 272-277.
- 16 Rogers, S. (2000). Interventions that facilitate socialization in children with autism. *Journal of*
17 *Autism and Developmental Disorders*, *30*(5), 399-409.
- 18 Rosenbaum, D. (2005). The Cinderella of psychology: The neglect of motor control in the
19 science of mental life and behavior. *American Psychologist*, *60*(4), 308-317. doi:
20 10.1037/0003-066X.60.4.308
- 21 Sparrow, S., Cicchetti, D., & Balla, D. (2005). *Vineland-II Adaptive Behavior Scales*. Circles Pines,
22 MN.: American Guidance Service.

Motor skills, adaptive behavior skills & ASD

- 1 Staples, K., MacDonald, M., & Zimmer, C. (2012). Assessment of motor behavior among children
2 and adolescents with autism spectrum disorder. *International review of research in*
3 *developmental disabilities*, 42, 179-214. doi: 10.1016/B978-0-12-394284-5.00007-3
- 4 Staples, K., & Reid, G. (2009). Fundamental movement skills and autism spectrum disorders.
- 5 Sutera, S., Pandey, J., Esser, E. L., & Rosenthal, M. A. (2007). Predictors of optimal outcome in
6 toddlers diagnosed with autism spectrum disorders. *Journal of Autism and*
7 *Developmental Disorders*, 37(1), 98-107. doi: 10.1007/s10803-006-0340-6
- 8 Swiezy, N. (2008). Bridging for success in Autism: Training and collaboration across medical,
9 educational, and community systems. *Child and adolescent psychiatric clinics of North*
10 *America*, 17(4).
- 11 Teittelbaum, P., Teittelbaum, O., Nye, J., Fryman, J., & Maurer, R. G. (1998). Movement analysis
12 in infancy may be useful for early diagnosis of autism. *Psychology*, 95, 13982-13987.
- 13 Teittelbaum, P., Teittelbaum, O., Nye, J., Fryman, J., & Maurer, R. (1998). Movement analysis in
14 infancy may be useful for early diagnosis of autism. *Psychology*, 95, 13982-13987.
- 15 Vernazza-Martin, S., Martin, N., Vernazza, A., Lepellec-Muller, A., Rufo, M., Massion, J., &
16 Assaiante, C. (2005). Goal directed locomotion and balance control in autistic children.
17 *Journal of Autism and Developmental Disorders*, 35(1), 91-102.
- 18 Wong, V. C. N., & Kwan, Q. K. (2010). Randomized controlled trial for early intervention for
19 autism: A pilot study of the autism 1-2-3 Project. *Journal of Autism and Developmental*
20 *Disorders*, 40, 677-688.
- 21 Yirmiya, N., & Ozonoff, S. (2007). The very early autism phenotype. *Journal of Autism and*
22 *Developmental Disabilities*, 37(1), 1-11. doi: 10.1007/s10803-006-0329-1

Motor skills, adaptive behavior skills & ASD

1

2

Table 1.

Descriptive characteristics of the participants

Variable	ASD (n=172)	PDD-NOS (n=22)	Non-ASD (n=39)
Age (months)	30.3 (5.7)	31.1 (7.6)	30.0 (5.9)
Gender	M=141, F= 31	M=16, F=6	M=27, F=12
Race/Ethnicity	Caucasian= 113, Other=59	Caucasian=14, Other=8	Caucasian=27, Other=9
Maternal Education	Grad/ professional=30 College degree=46 Some college=51 High school=26 Some high school= 3 Unknown=5	Grad/professional=3 College degree=5 Some college=4 High school=5 Some high school=0 Unknown=5	Grad/professional=6 College degree=6 Some college=11 High school=11 Some high school=1 Unknown=4
Gross Motor Age Equivalent	21.7 (6.2)	20.9 (7.9)	25.3 (7.4)
Fine Motor Age Equivalent	18.4 (6.1)	21.1 (7.2)	23.5 (7.0)
Gross motor difference variable (months)	8.8(7.6)*	7.6(7.5)	6.5(7.6)
Fine motor difference variable (months)	12.0(7.2)	8.6(6.4)	8.4(8.0)
Visual Reception Age Equivalent	19.2 (6.4)	22.4 (7.7)	25.9 (7.4)
Receptive Language Age Equivalent	8.5 (6.0)	16.5 (6.7)	23.3 (9.1)
Expressive Language Age Equivalent	10.5 (5.8)	13.4 (7.1)	20.7 (10.3)

Vineland Fine Motor Age Equivalent	18.1 (4.9)	19.6 (6.9)	20.9 (6.6)
Vineland Gross Motor Age Equivalent	20.6 (5.6)	21.5 (11.1)	21.6 (7.8)
Vineland Overall Motor Age Equivalent	19.5 (5.1)	20.7 (7.9)	21.0 (7.1)
Vineland Standard Motor Score	73.2 (12.6)	75.3 (19.2)	78.0 (16.1)
Ratio Verbal IQ	31.5 (18.2)	48.8 (17.0)	73.4 (26.6)
Ratio Non-Verbal IQ	63.2 (19.9)	72.2 (24.6)	83.2 (18.4)

*Mean (standard deviation)

Table 2.

Multiple regression analysis testing fine motor skill on the adaptive behavior composite, daily living skills adaptive social and adaptive communicative skills.

Variable	Adaptive Behavior Composite			Daily Living Skills		
	<i>B</i>	<i>SE B</i>	<i>β</i>	<i>B</i>	<i>SE B</i>	<i>β</i>
Fine motor skills	.497	0.08	.352*	1.92	0.34	.34
Calibrated severity	-.956	0.19	-.304*	-4.282	0.76	-.33
<i>R</i> ²	.19			.29		
<i>F</i>	46.7*			46.7		

p* < .001 *p* < .05.

Adaptive Social Skills

Adaptive Communicative Skills

Variable	<i>B</i>	<i>SE B</i>	<i>β</i>	<i>B</i>	<i>SE B</i>	<i>β</i>
Fine motor skills	.196	0.09	.124**	.358	0.087	.231
Calibrated severity	-1.9	0.20	-.537*	-1.70	0.194	-.49
<i>R</i> ²	.35			.37		
<i>F</i>	59.0*			65.1		

Table 3.

Multiple regression analysis testing gross motor skill on the adaptive behavior composite, daily living skills adaptive social and adaptive communicative skills.

Variable	Daily Living Skills		
	<i>B</i>	<i>SE B</i>	<i>β</i>
Gross motor skills	1.6	.32	.29*
Calibrated severity	-5.63	0.768	-.439*
<i>R</i> ²	.27		
<i>F</i>	42.5*		

**p* < .001