

History of Entomopathogenic Nematology

G. O. POINAR JR.,¹ P. S. GREWAL²

Abstract: The history of entomopathogenic nematology is briefly reviewed. Topic selections include early descriptions of members of *Steinernema* and *Heterorhabditis*, how only morphology was originally used to distinguish between the species; descriptions of the symbiotic bacteria and elucidating their role in the nematode-insect complex, including antibiotic properties, phase variants, and impeding host defense responses. Other topics include early solutions regarding production, storage, field applications and the first commercial sales of entomopathogenic nematodes in North America. Later studies centered on how the nematodes locate insect hosts, their effects on non-target organisms and susceptibility of the infective juveniles to soil microbes. While the goals of early workers was to increase the efficacy of entomopathogenic nematodes for pest control, the increasing use of *Heterorhabditis* and *Photorhabdus* as genetic models in molecular biology is noted.

Key words: History, entomopathogenic nematodes, *Heterorhabditis* spp., *Steinernema* spp.

Entomopathogenic nematodes of the families Steinernematidae and Heterorhabditidae are symbiotically associated with bacteria in the genera *Xenorhabdus* and *Photorhabdus*, respectively. When an infective juvenile enters the body cavity of a susceptible host, the bacteria are released, multiply and host death occurs within two days, hence the term, entomopathogenic. The nematodes develop and reproduce within the insect cadaver, feeding on the symbiotic bacteria and degraded host tissues.

The first entomopathogenic nematode was described by Steiner as *Aplectana kraussei* (now *Steinernema kraussei*) in 1923 and at that time was considered no more than a curiosity whose systematic position was problematic. The systematic position of the second entomopathogenic nematode, *Neoalectana glaseri* Steiner (1929) from material isolated by Glaser and Fox (1930), was still not certain and Steiner placed it in the family Oxyuridae. The early history of this nematode will be discussed later but after its use against the Japanese beetle, it ceased to be of interest and all cultures were lost.

It was not until Jaroslav Weiser (1955) (Fig. 1) described a European population of *Neoalectana carpocapsae* from codling moth larvae and Dutky and Hough (1955) isolated the DD-136 strain of an undescribed steinernematid from codling moth larvae in Eastern North America that serious studies on the pathogenicity and life history of entomopathogenic nematodes began. In 1965, cultures of *S. carpocapsae* were obtained from Weiser and using morphology and hybridization studies, it was shown that the Czechoslovakian strain of *S. carpocapsae* and the North American DD-136 nematode were conspecific (Poinar, 1967).

The symbiotic bacterium (under the name, *Achromobacter nematophilus*) associated with *S. carpocapsae* was described by Poinar and Thomas (1965). The location of the bacteria in the infective stage juveniles using light

microscopy and later electron microscopy was then demonstrated (Poinar, 1966; Poinar and Leutenegger, 1968). The role of the bacterium in the development of the nematode and death of the host was elucidated (Poinar and Thomas, 1966, 1967) and the bacterium was later transferred to a new genus, *Xenorhabdus* (Thomas and Poinar, 1979) (Fig. 2).

At first, morphology of the male tail or features of the infective stage juveniles alone could be used to differentiate between the various species of *Steinernema* (Poinar, 1986; Wright, 1990). Also, using the biological species concept, which is well suited for *Steinernema*, cross-breeding could be performed to determine specific or intra-specific status of the numerous geographical strains that were discovered (Poinar and Veremtschuk, 1970; Poinar, 1986, 1990). As more isolates were unearthed, (there are now some 36 species of *Steinernema* (Stock and Hunt, 2005), measurements between them overlapped and genomic analysis also was used to determine their specific status (Liu et al., 2000; Ciche, 2007).

The genus *Heterorhabditis* was described in 1976 and the symbiotic bacterium of *H. bacteriophora* was characterized as *Xenorhabditis luminescence* in 1979 (Poinar, 1976; Thomas and Poinar, 1979). The fascinating character of the symbiotic bacteria of *Heterorhabditis* spp. was its ability to fluoresce, so much so that the entire infected insect cadaver glowed in the dark (Fig. 3) and light even could be detected in a single infective stage (Poinar et al., 1980a). This bacterial species was later transferred to the genus *Photorhabdus* (Boemare et al., 1993). The location of the bacterial cells in the infective stage was shown with electron microscopy (Poinar et al., 1977) (Fig. 4) and aspects of its behavior were elucidated by Milstead (1977). As with *Steinernema*, there are many geographic species and strains of *Heterorhabditis* (Poinar, 1990; Stock and Hunt, 2005) and the global distribution of both *Heterorhabditis* and *Steinernema* indicates that their lineages were present when all land masses were combined as the Pangaea supercontinent. It is interesting to note that a genetic analysis showed that *Heterorhabditis* is a sister group to the vertebrate-parasitic strongyloids and that both groups arose independently

Received for publication February 10, 2012.

¹Department of Zoology, Oregon State University, Corvallis, OR 97331.

²Ohio State University, Wooster, OH 44691.

Email: poinarg@science.oregonstate.edu

This paper was edited by Nancy Kokalis-Burelle.



FIG. 1. Jaroslav Weiser described *Steinernema carpocapsae* in 1955. At the International Colloquium of Insect Pathology and Microbial Control in Wageningen, the Netherlands in 1966. Photo by G. Poinar.

from the free-living *Rhabditis* group (Kiontke et al., 2007). The obligate association of *Heterorhabditis* with a unique genus of luminescent symbiotic bacteria, its ability to enter the body of healthy insects, the alternating



FIG. 2. Gerard Thomas (left) and George Poinar at UC Berkeley described the symbiotic bacterium associated with *Steinernema carpocapsae* in 1965 and *Heterorhabditis bacteriophora* in 1979. They also revealed the significance of the bacteria in the development of the nematodes. Photo by Roberta Poinar.




FIG. 3. One character that distinguished *Heterorhabditis* from other rhabditids was its ability to transmit a luminescent bacterium. Here are cadavers of the wax moth glowing in the dark after being infected with *Heterorhabditis bacteriophora* 48 hours earlier. Photo by G. Thomas.

of sexual and hermaphroditic generations and the unique morphology of the parasitic adults explains why this genus was assigned family status. One unique feature of the infective juveniles of *Heterorhabditis* that is lacking in *Steinernema* and other rhabditids is the presence of a dorsal “hook” on the tip of the head. This structure allows the infectives to enter the body cavity through the outer tegument of potential hosts, as well as through the trachea and gut wall (Bedding and Molyneux, 1982; Poinar and Georgis, 1990). The ancient age of the *Heterorhabditis* clade was recently shown when a 100 million year old fossil (*Proheterorhabditis burmanicus*) was discovered in Early Cretaceous Burmese amber (Poinar, 2011).

It is now evident that all *Steinernema* species have mutualistic associations with species of *Xenorhabdus* while all *Heterorhabditis* species have symbiotic associations with *Photorhabdus* species (Akhurst and Boemare, 1990; Boemare, 2002). The elucidation of the symbiosis between entomopathogenic nematodes and their associated bacteria was a major turning point in the development of the nematodes as commercial biological control agents.



FIG. 4. Electron micrograph of *Photorhabdus luminescens* in the anterior portion of the intestine of the infective stage of *Heterorhabditis bacteriophora*. Photo by Roberta Poinar.

Dutky (1959) first noticed the antibiotic properties of the bacterium associated with *S. carpocapsae*, which explained how it could destroy foreign bacteria that invaded the insect cadaver containing the developing nematodes. Since then several antibiotics, including xenorhabdins, xenocaumacins, hydroxystilbenes, indole derivatives, and anthraquinone derivatives, were recovered from cultures of *Xenorhabdus* and *Photorhabdus* (Webster et al., 2002).

Akhurst (1980) discovered that in culture, *Xenorhabdus* existed in two (or more) genetically identical phase variants that differed in colony morphology, colony color and antimicrobial activity. The primary phase, which is carried by naturally occurring infective stages, provided maximum nematode growth and antibiotic production. But the primary phase would suddenly revert to a secondary phase, which was much less supportive of nematode growth and had limited antibody production. This shift was a major hindrance in the commercial production of the nematodes. The cause of this sudden phase shift was unknown until a bacteriophage of *Heterorhabditis* was discovered that only attacked the primary phase. Shifting to the secondary phase would then be a survival response of the *Xenorhabdus* primaries (Poinar et al., 1980b; Poinar et al., 1989).

Insects have a barrage of defense reactions to invading parasites and the most significant against nematodes is melanization and encapsulation. Normally, the bacteria will kill the host before a lethal response can be effected. However, in some experimental hosts such as mosquitoes, a rapid melanization reaction will kill the infectives before they can release their symbiotic bacteria (Bronskill, 1962; Welch and Bronskill, 1962). Also, if the infectives of *S. carpocapsae* lack cells of their symbiotic bacterium, the developing stages can be encapsulated and killed, even in larvae of *Galleria mellonella*, which is the most common host used for raising entomopathogenic nematodes (Poinar, 1969).

There were also natural enemies of the infective stages to consider, especially protozoa and fungi. When caterpillars already infected with microsporidians were later invaded by *S. carpocapsae*, the protozoan infections were transferred to the nematodes (Veremchuk and Issi, 1970). Naturally occurring populations of entomopathogenic nematodes can also be infected with microsporidians (Poinar, 1988). Infective stages of entomopathogenic nematodes are also susceptible to infection by several common soil fungi (Poinar and Jansson, 1986a, 1986b) showing that before applying the nematodes to soils rich in humus, it is prudent to make a survey of nematophagous fungi that might be present.

While the first entomopathogenic nematode, *S. kraussei* Steiner, was described in 1923, the biocontrol potential of these nematodes was first investigated by Glaser and his colleagues who investigated *S. glaseri* to control populations of the Japanese beetle that had

recently invaded New Jersey. Growth of the nematodes was tested on different types of artificial media for mass production (Fig. 5). While *S. glaseri* is associated with a symbiotic bacterium, Glaser and his group were unaware of its presence and this bacterium became lost during the sterilization processes when the nematodes were transferred to artificial media. It was fortunate for the New Jersey team that *S. glaseri* is one of the most catholic nematodes in the genus regarding the ability to develop on other bacteria as well as yeast in the absence of its symbiont. While production is much lower than with the natural-occurring symbiotic bacterium, the nematodes can still kill, invade, destroy and reproduce inside insects (Poinar, 1969). Mass-production on artificial media was achieved and large-scale field releases of *S. glaseri* were conducted with the nematodes dispersed from a motor driven tank (Glaser, 1932; Glaser and Farrell, 1935). Details of the field trials are summarized in Poinar (1979). The symbiotic bacterium was later discovered in a population of *S. glaseri* from North Carolina (Poinar and Brooks, 1977; Poinar, 1978).

Living insects were used to produce the first entomopathogenic nematodes for field testing. Nutrilite Corporation in Lakeview, CA used larvae of the wax moth, *Galleria mellonella* to produce Biotrol NCS-DD-136 in 1970 for experimental use. In 1981, "The Nematode Farm" in Berkeley produced several entomopathogenic nematodes (*S. carpocapsae*, *S. glaseri* and *H. bacteriophora*) in *Galleria mellonella* for commercial use against garden pests. Also in 1981, BR Supply in Exeter, CA raised *S. carpocapsae* on crickets and packaged a product called Neocide for use against the carpenter worm. In 1982, Biosys, in Palo Alto, CA, (previously established as "The California Nematode Laboratories" in Emeryville, CA, and for a brief period "Biosis" in Palo Alto, CA), was the first to use a fermentation process to mass produce *Steinernema* spp. and their commercial products, Bio-Safe, BioVector, etc., were aimed at lawn and garden insects. In 1983, Biotechnology Australia produced



FIG. 5. Rudolf Glaser (right) and co-worker examine culture plates with *S. glaseri* in their New Jersey laboratory in the early 1930s.

nematodes on particles of sponge impregnated with an artificial diet, based on a method developed earlier by Bedding (1981). This technique used polyether polyurethane sponge as a three-dimensional support that allowed the nematodes to move through the matrix and provide air exchange. Their product, Otinem, was aimed at black vine weevils in Australia and Europe. Commercial production of entomopathogenic nematodes in liquid culture was later perfected by a team of researchers led by Friedman (1990) at Biosys Inc. A number of additional smaller companies, some as cottage industries, appeared in the mid- 80s.

One of the serious problems in the commercialization of entomopathogenic nematodes, aside from mass production, was storage under conditions that maintained high viability together with high infectivity. Refrigeration was a suitable method but not practical for retailers and growers who wanted to sell or apply the nematodes over a period of several weeks or even days. Studies were then initiated on the possibility of desiccating the nematodes so they could be stored at room temperatures. Simons and Poinar (1973) discovered that if the infective stages of *S. carpocapsae* were desiccated slowly, they entered a partial anhydrobiosis and could be quickly revived with water and still retain their infectivity. Based on these findings, Bedding (1988) later developed a "clay sandwich" formulation in which nematodes were placed in layers of clay to remove surface water and induce partial anhydrobiosis. Anhydrobiosis was further used to enhance the ambient storage stability of entomopathogenic nematodes (Grewal, 2000). Scientists at Biosys (see Georgis, 1990) developed an alginate formulation in which sheets of calcium alginate spread over plastic screens were used to entrap and maintain nematodes. Bedding and Butler (1994) reported a formulation in which the nematode slurry was mixed with a powder of anhydrous polyacrylamide to achieve a water activity between 0.800 and 0.995. Capinera and Hibbard (1987) developed a granular formulation in which nematodes were partially encapsulated in lucerne meal and wheat flour. Later Connick et al. (1993) described an extruded granule in which nematodes were distributed throughout a wheat gluten matrix. This "pesta" formulation included a filter and humectant to enhance nematode survival.

A major leap in the development of nematode formulations was reported by Silver et al. (1995) who developed a water dispersible granular formulation in which the nematodes were encased in 10-20 mm diameter granules consisting of a mixture of various types of silica, clay, cellulose, lignin, and starches. With this formulation, the shelf-life of commercially produced *S. carpocapsae* was extended to 7 months at ambient temperatures (Gaugler et al., 2000). Applications against aerial pests also posed a problem since if the nematodes desiccated too rapidly, their effectiveness was greatly reduced. Webster and Bronskill (1968) used an evaporation

retardant to prolong the life of the nematodes used against foliar pests.

The infective stages of entomopathogenic nematodes can be easily applied using conventional pesticide application equipment, however, applying the nematodes simultaneously with other agents saves labor costs. Rao et al. (1975) showed that *S. carpocapsae* can be tank mixed with certain insecticides and in their trials against the corn rootworm, Poinar et al. (1983) applied the infective stages of *S. carpocapsae* with liquid fertilizer. These initial results were followed by a series of comprehensive studies on the effects of combining the infective stages of *Steinernema* and *Heterorhabditis* with pesticides (Rovesti and Deseo, 1989, 1990, 1991). As compatibility information is critical for implementation of nematodes in integrated pest management systems, a comprehensive review was published by Koppenhöfer and Grewal (2005). Kaya and Nelson (1985) investigated the application of infective stages in alginate gels for increased persistence and this practice was commercialized in the application of nematodes to tree trunks. Since application problems were often experienced, Lello et al. (1996) and Fife et al. (2003, 2004, 2005) systematically evaluated the influence of droplet size, pressure differentials, hydraulic nozzles, contraction flow fields and agitation, on nematode viability and virulence. Masson et al. (1999) proposed the potential utility of spinning disc technology for the application of nematodes against foliar pests. Reed et al. (1986) were the first to apply nematodes through trickle irrigation, Wright et al. (1993) through center-pivot irrigation, and Cabanilas and Raulston (1996) through furrow irrigation systems. Subsurface application of nematodes with an adapted seed-driller was reported by Shetlar (1993). Bari (1992) developed a method of soaking plant cuttings in nematode suspensions to control the artichoke plume moth while Pye and Pye (1985) proposed a root dip method to economize nematode application rates. A slow release formulation using an absorbent gel was used to apply nematodes in citrus (Georgis, 1990) and a similar formulation (tea bag) was used in oilseed rape (Menzler-Hokkanen and Hokkanen, 2004). Infected insect cadavers can also serve as slow release systems for nematodes (Jansson and Lecrone, 1994) and Shapiro-Illan et al (2001) improved this method by for application by formulating nematode infected cadavers with powered starch to reduce their stickiness.

Miller (1989) developed a pathogenicity assessment technique to determine the virulence of commercially produced *S. carpocapsae*. This method was later called the one-on-one *Galleria mellonella* bioassay (Converse and Miller, 1999). Grewal et al. (1999) developed the sandwell method, which is suitable for routine quality assessment of most entomopathogenic nematode species at low concentrations. Many other methods serve as indicators of infectiousness, pathogenicity or the general

quality of the nematodes (see Grunder et al., 2005). Gaugler et al. (2000) assessed the quality of commercially produced nematodes, thus raising awareness about the importance of effective quality control during commercialization. Georgis et al., (2006) discussed the success and failures of entomopathogenic nematodes used as biological control agents.

Attention then turned to the behavior of the infective stages and how they were able to locate insect hosts. Reed and Wallace (1965) described three types of movement of the infective stages of *S. carpocapsae*; gliding, bridging and leaping. The infectives used a gliding motion to reach the soil surface. The bridging motion involved the nematodes standing on their tails and waving their anterior ends (nictating). Such behavior had already been noted in free-living rhabditids that had phoretic relationships with insects. The amazing leaping behavior consisted of the infectives coiling their bodies around a water droplet (which produced a tension force), suddenly releasing the droplet by uncoiling and being propelled horizontally across the substrate by the tension force. The actual leap was too rapid to be followed with the naked eye. It is obvious that various clues are used by the infective stages to locate their hosts. Byers and Poinar (1982) showed that *S. carpocapsae* infectives could locate hosts by minute temperature gradients while Lewis et al. (1992) showed *S. carpocapsae* and *S. glaseri* responded to long-range volatile cues.

Differences in the vertical and horizontal dispersal of *Steinernema* infectives were determined to be related to host-finding behavior by Moyle and Kaya (1981). These authors noted that while *S. carpocapsae* did not shift its position significantly from the site of application, *S. glaseri* moved horizontally long distances. Georgis and Poinar (1983a, 1983b) showed how soil texture affected the distribution and infectivity of *S. carpocapsae* and *S. glaseri*, respectively and Poinar and Hom (1986) studied the survival and horizontal movement of the infective stages of *S. carpocapsae*. Vertical migration of *Heterorhabditis* spp. in soil was investigated by Georgis and Poinar (1983c).

A series of studies conducted in Gaugler's laboratory described two host finding behaviors of the infective stages; ambushers and cruisers. Nematodes using the ambush type of foraging are better adapted at locating highly mobile hosts on the soil surface while nematode species that use the cruiser type are more adapted for sedentary hosts in the soil. Gaugler and Campbell (1991) proposed that the ambushing type of host-finding behavior may explain the limited movement of some entomopathogenic nematode species in the soil. Grewal et al. (1994a, 1994b) demonstrated that *S. feltiae* and *S. riobrave* use an intermediate type of foraging behavior between the ambushing and cruising exhibited by *S. carpocapsae* and *S. glaseri*, respectively. These studies showed researchers that it was possible to

match the host-finding behavior of nematodes with the life history parameters of target pests.

Poinar (1979) summarized early reports on challenging non-insect invertebrates and vertebrates with *S. carpocapsae* and Georgis et al. (1991) later demonstrated the safety of entomopathogenic nematodes to soil invertebrates. Aside from the negative report of *S. carpocapsae* killing adult honey bees (Hackett and Poinar, 1973), these nematodes have a minimal effect on non-target invertebrates. However, apart from their use for controlling a wide range of insects, the ability of entomopathogenic nematodes to infect some non-insect groups was a bonus. Samish and Glazer (1991) discovered that entomopathogenic nematodes are capable of killing engorged female cattle ticks. Ticks of the genera *Amblyomma*, *Argas*, *Boophilus*, *Dermacentor*, *Hylomma*, and *Rhipicephalus* have now been found to be susceptible to entomopathogenic nematodes (Glazer et al., 2005). Although the infective nematodes can invade and kill ticks and thus have potential for their control, there is no evidence of nematode reproduction in these arachnids.

Some insect vectors of human diseases are also susceptible to entomopathogenic nematodes. The susceptibility of fleas was first demonstrated with the cat flea, *Ctenocephalides felis* (Silverman et al., 1982). Biosys developed a biocontrol product with *S. carpocapsae* for the control of fleas in home lawns as a part of an integrated control program (Manweiler, 1994). Susceptibility of the body louse *Pediculus humanus humanus* to entomopathogenic nematodes was first demonstrated by Weiss et al. (1993). They reported that *S. carpocapsae* and *S. glaseri* caused over 85% mortality of female lice within 24 h. Doucet et al. (1998) showed that the head louse *P. humanus capitis* also is susceptible to entomopathogenic nematodes and *H. bacteriophora* was most effective of those tested. Larvae of phlebotomine flies, the vectors of leishmaniasis, are also susceptible to infection by *Steinernema* and *Heterorhabditis* (Poinar et al., 1993).

There are some reports of vertebrate infections by entomopathogenic nematodes. The infectives of *S. carpocapsae* were able to kill tadpoles of the Antillean toad, *Bufo marinus* (Kermarrec and Mauléon, 1985) and *Heterorhabditis* and *Steinernema* infectives caused mortality of frog tadpoles (Poinar and Thomas, 1988). While runoff could carry the infectives into standing water where tadpoles occurred, the likelihood of them making contact with amphibian larvae would be minimal.

Although natural populations of entomopathogenic nematodes are well-adapted to their native habitat and hosts through natural selection, additional useful traits could be established in their genome to make them even more efficient against other hosts in different environments. Poinar (1991) discussed some desirable traits that could be introduced into nematodes through recombinant DNA technology such as microinjection,



FIG. 6. Harry Kaya (left) and Randy Gaugler at the “International Symposium on Entomopathogenic nematodes in Biological Control” that they organized in Asilomar, CA, August 20-22, 1989. Photo by G. Poinar.

gene transplanted, mutagenesis and selective breeding. Already in 1980, Burman and Pye had developed a temperature-selective strain of *S. carpocapsae* whose infectives gravitate toward the temperature at which they underwent development. Fodor et al (1989) were able to obtain mutants of *S. carpocapsae* that were resistant to anthelmintics. Gaugler et al. (1989) demonstrated the use of selective breeding for enhanced host-finding in *S. carpocapsae*. Hashmi et al. (1995) used microinjection to produce a transgenic strain of *H. bacteriophora* that incorporated a heat shock protein gene from *Caenorhabditis elegans* and Sandhu et al. (2006) reported expressed-sequenced tags (ESTs) of *H. bacteriophora*. Although the benefits of genetic selection were demonstrated, the only field application of a selected strain was the Kapow strain of *S. carpocapsae* created by Jim Lindegren. This strain was developed by using the first juveniles to leave the host to infect subsequent hosts. The infectives of the Kapow strain were more active than the source strain and were used in field trials against navel orangeworms in California almond orchards (Agudelo-Silva et al., 1987). Genetic manipulation of the symbiotic bacteria is also feasible and studies have been undertaken to examine the genome of *Photorhabdus luminescens* (Duchaud et al., 2003).



FIG. 7. Infective stage of the HP88 strain of *Heterorhabditis bacteriophora*. Aside from their role as biocontrol agents, members of this genus and their associated bacteria are also serving as genetic models in molecular biology. Photo by G. Poinar.

This brief synopsis of the history of entomopathogenic nematodes necessitated a certain brevity, which did not allow us to include the works of many others who contributed to this field. For instance, the “International Symposium on Entomopathogenic nematodes” in Biological Control that Harry Kaya and Randy Gaugler organized at Asilomar, CA in 1989 (Fig. 6) provided a chance for researchers around the world to discuss common problems. It is easy to forget the many past hurdles that had to be overcome in order to reach where we are today. One crucial obstacle was obtaining government registration of not just the nematodes, but also their symbiotic bacteria. Educating the public by making it clear that entomopathogenic nematodes would not infect plants and humans was another early issue. There are still many ways in which the productivity and use of entomopathogenic nematodes can be improved (Grewal et al., 2005), however an interesting sidelight regarding *Heterorhabditis* and its associated bacterium is their increasing use as genetic models in molecular biology (Cliche, 2007; Duchaud et al., 2003) (Fig. 7).

Acknowledgments. The senior author thanks Roberta Poinar for comments on earlier drafts of this work.

LITERATURE CITED

- Agudelo-Silva, F., Lindegren, J. E., and Valero, K. A. 1987. Persistence of *Neoalectana carpocapsae* (Kapow selection) infectives in almonds under field conditions. *Florida Entomologist* 70:288–291.
- Akhurst, R. J. 1980. Morphological and functional dimorphism in *Xenorhabditis* spp., bacteria symbiotically associated with the insect pathogenic nematodes *Neoalectana* and *Heterorhabditis*. *Journal of General Microbiology*, 121:303–309.
- Akhurst, R. J., and Boemare, N. E. 1990. Biology and Taxonomy of *Xenorhabdus*, Pp. 75–90 in R. Gaugler, and H. K. Kaya, eds. *Entomopathogenic nematodes in Biological Control*. Boca Raton: CRC Press.
- Bari, M. A. 1992. Disinfestation of artichoke stumps with entomopathogenic nematode against the artichoke plume moth. *Proceedings of the International Congress of Entomology*, Beijing, China. 19:319.

- Bedding, R. A. 1981. Low cost in vitro mass production of *Neoalectana* and *Heterorhabditis* species (Nematoda) for field control of insect pests. *Nematologica* 27:109–114.
- Bedding, R. A. 1988. Storing third stage infective nematode juveniles by mixing with clay, placing between layers of clay or contacting with adsorbent. International Patent WO 88/08668.
- Bedding, R. A., and Akhurst, R. J. 1975. A simple baiting technique for the detection of insect parasitic rhabditid nematodes in soil. *Nematologica* 21:109–110.
- Bedding, R. A., and Molyneux, A. S. 1982. Penetration of insect cuticle by infective juveniles of *Heterorhabditis* spp. (Heterorhabditidae: Nematoda). *Nematologica* 28:354–359.
- Bedding, R. A., and Butler, K. L. 1994. Storage and transport of entomopathogenic nematodes. Australian Patent No. 608852.
- Boemare, N. E. 2002. Biology, taxonomy and systematics of *Photorhabdus* and *Xenorhabdus*. Pp. 35–56 in R. Gaugler, ed. *Entomopathogenic Nematology*. Wallingford, UK: CABI Publishing.
- Boemare, N. E., Akhurst, R. J., and Mourant, R. G. 1993. DNA relatedness between *Xenorhabdus* spp. (Enterobacteriaceae), symbiotic bacteria of entomopathogenic nematodes, and a proposal to transfer *Xenorhabdus luminescens* to a new genus, *Photorhabdus* gen. nov. *International Journal of Systematic Bacteriology* 43:249–255.
- Bronskill, J. F. 1962. Encapsulation of rhabditoid nematodes in mosquitoes. *Canadian Journal of Zoology* 40:1269–1275.
- Burman, M., and Pye, A. E. 1980. *Neoalectana carpocapsae*: Movements of nematode populations on a thermal gradient. *Experimental Parasitology* 48:258–265.
- Byers, J. A., and Poinar, G. O., Jr. 1982. Location of insect hosts by the nematode, *Neoalectana carpocapsae*, in response to temperature. *Behaviour* 79:1–10.
- Cabanillas, H. E., and Raulston, J. R. 1996. Effects of furrow irrigation on the distribution and infectivity of *Steinernema riobravii* against corn earworm in corn. *Fundamental and Applied Nematology* 19:273–281.
- Capinera, J. L., and Hibbard, B. E. 1987. Bait formulations of chemical and microbial insecticides for suppression of crop-feeding grasshoppers. *Journal of Agricultural Entomology* 4:337–344.
- Ciche, T. 2007. The biology and genome of *Heterorhabditis bacteriophora*. The biology and genome of *Heterorhabditis bacteriophora* (February 20, 2007), *WormBook*, ed. The *C. elegans* Research Community, WormBook, Doi/10.1895/wormbook.1.135.1, <http://www.wormbook.org>.
- Connick, W. J., Nickle, W. R., and Vinyard, B. T. 1993. Pesta: new granular formulations for *Steinernema carpocapsae*. *Journal of Nematology* 25:198–203.
- Converse, V., and Miller, R. W. 1999. Development of the one-on-one quality assessment assay for entomopathogenic nematodes. *Journal of Invertebrate Pathology* 74:143–148.
- Doucet, M. A., Miranda, M. B., and Bertolotti, M. A. 1998. Infectivity of entomogenous nematodes (Steinernematidae and Heterorhabditidae) to *Pediculus humanus capitis* De Geer (Anoplura: Pediculidae). *Fundamental and Applied Nematology* 21:13–16.
- Duchaud, E., Rusniok, C., Frangeul, L., Buchrieser, C., Givaudan, A., Taourit, S., Bocs, S., Boursaux-Eude, C., Chandler, M., and Charles, J. F. 2003. The genome sequence of the entomopathogenic bacterium *Photorhabdus luminescens*. *Nature Biotechnology* 21:1307–1313.
- Dutky, S. R. 1959. *Insect Microbiology*. *Advances in Applied Microbiology* 1:175–200.
- Dutky, S. R., and Hough, W. S. 1955. Note on a parasitic nematode from codling moth larvae, *Carpocapsae pomonella*. *Proceedings of the Entomological Society of Washington* 57:244.
- Fife, J. P., Derksen, R. C., Ozkan, H. E., and Grewal, P. S. 2003. Effects of pressure differentials on the viability and infectivity of entomopathogenic nematodes. *Biological Control*, 27:65–72.
- Fife, J. P., Derksen, R. C., Ozkan, H. E., Grewal, P. S., Chalmers, J. J., and Krause, C. R. 2004. Evaluation of a contraction flow field on hydrodynamic damage to entomopathogenic nematodes—a biological pest control agent. *Biotechnology and Bioengineering* 86:96–107.
- Fife, J. P., Ozkan, H. E., Derksen, R. C., Grewal, P. S., and Krause, C. R. 2005. Viability of a biological pest control agent through hydraulic nozzles. *Transactions of the American Society of Agricultural Engineers* 48:45–54.
- Fodor, A., Séringer, G., and Georgis, R. 1989. Application of insect pathogenic nematodes (*Neoalectana carpocapsae* and *Heterorhabditis* spp.) against Colorado potato beetle (*Leptinotarsa decemlineata*) larvae: Small plot experiments and laboratory trials. *Novenyved* 25:215.
- Friedman, M. J. 1990. Commercial production and development. Pp. 153–172 in R. Gaugler, and H. K. Kaya, eds. *Entomopathogenic nematodes in Biological Control*. Boca Raton, CRC Press.
- Gaugler, R., and Campbell, J. F. 1991. Selection for enhanced host-finding of scarab larvae (Coleoptera: Scarabaeidae) in an entomopathogenic nematode. *Environmental Entomology* 20:700–706.
- Gaugler, R., Campbell, J. F., and McGuire, T. R. 1989. Selection for host-finding in *Steinernema feltiae*. *Journal of Invertebrate Pathology* 54:363–372.
- Gaugler, R., Grewal, P., Kaya, H., and Smith-Fiola, D. 2000. Quality assessment of commercially produced entomopathogenic nematodes. *Biological Control* 17:100–109.
- Georgis, R. 1990. Formulation and application technology. Pp. 173–191 in R. Gaugler, and H. K. Kaya, eds. *Entomopathogenic Nematodes in Biological Control*. Boca Raton: CRC Press.
- Georgis, R., and Poinar, G. O., Jr. 1983a. Effect of soil texture on the distribution and infectivity of *Neoalectana carpocapsae* (Nematoda: Steinernematidae). *Journal of Nematology* 15:308–311.
- Georgis, R., and Poinar, G. O., Jr. 1983b. Effect of soil texture on the distribution and infectivity of *Neoalectana glaseri* (Nematoda: Steinernematidae). *Journal of Nematology* 15:329–332.
- Georgis, R., and Poinar, G. O., Jr. 1983c. Vertical migration of *Heterorhabditis bacteriophora* and *H. heliothidis* (Nematoda: Heterorhabditidae). *Journal of Nematology* 15:652–654.
- Georgis, R., Kaya, H. K., and Gaugler, R. 1991. Effect of steinernematid and heterorhabditid nematodes (Rhabditida: Steinernematidae and Heterorhabditidae) on non-target arthropods. *Environmental Entomology* 20:815–822.
- Georgis, R., Koppenhöfer, A. M., Lacey, L. A., Bélair, G., Duncan, L. W., Grewal, P. S., Samish, M., Tan, L., Torr, P., and van Tol, R. W. H. M. 2006. Successes and failures in the use of parasitic nematodes for pest control. *Biological Control* 38:103–123.
- Glaser, R. W. 1932. Studies on *Neoalectana glaseri*, a nematode parasite of the Japanese beetle (*Popillia japonica*). New Jersey Department of Agriculture, Circular 211:1–34.
- Glaser, R. W., and Fox, H. 1930. A nematode parasite of the Japanese beetle (*Popillia japonica* Newm.). *Science* 71:16–17.
- Glaser, R. W., and Farrell, C. C. 1935. Field experiments with the Japanese beetle and its nematode parasite. *Journal of the New York Entomological Society* 43:345–371.
- Grewal, P. S. 2000. Enhanced ambient storage stability of an entomopathogenic nematode through anhydrobiosis. *Pest Management Science* 56:401–406.
- Grewal, P. S., Selvan, S., and Gaugler, R. 1994a. Thermal adaptation of entomopathogenic nematodes: niche breadth for infection, establishment, and reproduction. *Journal of Thermal Biology* 19:245–253.
- Grewal, P. S., Lewis, E. E., Campbell, J. F., and Gaugler, R. 1994b. Host finding behavior as a predictor of foraging strategy for entomopathogenic nematodes. *Parasitology* 108:207–215.
- Grewal, P. S., Converse, V., and Georgis, R. 1999. Influence of production and bioassay methods on infectivity of two ambush foragers (Nematoda: Steinernematidae). *Journal of Invertebrate Pathology* 73:40–44.
- Grewal, P. S., Ehlers, R. U., and Shapiro-Illan, D. I. 2005. Critical issues and research needs for expanding the use of nematodes in

- biocontrol. Pg. 479–489 in P. S. Grewal, R. U. Ehlers, and D. Shapiro-Ilan, eds. *Nematodes as Biocontrol Agents*. Wallingford, UK: CABI Publishing.
- Grunder, J. M., Ehlers, R.-U., and Jung, K. 2005. Quality Control of Entomopathogenic Nematodes. COST Action 819, Agroscope Faw, Wädenswil, Switzerland.
- Hackett, K. J., and Poinar, G. O., Jr. 1973. The ability of *Neoapectana carpocapsae* Weiser (Steinernematidae: Rhabditoidea) to infect adult honeybees (*Apis mellifera*, Apidae: Hymenoptera). *American Bee Journal* 113:100.
- Hashmi, S., Hasnmi, G., and Gaugler, R. 1995. Genetic transformation of an entomopathogenic nematode by microinjection. *Journal of Invertebrate Pathology* 66:293–296.
- Jansson, R. K., and Lecrone, S. H. 1994. Application methods for entomopathogenic nematodes (Rhabditida: Heterorhabditidae) aqueous suspensions versus infected cadavers. *Florida Entomologist* 77:281–284.
- Kaya, H. K., and Nelson, C. E. 1985. Encapsulation of steinernematod and heterorhabditid nematodes with calcium alginate; a new approach for insect control and other applications. *Environmental Entomology* 14:572–574.
- Kermarrec, A., and Mauléon, H. 1985. Potential noxiousness of the entomogenous nematode *Neoapectana carpocapsae* Weiser to the Antillean toad *Bufo marinus* L. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit (Gent)* 50:831–838.
- Kiontke, K., Barrière, A., Kolotuev, I., Podbilewicz, B., Sommer, R., Fitch, D. H. A., and Félix, M.-A. 2007. Trends, Stasis, and Drift in the evolution of nematode vulva development. *Current Biology* 17:1925–1937.
- Koppenhöfer, A. M., and Grewal, P. S. 2005. Interactions with other biological control agents and agrochemicals. Pg. 363–381 in P. S. Grewal, R. U. Ehlers, and D. Shapiro-Ilan, eds. *Nematodes as Biocontrol Agents*. Wallingford, UK, CABI Publishing.
- Lello, E. R., Patel, M. N., Matthews, G. A., and Wright, D. J. 1996. Application technology for entomopathogenic nematodes against foliar pests. *Crop Protection* 15:567–574.
- Lewis, E. E., Gaugler, R., and Harrison, R. 1992. Entomopathogenic nematode host finding: response to host contact cues by cruise and ambush foragers. *Parasitology* 105:309–319.
- Liu, J., Poinar, G. O., Jr., and Berry, R. E. 2000. Control of insect pests with entomopathogenic nematodes: The impact of molecular biology and phylogenetic reconstruction. *Annual Review of Entomology* 45:287–306.
- Manweiler, S. A. 1994. Development of the first cat flea biological control product employing the entomopathogenic nematode *Steinernema carpocapsae*. Pp.1005–1012 in *Proceedings of the Brighton Crop Protection Conference: Pest and Diseases*, Farnham, Surrey: British Crop Protection Council.
- Menzler-Hokkanen, I., and Hokkanen, H. M. T. 2004. Developing entomopathogenic nematode delivery systems for biological control of oilseed rape pests. *International organization for Biological and integrated control of noxious animals and plants, West Palaearctic Regional Section, Bulletin* 28:19–22.
- Miller, R. W. 1989. Novel pathogenicity assessment technique for *Steinernema* and *Heterorhabditis* entomopathogenic nematodes. *Journal of Nematology* 21:574.
- Milstead, J. E. 1977. The life cycle and pathobiology of *Heterorhabditis bacteriophora* Poinar (Rhabditoidea: Nematoda) in its lepidopteran hosts. Ph.D. dissertation. University of California, Berkeley, 113 pp.
- Moyle, P. L., and Kaya, H. K. 1981. Dispersal and infectivity of the entomogenous nematode, *Neoapectana carpocapsae* Weiser (Rhabditida: Steinernematidae), in sand. *Journal of Nematology* 13: 295–300.
- Poinar, G. O., Jr. 1966. The presence of *Achromobacter Nematophilus* Poinar and Thomas in the infective stage of a *Neoapectana* sp. (Steinernematidae: Nematoda). *Nematologica* 12:105–108.
- Poinar, G. O., Jr. 1967. Description and Taxonomic position of the DD-136 nematode (Steinernematidae, Rhabditoidea) and its relationship to *Neoapectana carpocapsae* Weiser. *Proceedings of the Helminthological Society of Washington* 34:199–209.
- Poinar, G. O., Jr. 1969. Arthropod immunity to worms. Pg. 173–210 in G. J. Jackson, and R. Herman, eds. *Immunity to Parasitic Animals*. Vol. 1. New York: Appleton-Century Crofts Co.
- Poinar, G. O., Jr. 1976. Description and biology of a new insect parasitic rhabditoid, *Heterorhabditis bacteriophora* n. gen., n. sp. (Heterorhabditidae, n. fam.) (Rhabditida [Oerley]). *Nematologica* 21:463–470.
- Poinar, G. O., Jr. 1978. Generation polymorphism in *Neoapectana glaseri* Steiner (Steinernematidae: Nematoda), redescribed from *Strigoderma arboricola* (Fab.) (Scarabaeidae: Coleoptera) in North Carolina. *Nematologica* 24:105–114.
- Poinar, G. O., Jr. 1979. *Nematodes for Biological Control of Insects*. CRC Press, Boca Raton. 277 pp.
- Poinar, G. O., Jr. 1986. Recognition of *Neoapectana* species (Steinernematidae: Rhabditida). *Proceedings of the Helminthological Society of Washington* 53:121–129.
- Poinar, G. O., Jr. 1988. A microsporidian parasite of *Neoapectana glaseri* (Steinernematidae: Rhabditida). *Revue de Nematologie* 11:359–361.
- Poinar, G. O., Jr. 1991. Genetic engineering of nematodes for pest control. Pg. 77–93 in K. Maramorosch, ed. *Biotechnology for Biological Control of Pests and Vectors*. Boca Raton: CRC Press.
- Poinar, G. O., Jr. 1990. Taxonomy and biology of Steinernematidae and Heterorhabditidae. Pg. 23–61 in R. Gaugler and H. K. Kaya, eds. *Entomopathogenic Nematodes in Biological Control*. Boca Raton: CRC Press.
- Poinar, G. O., Jr. 2011. *The Evolutionary History of nematodes*. Brill, Leiden, 439 pp.
- Poinar, G. O., Jr., and Thomas, G. M. 1965. A new bacterium, *Achromobacter Nematophilus* sp. nov. (Achromobacteriaceae Eubacteriales) associated with a nematode. *International Bulletin of Bacterial Nomenclature and Taxonomy* 15:249–252.
- Poinar, G. O., Jr., and Thomas, G. M. 1966. Significance of *Achromobacter nematophilus* (Achromobacteriaceae Eubacteriales) in the development of the nematode, DD-136 (*Neoapectana* sp. Steinernematidae). *Parasitology* 56:385–390.
- Poinar, G. O., Jr., and Thomas, G. M. 1967. The Nature of *Achromobacter nematophilus* as an insect pathogen. *Journal of Invertebrate Pathology* 9:510–514.
- Poinar, G. O., Jr., and Leutenegger, R. 1968. Anatomy of the infective and normal third stage juveniles of *Neoapectana carpocapsae* Weiser. *Journal of Parasitology* 54:340–350.
- Poinar, G. O., Jr., and Veremtshuk, G. V. 1970. A new strain of entomopathogenic nematodes and geographical distribution of *Neoapectana carpocapsae* Weiser (Rhabditida, Steinernematidae). *Zoological Zhurnal* 49:966–969.
- Poinar, G. O., Jr., and Brooks, W. M. 1977. Recovery of the entomogenous nematode, *Neoapectana glaseri* Steiner from a native insect in North Carolina. *International Research Communications System Medical Science* 5:473.
- Poinar, G. O., Jr., and Hom, A. 1986. Survival and horizontal movement of infective stage *Neoapectana carpocapsae* in the field. *Journal of Nematology* 18:34–36.
- Poinar, G. O., Jr., and Jansson, H. B. 1986a. Susceptibility of *Neoapectana* spp. and *Heterorhabditis heliothidis* to the endoparasitic fungus *Drechmeria coniospora*. *Journal of Nematology* 18:225–230.
- Poinar, G. O., Jr., and Jansson, H.-B. 1986b. Infection of *Neoapectana* and *Heterorhabditis* (Rhabditida: Nematoda) with the predatory fungi, *Monacrosporium ellipsosporum* and *Arthrobotrys oligospora* (Moniliales: Deuteromycetes). *Revue du Nematologie* 9:241–244.
- Poinar, G. O., Jr., and Thomas, G. M. 1988. Infection of frog tadpoles (Amphibia) by insect parasitic nematodes (Rhabditida). *Experientia* 44:528–531.

- Poinar, G. O., Jr., and Georgis, R. 1990. Characterization and field application of *Heterorhabditis bacteriophora* strain HP88 (Heterorhabditidae: Rhabditida). *Revue de Nématologie* 13:387–393.
- Poinar, G. O., Jr., Thomas, G. M., and Hess, R. 1977. Characteristics of the specific bacterium associated with *Heterorhabditis bacteriophora* (Heterorhabditidae: Rhabditida). *Nematologica* 23: 97–102.
- Poinar, G. O., Jr., Thomas, G., Haygood, M., and Neelson, K. H. 1980a. Growth and luminescence of the symbiotic bacteria associated with the terrestrial nematode, *Heterorhabditis bacteriophora*. *Soil Biology and Biochemistry* 12:5–10.
- Poinar, G. O., Jr., Hess, R. T., and Thomas, G. 1980b. Isolation of defective bacteriophages from *Xenorhabdus* spp. (Enterobacteriaceae). *IRCS Medical Science* 8:141.
- Poinar, G. O., Jr., Evans, J. S., and Schuster, E. 1983. Field test of the entomogenous nematode, *Neoaplectana carpocapsae*, for control of corn rootworm larvae (*Diabrotica* sp., Coleoptera). *Protection Ecology* 5:337–342.
- Poinar, G. O., Jr., Hess, R. T., Lanier, W., Kinney, S., and White, J. H. 1989. Preliminary observations of a bacteriophage infecting *Xenorhabdus luminescens* (Enterobacteriaceae). *Experientia* 45:191–192.
- Poinar, G. O., Jr., Ferro, C., Morales, A., and Tesh, R. B. 1993. *Anandranema phlebotophaga* n.gen., n.sp. (Allantonematidae: Tylenchida), a new nematode parasite of phlebotomine sand flies (Psychodidae: Diptera) with notes on experimental infections of these insects with parasitic rhabditoids. *Fundamental and Applied Nematology* 16:11–16.
- Pye, A. E., and Pye, N. L. 1985. Different applications of the insect parasitic nematode *Neoaplectana carpocapsae* to control the large pine weevil, *Hylobius abietis*. *Nematologica* 31:109–116.
- Rao, P. S. P., Das, P. K., and Pahdi, G. 1975. Note on compatibility of DD-136 (*Neoaplectana carpocapsae*) Dutky, an insect parasitic nematode with some insecticides and fertilizers. *Indian Journal of Agricultural Science*. 45:275–277.
- Reed, E. M., and Wallace, H. R. 1965. Leaping locomotion by an insect-parasitic nematode. *Nature* 206:210–211.
- Reed, D. K., Reed, G., and Creighton, C. S. 1986. Introduction of entomogenous nematodes into trickle irrigation systems to control striped cucumber beetle (Coleoptera, Chrysomelidae). *Journal of Economic Entomology* 79:1330–1333.
- Rovesti, L., and Deseö, K. V. 1989. Effect of neem kernel extract on steinernematode and heterorhabditid nematodes. *Nematologica* 35: 493–496.
- Rovesti, L., and Deseö, K. V. 1990. Compatibility of chemical pesticides with the entomopathogenic nematodes, *Steinernema carpocapsae* Weiser and *Steinernema feltiae* (Nematoda, Steinernematodae). *Nematologica* 36:237–245.
- Rovesti, L., and Deseö, K. V. 1991. Compatibility of pesticides with the entomopathogenic nematode, *Heterorhabditis heliothidis*. *Nematologica* 37:113–116.
- Samish, M., and Glazer, I. 1991. Killing ticks with parasitic nematodes of insects. *Journal of Invertebrate Pathology* 58:281–282.
- Sandhu, S. K., Jagdale, G. B., Hogenhout, S. A., and Grewal, P. S. 2006. Comparative analysis of the expressed genome of the infective juvenile entomopathogenic nematode, *Heterorhabditis bacteriophora*. *Molecular Biochemistry and Parasitology* 145:239–244.
- Shetlar, D. J. 1999. Application methods in different cropping systems. Pg. 31–36 in S. Polavarapu, ed. *Proceedings of the National Workshop on optimal use of insecticidal nematodes in pest management*. Chatsworth, NJ: Rutgers University.
- Shapiro-Ilan, D. I., Lewis, E. E., Behle, R. W., and McGuire, M. R. 2001. Formulation of entomopathogenic nematode-infected cadavers. *Journal of Invertebrate Pathology* 78:17–23.
- Silver, S. C., Dunlop, D. B., and Grove, D. I. 1995. Granular formulation of biological entities with improved storage stability. International Patent No. WO 95/05077.
- Silverman, J., Platzer, E. G., and Rust, M. K. 1982. Infection of the cat flea, *Ctenocephalides felis* (Bouche) by *Neoaplectana carpocapsae* Weiser. *Journal of Nematology* 14:394–397.
- Simons, W. R., and Poinar, F. O., Jr. 1973. The ability of *Neoaplectana carpocapsae* (Steinernematidae: Nematodea) to survive extended periods of desiccation. *Journal of Insect Pathology* 22:228–230.
- Steiner, G. 1923. *Aplectana krausseii* n. sp., einer in der Blattwespe *Lyda* sp. parasitierende Nematodenform, nebst Bemerkungen über das Seitenorgan der parasitischen nematoden. *Zentralblatt fuer Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene* 59:14–18.
- Steiner, G. 1929. *Neoaplectana glaseri* n. g., n. sp. (Oxyuridae) a new nematode parasite of the Japanese beetle (*Popillia japonica* Newm.). *Journal of the Washington Academy of Science*. 19:436–440.
- Stock, S. P., and Hunt, D. J. 2005. Morphology and systematics of Nematodes used in Biocontrol. In Grewal, P. S., Ehlers, R-U., and Shapiro-Ilan, D. I. *Nematodes as biocontrol agents*. Wallingford: CABI Publishing.
- Thomas, G. M., and Poinar, G. O., Jr. 1979. *Xenorhabdus* gen. nov., a genus of entomopathogenic, nematophilic bacteria of the family Enterobacteriaceae. *International Journal of Systematic Bacteriology* 29:352–360.
- Veremchuk, G. V., and Issi, I. V. 1970. The development of microsporidians of insects in the entomopathogenic nematode *Neoaplectana carpocapsae* (Nematoda: Steinernematodidae). *Parazitologiya* 4:3–7.
- Webster, J. M., and Bronskill, J. F. 1968. Use of Gelgard M and an evaporation retardant to facilitate control of larch sawfly by a nematode-bacterium complex. *Journal of Economic Entomology* 61:1370–1371.
- Webster, J. M., Chen, G., Hu, K., and Li, J. 2002. Bacterial metabolites. Pp. 99–114 in R. Gaugler, ed. *Entomopathogenic Nematology*. Oxon, UK: CAB International.
- Weiser, J. 1955. *Neoaplectana carpocapsae* n. sp. (Anguillata, Steinernematidae) nový cizopasník housenek obalece jablecneho, *Carpocapsae pomonella* L. *Vestník Československe Spolecnosti Zoologicke* 19:44–52.
- Weiss, M., Glazer, I., Mumcuoglu, K. Y., Elking, Y., and Galun, R. 1993. Infectivity of steinernematid and heterorhabditid nematodes for the human body louse *Pediculus humanus humanus* (Anoplura: Pediculidae). *Fundamental and Applied Nematology* 16:489–493.
- Welch, H. E., and Bronskill, J. F. 1962. Parasitism of mosquito larvae by the nematode, DD-136 (Nematoda: Neoaplectanidae). *Canadian Journal of Zoology* 40:1263–1268.
- Wright, L. C., Witkowski, J. F., Echtenkamp, G., and Georgis, R. 1993. Efficacy and persistence of *Steinernema carpocapsae* (Rhabditida: Steinernematidae) applied through center-pivot irrigation system against larval corn rootworms (Coleoptera: Chrysomelidae). *Journal of Economic Entomology* 86:148–1354.
- Wright, P. J. 1990. Morphological characterization of the entomogenous nematodes *Steinernema* spp. and *Heterorhabditis* spp. (Nematoda: Rhabditida). *New Zealand of Zoology* 17:577–586.