

Article (refereed)

Austnes, Kari; Evans, Chrisptoher D.; Eliot-Laize, Caroline; Naden, Pamela S.; Old, Gareth H.. 2010. Effects of storm events on mobilisaton and in-stream processing of dissolved organic matter (DOM) in a Welsh peatland catchment. *Biogeochemistry*, 99 (1-3). 157-173. [10.1007/s10533-009-9399-4](https://doi.org/10.1007/s10533-009-9399-4)

© Springer Science+Business Media B.V. 2009

This version available <http://nora.nerc.ac.uk/8246/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the authors and/or other rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.

www.springerlink.com

Contact CEH NORA team at
noraceh@ceh.ac.uk

Editorial Manager(tm) for Biogeochemistry
Manuscript Draft

Manuscript Number: BIOG1172R1

Title: Effects of storm events on mobilisation and in-stream processing of dissolved organic matter (DOM) in a Welsh peatland catchment

Article Type: Manuscript

Keywords: DOC; DOM quality; Fluorescence; In-stream processes; Peat stream; Storm events

Corresponding Author: Dr. Kari Austnes, Ph.D.

Corresponding Author's Institution:

First Author: Kari Austnes, Ph.D.

Order of Authors: Kari Austnes, Ph.D.; Christopher D Evans; Caroline Eliot-Laize; Pamela S Naden; Gareth H Old

Abstract: Peatlands are important contributors of dissolved organic matter (DOM) to downstream aquatic systems. We investigated the effects of storm events on dissolved organic carbon (DOC) concentrations and DOM quality in a stream draining a Welsh peatland catchment. Intensive stream samples were collected and analysed for pH, DOC, dissolved organic nitrogen (DON), absorbance and fluorescence. Soil water samples and samples of sphagnum pore water were also collected, and a simple end-member mixing model was applied to account for changes occurring during the events. Fluorescence data were interpreted using parallel factor analysis (PARAFAC). DOC concentrations increased and pH decreased during the storm events. The soil water data and the mixing model indicated that this was due to a change of flow paths and draining of the DOC-rich acrotelm. Absorbance indices and the DOC/DON ratio suggested that the DOM released during events was less degraded. There was a striking, inversely related diurnal pattern in absorbance and fluorescence after the discharge peak. The diurnal pattern and a lack of fit with the mixing model suggested that fluorescing DOM was mainly produced in-stream. Fluorescence has been found to peak in the morning and decline during day-time due to photo-bleaching. We hypothesise that the input of additional DOM during events causes a change in the diurnal pattern, giving a peak at mid-day, when the processing of the additional DOM is highest.

Response to Reviewers: Dear editor/reviewer

The manuscript "Effects of storm events on mobilisation and in-stream processing of dissolved organic matter (DOM) in a Welsh peatland catchment" has been revised according to the reviewer's comments.

The main issue addressed in the review was the potential iron interference of the spectrophotometric analyses. We have now explored this issue thoroughly, and our conclusion is that there may have been some iron interference, but that this does not appear to affect the interpretation - our main focus was temporal patterns throughout the events, not absolute levels, and these patterns were hardly affected by the iron correction. The issue has been addressed in the methods (lines 173-195 and results (lines 351-364+394-401) sections. For absorbance and sUVa, Fe corrected values were included in the revised paper, based on a published equation (Weishaar et al 2003). The fluorescence estimate was based on a similar regression, but it was more crude as we had to produce the equation ourselves

based on the data in the paper by Ohno et al. (2008). Hence, we did not feel it would be appropriate to present these data, but the effect was much the same as that for absorbance, and again did not alter the interpretation. Having accounted for Fe interference in our revised analysis, described this in the methods, and found that it did not alter the interpretation, we did not feel there was a need to alter or expand the discussion and conclusions further on this subject (cf specific comment 11). However, further detail could be included if requested by the editor.

Specific comments:

2. The filters were rinsed with both water and sample before proper filtering, to exclude the possibility of filter bleed. A description of this has been included in the text (lines 155-156 and 160).
3. The two different filters were used for two different aliquots of the sample. This has now been made clearer in the text (lines 157-158). Due to risk of contamination, samples to be analysed for fluorescence had to be filtered at CEH Wallingford according to their procedures. The different filter size was also according to their procedures. 1.2 μm is sufficient to remove substituents that can cause scattering. There is hardly any difference in fluorescence between 0.45 and 1.2 μm , as the important region is around 0.1 μm , where microbial and cellular material is removed from the solution.
4. sUVa and E2/E3 are not specific measures of particular properties of DOM, but are correlated with some properties. This is thoroughly explained at the beginning of the discussion on DOM quality. In our opinion it would be confusing and too complicated to include this information in the methods section. The interpretation of absorbance or fluorescence analyses are not explained either, for the same reason. It is merely stated in the introduction that spectrophotometric analysis gives information on DOM quality (lines 103-107).
5. The fluorescence exhibited by two of the sphagnum samples may indicate some protein-like material. The excitation wavelength fits with that of tyrosine or especially tryptophan, but the emission wavelength was somewhat lower. However, as explained (lines 284-285) this was not a significantly strong feature across all samples for a component in this region to be validated. PARAFAC extracts components in all regions of the EEM measured, so if this was a strong feature, it would have been expressed in a valid component. As stated in lines 371-374 some overlap with the fluorescence caused by contamination in some of the samples may have reduced the chances of validating a component in this region, but most likely not, as it was only a slight overlap between the regions of true and contaminant fluorescence. Hence, as a true component in this region was not validated, we did not include a thorough discussion of what this component, if validated, would have represented, even though it was most likely some protein-like material.
6. This has already been addressed.
7. Yes, we did measure blank samples from these tubes. This has now been stated in lines 370-371.
8. The point of the mixing model was merely to give an indication of the shape of the end-member variation through the event, not the exact proportions. The mixing model also highlights the strong deviation of fluorescence from anything resembling conservative mixing. We consequently think the mixing model serves as a good illustration, and would like to keep it. A couple of lines (439-441) on the reliability of the different models were included.
9. True, this is not really a good way to express this. The main reason to include E2/E3 is that it is a different type of parameter, and when both parameters indicate the same change in properties, this strengthens the conclusion. E2/E3 is in fact also a more robust parameter, because there is less risk of contamination when only using spectrophotometric analysis (a line on this has been included in the text, line 457). However, more readers will be familiar with the sUVa index, so it would not be a good alternative to use E2/E3 only. As explained above, there is no reason to believe that iron interference would have a strong impact on E2/E3.
10. The mechanism is not clear here, as it cannot be drawn from the data themselves. However, it is not likely to be related to sorption in the mineral soil as the catchment is strongly peat-dominated. Sorption to organic soil is more likely, but we have no evidence to support such a mechanism.
11. The discussion has been shortened wherever possible. The main changes have been made to the first section, Effects of events on DOC concentration. Here the last two paragraphs have been removed.

We found this discussion interesting, but removing them makes the manuscript more focused. Accordingly, some details have been removed from the results chapter, as these were mainly a basis for this discussion. No complete paragraphs have been removed from the last two sections of the discussion, as these are the most important parts. The third section has been left as it was, while the second section has been modified to make the discussion shorter and clearer.

Figures: Figure 7 is mainly an example, and can be removed. We would like to keep the rest of the figures, as they give important background, or serve as useful illustrations to the reader. The figure numbers are thus changed for figures 8-11, to 7-10. Some cosmetic changes have been made to figures 3-5+7.

Yours sincerely
Kari Austnes

1 **Effects of storm events on mobilisation and in-stream processing of dissolved**
2 **organic matter (DOM) in a Welsh peatland catchment**

3

4 K. Austnes^{1*}, C.D. Evans², C. Eliot-Laize³, P.S. Naden⁴ and G.H. Old⁴

5

6 1. Department of Plant and Environmental Sciences, Norwegian University of Life
7 Sciences, P.O. Box 5003, N-1432 Ås, Norway. Current address: Norwegian Institute
8 for Water Research, Gaustadalléen 21, N-0349 Oslo, Norway

9 2. Centre for Ecology and Hydrology Bangor, Environment Centre Wales, Deiniol
10 Road, Bangor, Gwynedd, LL57 2UW, UK

11 3. Centre for Ecology and Hydrology Wallingford, Maclean Building, Benson Lane,
12 Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK. Current address:
13 Premier Analytical Services, Microscopy, The Lord Rank Centre, Lincoln Road,
14 High Wycombe, Buckinghamshire, HP12 3QR, UK

15 4. Centre for Ecology and Hydrology Wallingford, Maclean Building, Benson Lane,
16 Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

17

18 *Corresponding author:

19 Tel.: +47 02348

20 Fax: +47 22185200

21 E-mail address: kari.austnes@niva.no

22

23 Running head: Storm events and DOM effects in a peatland catchment

24 Article type: Original paper

25

26 Key words: DOC, DOM quality, Fluorescence, In-stream processes, Peat stream,
27 Storm events

28

29 **Abstract**

30

31 Peatlands are important contributors of dissolved organic matter (DOM) to
32 downstream aquatic systems. We investigated the effects of storm events on dissolved
33 organic carbon (DOC) concentrations and DOM quality in a stream draining a Welsh
34 peatland catchment. Intensive stream samples were collected and analysed for pH,
35 DOC, dissolved organic nitrogen (DON), absorbance and fluorescence. Soil water
36 samples and samples of sphagnum pore water were also collected, and a simple end-
37 member mixing model was applied to account for changes occurring during the
38 events. Fluorescence data were interpreted using parallel factor analysis (PARAFAC).
39 DOC concentrations increased and pH decreased during the storm events. The soil
40 water data and the mixing model indicated that this was due to a change of flow paths
41 and draining of the DOC-rich acrotelm. Absorbance indices and the DOC/DON ratio
42 suggested that the DOM released during events was less degraded. There was a
43 striking, inversely related diurnal pattern in absorbance and fluorescence after the
44 discharge peak. The diurnal pattern and a lack of fit with the mixing model suggested
45 that fluorescing DOM was mainly produced in-stream. Fluorescence has been found
46 to peak in the morning and decline during day-time due to photo-bleaching. We
47 hypothesise that the input of additional DOM during events causes a change in the
48 diurnal pattern, giving a peak at mid-day, when the processing of the additional DOM
49 is highest.

50

51 **Introduction**

52

53 Peatlands are important contributors of dissolved organic matter (DOM) to
54 downstream aquatic systems. Mean dissolved organic carbon (DOC) fluxes in rivers
55 draining peat dominated areas are higher than in rivers draining most other landscape
56 types, in the order of 50-100 kg ha⁻¹ yr⁻¹ (Aitkenhead and McDowell 2000).
57 Percentage peat cover is found to be a good predictor of DOC concentration across
58 large spatial scales (Aitkenhead et al. 1999).

59 The concentration of DOC is commonly found to correlate positively with
60 discharge in streams and rivers in temperate and boreal catchments (e.g. Thurman
61 1985, p. 50; Hope et al. 1994; Soulsby et al. 2003). Likewise, DOC concentrations
62 have frequently been found to increase during storm events (e.g. Hinton et al. 1997;
63 Buffam et al. 2001). However, most studies of discharge-DOC concentration
64 relationships have been conducted in catchments with organo-mineral or mixed soils
65 (Clark et al. 2007a). Given the significant contribution of peat systems to DOC
66 concentrations and fluxes, it is important to investigate the relationship between
67 discharge and not only DOC (as a measure of DOM quantity) but also DOM quality
68 in these systems. Studying these relationships is especially important in a climate
69 change perspective, given that annual precipitation, as well as extremes of daily
70 precipitation, are likely to increase in northern Europe (Christensen et al. 2007).

71 In streams draining organo-mineral soils, the increased DOC concentration at
72 high discharge is commonly explained by changes in flow paths towards increased
73 lateral flow through the upper horizons, where DOC concentrations are higher (e.g.
74 McDowell and Likens 1988; Boyer et al. 1997; Hinton et al. 1998; Inamdar et al.
75 2006). Change in flow path can be an important control on DOC concentration in peat

76 streams as well, but seems to depend upon the hydrological connectivity between the
77 peat and the underlying bedrock/mineral soil (Clark et al. 2007a, b). Peat soils consist
78 of an upper horizon (the acrotelm) with roots and decomposing plant material, and a
79 lower horizon (the catotelm) with dense peat (Evans et al. 1999). The stored carbon in
80 the catotelm is hydrologically disconnected from the stream (Billett et al. 2006).
81 Where there is hydrological connectivity between the peat and the mineral soil, base
82 flow is characterised by alkaline DOC poor groundwater (Worrall et al. 2002; Clark et
83 al. 2007a). As the water table rises during an event, resulting in subsurface flow in the
84 acrotelm, stream DOC concentration increases due to the input of DOC rich soil water
85 (Evans et al. 1999; Worrall et al. 2002; Soulsby et al. 2003; Clark et al. 2007a, b).
86 However, further progress of the event may introduce low DOC, rain-like water
87 (Clark et al. 2007b), due to exhaustion of the acrotelm or the occurrence of saturated
88 overland flow or macropore flow (Evans et al. 1999; Worrall et al. 2002). Hence,
89 stream concentration can be described by mixing of water from three different source
90 areas, so-called end-members (Worrall et al. 2002). In other peat catchments, base
91 flow is chemically similar to the acidic DOC rich soil water in the acrotelm (Clark et
92 al. 2007b). In these systems stream water during events derives from two end-
93 members only: soil water and the rain-like water (Clark et al. 2007a). This causes a
94 decrease in stream DOC concentration during events (Clark et al. 2007a; Eimers et al.
95 2008).

96 Literature on the effects of storm events on DOM quality is sparse. If storm
97 events cause changes of flow paths and not merely dilution, one would assume a
98 change in the quality of the DOM released to the stream (Buffam et al. 2001). DO^{14}C
99 data from peat moorland in the upper Conwy catchment showed that base flow
100 releases old, soil-derived DOC, whereas high flow releases younger DOC, probably

101 derived from recent plant material (Evans et al. 2007). Probably, a difference in age
102 also implies different degree of degradation, which in turn affects DOM quality
103 (Qualls and Haines 1992; Kalbitz et al. 2003; Saadi et al. 2006). Spectrophotometric
104 measurement of absorbance and fluorescence can give information on DOM quality in
105 terms of chemical characteristics of the organic material and its bioavailability (e.g.
106 Senesi 1990; Peuravuori and Pihlaja 1997; Croué et al. 1999; McKnight et al. 2001;
107 Kalbitz et al. 2003; Fellman et al. 2008). To our knowledge fluorescence scanning of
108 high frequency event samples has not previously been conducted.

109 In the present study high frequency measurement of chemical and
110 spectrophotometric characteristics of high frequency stream water samples was
111 conducted in a peatland catchment in Wales. The objectives of the study were: 1) To
112 investigate the effect of storm events on DOC concentration, 2) to investigate the
113 effects of storm events on DOM quality, and 3) to explore the usefulness of
114 fluorescence analyses to assess short term changes in DOM quality.

115

116 **Methods**

117

118 *Field site*

119

120 The study site is a small (3.2 km²) peatland catchment in north Wales drained by the
121 stream Afon Ddu (Fig. 1). The catchment is a sub-catchment of the upper Conwy
122 catchment, and it is part of the large Migneint blanket peatland area. Catchment
123 altitude ranges from 440 to 540 m.a.s.l., with the main part being below 500 m.a.s.l..
124 Average temperature is 8°C and mean annual rainfall is 2300 mm. Hydrological
125 response to rainfall events is rapid (Billett et al. 2007). The bedrock is volcanic, mixed

126 rhyolitic and basaltic, of Ordovician origin. Peat soils (histosols) dominate, but at
127 elevated sites podzols occur. The soil depth is usually around 1-2 m, but may be up to
128 5 m. The vegetation is dominated by heather (*Calluna vulgaris* L.) and *Sphagnum*
129 spp. mosses, with scattered grasses and sedges. There is some drainage ditching in the
130 catchment dating from the 1930s-1960s.

131

132 *Water sampling*

133

134 Intensive high-flow event sampling in the Afon Ddu was conducted in the autumn of
135 2007 (sampling point at Ordnance Survey coordinates 277520,344920). Samples were
136 collected during autumn because DOC concentrations were expected to be high (Clark
137 et al. 2007b). Samples were collected during three events (September 17th-18th,
138 October 3rd-4th, October 26th-30th), hereafter referred to as events 1, 2 and 3. During
139 events 1 and 2, samples were collected for 24 hours (event 1 two-hourly, event 2
140 hourly), while during the larger event 3 samples were collected two-hourly for 96
141 hours. Samples were collected using a Xian 1000 autosampler (Hach Lange,
142 Germany).

143 Samples were collected from a number of nearby locations during each event,
144 to provide information on potential end-members. These were sphagnum pore water
145 samples collected by manual squeezing, and three different types of soil water
146 samples collected from blanket peat: bulked soil water collected at 5 cm using micro-
147 rhizon suction samplers and soil water collected at 5 and 10 cm using zero-tension
148 lysimeters.

149

150

151 *Analyses*

152

153 All samples were split in three for different analyses. The aliquot that was to be
154 analysed for carbon and nitrogen was filtered immediately (within maximum 2 days)
155 through Whatman sterile 0.45µm cellulose nitrate filters. The filters were rinsed with
156 deionised water and an aliquot of sample. All samples were stored at <3°C.

157 The event sample aliquots that were to be analysed for fluorescence and
158 absorbance were brought to CEH Wallingford as soon as possible (within 0-5 days).
159 Prior to analysis all samples were filtered through Whatman 1.2 µm glass microfibre
160 GFC filters, after rinsing with water and an aliquot of sample. All equipment used was
161 acid washed. The water used was purified with a NANOpure DIamond Analytical and
162 UV Systems from Barnstead. One cm quartz cells were used. Fluorescence analyses
163 were performed on a Varian Cary Eclipse instrument. Several blanks were run, to
164 ensure that the equipment used had no residual fluorescence. The samples were
165 scanned for emission from 280 to 500 nm at excitation 200 to 400 nm. Absorbance
166 was measured on a Varian Cary 50 instrument, measuring the spectrum from 200 to
167 800 nm. Whenever the absorbance was above 0.3 cm⁻¹, the samples were diluted to
168 below this level, and both analyses repeated, as inner-filtering correction (see below)
169 has been found to be insufficient above this level (Ohno 2002). The sUVa index was
170 calculated as absorbance measured at 254 nm divided by DOC (mg l⁻¹) (Vogt et al.
171 2004) and the E2/E3 index as absorbance measured at 250 nm divided by absorbance
172 measured at 365 nm (Peuravuori and Pihlaja 1997).

173 Both absorbance and fluorescence may be affected by iron interference.
174 Absorbance may be overestimated, as Fe ions absorb light (Weishaar et al. 2003).
175 Fluorescence may be underestimated, due to quenching of the fluorescence signal

176 caused by Fe-DOM complexation (Zepp et al. 2004; Ohno et al. 2008). To investigate
177 possible iron interference, data from biweekly sampling (November 2006 to January
178 2008) at the same spot were used, as Fe concentration was not analysed in the event
179 samples. Fe in the biweekly samples were analysed by ICP-OES. Preliminary
180 analyses showed that the Fe concentration was positively linearly related to DOC
181 concentration ($R^2 = 0.59$). This is in line with Neal et al. (2008), who observed strong
182 correlation between Fe and DOC concentration across a range of UK sites. However,
183 there was also a negative linear relationship with $\log(\text{flow})$ (Environment Agency
184 discharge data for the River Conwy at Cwm Llanerch
185 (http://www.nwl.ac.uk/ih/nrfa/station_summaries/066/011.html) ($R^2 = 0.30$). Hence,
186 Fe concentration could be reasonably well explained ($R^2 = 0.78$) using multiple linear
187 regression with DOC concentration and $\log(\text{flow})$ as explanatory variables. A linear
188 relationship ($R^2 = 0.82$) was established between Cwm Llanerch flow and Afon Ddu
189 stage (data from August to November 2007). Fe concentration in the event samples
190 was estimated using the multiple linear regression equation with event sample DOC
191 concentration and Cwm Llanerch flow (calculated from Afon Ddu stage) as inputs.
192 The contribution of Fe to absorbance at 254 nm was estimated using the equation for a
193 pure Fe^{3+} solution in Weishaar et al. (2003). Weishaar et al. showed that the
194 absorbances of Fe and DOM are additive. A crude estimate of Fe quenching of
195 fluorescence was done based on data in Ohno et al. (2008).

196 The remaining analyses on event samples were conducted at CEH Bangor.
197 Unfiltered samples were analysed for pH using a Metrohm SM 702 Titrino. Filtered
198 samples were analysed for total dissolved nitrogen (TDN) and non purgeable organic
199 carbon (NPOC) by elemental analysis using a Thermalox TOC/TN Analyser. The
200 purging prior to total carbon analysis removes inorganic carbon as CO_2 (by addition

201 of 11 μl of 1M HCl and purging with oxygen for 90 sec), so that only dissolved
202 organic carbon (DOC) is determined. Nitrate-N ($\text{NO}_3\text{-N}$) and ammonium-N ($\text{NH}_4\text{-N}$)
203 were analysed by autoanalyser (Skalar SA-40). $\text{NO}_3\text{-N}$ was analysed by the
204 sulphaniamide/NEDA/Cd/Cu reduction method, with extinction at 540nm, and $\text{NH}_4\text{-}$
205 N by the Indol-phenol blue method, with extinction at 660nm. Dissolved organic
206 nitrogen (DON) was calculated by subtracting $\text{NO}_3\text{-N}$ and $\text{NH}_4\text{-N}$ from TDN.

207

208 *Discharge*

209

210 Discharge at the Afon Ddu lower site was measured by a Starflow Ultrasonic Doppler
211 instrument with an integrated micrologger (Unidata, Western Australia). Due to
212 instrument failure, the velocity measurements were not correct, so accurate discharges
213 could not be calculated. However stage data were reliable, and provided an effective
214 proxy for the discharge changes associated with each event.

215 Rain data used were from the Snowdon Environmental Change Network site,
216 about 20 km north-west of the Afon Ddu catchment (Countryside Council for Wales
217 2008).

218

219 *Data analysis*

220

221 The event data are described quantitatively using different approaches. Base flow
222 composition was defined as the average stream water composition of samples taken
223 when there was minimal change in stream stage (the first 3, 5 and 11 samples in event
224 1, 2 and 3, respectively; average for pH calculated via H^+). In event 1 the stable stage
225 level did not represent a true base flow level, as it closely followed an earlier event.

226 To account for changes in the different parameters during the events, the percentage
227 change from the base flow to a maximum or minimum was calculated. Base flow and
228 average soil water (the micro-rhizon and the two zero-tension lysimeter samples) was
229 compared for each event using two sample t-test. Results for sphagnum pore water
230 could not be compared statistically to base flow or soil water, as there was only one
231 sample per event. Correlation of parameters was done using Pearson's r. Results were
232 considered significant for $P < 0.05$. Minitab Release 14 was used as statistical
233 software. Regression analyses for the Fe-estimation were performed with JMP 7.0.1.

234 A simple two end-member mixing model was applied to each event and
235 parameter (H^+ and DOC concentration, absorbance at 254 nm and fluorescence
236 intensity), using the average base flow as one end-member and average soil water as
237 the other end-member. This corresponds to the end-member model described by
238 Worrall et al. (2002) but omits the rain end-member (dilution). The model calculates
239 the proportion of each of the two end-members contributing to stream water, starting
240 from the first non-base flow sample. It is a crude model, as there were probably other
241 end-members; base flow is not a true end-member; and none of the parameters are
242 conservative. The intention was simply to assess whether the proposed two end-
243 member system could consistently explain observed variations in different
244 parameters, i.e. that the contribution of the two end-members was consistent for
245 different parameters throughout the event.

246 Fluorescence data were analysed using parallel factor analysis (PARAFAC) in
247 MATLAB (version R2007b), according to the procedure described by Stedmon and
248 Bro (2008). PARAFAC decomposes the complex set of sample emission-excitation
249 matrices (EEMs) and extracts specific components (Stedmon et al. 2003),

250 representing groups of fluorophores with similar fluorescence characteristics
251 (Stedmon and Markager 2005). The model is defined as

$$252 \quad x_{ijk} = \sum_{f=1}^F a_{if} b_{jf} c_{kf} + \varepsilon_{ijk}$$

253 $i = 1, \dots, I; \quad j = 1, \dots, J; \quad k = 1, \dots, K$ (1)

254 where x_{ijk} is the fluorescence intensity of the i th sample at emission wavelength j and
255 excitation wavelength k . a_{if} is directly proportional to the concentration of the f th
256 component in sample i . b_{jf} is linearly related to the quantum efficiency (fraction of
257 absorbed energy emitted as fluorescence) of the f th component at emission
258 wavelength j . c_{kf} is linearly proportional to the specific absorption coefficient of the
259 f th component at excitation wavelength k . The residual matrix ε_{ijk} represents the
260 variability not accounted for by the model, and the model is found by minimising the
261 sum of squared residuals (Stedmon et al. 2003).

262 The raw data were corrected for inner-filtering effects according to Parker and
263 Barnes (1957). Inner filtering reduces the fluorescence intensity due to the absorption
264 of the excitation beam and of emitted light after excitation (Lakowicz 1983, p. 44).
265 The correction suggested by Lakowicz (1983, p. 44) and applied by e.g. McKnight et
266 al. (2001) and Ohno (2002) gave nearly identical results as the Parker and Barnes
267 (1957) correction, but the latter correction was selected because it takes account of the
268 dimensions of the excitation and emission beams. The matrices were then corrected
269 for instrument bias, i.e. the effects of lamp output and instrument sensitivity/response.
270 A blank, measured on a sealed cell of Milli-Q water, was subtracted to remove/reduce
271 the Raman line (Hudson et al. 2007). Finally a mask was applied to the data matrix,
272 setting all data in the regions without fluorescence (excitation wavelength exceeds
273 emission wavelength) to zero, and replacing all data in the regions greatly influenced

274 by the two Rayleigh-Tyndall lines (Hudson et al. 2007) with missing values (NaN in
275 MATLAB).

276 A series of PARAFAC models with three to seven components were fitted to
277 the data. As some of the components gave negative excitation or emission loadings,
278 non-negativity constraints were applied. The PARAFAC model could be split half
279 validated (i.e. separate modelling of two halves of the samples gave statistically
280 identical results, cf. Stedmon, C. A. and Bro, R. 2008) for up to three components.
281 Initial exploration revealed that the sphagnum pore water samples from event 1 and 2
282 showed a high residual fluorescence at low excitation and emission wavelength. Some
283 other samples, especially soil water samples, also exhibited some fluorescence in this
284 region. However, overall this feature was present in too few samples for a model
285 including a component in this region to be validated. Thus, the two sphagnum pore
286 water samples were considered as outliers, and removed prior to the final modelling.
287 The final model was the three component model giving the smallest residual error out
288 of ten models fitted following random initialisation of the model. The maximum
289 fluorescence intensity of the different components (F_{max}) for individual samples was
290 corrected for dilution by multiplying by the dilution factor.

291

292 **Results**

293

294 *Hydrology*

295

296 The rainfall during event 1, 2 and 3 was 16, 17 and 114 mm, respectively (Fig. 2).
297 Event 1 was the second in a series of events following an 11 day dry period. There
298 was a 26 mm event just prior to event 1, and the stage was 198 mm at the onset of the

299 event. During the event, stage rose to 334 mm. Event 2 was preceded by 7 dry days,
300 with a series of large events before that (total precipitation 347 mm). The stream stage
301 was 138 mm at the onset of the event and maximum stage during the event was 211
302 mm. Event 3 was preceded by a 10 day dry period. The stream stage was at 131 mm
303 at the onset. During the event, stage rose to 1045 mm. Event 3 was clearly the largest
304 of the three events, and one of the largest of the whole autumn.

305

306 *pH and DOC*

307

308 The pH decreased during all events (Fig. 3). The largest decrease (2.3 pH units from
309 base flow levels) occurred in event 3, compared to 0.9 and 1.1 pH units in events 1
310 and 2, respectively. Base flow pH was significantly higher than that of soil water in all
311 events. pH of sphagnum pore water was at a similar level as that of soil water.

312 DOC concentrations increased during all events (Fig. 3), by 16%, 44% and
313 89% relative to base flow in events 1, 2 and 3, respectively. The DOC concentration
314 was significantly and negatively correlated with pH (r^2 0.59, 0.81 and 0.86 for event
315 1, 2 and 3, respectively). Soil water DOC concentrations were higher than those of
316 base flow, but the difference was only significant for event 2 and 3. DOC
317 concentration in sphagnum pore water was higher than in soil water in event 1 and 2,
318 and at the same level as in soil water in event 3.

319

320 *DOC/DON ratio*

321

322 The DOC/DON ratio data (Fig. 4) were noisy, but there was an upward trend in all
323 events. Maximum values were 58, 57 and 103% higher than base flow values in event

324 1, 2 and 3, respectively. There was no significant difference between soil water and
325 base flow DOC/DON ratios in any of the events. The DOC/DON ratios of the
326 sphagnum samples were far lower than those of both base flow and soil water
327 samples.

328

329 *Absorbance*

330

331 Absorbance at 254 nm increased during events 2 and 3 (Fig. 5), with 30 and 53%
332 maximum increases from base flow, respectively. In these events, absorbance was
333 significantly and positively correlated with DOC concentration (r^2 0.88 and 0.92,
334 respectively). In event 3 there was a striking diurnal pattern in absorbance after the
335 initial peak, with consistent maxima at 7 am and minima at 1 pm. Soil water
336 absorbance was significantly higher than that of base flow in events 2 and 3.
337 Absorbance of sphagnum pore water was generally lower than that of both base flow
338 and soil water.

339 The sUVa index (Fig. 5) decreased from base flow levels during event 1 (16%
340 change), and more clearly during event 3 (28% change). There was no apparent
341 pattern in event 2. The four deviating samples in event 2 with relatively low sUVa
342 values correspond to the four samples with relatively low DOC, and appears to be
343 caused by a DOC measurement error, as this deviation was not observed for
344 absorbance. The reason for this error could not be established. Base flow sUVa was
345 significantly higher than that of soil water in all events. sUVa in sphagnum pore water
346 was far lower than sUVa in both base flow and soil water.

347 There was a rising trend in E2/E3 (Fig. 5) for all events, with a change from
348 base flow to maximum levels of 5, 3 and 27% for events 1, 2 and 3, respectively. Soil

349 water E2/E3 was significantly higher than that of base flow in all events. E2/E3 in
350 sphagnum pore water was at a similar level as in soil water.

351 The average estimated Fe concentration was 2.2, 1.6 and 1.5 mg l⁻¹ for events
352 1, 2 and 3, respectively, i.e. above the level (0.5 mg l⁻¹) where iron interference should
353 be considered (Weishaar et al. 2003). The estimated Fe concentration followed closely
354 the DOC concentration, and was only slightly modified by flow. Fig. 5 shows the
355 effect of correcting for Fe absorbance on absorbance and sUVa. The effect is
356 pronounced, and the resulting sUVa values are more realistic (cf. Weishaar et al.
357 2003, table 1). However, the temporal patterns are preserved to a high degree. As the
358 following discussion is mainly focused on the relative changes and temporal patterns,
359 the issue of iron interference will not be further discussed, but it should be kept in
360 mind that the absorbance and sUVa values are overestimated. E2/E3 is probably not
361 strongly affected by iron interference. Weishaar et al. (2003) observed similar iron
362 interference at 280 nm as 254 nm, so the interference at 365 nm is not believed to be
363 very different. Moreover, the E2/E3 values obtained are realistic (cf. Peuravuori and
364 Pihlaja 1997, table 2).

365

366 *Fluorescence*

367

368 Component 3 from the PARAFAC analysis was considered to represent
369 contamination, as it only appeared in the samples where a certain plastic tube was
370 used for dilution. Blank samples using the same plastic tube exhibited the same
371 feature. This component is consequently not discussed in the following. However, this
372 component overlapped with the region that was poorly modelled for the sphagnum

373 samples, and the presence of this contamination component may have added to the
374 difficulty of validating a model with a true component in this region.

375 The two remaining components are shown in Fig. 6. Both components were
376 double-peaked. Component 1 (C1) had maximum excitation at <250 nm with a
377 secondary peak at 310 nm, and maximum emission at 440 nm. Component 2 (C2) had
378 a maximum at excitation 260 nm with a secondary peak at 360 nm, and maximum
379 emission at 474 nm. The temporal variation in the fluorescence intensity (Fig. 7) did
380 not seem to coincide with the changes in stream stage. Throughout the sampling
381 periods the intensity levels were either stable or decreasing. C1 had the clearest
382 decreasing trend. This resulted in a decrease in the C1/C2 ratio in all events, with a
383 levelling off (event 1 and 2) or increase (event 3) towards the end of the events (Fig.
384 7). The decrease from base flow levels to minimum levels was 7, 18, and 17% in
385 events 1, 2 and 3, respectively. C1 was higher in base flow than soil water, but this
386 was only significant for event 2. C2 was higher in soil water, and this was significant
387 for event 2 and 3. Both C1 and C2 were lower in sphagnum pore water than in base
388 flow and soil water. For C1/C2, base flow was significantly higher than soil water in
389 all events. C1/C2 of sphagnum pore water was at the same level as for base flow.

390 In event 3, there was a clear diurnal pattern in C1 and C2, especially observed
391 after the stage peak. The fluorescence peaked at 1 pm and was lowest during the night
392 (9 pm to 7 am). The pattern coincided with the in situ temperature variation, while it
393 was inversely related to the pattern observed for absorbance (Fig. 8).

394 Ohno et al. (2008) investigated Fe quenching of PARAFAC-derived
395 components. Their results could be used as basis for a crude estimate of fluorescence
396 quenching, as their components 1 and 2 corresponded well with C1 and C2,
397 respectively. The effect on overall levels was pronounced, but the temporal patterns

398 were preserved to a high degree. Ohno et al. (2008) showed that component 1 was
399 more strongly quenched than component 2. However, the estimated effect of
400 quenching on C1/C2 was minor. Hence, Fe quenching seems unimportant to the
401 temporal patterns in fluorescence, and will not be further discussed.

402

403 *Mixing model*

404

405 Mixing models for event 1 and 2 did not show any consistent patterns between
406 parameters (data not shown). For event 3 (Fig. 9), there was a consistent, although not
407 identical, pattern in H⁺ concentration, DOC concentration and absorbance at 254 nm.
408 The curves showed a decrease in the proportion of base flow-type water at the start of
409 the event and a subsequent increase coinciding with the decrease in stream stage. The
410 models based on H⁺ and absorbance indicated lower and higher contribution of soil
411 water at peak flow compared to the model based on DOC concentration, respectively.
412 Mixing models for event 3 based on C1 and C2 were not consistent with those of the
413 other three parameters (Fig. 9).

414

415 **Discussion**

416

417 *Effects of events on DOC concentration*

418

419 The higher pH and lower DOC concentration in base flow compared to soil water
420 indicates that the peat system in Afon Ddu is not like the system described by Clark et
421 al. (2007a, b), where base flow was chemically similar to soil water. Like DOC
422 concentration, absorbance at 254 nm was also lower in base flow than in soil water.

423 Absorbance is generally known to be closely correlated with DOC concentration (e.g.
424 Dobbs et al. 1972; Brandstetter et al. 1996; Korshin et al. 1997). Relatively high
425 baseflow DOC concentration indicates a significant input of drainage from the peat,
426 whilst the high baseflow pH demonstrates some hydrological connectivity with
427 underlying base-rich bedrock and/or mineral soil patches within the catchment.

428 The increased DOC concentration, increased absorbance and decreased pH at
429 high flow can be explained by an increased contribution of soil water, due to a rising
430 water table within the acrotelm and subsurface flow (Evans et al. 1999; Worrall et al.
431 2002). A higher similarity between soil water and stream water at high flow with
432 respect to DOC concentrations was also observed by Billett et al. (2006). The change
433 in stream chemistry could not solely be caused by an increased input of water from
434 the sphagnum layer, as this could not have explained the trend in absorbance. There
435 was little evidence of dilution at peak flow. The process could thus be reasonably well
436 described for event 3 by the applied two end-member mixing model, rather than a
437 three end-member model as suggested in Worrall et al. (2002). The discrepancies
438 observed between the DOC concentration, H^+ concentration and absorbance at 254
439 nm were most likely due to the lack of conservative mixing. The most reliable model
440 is probably the one for DOC, as H^+ concentration is unlikely to be conservative, and
441 absorbance behaves abnormally at peak flow. The inconsistency between parameters
442 in the models for event 1 and 2 can probably be explained by the limited size of the
443 events and small changes in stream chemistry, giving too much noise in such a crude
444 model.

445

446 *Effects of events on DOM quality*

447

448 The sUVa index provides a measure of the change in DOM quality expressed by
449 deviation from the correlation between DOC concentration and absorbance. sUVa has
450 been shown to be positively correlated with aromaticity and molecular weight (Croué
451 et al. 1999; Weishaar et al. 2003; Hood et al. 2005). Both sUVa and aromaticity
452 increase upon biodegradation (Kalbitz et al. 2003; Saadi et al. 2006). A high sUVa is
453 also associated with lower bioavailability (Fellman et al. 2008). Thus, the decrease in
454 sUVa at high flow indicates that the DOM released at high flow is less degraded and
455 aromatic and of lower molecular weight. This is supported by the increase in E2/E3,
456 as E2/E3 is negatively correlated to aromaticity and molecular weight (Peuravuori and
457 Pihlaja 1997). E2/E3 is a more robust parameter than sUVa, giving a smoother trend.

458 DOC/DON data from this study are noisy, but clearly show increases in all
459 three events. Increasing DOC/DON may be indicative of increasing aromaticity
460 (McKnight et al. 1997; Hood et al. 2005), because algal and microbial material has
461 low DOC/DON (McKnight et al. 1994) and low aromaticity (McKnight et al. 2001).
462 However, even the lowest DOC/DON values observed are much higher than those
463 associated with algal or microbial material (McKnight et al. 1994). Moreover, the
464 highest DOC/DON values observed correspond to the lower range of values observed
465 for different litter leachates (Magill and Aber 2000). Hence, in this case it is more
466 likely that increasing DOC/DON indicates decreasing degree of degradation (Melillo
467 et al. 1982; Qualls and Haines 1992; Currie et al. 1996; Yano et al. 2004). Thus both
468 absorbance and DOC/DON indicate that base flow is characterised by older and
469 probably more degraded DOM, while soil water comprises younger, less degraded
470 DOM (Evans et al. 2007).

471 With increased absorbance at high flow, one might expect to see a similar
472 response in fluorescence. However, unlike absorbance, fluorescence does not

473 invariably correlate well with DOC, and fluorescence relative to absorbance may vary
474 between different sources (Baker et al. 2008). In addition, fluorescing structures
475 constitute only a minor part of humic molecules (Miano et al. 1988; Senesi et al.
476 1991). Thus, it is possible for fluorescence to decrease even if absorbance, as a bulk
477 parameter, increases.

478 The mixing model for event 3 indicates that stream fluorescence during the
479 event was not a result of mixing of soil water and base flow water, as it was for
480 absorbance. For C1, soil water fluorescence was not sufficiently low to explain the
481 decrease, and C2 remained approximately at base flow levels throughout the event
482 despite the higher fluorescence intensity in soil water. The effect could not be
483 explained by increased contribution of low fluorescence sphagnum pore water, as this
484 does not fit with the observed changes in absorbance, DOC/DON and C1/C2 during
485 the events. Rather, the lack of consistency between fluorescence parameters and the
486 other parameters in the mixing models, the diurnal variation in fluorescence, and the
487 co-variation in fluorescence and in situ stream temperature observed in event 3,
488 collectively suggest that stream fluorescence may be governed by in-stream microbial
489 processes.

490 In-stream processing of DOM has been found to be an important control on
491 stream DOC in a similar catchment (Dawson et al. 2001). The microbial processing
492 probably occurs in biofilms, as it is generally believed that this is where most
493 microorganisms in natural systems exist (Sutherland 2001). Biofilms on the stream
494 bed, rather than suspended aggregates in the water column tend to dominate
495 ecosystems with high downstream transport and sediment-surface-area to water-
496 volume ratio, as in headwaters (Battin et al. 2008). Lock and Hynes (1976) showed
497 that the stream bottom, not the water itself is responsible for the major part of DOC

498 removal, and biofilms on the stream bed have been shown to remove a substantial
499 amount of DOC (Fiebig et al. 1990; Fiebig and Lock 1991). In the Afon Ddu there
500 were no indications of in-stream processing of bulk DOC. This either confirms that
501 only a minor part of the DOM released is subject to such processing, or that the
502 spectrophotometric methods, introducing less uncertainty than the DOC analysis, are
503 better for detecting short-term variation in DOM. The in-stream processing will be
504 further covered in the next section.

505 Microbial processing of DOM is limited by microbial capacities, i.e. reaction
506 rates (Battin et al. 2008). Hence, one can expect increased flow to give dilution of
507 stream fluorescence. The trends in C1 and C2 can thus be explained by dilution as the
508 main control, and only some compensation by fluorescing material from the soil. This
509 indicates that highly fluorescing DOM in soil water is not as mobile as less
510 fluorescing DOM. The compensation by soil water was highest for C2, where the
511 levels were higher in soil water than in base flow. The C1/C2 ratio would not be
512 affected by dilution, so the downwards trend was due to the small contribution of
513 fluorescing material from the soil.

514 A decreased C1/C2 represents a shift in peak position to higher wavelengths,
515 and this has been associated with higher density of aromatic rings, higher degree of
516 aromatic substitution and conjugation, higher molecular weight, and higher
517 hydrophobicity (Senesi 1990; Coble et al. 1998; Sharma and Schulman 1999, p. 20-
518 21; Wu et al. 2003). A peak shift to higher wavelengths may also signify an increase
519 in the importance of humic compared to fulvic acids (Senesi 1990). The C1/C2 ratio
520 thus suggests that the aromaticity and molecular weight of DOM was higher in soil
521 water compared to base flow, which is opposite to the conclusion based sUV_a, E2/E3
522 and DOC/DON. However, this fits in well with the idea that the fluorescing material

523 is mainly produced in-stream, which would make base flow fluorescing material
524 relatively younger than that deriving from soil water, as opposed to what is the case
525 for the bulk material. The overall conclusion is still that DOM released at high flow is
526 generally less degraded, as the fluorescing part constitutes only a minor part of the
527 DOM.

528

529 *Effects of events on in-stream processing of DOM*

530

531 As discussed above, the diurnal pattern in fluorescence observed in event 3 appears to
532 reflect diurnal variation in in-stream production of fluorescing DOM. Both
533 temperature and light are environmental factors that vary on a daily timescale.
534 Temperature has a positive effect on fluorescence, as the activity of the DOM-
535 producing microorganisms is stimulated by increasing temperature (Christ and David
536 1996; Gødde et al. 1996; Freeman et al. 2001). UV light is known to cause a decrease
537 in fluorescence, through so-called photo-bleaching (e.g. Skoog et al. 1996; Moran et
538 al. 2000; Patel-Sorrentino et al. 2004), and can be expected to co-vary with stream
539 temperature, peaking at mid-day.

540 To our knowledge a diurnal pattern in fluorescence has only been observed by
541 Spencer et al. (2007). They measured in situ fluorescence during base flow by means
542 of a fluorometer measuring at 370 nm excitation and over a broad emission band
543 centred at 460 nm. However, whereas fluorescence in event 3 peaked at mid-day and
544 was positively related to temperature, the diurnal pattern reported by Spencer et al.
545 peaked in the early morning (around 9 am), had a minimum in late afternoon (around
546 6 pm), and was negatively related to temperature. According to Spencer et al., the
547 diurnal pattern was related to day-time photo-bleaching of the DOM. This is

548 confirmed by base flow data from similar in situ measurements (excitation 330 nm,
549 emission 450 nm) from Afon Ddu (data not shown) and another stream (Nant y
550 Brwyn), 2 km away on the same area of blanket peat (Fig. 10), which showed a
551 similar diurnal pattern.

552 We hypothesise that during an event the diurnal fluorescence pattern is shifted,
553 due to the extra input of soil water DOM represented by absorbing material. Nieto-
554 Cid et al. (2006) showed that fluorescent humic substances could be produced by
555 microbial degradation of DOM on a short (<1 day) timescale, and that the magnitude
556 of production was related to microbial activity. If even a small proportion of the DOM
557 transported from the soil can be used by biofilm microbial communities, this could
558 explain the rapid, diurnal production of fluorescent DOM. Degradation would be
559 expected to peak at mid-day due to the higher temperatures, which could explain the
560 shift in the diurnal cycle, and also the negative relationship between fluorescence and
561 absorbance in event 3. This positive effect of temperature apparently surpasses any
562 negative effect of UV light at mid-day, perhaps because the cloud cover in the period
563 during and after the storm event was high, and UV intensity consequently low.
564 Alternatively, or additionally, given the high input of DOM, photo-bleaching could
565 have had a positive impact on production of fluorescing DOM, as UV radiation is
566 found to increase the substrate quality, and thus the degradability, of terrestrially
567 derived DOM (Moran et al. 2000; Tranvik and Bertilsson 2001; Anesio et al. 2005).

568 The hypothesis appears to be supported by event data from the Nant y Brwyn
569 (Fig. 10). At base flow the in situ fluorescence minimum was at about 4-7 pm and
570 peaked in the early morning (6-8 am). In the first day of the event shown (June 18th),
571 the pattern was shifted, and after the peak in discharge (June 19th) the in situ
572 fluorescence remained high all day until a minimum at 8 pm. In the following few

573 days in situ fluorescence continued to stay high throughout a longer part of the day,
574 with a delayed minimum compared to base flow, until June 24th, when the maximum
575 was again at 8 am and the minimum at 4 pm. There was more noise in these data than
576 the event data for the Afon Ddu, as the PARAFAC model involves smoothing. To
577 compare, the intensities at the wavelength measured at Nant y Brwyn were extracted
578 from the event 3 EEMs (data not shown), and although they exhibited more noise, the
579 diurnal pattern could still be distinguished. If the hypothesis is correct, then at least
580 part of the absorbing, humic DOM released during events must be rapidly processed
581 close to where it was released. However, more data and a more quantitative approach
582 are needed to confirm this. The hypothesis implies that the production of fluorescing
583 DOM not only peaks at a different time of the day during and after an event, but that
584 there is a general increase in fluorescence intensity. Dilution may explain why this
585 was not observed.

586

587 **Conclusions**

588

589 In peatlands where base flow is mainly high pH/low DOC ground water, events result
590 in increased DOC concentration due to flushing of the shallow acrotelm. The increase
591 in DOC is accompanied by decreased pH. The DOM released during events appears
592 to be less degraded, less aromatic and more bioavailable. An increased input of this
593 material is believed to have caused stimulation of in-stream microbial communities,
594 producing fluorescing metabolites.

595

596 Absorbance proved to be a good indicator of changes in DOM quality and an
597 identification of the shift of flow paths and sources throughout an event. Fluorescence

598 was less useful in that respect, due to the apparent dominance of in-stream production.
599 However, absorbance and fluorescence data combined gave indications of short term
600 changes in in-stream processing of DOM during events which have not previously
601 been observed.

602

603 **Acknowledgements**

604

605 This work was partly funded by the EU Eurolimpacs project (the Commission of
606 European Communities GOCE-CT-2003-505540). We thank David Williams, Simon
607 Grant, Annette Burden and Timothy Jones for assistance in the field; Annie Brittain
608 and Steve Hughes for help with the chemical analyses; David Norris for drawing the
609 catchment map; Colin Stedmon for assistance in the PARAFAC analysis; and
610 professor Jan Mulder for valuable comments on the manuscript.

611

612 **References**

613

- 614 Aitkenhead JA, McDowell WH (2000) Soil C:N ratio as a predictor of annual riverine
615 DOC flux at local and global scales. *Glob Biogeochem Cycles* 14(1):127-138
- 616 Aitkenhead JA, Hope D, Billett MF (1999) The relationship between dissolved
617 organic carbon in stream water and soil organic carbon pools at different
618 spatial scales. *Hydrol Process* 13(8):1289-1302
- 619 Anesio AM, Graneli W, Aiken GR, Kieber DJ, Mopper K (2005) Effect of humic
620 substance photodegradation on bacterial growth and respiration in lake water.
621 *Appl Environ Microbiol* 71(10):6267-6275

622 Baker A, Bolton L, Newson M, Spencer RGM (2008) Spectrophotometric properties
623 of surface water dissolved organic matter in an afforested upland peat
624 catchment. *Hydrol Process* 22(13):2325-2336

625 Battin TJ, Kaplan LA, Findlay S, Hopkinson CS, Marti E, Packman AI, Newbold JD,
626 Sabater F (2008) Biophysical controls on organic carbon fluxes in fluvial
627 networks. *Nature Geosci* 1(2):95-100

628 Billett MF, Deacon CM, Palmer SM, Dawson JJC, Hope D (2006) Connecting
629 organic carbon in stream water and soils in a peatland catchment. *J Geophys*
630 *Res* 111(G2):G02010. DOI 10.1029/2005JG000065

631 Billett MF, Garnett MH, Harvey F (2007) UK peatland streams release old carbon
632 dioxide to the atmosphere and young dissolved organic carbon to rivers.
633 *Geophys Res Lett* 34(23):L23401. DOI 10.1029/2007GL031797

634 Boyer EW, Hornberger GM, Bencala KE, McKnight DM (1997) Response
635 characteristics of DOC flushing in an alpine catchment. *Hydrol Process*
636 11(12):1635-1647

637 Brandstetter A, Sletten RS, Mentler A, Wenzel WW (1996) Estimating dissolved
638 organic carbon in natural waters by UV absorbance (254 nm). *J Plant Nutr*
639 *Soil Sci* 159(6):605-607

640 Buffam I, Galloway JN, Blum LK, McGlathery KJ (2001) A stormflow/baseflow
641 comparison of dissolved organic matter concentrations and bioavailability in
642 an Appalachian stream. *Biogeochem* 53(3):269-306

643 Christ MJ, David MB (1996) Temperature and moisture effects on the production of
644 dissolved organic carbon in a Spodosol. *Soil Biol Biochem* 28(9):1191-1199

645 Christensen JH, Hewitson B, Busuioc A, Chen A, Gao X, Held I, Jones R, Kolli RK,
646 Kwon W-T, Laprise R, Magaña Rueda V, Mearns L, Menéndez CG, Räisänen

647 J, Rinke A, Sarr A, Whetton P (2007) Regional climate projections. In:
648 Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M,
649 Miller HL (eds) Climate change 2007: The physical science basis.
650 Contribution of working group I to the fourth assessment report of the
651 intergovernmental panel on climate change. Cambridge University Press,
652 Cambridge, United Kingdom and New York, NY, USA

653 Clark JM, Lane SN, Chapman PJ, Adamson JK (2007a) Export of dissolved organic
654 carbon from an upland peatland during storm events: Implications for flux
655 estimates. *J Hydrol* 347(3-4):438-447

656 Clark JM, Lane SN, Chapman PJ, Adamson JK (2007b) Link between DOC in near
657 surface peat and stream water in an upland catchment. *Sci Total Environ* (in
658 press). DOI 10.1016/j.scitotenv.2007.11.002

659 Coble PG, Del Castillo CE, Avril B (1998) Distribution and optical properties of
660 CDOM in the Arabian Sea during the 1995 Southwest Monsoon. *Deep-Sea*
661 *Res Part II* 45(10-11):2195-2223

662 Countryside Council for Wales (2008) Yr Wyddfa – Snowdon ECN Site.
663 <https://weather.ccw.gov.uk/>. Cited 25 Jan 2008

664 Croué J-P, Violleau D, Bodaire C, Legube B (1999) Removal of hydrophobic and
665 hydrophilic constituents by anion exchange resin. *Water Sci Technol*
666 40(9):207-214

667 Currie WS, Aber JD, McDowell WH, Boone RD, Magill AH (1996) Vertical
668 transport of dissolved organic C and N under long-term N amendments in pine
669 and hardwood forests. *Biogeochem* 35(3):471-505

670 Dawson JJC, Bakewell C, Billett MF (2001) Is in-stream processing an important
671 control on spatial changes in carbon fluxes in headwater catchments? *Sci Total*
672 *Environ* 265(1-3):153-167

673 Dobbs RA, Dean RB, Wise RH (1972) Use of ultraviolet absorbance for monitoring
674 total organic carbon content of water and wastewater. *Water Res* 6(10):1173-
675 1180

676 Eimers MC, Buttle J, Watmough SA (2008) Influence of seasonal changes in runoff
677 and extreme events on dissolved organic carbon trends in wetland- and
678 upland-draining streams. *Can J Fish Aquat Sci* 65(5):796-808

679 Evans CD, Freeman C, Cork LG, Thomas DN, Reynolds B, Billett MF, Garnett MH,
680 Norris D (2007) Evidence against recent climate-induced destabilisation of
681 soil carbon from ^{14}C analysis of riverine dissolved organic matter. *Geophys*
682 *Res Lett* 34(7):L07407. DOI 10.1029/2007GL029431

683 Evans MG, Burt TP, Holden J, Adamson JK (1999) Runoff generation and water table
684 fluctuations in blanket peat: evidence from UK data spanning the dry summer
685 of 1995. *J Hydrol* 221(3-4):141-160

686 Fellman JB, D'Amore DV, Hood E, Boone RD (2008) Fluorescence characteristics
687 and biodegradability of dissolved organic matter in forest and wetland soils
688 from coastal temperate watersheds in southeast Alaska. *Biogeochem*
689 88(2):169-184

690 Fiebig DM, Lock MA (1991) Immobilization of dissolved organic matter from
691 groundwater discharging through the stream bed. *Freshw Biol* 26(1):45-55

692 Fiebig DM, Lock MA, Neal C (1990) Soil water in the riparian zone as a source of
693 carbon for a headwater stream. *J Hydrol* 116(1-4):217-237

694 Freeman C, Evans CD, Monteith DT, Reynolds B, Fenner N (2001) Export of organic
695 carbon from peat soils. *Nature* 412(6849):785-785

696 Gödde M, David MB, Christ MJ, Kaupenjohann M, Vance GF (1996) Carbon
697 mobilization from the forest floor under red spruce in the northeastern U.S.A.
698 *Soil Biol Biochem* 28(9):1181-1189

699 Hinton MJ, Schiff SL, English MC (1997) The significance of storms for the
700 concentration and export of dissolved organic carbon from two Precambrian
701 Shield catchments. *Biogeochem* 36(1):67-88

702 Hinton MJ, Schiff SL, English MC (1998) Sources and flowpaths of dissolved organic
703 carbon during storms in two forested watersheds of the Precambrian Shield.
704 *Biogeochem* 41(2):175-197

705 Hood E, Williams MW, McKnight DM (2005) Sources of dissolved organic matter
706 (DOM) in a Rocky Mountain stream using chemical fractionation and stable
707 isotopes. *Biogeochem* 74(2):231-255

708 Hope D, Billett MF, Cresser MS (1994) A review of the export of carbon in river
709 water - fluxes and processes. *Environ Pollut* 84(3):301-324

710 Hudson N, Baker A, Reynolds D (2007) Fluorescence analysis of dissolved organic
711 matter in natural, waste and polluted waters - A review. *River Res Appl*
712 23(6):631-649

713 Inamdar SP, O'Leary N, Mitchell MJ, Riley JT (2006) The impact of storm events on
714 solute exports from a glaciated forested watershed in western New York,
715 USA. *Hydrol Process* 20(16):3423-3439

716 Kalbitz K, Schwesig D, Schmerwitz J, Kaiser K, Haumaier L, Glaser B, Ellerbrock R,
717 Leinweber P (2003) Changes in properties of soil-derived dissolved organic
718 matter induced by biodegradation. *Soil Biol Biochem* 35(8):1129-1142

719 Korshin GV, Li CW, Benjamin MM (1997) Monitoring the properties of natural
720 organic matter through UV spectroscopy: A consistent theory. *Water Res*
721 31(7):1787-1795

722 Lakowicz JR (1983) *Principles of fluorescence spectroscopy*. Plenum Press, New
723 York

724 Lock MA, Hynes HBN (1976) The fate of "dissolved" organic carbon derived from
725 autumn-shed maple leaves (*Acer saccharum*) in a temperate hard-water
726 stream. *Limnol Oceanogr* 21(3):436-443

727 Magill AH, Aber JD (2000) Dissolved organic carbon and nitrogen relationships in
728 forest litter as affected by nitrogen deposition. *Soil Biol Biochem* 32(5):603-
729 613

730 McDowell WH, Likens GE (1988) Origin, composition, and flux of dissolved organic
731 carbon in the Hubbard Brook valley. *Ecol Monogr* 58(3):177-195

732 McKnight DM, Andrews ED, Spaulding SA, Aiken GR (1994) Aquatic fulvic acids in
733 algal-rich antarctic ponds. *Limnol Oceanogr* 39(8):1972-1979

734 McKnight DM, Harnish R, Wershaw RL, Baron JS, Schiff S (1997) Chemical
735 characteristics of particulate, colloidal, and dissolved organic material in Loch
736 Vale Watershed, Rocky Mountain National Park. *Biogeochem* 36(1):99-124

737 McKnight DM, Boyer EW, Westerhoff PK, Doran PT, Kulbe T, Andersen DT (2001)
738 Spectrofluorometric characterization of dissolved organic matter for indication
739 of precursor organic material and aromaticity. *Limnol Oceanogr* 46(1):38-48

740 Melillo JM, Aber JD, Muratore JF (1982) Nitrogen and lignin control of hardwood
741 leaf litter decomposition dynamics. *Ecol* 63(3):621-626

742 Miano TM, Sposito G, Martin JP (1988) Fluorescence spectroscopy of humic
743 substances. *Soil Sci Soc Am J* 52(4):1016-1019

744 Moran MA, Sheldon WM, Zepp RG (2000) Carbon loss and optical property changes
745 during long-term photochemical and biological degradation of estuarine
746 dissolved organic matter. *Limnol Oceanogr* 45(6):1254-1264

747 Neal C, Lofts S, Evans CD, Reynolds B, Tipping E, Neal M (2008) Increasing iron
748 concentrations in UK upland waters. *Aquat Geochem* 14 (3): 263-288

749 Nieto-Cid M, Álvarez-Salgado XA, Pérez FF (2006) Microbial and photochemical
750 reactivity of fluorescent dissolved organic matter in a coastal upwelling
751 system. *Limnol Oceanogr* 51(3):1391-1400

752 Ohno T (2002) Fluorescence inner-filtering correction for determining the
753 humification index of dissolved organic matter. *Environ Sci Technol*
754 36(4):742-746

755 Ohno T, Amirbahman A, Bro R (2008) Parallel factor analysis of excitation-emission
756 matrix fluorescence spectra of water soluble soil organic matter as basis for
757 the determination of conditional metal binding parameters. *Environ Sci*
758 *Technol* 42 (1): 186-192

759 Parker CA, Barnes WJ (1957) Some experiments with spectrofluorimeters and filter
760 fluorimeters. *Analyst* 82(978):606-618

761 Patel-Sorrentino N, Mounier S, Lucas Y, Benaim JY (2004) Effects of UV-visible
762 irradiation on natural organic matter from the Amazon basin. *Sci Total*
763 *Environ* 321(1-3):231-239

764 Peuravuori J, Pihlaja K (1997) Molecular size distribution and spectroscopic
765 properties of aquatic humic substances. *Anal Chim Acta* 337(2):133-149

766 Qualls RG, Haines BL (1992) Biodegradability of dissolved organic matter in forest
767 throughfall, soil solution, and stream water. *Soil Sci Soc Am J* 56(2):578-586

768 Saadi I, Borisover M, Armon R, Laor Y (2006) Monitoring of effluent DOM
769 biodegradation using fluorescence, UV and DOC measurements.
770 Chemosphere 63(3):530-539

771 Senesi N (1990) Molecular and quantitative aspects of the chemistry of fulvic acid
772 and its interactions with metal ions and organic chemicals: Part II. The
773 fluorescence spectroscopy approach. Anal Chim Acta 232(1):77-106

774 Senesi N, Miano TM, Provenzano MR, Brunetti G (1991) Characterization,
775 differentiation, and classification of humic substances by fluorescence
776 spectroscopy. Soil Sci 152(4):259-271

777 Sharma A, Schulman SG (1999) Introduction to fluorescence spectroscopy. Wiley,
778 New York

779 Skoog A, Wedborg M, Fogelqvist E (1996) Photobleaching of fluorescence and the
780 organic carbon concentration in a coastal environment. Mar Chem 55(3-
781 4):333-345

782 Soulsby C, Rodgers P, Smart R, Dawson J, Dunn S (2003) A tracer-based assessment
783 of hydrological pathways at different spatial scales in a mesoscale Scottish
784 catchment. Hydrol Process 17(4):759-777

785 Spencer RGM, Pellerin BA, Bergamaschi BA, Downing BD, Kraus TEC, Smart DR,
786 Dahgren RA, Hernes PJ (2007) Diurnal variability in riverine dissolved
787 organic matter composition determined by in situ optical measurement in the
788 San Joaquin River (California, USA). Hydrol Process 21(23):3181-3189

789 Stedmon CA, Markager S (2005) Resolving the variability in dissolved organic matter
790 fluorescence in a temperate estuary and its catchment using PARAFAC
791 analysis. Limnol Oceanogr 50(2):686-697

792 Stedmon, C. A. and Bro, R. 2008 Characterizing dissolved organic matter
793 fluorescence with parallel factor analysis: a tutorial. *Limnology and*
794 *Oceanography: Methods* (in press)

795 Stedmon CA, Markager S, Bro R (2003) Tracing dissolved organic matter in aquatic
796 environments using a new approach to fluorescence spectroscopy. *Mar Chem*
797 82(3-4):239-254

798 Sutherland IW (2001) The biofilm matrix - an immobilized but dynamic microbial
799 environment. *Trends Microbiol* 9(5):222-227

800 Thurman EM (1985) *Organic geochemistry of natural waters*. Nijhoff, Dordrecht

801 Tranvik LJ, Bertilsson S (2001) Contrasting effects of solar UV radiation on dissolved
802 organic sources for bacterial growth. *Ecol Lett* 4(5):458-463

803 Vogt RD, Akkanen J, Andersen DO, Bruggemann R, Chatterjee B, Gjessing E,
804 Kukkonen JVK, Larsen HE, Luster J, Paul A, Pflugmacher S, Starr M,
805 Steinberg CEW, Schmitt-Kopplin P, Zsolnay A (2004) Key site variables
806 governing the functional characteristics of Dissolved Natural Organic Matter
807 (DNOM) in Nordic forested catchments. *Aquat Sci* 66(2):195-210

808 Weishaar JL, Aiken GR, Bergamaschi BA, Fram MS, Fujii R, Mopper K (2003)
809 Evaluation of specific ultraviolet absorbance as an indicator of the chemical
810 composition and reactivity of dissolved organic carbon. *Environ Sci Technol*
811 37(20):4702-4708

812 Worrall F, Burt TP, Jaeban RY, Warburton J, Shedden R (2002) Release of dissolved
813 organic carbon from upland peat. *Hydrol Process* 16(17):3487-3504

814 Wu FC, Evans RD, Dillon PJ (2003) Separation and characterization of NOM by
815 high-performance liquid chromatography and on-line three-dimensional

816 excitation emission matrix fluorescence detection. Environ Sci Technol
817 37(16):3687-3693
818 Yano Y, Lajtha K, Sollins P, Caldwell BA (2004) Chemical and seasonal controls on
819 the dynamics of dissolved organic matter in a coniferous old-growth stand in
820 the Pacific Northwest, USA. Biogeochem 71(2):197-223
821 Zepp RG, Sheldon WM, Moran MA (2004) Dissolved organic fluorophores in
822 southeastern US coastal waters: correction method for eliminating Rayleigh
823 and Raman scattering peaks in excitation-emission matrices. Mar Chem 89 (1-
824 4): 15-36

825

826

827

828

829

830

831

832

833

834

835

836 **Figure captions**

837

838 **Fig. 1** Location of the field site (left) and map of the Afon Ddu catchment (right).

839

840 **Fig. 2** Stream stage and precipitation for the whole autumn (top) and for single events
841 (bottom). Markers in top panel indicate time of sampling. Note different time scale for
842 event 3.

843

844 **Fig. 3** pH and dissolved organic carbon (DOC) concentration in stream samples,
845 sphagnum pore water and soil water from the three events. BF = average base flow;
846 Sph = sphagnum pore water; MR = soil water collected by micro-rhizons; ZT5 = soil
847 water collected by zero-tension lysimeter at 5 cm; ZT10 = soil water collected by
848 zero-tension lysimeter at 10 cm. Note different time scale for event 3.

849

850 **Fig. 4** Dissolved organic carbon/dissolved organic nitrogen (DOC/DON) ratio in
851 stream samples, sphagnum pore water and soil water from the three events. BF =
852 average base flow; Sph = sphagnum pore water; MR = soil water collected by micro-
853 rhizons; ZT5 = soil water collected by zero-tension lysimeter at 5 cm; ZT10 = soil
854 water collected by zero-tension lysimeter at 10 cm. Note different time scale for event
855 3.

856

857 **Fig. 5** Absorbance at 254 nm, sUVa (absorbance at 254 nm/DOC concentration) and
858 E2/E3 (absorbance at 250 nm/absorbance at 365 nm) in stream samples, sphagnum
859 pore water and soil water from the three events. BF = average base flow; Sph =
860 sphagnum pore water; MR = soil water collected by micro-rhizons; ZT5 = soil water
861 collected by zero-tension lysimeter at 5 cm; ZT10 = soil water collected by zero-
862 tension lysimeter at 10 cm. Note different time scale for event 3.

863

864 **Fig. 6** Excitation-emission matrices of the two parallel factor analysis (PARAFAC)
865 generated components C1 (top) and C2 (bottom). The scale is emission
866 loading*excitation loading (i.e. $b_j c_k$ in equation 1).

867

868 **Fig. 7** Maximum fluorescence intensity (F_{max}) of component C1 and C2 and the ratio
869 of the two in stream samples, sphagnum pore water and soil water from the three
870 events. BF = average base flow; Sph = sphagnum pore water; MR = soil water
871 collected by micro-rhizons; ZT5 = soil water collected by zero-tension lysimeter at 5
872 cm; ZT10 = soil water collected by zero-tension lysimeter at 10 cm. Note different
873 time scale for event 3.

874

875 **Fig. 8** Maximum fluorescence intensity (F_{max}) of component C1 and C2 in event 3
876 plotted with in situ temperature (top) and absorbance at 254 nm (bottom).

877

878 **Fig. 9** Two end-member mixing model for event three, showing the proportion of base
879 flow type water needed to explain stream concentration when the other end-member is
880 soil water (average of the micro-rhizon sample and the two zero-tension lysimeter
881 samples). Mixing model excluding (left) and including (right) C1 and C2.

882

883 **Fig. 10** Fluorescence measured in situ (excitation 330 nm, emission 450 nm) and
884 discharge in Nant y Brwyn.

885

Figure 1
[Click here to download high resolution image](#)




Figure 2

[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 4

[Click here to download high resolution image](#)




Figure 5
[Click here to download high resolution image](#)




Figure 6
[Click here to download high resolution image](#)





Figure 7

[Click here to download high resolution image](#)




Figure 8
[Click here to download high resolution image](#)





Figure 9
[Click here to download high resolution image](#)





Figure 10
[Click here to download high resolution image](#)

