

11-2023

## Literature Review on the Effectiveness of POC Lactate Tests for Post-Operative Sepsis Discharge

Blaire Adler

Paarth Jain

Nathan Rim

Rachel Monane

Follow this and additional works at: [https://jdc.jefferson.edu/si\\_des\\_2025\\_phase1](https://jdc.jefferson.edu/si_des_2025_phase1)



Part of the [Art and Design Commons](#), and the [Medicine and Health Sciences Commons](#)

**[Let us know how access to this document benefits you](#)**

---

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

**Project title:**

**Literature Review on the Effectiveness of POC Lactate Tests for Post-Operative Sepsis**

**Discharge**

**Author(s):**

**Blaire Adler, BS\*\*, Paarth Jain, BS\*\*, Nathan Rim, BS\*\*, Rachel Monane, BBA\*\***

**Background:**

Every year in the USA, there are at least two million surgery-related infections; around 11% are derived from decompression spinal cord surgery. Generally, later detection of sepsis leads to significant burdens on the healthcare system; this manifests financially, temporally, and in poorer patient outcomes. “SepsisCheckToGo” is a toolkit aimed to minimize the number of sepsis complications in spinal surgery patients postoperatively; the kit would include a pulse oximeter, thermometer, and fingertip POC lactate tests to detect early sepsis signs after discharge and before the follow-up visit (which would occur around one month after initial discharge).

**Methods:**

Our team did a literature review on NIH PubMed; we aimed to gather existing information on the effectiveness of point-of-care lactate tests. After compiling relevant data, our team analyzed each article for statistics on the effectiveness of point-of-care lactate tests in the context of post-surgical usage. These findings were analyzed for any relevant information on sepsis prevention.

**Results:**

Data from many different studies showed that point-of-care finger lactate tests have reliable accuracy in blood lactate values. The tools were compared to gold-standard tests and maintained accuracy while being easy to use. However, there is a significant lack of evidence for the contribution of point-of-care lactate testing in improving rates of post-operative sepsis.

**Conclusion:**

Point-of-care lactate tests are simple for healthcare providers to use in a variety of different contexts; thus, they are reliable for quick measurements of lactate levels. Unfortunately, there is lack of evidence that this has much utility in patient diagnosis/management to cause any significant changes in patient outcome, particularly in the detection/prevention of genuine sepsis. Therefore, this tool may not be useful for the “SepsisCheckToGo” kit, as it could increase healthcare costs and add unnecessary steps to patients’ post-surgery steps without real evidence of alleviated outcomes.