





RESEARCH ARTICLE

REVISED Evaluating the effect of digital technology on the learning of orthodontic cephalometric measurement [version 3; peer review: 2 approved]

Previously titled: Evaluating the effectiveness of integrating digital technology into orthodontic cephalometric measurement teaching

Xin Yu^{1-3*}, Yu Tian ^{1,2*}, Dandan Li^{1,2*}, Wen Sun^{1,2}, Hua Wang^{1,2}, Siyang Yuan⁴, Bin Yan^{1,2}, Lin Wang^{1,2}, Yongchu Pan ^{1,2}

¹Department of Orthodontics, Affiliated Nantong Stomatological Hospital of Nantong University, Nantong, 226006, China

²Department of Orthodontics, The Affiliated Stomatological Hospital of Nanjing Medical University, Nanjing, 210029, China

³Jiangsu Key Laboratory of Oral Diseases, Nanjing, 210029, China

⁴School of Dentistry, University of Dundee, Dundee, DD1 4HN, UK

* Equal contributors

V3 First published: 18 Mar 2022, 11:328
<https://doi.org/10.12688/f1000research.109876.1>

Second version: 12 Dec 2022, 11:328
<https://doi.org/10.12688/f1000research.109876.2>

Latest published: 13 Jun 2023, 11:328
<https://doi.org/10.12688/f1000research.109876.3>

Abstract









Background: This study aimed to evaluate the effect of applying digital technology in orthodontic learning, students' new skills and satisfaction towards it.



Methods: In total, 94 undergraduates of stomatology were recruited and randomly allocated to two groups. According to the cross-over design, both groups completed cephalometric measurements through the traditional hand-drawn method and digital technology (the Dolphin software) in different orders. By traditional hand-drawn method, students need to depict the outlines of the craniofacial anatomical structures on the sulfuric transfer paper first, then marked the measurement points and completed the measurement of line spacings and angles. By digital technology, they should mark the points in the software and adjust the automatically generated outlines of the structures and obtained the results. Besides, an online questionnaire was designed to investigate students' attitudes toward the digital technology. Two professional orthodontists were invited as instructors. They measured a lateral cranial radiograph by two methods with one week's interval, and their intra- and inter-class correlation coefficient were measured. The means of their measurements were set as standards.

Results: The inter- and intra-ICC of two instructors surpassed 90%,

Open Peer Review

Approval Status  

	1	2
version 3 (revision) 13 Jun 2023		 view
		
version 2 (revision) 12 Dec 2022	 view	 view
		
version 1 18 Mar 2022	 view	 view

- Cinzia Maspero** , University of Milan, Milan, Italy
Ospedale Maggiore Policlinico, Milan, Italy
- Ana Corte-Real** , University of Coimbra, Coimbra, Portugal
University of Coimbra, Coimbra, Portugal

Any reports and responses or comments on the

and there were no significant differences between their measurements, and the measurements by two methods. There were significant differences of students' measurements ($P_{1-SNA} < 0.01$, $P_{1-SNB} = 0.01$ and $P_{1-L1-NB(mm)} < 0.01$) between the traditional method and digital technology. Besides, the most results of digital technology were closer to the standards than those of traditional method, including five items with statistical significance ($P_{2-SNB} < 0.05$, $P_{2-L1-NB(mm)} < 0.01$, $P_{2-FMA} < 0.05$, $P_{2-FMIA} < 0.05$, $P_{2-IMPA} < 0.01$), while three items were the opposite ($P_{2-SNA} < 0.05$, $P_{2-ANB(mm)} < 0.01$, $P_{2-NA-PA} < 0.01$). The questionnaire showed more students preferred digital technology compared with traditional method and 72% of participants mastered 50-80% of cephalometric knowledge after the course.

article can be found at the end of the article.

Keywords

orthodontic teaching; cephalometric measurement; traditional hand-drawn method; digital technology; the Dolphin software

Corresponding authors: Bin Yan (byan@njmu.edu.cn), Yongchu Pan (panyongchu@njmu.edu.cn)

Author roles: **Yu X:** Conceptualization, Visualization, Writing – Original Draft Preparation; **Tian Y:** Investigation, Visualization, Writing – Original Draft Preparation; **Li D:** Formal Analysis, Methodology, Writing – Review & Editing; **Sun W:** Formal Analysis, Project Administration, Writing – Review & Editing; **Wang H:** Data Curation, Resources, Writing – Review & Editing; **Yuan S:** Investigation, Visualization, Writing – Review & Editing; **Yan B:** Data Curation, Supervision, Writing – Review & Editing; **Wang L:** Project Administration, Supervision, Writing – Review & Editing; **Pan Y:** Conceptualization, Funding Acquisition, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by the Educational Project of Nanjing Medical University (2019ZC027 and ZC32015JG01), the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD-2018-87), and the Jiangsu Provincial Key Medical Discipline (ZDXKA2016026).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2023 Yu X *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Yu X, Tian Y, Li D *et al.* **Evaluating the effect of digital technology on the learning of orthodontic cephalometric measurement [version 3; peer review: 2 approved]** F1000Research 2023, 11:328

<https://doi.org/10.12688/f1000research.109876.3>

First published: 18 Mar 2022, 11:328 <https://doi.org/10.12688/f1000research.109876.1>

REVISED Amendments from Version 2

To make the article more easily to understand, some minor mistakes and phrases have been revised.

Any further responses from the reviewers can be found at the end of the article

Introduction

Orthodontics, as a discipline of stomatology, aims to study various kinds of malocclusion, including deformities of teeth, jaws and the craniofacial region.¹ Diagnosis is the most important part of orthodontic clinical work, among which cephalometric measurement is an essential procedure.² Nevertheless, cephalometric measurement indicators are numerous and complicated. In the past, the traditional hand-drawn method was applied, which required reading lamps, sulfuric acid transfer paper, dividers and so on. In addition, previous studies suggested that undergraduates tended to show less confidence in reading and measuring lateral cranial radiographs.³

As continuous developments are seen in digital technology, such technology has been applied in the education for undergraduates in recent decades. Digital technology possesses vivid images and operable processes, making it more intuitive, interactive and understandable than simply imparting theoretical knowledge to students, which may help improve the effect of teaching practice.⁴⁻⁶ Buchanan JA *et al.* found that before starting clinical work, students could better master theoretical knowledge through simulated operation or computer-aided learning method.^{7,8} It was also reported that students' attitudes towards computer-aided learning and digital technique were positive.⁹⁻¹³ Digital technology has offered great potential for dental education as well.¹⁴ For instance, Nagy ZA *et al.* reported that the Dental Teacher software could help students more efficiently learn the preparation technique of onlay restorations and facilitate their individual performances.¹⁵ Liu L *et al.* also found the digital training system might be a good alternative to the traditional training method in the preclinical practice of tooth preparation.¹⁶

The digital cephalometric analysis system, widely incorporated in intelligent software, was developed to computerize the manual tasks and output the specific results automatically. It was reported to be more time-saving than traditional measurement method and helpful for reducing unnecessary errors during the measurement process.¹⁷ Farooq *et al.* also found that the accuracy of cephalometric measurement by digital tracing with FACAD[®] was similar with the manual method. Furthermore, its advantages of digital imaging, such as quality improvement, file transmission and archiving made digitalized cephalometric analysis preferable in daily use.¹⁸

The Dolphin software[®] (Dolphin Imaging & management solution, America) is widely applied in the field of orthodontics, possessing functions like storage and management of patients' information and images. After users upload computerized tomography photographs, it can also achieve three-dimensional imaging, cephalometric measurement and treatment effect prediction. It incorporates more than 400 cephalometric analytical methods. This software has been reported to have the potential as an animation textbook for medical college students.¹⁹ Although it was assumed to exert some positive effects in the teaching process, there is still lack of research investigating the effectiveness of applying it in orthodontic cephalometric teaching process. Thus, the aim of this study was to examine the effectiveness of applying digital technology (the Dolphin software[®]) in teaching cephalometric measurement.

Methods**Ethical considerations**

This study followed the guidelines of the Nanjing Medical University ethics review committee and received the approval of the committee (approval number: PJ2019-053-001). All participants gave written informed consent.

Study size

G*Power software version 3.1.9.7 (RRID:SCR_013726) was used to estimate the required sample size for this study. This study used two independent *t* tests to calculate the number of students needed. The study power was set at 90% and alpha value set at 0.05. Based on these, a minimum sample of 86 subjects was required.

Participants

This study approached fourth-year undergraduate students of Stomatology in Nanjing Medical University by inviting them to attend this course, followed by their voluntary registration. All the students, consisting of 63 female and 31 male students (around 21-23 years old), agreed to participate in this pedagogical experiment and signed the informed consent form. All the students have neither learned about the method of measuring the lateral cranial radiographs nor used the Dolphin software before. We also invited two orthodontist faculty members with over 5 years of clinical experience as instructors.




Figure 1. Flow diagram of the teaching experiment.

Study design

This study was conducted from May 12, 2020 to June 16, 2020. The flow diagram of this pedagogical study was shown in [Figure 1](#). For students, it was divided into pre-test learning, group and cephalometric measurement, and questionnaire survey. For instructors, it was divided into monitoring activity, cephalometric measurement and inter- and intra-class correlation coefficient of their measurements.

Part of students

Pre-test learning

The total teaching period consisted of 7 credit hours, including 4 credit hours of theoretical class and 3 credit hours for practical instruction. During the theoretical class, one instructor imparted relevant knowledge to the students in detail, including the positions of anatomical markers and the meanings of commonly used measurement items.

In the practical instruction class, another instructor guided 94 students to review the basic knowledge and showed them how to complete cephalometric measurement by the traditional method and digital technology (the Dolphin software[®]). For the traditional method, the sulfuric acid transfer paper was fixed to the radiograph with a clip. Then, the patient's soft tissue profile and hard tissue anatomical structures were depicted on the reading lamp. Finally, the commonly used anatomical points were identified on the sulfuric acid transfer paper and the measurement was completed with the ruler and protractor.

While using digital technology, the instructor adjusted results of line spacing on the lateral cranial radiograph to their actual size at first. Then, the instructor accomplished the measurement by adjusting the gray contrast value and other auxiliary methods. After learning the relevant knowledge of commonly used cephalometric measurement points and items, students were encouraged to review relevant contents after class.

Group and cephalometric measurement

One week after the end of the pre-test instruction, 94 students were randomly allocated into two groups through the RAND function in Excel software (Microsoft® Excel 2019MSO (2201 Build 16.0.14827.20198 version for 64 bit) and were required to complete the same cephalometric measurement by both traditional method and digital technology (the Dolphin Imaging® 11.8).

According to the cross-over design, one group took the traditional method first to complete the measurement and then used the Dolphin software®, while the other group completed in the opposite order. A total of 15 cephalometric items (Figure 2) were measured, such as the angle between Sella-Nasion plane and the Nasion-Subspinale plane (SNA), the angle between Sella-Nasion plane and the Nasion-Supramental plane (SNB), the angle between the Nasion-Subspinale plane and the Nasion-Supramental plane (ANB). The measurements of the traditional method and Dolphin software® were recorded, respectively. The collection, input and check of these data were completed by three authors.

Questionnaire survey

In order to survey the effectiveness of applying the digital software in cephalometric teaching and students' attitudes toward it (the Dolphin software®), we designed an online questionnaire and collected results by Wenjuanxing (Ranxing Co. Ltd., Changsha, China). For example, to investigate how difficult students considered cephalometry is, we set three options ranging from "very difficult", "kind of tough" to "easy". As to the mastery degree of students after the course, the options were "50-80%", "20-50%" to "0-20%". Gender of the participants was recorded from the university records. The detailed questionnaire list and corresponding options are shown in Table 1.

Part of instructors

Monitoring activity

During this pedagogical experiments, two instructors took responsibility for guiding students with cephalometric measurements, assuring the measurement order of two student groups, monitoring the process of giving out and recollecting the questionnaire, as well as assistance in the data analysis.

Cephalometric measurement & Inter- and intra-class correlation coefficient test

Two instructors also measured the same radiograph by two methods. One week later, they measured the same radiograph by two methods again. The measurements by one instructor at one-week's interval were used for calculating intra-class correlation coefficient (ICC_{intra}), and measurements by two instructors were used for calculating inter-class correlation coefficient (ICC_{inter}).

Statistical analysis

The quantitative data of cephalometric measurements were analyzed by the statistical software SPSS 18.0 (IBM Corporation, Armonk, NY, RRID:SCR_016479). The measurements of students by two methods were compared using the independent sample *t* test, as well as comparing them with corresponding standards, respectively, with the level of significance set as $P < 0.05$.

As for questionnaire data, we obtained the statistical data through the built-in function on the online questionnaire platform, as it provided the constituent ratio of each option and participants list. Then we performed a descriptive analysis of these results.

Anatomical points calibration

Before the course, the standard of anatomical markers on lateral cranial radiographs were calibrated. Firstly, two orthodontists reviewed the basic definitions and meanings of common anatomical points and items. Then, they measured a lateral cranial radiograph on paper and by software, followed by discussing and unifying the anatomical position standard. Subsequently, they respectively completed cephalometric measurement of another lateral cranial radiograph. The intraclass correlation coefficient (ICC) was greater than 90%, showing consistency of cephalometric measurement between them.




Figure 2. Illustration of 15 commonly used measurement items. 1. SNA: the angle between the sella-nasion plane and the nasion-subspinale plane; 2. SNB: the angle between sella-nasion plane and the nasion-supramental plane; 3. ANB: the angle between the nasion-subspinale plane and the nasion-supramental plane; 4. NP-FH: the posterior-inferior angle between the facial plane and the Frankfort horizontal plane; 5. NA-PA: the angle between the nasion-subspinale plane and the pogonion-subspinale plane; 6. U1-NA (mm): the distance from the upper central incisor tip to the nasion-subspinale plane; 7. U1-NA: the angle between the upper central incisor and the nasion-subspinale plane; 8. L1-NB (mm): the distance from the lower central incisor tip to the nasion-supramental plane; 9. L1-NB: the angle between the lower central incisor and the nasion-supramental plane; 10. U1-L1: the angle between the upper central incisor and the lower central incisor; 11. Y axis angle: the anterior-inferior angle between the Frankfort horizontal plane and the sella-pogonion plane; 12. Po-NB (mm): the distance from the pogonion point to the nasion-supramental plane; 13. FMA: the angle between the Frankfort horizontal plane and the mandibular plane; 14. FMIA: the angle between the Frankfort horizontal plane and the long axis of the lower central incisor; 15. IMPA: the angle between the long axis of the lower central incisor and the mandibular plane. The profile tracing was completed with the Uceph software (Uceph, Chengdu, China).

Table 1. Students' attitudes towards the application of digital technology.

Questionnaire list	Option list		
Are you interested in studying cephalometric measurement?	A: very interested (55%)	B: a little interested (43%)	C: not interested at all (2%)
How helpful do you think cephalometric measurement is to the diagnosis of malocclusion?	A: very helpful (77%)	B: a little helpful (21%)	C: not helpful at all (2%)
How tough do you think cephalometric measurement is?	A: very difficult (21%)	B: kind of tough (66%)	C: easy (13%)
Which do you prefer as the better teaching method?	A: digital technology (33%)	B: traditional method (2%)	C: both are acceptable (65%)
How well do you think you master the cephalometric measurement analysis?	A: 50%-80% (72%)	B: 20%-50% (28%)	C: 0-20% (0%)

Results

Standards of cephalometric measurement

As shown in **Supplementary Table 1 and 2**, the inter- and intra-ICC were all surpassed 90%, and there were no significant differences between two instructors' measurements, and the measurements by two methods. As the result, the means of two instructors' measurements were set as the standards.

Table 2. Comparison of students' measurements and standard values by the digital technology and traditional method.

Measurement items	Digital technology			Traditional method			P ₁	P ₂
	Test value (X±SD)	Standard value	d ₁	Test value (X±SD)	Standard value	d ₂		
SNA (°)	92.46±1.88	92.00	1.53	91.41±1.47	92.10	1.19	<0.01	<0.05
SNB (°)	88.47±0.86	88.30	0.69	88.03±1.35	88.30	0.94	0.01	<0.05
ANB (°)	3.76±1.83	3.70	1.43	3.50±0.86	3.80	0.75	0.25	<0.01
NP-FH (°)	89.32±1.62	90.70	1.83	88.95±1.61	90.90	1.89	0.13	0.76
NA-PA (°)	5.99±3.60	6.40	2.79	5.98±1.86	6.50	1.56	0.98	<0.01
U1-NA (mm)	5.07±1.86	6.20	1.71	5.20±1.41	6.30	1.35	0.61	0.06
U1-NA (°)	21.26±3.77	19.70	3.41	21.69±4.75	19.90	3.88	0.50	0.26
L1-NB (mm)	6.09±0.48	6.70	0.73	5.76±0.75	6.80	1.00	<0.01	<0.01
L1-NB (°)	25.10±2.24	26.45	2.19	25.69±3.40	26.60	2.68	0.16	0.09
U1-L1 (°)	129.68±4.67	130.25	4.04	130.16±5.97	130.50	4.82	0.54	0.08
Y axis (°)	64.45±0.99	62.80	1.79	64.22±1.93	62.70	1.91	0.32	0.52
Po-NB (mm)	1.13±0.46	1.05	0.37	1.21±0.44	1.02	0.35	0.21	0.54
FMA (°)	28.89±2.05	28.25	1.71	29.31±2.51	28.50	2.16	0.22	<0.05
FMIA (°)	63.46±2.78	63.75	2.27	63.36±3.52	63.80	2.90	0.83	<0.05
IMPA (°)	87.94±2.78	88.00	2.31	87.80±4.03	88.20	3.30	0.79	<0.01

X̄: Mean of students' measurements; SD: Standard deviation of students' measurements; Standard value: Mean of two orthodontists' measurements by traditional method or digital technology; d₁: Mean of absolute values of differences between students' results and standard values by digital software; d₂: Mean of absolute values of differences between students' results and standard values by traditional method; P₁: The P value of the independent sample t test on students' measurements by two methods; P₂: The P value of the independent sample t test on absolute values of differences between students' results and the standards by two methods.

SNA: the angle between the sella-nasion plane and the nasion-subspinale plane; SNB: the angle between sella-nasion plane and the nasion-supramental plane; ANB: the angle between the nasion-subspinale plane and the nasion-supramental plane; NP-FH: the posterior-inferior angle between the facial plane and the Frankfort horizontal plane; NA-PA: the angle between the nasion-subspinale plane and the pogonion-subspinale plane; U1-NA (mm): the distance from the upper central incisor tip to the nasion-subspinale plane; U1-NA: the angle between the upper central incisor and the nasion-subspinale plane; L1-NB (mm): the distance from the lower central incisor tip to the nasion-supramental plane; L1-NB: the angle between the lower central incisor and the nasion-supramental plane; U1-L1: the angle between the upper central incisor and the lower central incisor; Y axis angle: the anterior-inferior angle between the Frankfort horizontal plane and the sella-pogonion plane; Po-NB (mm): the distance from the pogonion point to the nasion-supramental plane; FMA: the angle between the Frankfort horizontal plane and the mandibular plane; FMIA: the angle between the Frankfort horizontal plane and the long axis of the lower central incisor; IMPA: the angle between the long axis of the lower central incisor and the mandibular plane.

Accuracy of students' cephalometric measurement

Statistically significant differences were observed in measurements of SNA, SNB and the distance from the lower central incisor tip to the nasion-supramental plane (L1-NB (mm)) between the traditional method and digital technology ($P_{1-SNA}<0.01$, $P_{1-SNB}=0.01$, $P_{1-L1-NB (mm)}<0.01$) (Table 2), while other items showed no significant differences. Besides, the measurements by digital technology were closer to the standard values than those by the traditional method. The accuracy of five items measurements using digital technology was significantly higher, including SNB, L1-NB (mm), the angle between the Frankfort horizontal plane and the mandibular plane (FMA), the angle between the Frankfort horizontal plane and the long axis of the lower central incisor (FMIA), and the angle between the long axis of the lower central incisor and the mandibular plane (IMPA) ($P_{2-SNB}<0.05$, $P_{2-L1-NB (mm)}<0.01$, $P_{2-FMA}<0.05$, $P_{2-FMIA}<0.05$, $P_{2-IMPA}<0.01$). However, five items presented the opposite result, among which three items were statistically significant (SNA, ANB and the angle between the Nasion-Subspinale plane and the pogonion-subspinale plane (NA-PA) ($P_{2-SNA}<0.05$, $P_{2-ANB}<0.01$, $P_{2-NA-PA}<0.01$).

Attitudes of students towards the digital technology

We assigned the questionnaires to all the participants with 82 of 94 students filling out the questionnaire and the response rate was 87%. The statistical results of the questionnaire are shown in Table 1. Among the respondents, 66% thought studying cephalometry was very difficult and 21% thought it was kind of tough. After instruction, review and practice, 72% of them considered they had mastered 50-80% of relevant knowledge and a few students thought they had mastered 20-50%. About 33% of students preferred the digital technology than traditional method (2%) as a better teaching method and 65% held that both were acceptable, which indicated good acceptance by students of digital technology applied in the teaching process. In addition, 98% of participants expressed their interest in studying cephalometry and considered cephalometric analysis helpful to diagnosis of malocclusion.

Discussion

Cephalometric measurement is essential for diagnosis and treatment plan design of patients with malocclusions, which are the most significant procedures in orthodontic clinical work. Orthodontic educators put forward that orthodontic teaching for undergraduates should focus on diagnosis and recognition of problems.²⁰ However, there is not a generally accepted teaching method, and a wide variation of course durations and contents exist in different dental colleges and faculties.²¹ How to arouse students' interest and achieve better teaching effects is a major problem faced by orthodontic educators. Since only a few reports explored this aspect, we designed this didactical experiment.

In our study, three cephalometric items measured by two methods were statistically different, while other items were basically similar. These results implied that digital technology could achieve similar results to the traditional method during cephalometric measurement. Additionally, the majority of items measured by digital technology were closer to the standards, including five statistically significant items, which suggested students could achieve more accurate results by digital technology. This may be attributed to the function of digital software to adjust the gray contrast value of X-ray films (Supplementary figure 1), making it easier to identify the unclear points on the printed paper. In addition, the automatic generation of results also helped to avoid evitable errors during manual measurement. These results were in accordance with previous studies, which suggested that the accuracy of digital measurement on 3-dimensional cone beam computed tomography images was basically similar to or even higher than that of manual measurement.^{22,23}

However, in spite of the convenience it provided, the digital technology may lead to a lack of deep understanding of corresponding contents, such as definitions and meanings of these measurement items. The traditional method could better cultivate practical abilities of students and enhance their memory of relevant knowledge. Besides, although the accuracy of some items obtained by digital technology were significantly higher, a few items showed the opposite result (SNA, ANB and NA-PA). The subspinale point (the A point) was associated with these three items, which suggested that the traditional method was more accurate than digital technology in positioning the subspinale point on the lateral cranial radiograph. The subspinale point is the most concave point of the arc from the anterior nasal spine point (the ANS point) to the superior prosthion point (the Spr point). Compared with digital measurement, the advantage of manual measurement is that the arc can be traced on paper, and some auxiliary instruments like ruler and protractor can help to locate the subspinale point (Supplementary figure 2). These results indicated abundant experience was required to identify the subspinale point, reminding both orthodontic educators and students to devote more time and energy to deep learning it.

Before the course, the majority of students showed fear towards abstract and complex concepts, assuming cephalometric measurement was difficult to master. However, with instruction and practice, most students could master 50-80% of relevant knowledge, which could result from digital technology realizing visualization of numerous and complicated anatomical markers. Previous studies found that visualization was extremely attractive to young students, and significantly aroused their interest and sense of participation.^{24,25} Our survey confirmed that more students preferred digital

technology (33%) as a better teaching method than the traditional method. As a result, the application of digital technology in teaching cephalometric measurement was widely accepted by students and contributed to favorable teaching results.

This study still had some limitations. For example, the sample size could be further enlarged. Secondly, the measurement time for students were not strictly required, which may have resulted in the underperformances or supernormal performances of students. Thirdly, exploration of student's attitudes towards digital technology being applied in this course was not sufficient. The mentioned issues needed to be improved may interrupt us from to accurately assessing the real advantages and disadvantages of the appliance of this digital technology. These findings could guide and encourage university orthodontic teachers to apply this technology in cephalometric teaching and pay more attention to considering the position identification of the subspinale point.

Conclusion

This study investigated the effect of applying the digital technology (the Dolphin software®) in cephalometric learning, demonstrating better effect and acceptance of this technology.

Data availability

Extended data

Figshare: Evaluating the effect of digital technology on the learning of orthodontic cephalometric measurement, <https://doi.org/10.6084/m9.figshare.21550599>.²⁶

This project contains the following extended data:

- Supplementary File.docx (Supplementary figure 1, 2; Supplementary table 1, 2)
- The results of students' measurements.xlsx
- Statistical results of the questionnaire.xlsx

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/) (CC-BY 4.0).

References

1. Ditmarov A: **Orthodontics: Orthodontics vs orthodontiya.** *Br. Dent. J.* Jul 13 2018; **225**(1): 2. [PubMed Abstract](#) | [Publisher Full Text](#)
2. Jheon AH, Oberoi S, Solem RC, *et al.*: **Moving towards precision orthodontics: An evolving paradigm shift in the planning and delivery of customized orthodontic therapy.** *Orthod. Craniofac. Res.* Jun 2017; **20**(Suppl 1): 106–113. [PubMed Abstract](#) | [Publisher Full Text](#)
3. Drage NA, Atkin PA, Farnell DJJ: **Dental and maxillofacial radiology: confidence, knowledge and skills in the newly graduated dentist.** *Br. Dent. J.* Apr 2020; **228**(7): 546–550. [Publisher Full Text](#)
4. Zitzmann NU, Matthisson L, Ohla H, *et al.*: **Digital Undergraduate Education in Dentistry: A Systematic Review.** *Int. J. Environ. Res. Public Health.* May 7 2020; **17**(9). [PubMed Abstract](#) | [Publisher Full Text](#)
5. Inquimbert C, Tramini P, Romieu O, *et al.*: **Pedagogical Evaluation of Digital Technology to Enhance Dental Student Learning.** *Eur. J. Dent.* Feb 2019; **13**(1): 053–057. [PubMed Abstract](#) | [Publisher Full Text](#)
6. Silveira MS, Cogo ALP: **The contributions of digital technologies in the teaching of nursing skills: an integrative review.** *Rev. Gaucha Enferm.* Jul 13 2017; **38**(2): e66204. Contribuições das tecnologias educacionais digitais no ensino de habilidades de enfermagem: revisão integrativa. [PubMed Abstract](#) | [Publisher Full Text](#)
7. Buchanan JA: **Use of simulation technology in dental education.** *J. Dent. Educ.* Nov 2001; **65**(11): 1225–1231. [Publisher Full Text](#)
8. Urbankova A: **Impact of computerized dental simulation training on preclinical operative dentistry examination scores.** *J. Dent. Educ.* Apr 2010; **74**(4): 402–409. [PubMed Abstract](#) | [Publisher Full Text](#)
9. Al-Jewair TS, Qutub AF, Malkhassian G, *et al.*: **A systematic review of computer-assisted learning in endodontic education.** *J. Dent. Educ.* Jun 2010; **74**(6): 601–611. [Publisher Full Text](#)
10. Abbey LM, Arnold P, Halunko L, *et al.*: **CASE STUDIES for Dentistry: development of a tool to author interactive, multimedia, computer-based patient simulations.** *J. Dent. Educ.* Dec 2003; **67**(12): 1345–54.
11. Al-Jewair TS, Azarpazhooh A, Suri S, *et al.*: **Computer-assisted learning in orthodontic education: a systematic review and meta-analysis.** *J. Dent. Educ.* Jun 2009; **73**(6): 730–739. [PubMed Abstract](#) | [Publisher Full Text](#)
12. Rosenberg H, Grad HA, Matear DW: **The effectiveness of computer-aided, self-instructional programs in dental education: a systematic review of the literature.** *J. Dent. Educ.* May 2003; **67**(5): 524–532. [PubMed Abstract](#) | [Publisher Full Text](#)
13. Schott TC, Arsalan R, Weimer K: **Students' perspectives on the use of digital versus conventional dental impression techniques in orthodontics.** *BMC Med. Educ.* Mar 12 2019; **19**(1): 81. [PubMed Abstract](#) | [Publisher Full Text](#)
14. Ren Q, Wang Y, Zheng Q, *et al.*: **Survey of student attitudes towards digital simulation technologies at a dental school in China.** *Eur. J. Dent. Educ.* Aug 2017; **21**(3): 180–186. [Publisher Full Text](#)
15. Nagy ZA, Simon B, Tóth Z, *et al.*: **Evaluating the efficiency of the Dental Teacher system as a digital preclinical teaching tool.** *Eur.*

- J. Dent. Educ.* Aug 2018; **22**(3): e619–e623.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Liu L, Li J, Yuan S, *et al.*: **Evaluating the effectiveness of a preclinical practice of tooth preparation using digital training system: A randomised controlled trial.** *Eur. J. Dent. Educ.* Nov 2018; **22**(4): e679–e686.
[PubMed Abstract](#) | [Publisher Full Text](#)
 17. Chen SK, Chen YJ, Yao CC, *et al.*: **Enhanced speed and precision of measurement in a computer-assisted digital cephalometric analysis system.** *Angle Orthod.* Aug 2004; **74**(4): 501–507.
[Publisher Full Text](#)
 18. Farooq MU, Khan MA, Imran S, *et al.*: **Assessing the Reliability of Digitalized Cephalometric Analysis in Comparison with Manual Cephalometric Analysis.** *J. Clin. Diagn. Res.* Oct 2016; **10**(10): Zc20–zc23.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Hongyu Chen CW: **Wenli Lai The introduction of Dolphin software and its clinical application in orthodontics.** *Journal of Chinese Physician.* 2015; **17**(04): 611–613.
 20. Rock WP, O'Brien KD, Stephens CD: **Orthodontic teaching practice and undergraduate knowledge in British dental schools.** *Br. Dent. J.* Mar 23 2002; **192**(6): 347–351.
[PubMed Abstract](#) | [Publisher Full Text](#)
 21. Derringer KA: **Undergraduate orthodontic teaching in UK dental schools.** *Br. Dent. J.* Aug 27 2005; **199**(4): 224–232.
[PubMed Abstract](#) | [Publisher Full Text](#)
 22. Shahidi S, Oshagh M, Gozin F, *et al.*: **Accuracy of computerized automatic identification of cephalometric landmarks by a designed software.** *Dentomaxillofac. Radiol.* 2013; **42**(1): 20110187.
[PubMed Abstract](#) | [Publisher Full Text](#)
 23. Gupta A, Kharbanda OP, Sardana V, *et al.*: **Accuracy of 3D cephalometric measurements based on an automatic knowledge-based landmark detection algorithm.** *Int. J. Comput. Assist. Radiol. Surg.* Jul 2016; **11**(7): 1297–1309.
[PubMed Abstract](#) | [Publisher Full Text](#)
 24. Triepels CPR, Smeets CFA, Notten KJB, *et al.*: **Does three-dimensional anatomy improve student understanding?** *Clin. Anat.* Jan 2020; **33**(1): 25–33.
[PubMed Abstract](#) | [Publisher Full Text](#)
 25. Amer RS, Denehy GE, Cobb DS, *et al.*: **Development and evaluation of an interactive dental video game to teach dentin bonding.** *J. Dent. Educ.* Jun 2011; **75**(6): 823–831.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. Tian, Pan Y: **Evaluating the effect of digital technology on the learning of orthodontic cephalometric measurement.** figshare. Dataset. 2022.
[Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 3

Reviewer Report 24 November 2023

<https://doi.org/10.5256/f1000research.144525.r178260>

© 2023 Corte-Real A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Ana Corte-Real 

¹ Faculty of Medicine, University of Coimbra, Coimbra, Portugal

² Forensic Dentistry Laboratory, Faculty of Medicine, University of Coimbra, Coimbra, Portugal

The authors correspond positively to the suggestions.

Competing Interests: No competing interests were disclosed.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 2

Reviewer Report 11 January 2023

<https://doi.org/10.5256/f1000research.140872.r157996>

© 2023 Corte-Real A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Ana Corte-Real 

¹ Faculty of Medicine, University of Coimbra, Coimbra, Portugal

² Forensic Dentistry Laboratory, Faculty of Medicine, University of Coimbra, Coimbra, Portugal

The authors improved the manuscript. However, some suggestions should be pointed out in the Methods section.

- "All the students use (/used?) the Dolphin software before"... is it really what the authors intended to write?

- The original information of the manuscript is its impact on orthodontic learning and students' new skills and satisfaction (instead of "student's acceptance towards it"). It is difficult to understand the sentence, "The collection and input of these data were completed by three postgraduates, with two postgraduates responsible for the inputting and the third one in charge of checking." It should be clarified. The student's contribution to the cephalometric analysis and their performance and satisfaction evaluations should be clear.
- Intra and inter-class correlations were based on the instructors' data. What is its impact on the present study?

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medical and Dentistry Higher Education and Health Quality

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 20 Feb 2023

Yu Tian

The authors improved the manuscript. However, some suggestions should be pointed out in the Methods section.

Response: We are grateful for the reviewer's helpful suggestions. As shown below, we have addressed the reviewer's concerns, and provided point-by-point responses. We hope that the changes meet your approval.

1. "All the students use (/used?) the Dolphin software before"... is it really what the authors intended to write?

Response: Thank you for pointing out this mistake. We have revised the sentence to "All the students have neither learned about the method of measuring the lateral cranial radiographs nor used the Dolphin software before."

2. The original information of the manuscript is its impact on orthodontic learning and students' new skills and satisfaction (instead of "student's acceptance towards it").

Response: Thank you for your suggestion. According to your advice, we have revised the phrase to "orthodontic learning, students' new skills and satisfaction towards it".

3. It is difficult to understand the sentence, "The collection and input of these data were completed by three postgraduates, with two postgraduates responsible for the inputting and the third one in charge of checking." It should be clarified. The student's contribution to the cephalometric analysis and their performance and satisfaction evaluations should be clear.

Response: Thank you for your suggestion. To make it clear, we have revised the sentence to “The collection, input and check of these data were completed by three authors.”

4. Intra and inter-class correlations were based on the instructors' data. What is its impact on the present study?

Response: Thank you for your question. In order to evaluate students' performances, we needed standards. Thus, we invited the experienced instructors and set their measurement results as standards after checking the stability and consistency of their measurements.

Competing Interests: No competing interests existed.

Reviewer Report 04 January 2023

<https://doi.org/10.5256/f1000research.140872.r157997>

© 2023 Maspero C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Cinzia Maspero 

¹ Department of Orthodontics, Faculty of Medicine, University of Milan, Milan, Italy

² Fondazione IRCCS Cà Granda, Ospedale Maggiore Policlinico, Milan, Italy

All my requirements have been addressed. The manuscript has been improved and, in my opinion, it is suitable for indexing.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Orthodontic, Orthognati surgery, cephalometric, maxillary expansion, interceptive orthodontics, fixed appliances, clear aligners, pediatric dentistry

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 07 September 2022

<https://doi.org/10.5256/f1000research.121428.r135780>

© 2022 Corte-Real A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Ana Corte-Real 

¹ Faculty of Medicine, University of Coimbra, Coimbra, Portugal

² Forensic Dentistry Laboratory, Faculty of Medicine, University of Coimbra, Coimbra, Portugal

The topic is interesting for assessing learning from the student's perspective, but the title does not correspond to the content. The methodology section is the weakest part of the work. The authors should adjust the study design, including the analysis of intra and inter-observer accuracy and monitoring activities during the study. The selection of participants should include the inclusion and exclusion criteria as the previous use of digital technology for cephalometric measurements impacts the students' skills.

Is the work clearly and accurately presented and does it cite the current literature?

No

Is the study design appropriate and is the work technically sound?

No

Are sufficient details of methods and analysis provided to allow replication by others?

No

If applicable, is the statistical analysis and its interpretation appropriate?

No

Are all the source data underlying the results available to ensure full reproducibility?

No

Are the conclusions drawn adequately supported by the results?

No

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Medical and Dentistry Higher Education and Health Quality

I confirm that I have read this submission and believe that I have an appropriate level of expertise to state that I do not consider it to be of an acceptable scientific standard, for reasons outlined above.

Author Response 07 Nov 2022

Yu Tian

Response: We are grateful for the reviewer's helpful suggestions. As shown below, we have

addressed the reviewer's concerns, and provided point-by-point responses. We hope that the changes meet your approval.

1. The topic is interesting for assessing learning from the student's perspective, but the title does not correspond to the content.

Response: Thank you for your valuable suggestions. We fully understand and agree with your suggestion about the relationship between the title and content of this article. We've revised the title to make it more suitable for this study "Evaluating the effect of digital technology on the learning of orthodontic cephalometric measurement".

2. The methodology section is the weakest part of the work. The authors should adjust the study design, including the analysis of intra and inter-observer accuracy and monitoring activities during the study.

Response: Thank you for your valuable suggestion. We added in **Methods** section that "The measurements by one instructor at one-week's interval were used for calculating intra-class correlation coefficient (ICC_{intra}), and measurements by two instructors were used for calculating inter-class correlation coefficient (ICC_{inter}).", and in **Standards of cephalometric measurement** section that "The inter- and intra-ICC were all surpassed 90%, and there were no significant differences between two instructors' measurements, and the measurements by two methods".

As regard to the monitoring activity, we've added that "During this pedagogical experiments, two instructors took responsibility for guiding students with cephalometric measurements, assuring the measurement order of two student groups, monitoring the process of giving out and recollecting the questionnaire, as well as assistance in the data analysis." in the **Monitoring activity** section in page 8.

3. The selection of participants should include the inclusion and exclusion criteria as the previous use of digital technology for cephalometric measurements impacts the students' skills.

Response: Thank you for your suggestions. We agree with your advice on the selection of participant students. We've added in "Participants" section in line 5, page 6 that "...All the students did not learn about the method of measuring the lateral cranial radiographs and used the Dolphin software before".

Competing Interests: No competing interests were disclosed.

Reviewer Report 07 September 2022

<https://doi.org/10.5256/f1000research.121428.r148740>

© 2022 Maspero C. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? Cinzia Maspero 

¹ Department of Orthodontics, Faculty of Medicine, University of Milan, Milan, Italy

² Fondazione IRCCS Cà Granda, Ospedale Maggiore Policlinico, Milan, Italy

Thank you for submitting your research. The manuscript is interesting but some changes are necessary before taking it into consideration for publication. Here are my concerns:

- ABSTRACT: "Besides, the results of most items by digital technology were closer to the **standards** than those by the traditional method" What do you mean with standards?
- INTRODUCTION: replace the word subdiscipline with discipline.
- "These factors may not only affect accuracy of measurements, but also contribute to destruction of students' enthusiasm for further learning." Are you sure with this sentence?
- STUDY DESIGN: pre-test instruction. Please better explain what do you mean and what instruction have been given.
- FLOW CHART: anatomical point calibration: calibrated their standard. What do you mean? Specify which standard.
- CEFALOMETRIC MEASUREMENTS: it is not clear if the same lateral telerradiography have been used for all the students. Better specify.
- Illustration of 15 commonly used measurement items: which kind of tracing have been used?
- Please better specify the aim of this study.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Orthodontic, Orthognati surgery, cephalometric, maxillary expansion, interceptive orthodontics, fixed appliances, clear aligners, pediatric dentistry

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 07 Nov 2022

Yu Tian

Thank you for submitting your research. The manuscript is interesting but some changes are necessary before taking it into consideration for publication. Here are my concerns:

Response: We are grateful for the reviewer's helpful suggestions. As shown below, we have addressed the reviewer's concerns, and provided point-by-point responses. We hope that the changes meet your approval.

1. ABSTRACT: "Besides, the results of most items by digital technology were closer to the standards than those by the traditional method" What do you mean with standards?

Response: Thank you for your valuable suggestion. The means of two instructors' measurements were set as the standards, and we've added this in the abstract (line 4 in Results, page 2).

2. INTRODUCTION: replace the word subdiscipline with discipline.

Response: Thank you for your suggestion. According to your advice, we've used the word "discipline" in line 1, page3.

3. "These factors may not only affect accuracy of measurements, but also contribute to destruction of students' enthusiasm for further learning." Are you sure with this sentence?

Response: Thank you for your advice. We've deleted the sentence.

4. STUDY DESIGN: pre-test instruction. Please better explain what do you mean and what instruction have been given.

Response: Thank you for your suggestion. We've checked the manuscript, and described the pre-test instruction in detail in page 6, which included 7 credit hours and let students learn about the 15 cephalometric items and two measurement methods.

5. FLOW CHART: anatomical point calibration: calibrated their standard. What do you mean? Specify which standard.

Response: Thank you for your valuable suggestions. We revised the subtitle and described in detail that "Before the course, the inter- and intra-class correlation coefficient of two instructors' measurements were tested. Firstly, two orthodontists reviewed and discussed the locations of anatomical points. Then, they measured a lateral cranial radiograph by traditional and software methods. One week later, they measured the same radiograph by two methods again. The measurements by one instructor at one-week's interval were used

for calculating intra-class correlation coefficient (ICC_{intra}), and measurements by two instructors were used for calculating inter-class correlation coefficient (ICC_{inter})” in **Inter- and intra-class correlation coefficient of two instructors’ measurements** section.

6. CEPHALOMETRIC MEASUREMENTS: it is not clear if the same lateral teleradiography have been used for all the students. Better specify.

Response: Thank you for your suggestion. We’ve added in line 2, page 7 that “One week after the end of the pre-test instruction, 94 students were randomly allocated into two groups through the RAND function in Excel software (Microsoft® Excel® 2019MSO (2201 Build 16.0.14827.20198 version for 64 bit) and were required to complete cephalometric measurement of the same lateral cranial radiograph by both traditional method and digital technology (the Dolphin Imaging® 11.8).”.

7. Illustration of 15 commonly used measurement items: which kind of tracing have been used?

Response: Thank you for your question. The tracing was completed with the Uceph software, and we’ve added “The profile tracing was completed with the Uceph software (Uceph, Chengdu, China).” in the illustration of **Figure 2**.

8. Please better specify the aim of this study.

Response: Thank you for your suggestion. We’ve revised in abstract and introduction section that “This study aimed to evaluate the effect of applying digital technology in cephalometric measurement teaching and students’ acceptance towards it”.

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research