

Polish Heart Journal

The Official Peer-reviewed Journal of the Polish Cardiac Society since 1957

Online first

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon

ISSN 0022-9032 e-ISSN 1897-4279

Sudden cardiac arrest in the setting of coronary artery ectasia: Mechanistic and clinical perspectives. Author's reply

Authors: Małgorzata Zalewska-Adamiec, Maciej Południewski, Hanna Bachórzewska-

Gajewska, Sławomir Dobrzycki

Article type: Letter to the Editor

Received: December 23, 2023

Accepted: December 27, 2023

Early publication date: December 28, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Sudden cardiac arrest in the setting of coronary artery ectasia: Mechanistic and clinical

perspectives. Author's reply

Małgorzata Zalewska-Adamiec, Maciej Południewski, Hanna Bachórzewska-Gajewska,

Sławomir Dobrzycki

Department of Invasive Cardiology, Medical University in Bialystok, Białystok, Poland

Short title: Sudden cardiac arrest in coronary artery ectasia.

Correspondence to:

Małgorzata Zalewska-Adamiec, MD,

Department of Invasive Cardiology,

Medical University in Bialystok,

Skłodowskiej-Curie 24A, 15–276 Białystok,

phone: +48 603 784 468,

e-mail: mzalewska5@wp.pl

We would like to thank Yalta et al. [1] for your interest in our report on sudden cardiac arrest

(SCA) during diagnostic coronary angiography in a patient with coronary artery ectasias (CAE)

[2].

We believe that coronary artery ectasias constitute a very important clinical problem, both

diagnostic and therapeutic, therefore the occurrence of such a serious complication as SCA in

our patient with CAE motivated us to describe this case. Unfortunately, the limited number of

words in the clinical vignette prevents a thorough discussion of all clinical aspects. Therefore,

we are especially grateful for all the valuable comments of Yalta et al., to which we can respond

here.

Yalta et al. presented possible causes of SCA in patients with CAE in clinical implications:

• Myocardial ischemia due to slowed flow at the macrovascular level. We consider this

aspect to be the most likely cause of the angina pain reported by our patient.

• Severe dysfunction of coronary microcirculation responsible for myocardial ischemia.

We cannot rule out microcirculation disorders in our patient, but we have not currently

2

planned additional tests, including positron emission tomography. Further diagnosis of

the causes of ischemia in the patient depends on the further clinical course.

Possible vasospastic component requiring appropriate pharmacological treatment [3].

In our patient, we did not find a typical history of vasospastic angina. The patient

received typical pharmacological treatment (acetylsalicylic acid 75 mg/day, cilazapril 5

mg/day, amlodipine 5 mg/day, bisoprolol 3.75 mg/day and rosuvastatin 10 mg/day).

Trimetazidine 2×35 mg/day was added to the treatment.

Occurrence of acute coronary syndromes as a result of peripheral embolism of the distal

sections of coronary arteries. Our patient has not had any acute coronary syndrome to

date.

Mechanical complications of ectatically dilated arteries (rupture, fistulas).

Percutaneous and cardiac surgical interventions in patients with advanced coronary

artery ectasias resistant to pharmacological treatment. Qualifying these patients for

interventional treatment is extremely difficult and requires decision-making within the

Heart Team and often additional hemodynamic tests, e.g. fractionated flow reserve [4].

Analyzing all possible mechanisms of SCA in our patient, we considered the most likely cause

to be slow flow of the injected contrast agent in the ectatically dilated left coronary artery.

We referred our patient for further outpatient cardiological care with the recommendation of

regular electrocardiography monitoring using the Holter method. However, the in-depth

diagnostics (positron emission tomography, imaging of coronary arteries) depend on the

patient's clinical condition.

To sum up, the presented case of this patient and the demonstrated clinical implications related

to coronary artery ectasias indicate the need for special cardiological care for these patients,

taking into account various diagnostic tests and therapeutic methods. However, maintaining

registries and long-term observational studies of patients with CAE would allow for the

development of recommendations for the management of these patients in long-term care.

Article information

Conflict of interest: None declared.

Funding: None.

Open access: This article is available in open access under Creative Common Attribution-Non-

Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, which allows

downloading and sharing articles with others as long as they credit the authors and the publisher,

3

but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl

REFERENCES

- 1. Yalta K, Palabıyık O. Sudden cardiac arrest in the setting of coronary artery ectasia: Mechanistic and clinical perspectives. Kardiol Pol. 2023.
- Zalewska-Adamiec M, Południewski M, Bachórzewska-Gajewska H, et al. Slow flow in ectatic dilated coronary arteries as the cause of sudden cardiac arrest during diagnostic coronary angiography. Kardiol Pol. 2023 [Epub ahead of print], doi: 10.33963/v.kp.97726, indexed in Pubmed: 37997829.
- 3. Khedr A, Neupane B, Proskuriakova E, et al. Pharmacologic management of coronary artery ectasia. Cureus. 2021; 13(9): e17832, doi: 10.7759/cureus.17832, indexed in Pubmed: 34660041.
- Zalewska-Adamiec M, Kuzma L, Bachorzewska-Gajewska H, et al. Fractional flow reserve in the diagnosis of ischemic heart disease in a patient with coronary artery ectasia. Diagnostics (Basel). 2021; 12(1): 17, doi: 10.3390/diagnostics12010017, indexed in Pubmed: 35054184.