

Scotland's Rural College

Cryopreservation of a soil microbiome using a Stirling 1 cycle approach - a genomic assessment

Cafa, Giovanni; Bonnin, Juan Miguel; Holden, Nicola; Malone, Jacob G.; Mauchline, Tim H.; Clark, Ian; Taketani, Rodrigo; Ryan, Matthew J.

DOI:

[10.31220/agrirxiv.2021.00066](https://doi.org/10.31220/agrirxiv.2021.00066)

Print publication: 16/07/2021

Document Version

Early version, also known as preprint

[Link to publication](#)

Citation for published version (APA):

Cafa, G., Bonnin, J. M., Holden, N., Malone, J. G., Mauchline, T. H., Clark, I., Taketani, R., & Ryan, M. J. (2021). *Cryopreservation of a soil microbiome using a Stirling 1 cycle approach - a genomic assessment*. (agriRxiv). <https://doi.org/10.31220/agrirxiv.2021.00066>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Cryopreservation of a soil microbiome using a Stirling Cycle approach – a genomic assessment

Author names

Giovanni Cafà (<https://orcid.org/0000-0003-0819-2126>)¹, Juan Miguel Bonnin (<https://orcid.org/0000-0002-5095-6719>)¹, Nicola Holden (<https://orcid.org/0000-0002-7904-4529>)², Jacob G Malone^{3,4} (<https://orcid.org/0000-0003-1959-6820>), Tim H Mauchline (<https://orcid.org/0000-0002-8590-2614>)⁵, Ian Clark (<https://orcid.org/0000-0002-1589-5420>)⁵, Rodrigo Taketani⁵, Matthew J Ryan (<https://orcid.org/0000-0002-3689-6609>)^{1*}

Affiliation

¹ CABI, Egham, TW20 9TY

² SRUC, Department of Rural Land Use, Craibstone Estate, Aberdeen, AB21 9YA

³ Molecular Microbiology, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH

⁴ University of East Anglia, Norwich, NR4 7TJ

⁵ Sustainable Agriculture Sciences, Rothamsted Research, Harpenden, AL5 2JQ

Corresponding author

* Matthew J Ryan m.ryan@cabi.org

Abstract

Soil microbiomes are dynamic systems that respond to biotic and abiotic environmental factors such as those presented at seasonal scales or due to long-term anthropogenic regime shifts. These can affect the composition and function of microbiomes. Investigation of microbiomes can uncover hidden microbial roles in health and disease and discover microbiome-based interventions. Collections of soil samples are kept by various institutions in either a refrigerated or occasionally frozen state, but conditions are not optimised to ensure the integrity of soil microbiome. In this manuscript, we describe cryopreservation with a controlled rate cooler and estimate the genomic content of an exemplar soil sample before and after cryopreservation. The first hypothesis was to test the genomic integrity of the microbiome. We also enriched the soil sample with a liquid medium to estimate the growth of bacteria and compared their growth before and after cryopreservation. Sequence-based rRNA metabarcoding was used to demonstrate that the controlled rate cooler maintains intact the DNA content of the microbiome. Two methods of cryopreservation were applied and compared with control aliquots of soil. An optimised cryopreservation of soil samples is

34 essential for the development of microbiome research in order to retain stable, functionally intact
35 microbiomes. Our results showed that metabarcoding of 16S and ITS rRNA were useful methods to
36 estimate successful cryopreservation. The soil microbiome after enrichment with liquid medium
37 exhibited a similar response of cryopreserved soil and this was estimated with the comparison of the
38 ten most abundant bacterial taxa. These findings support a successful process of cryopreservation
39 and are promising for future use of this technology. To the best of our knowledge, this study is the
40 first report of cryopreservation of soil using a Stirling cycle cooling approach.

41

42 **Introduction**

43

44 Soil microbiomes are dynamic systems that respond to biotic and abiotic environmental factors in a
45 temporally and spatially dependent manner, such as those presented at seasonal scales or due to
46 long-term anthropogenic regime shifts, affecting their composition and function and their
47 environment (1,2). They offer great potential to contribute to the sustainable intensification of
48 agriculture and their application could reduce our dependence on the use of toxic chemicals in
49 agriculture and stimulate a more sustainable application of environmental resources (3,4).
50 Microbiome research and application may impact anthropogenic-driven climate change (5) and
51 developing such an understanding is essential to tackle challenges facing human society today, such
52 as the management of natural ecosystems and the mitigation of climate change (6).

53 To ensure robust application of whole beneficial microbiomes, methods for their preservation are
54 required. Ideally, such methods will include the preservation of biotic components, abiotic factors as
55 well as the physical environment where organisms function, such as the space where chemical
56 reactions occur. Altering these factors will likely influence the integrity and function of preserved
57 microbiomes when are subsequently resuscitated. For example, the removal of keystone organisms
58 encoding functional genes due to the use of a sub-optimal storage methodology could irreversibly
59 and negatively affect biological networks if these genes encode critical functions in a given
60 microbiome system (7). As such, preservation of the community complex is critical (8). Storage
61 capacity is a significant challenge to this approach. It is not practical to store large amounts of
62 material such as soil, hence an understanding of the amount of sample required to be representative
63 of the microbiome is required. For example, in agriculture a single field encompasses multiples of
64 localized microbiomes (9).

65 The development of preservation methods to conserve samples as representative 'snap shots' in
66 time is becoming a crucial part of microbiome research, as the requirement for reference material to
67 repeat and validate research outcomes and its utilisation as a source of material for potential
68 commercial use increases (7). Proving that the functionality of the microbiomes remain intact after
69 cryopreservation is therefore critical. There is evidence that only a fraction of the microbiome can
70 survive traditional cryopreservation and many organisms must therefore be isolated and cultured
71 under specific conditions before freezing. As such, there is a need to improve microbiome
72 cryopreservation methods.

73 The rapid advancement of nucleic acid sequencing technologies has precipitated an urgent
74 assessment of the role of biobanks to preserve microbiomes and underpin research to unlock their
75 functional potential (7). Preservation of metagenomes from these samples allows microbiomes to be
76 described in unprecedented detail and the functional potential of a given microbiome to be
77 deciphered. However, the analysis of nucleic acids does not provide information on the viability of
78 organisms, even though RNA analysis does give insights into microbial activity at the time of
79 preservation.

80 When considering microbiome preservation there are two questions that need to be answered: (i)
81 what should be preserved, and (ii) what is the best way of preserving it? (7). Historically, soil samples
82 have been frozen and stored in mechanical freezers without any control of the rate of freezing. The
83 application of a sub-optimal approach can compromise the microbial communities present as some
84 will not survive the freezing process. The use of a Stirling cycle freezer for cryopreservation is
85 considered to have significant advantages over traditional methodologies including nitrogen free
86 operation, application of low cooling rates, reduction of sample contamination risks and control of
87 ice nucleation (10). For the majority of organisms, applications of controlled cooling techniques
88 reduce the prospect of ice damage that would otherwise compromise membrane integrity and cell
89 viability. Assessment of cryopreservation regimes have included several tests to analyse viability of
90 microorganisms in addition to the relative stability of genomic profiles. Stirling cycle cooler has
91 shown recovery of fungi that was 97% effective, fungi from all major groups that were recovered,
92 including recognised cryopreservation-recalcitrant strains (10). The control of cooling is much more
93 finite with the Stirling cooler, reducing the variation in ice nucleation between replicates (10).

94 In this manuscript, we describe cryopreservation with a controlled rate cooler and estimate the
95 genomic content of a selected soil sample before and after cryopreservation. Two methods of
96 cryopreservation were used and compared with control aliquots of soil. Metabarcoding was used to
97 estimate the genomic content of a soil microbiome. We selected a representative soil sample from
98 the Rothamsted collection and used it to estimate the genomic integrity after cryopreservation. The
99 first hypothesis was to confirm the genomic integrity of the microbiome. We also treated the soil
100 sample with a liquid medium to estimate the growth of bacteria and compared their growth before
101 and after cryopreservation. We used PCR-based rRNA metabarcoding to demonstrate that the
102 controlled rate cooler maintains the DNA content of the microbiome.

103 **Methods**

104

105 *Soil type.* Bulk soil samples were collected from a permanently maintained Bare-Fallow (Stackyard,
106 Woburn Experimental Farm, Bedfordshire, UK.) in June 2020. GPS and soil classification data can be
107 found in Table 1.

108

109 *Sample preparation and cryopreservation.* Soil was mixed aseptically with a flame and ethanol
110 sterilised spatula to ensure even distribution of microbial communities, before taking 250 mg
111 aliquots for each sample. Soil samples were cryopreserved with two methods: controlled rate

112 cooling (CRC, 'rate'), and plunge cooling (PC, 'plunge'). CRC samples were cooled with a Stirling cycle
113 cooler, Via Freeze Duo (VFD 30006, Cytiva, Amersham, UK) under a standard IMI / CABI culture
114 collection cooling profile of 1°C min⁻¹ (between 5°C to -30°C) outside this range the cooling rate was
115 2°C min⁻¹ and were held at -80°C. Plunge cooled samples were fully submerged in dewars filled with
116 liquid nitrogen (LN), samples were taken out once bubbling of LN subsided to pre-sample intensities.
117 When each treatment was completed it was immediately stored in LN vapour phase ultra-cold
118 storage (<175°C) for 72 hours. When required, both soil sample treatments were thawed by rapid
119 warming in a water bath set to 37°C for 5 minutes. Samples were annotated based on their
120 cryopreservation method. Control: no cryopreservation; Plunge: rapidly cooled via PC in LN; Rate:
121 CRC with Stirling cycle cooler (VFD30006).

122 *Enrichment method.* Methodology for enrichment was adapted from Yang 2015 (11), 250 mg of soil
123 was added to a falcon tube containing 2.5 ml of 1:10 trypticase soy broth (TSB; Oxoid; 17.0 g
124 pancreatic digest of casein, 3.0 soya bean, 5.0 g NaCl, 2.5 g dipotassium hydrogen phosphate, 2.5 g
125 glucose). This solution was vortexed for 30 seconds, sealed with tape and incubated at 30°C at 150
126 rpm. for 70 hours. Enriched solutions were vacuum filtered using Nalgene™ Sterile Analytical Filter
127 Units (Thermo Fisher Scientific). Filter paper was used for downstream DNA extractions.

128 *DNA extraction, PCR, and NGS library preparation.* DNA was isolated from aliquots of 250 mg of the
129 soil sample with DNeasy PowerSoilKit (Qiagen, Manchester, UK) according to the manufacturers'
130 instruction. For the enrichment, a preliminary filtration of the liquid medium was undertaken with
131 nylon Whatman membrane filters with pore size of 0.2 µm (Merck Life Science UK Ltd, Gillingham,
132 UK). DNA was quantified using Qubit 3.0 Fluorometer (Thermo Fisher Scientific) and normalised to 5
133 ng/µl before PCR reactions. Genomic DNA was amplified with two primer pairs for 16S and ITS rRNA
134 genes, for bacterial and fungal amplicons respectively, with primers modified with the Illumina
135 adapter overhang sequences to the 5'-end. Metabarcoding was performed on the V3–V4 16S rRNA
136 region for bacteria with V3F (5'-CCTACGGGNGGCWGCAG-3') and V3R (5'-
137 GACTACHVGGGTATCTAATCC-3') (12). For the ITS rRNA, the primer were: ITS1 F12 (5'-
138 GAACCGCGGARGGATCA-3') (13) and ITS2 (5'-GCTGCGTTCTTCATCGATGC-3') (14) (15). PCR was
139 undertaken in a Bioer TC1300 LifeECO Thermal cycler (Alpha Laboratories, Eastleigh, UK) with a
140 reaction mix containing 5 µl of each primer at the concentration of 1 µM, 2.5 µl of template DNA at
141 the concentration of 5 ng/µl, and 12.5 µl of KAPA HiFi HotStart ReadyMix (Roche Life Sciences,
142 Welwyn, UK) to a final volume of 25 µl with PCR grade water. PCR reactions were preincubated for 3
143 min at 95°C followed by 25 cycles of 30 sec at 95°C, 30 sec at 55°C, and 30 sec at 72°C. Samples were
144 finally incubated for 5 min at 72°C, followed by chilling to 10°C. Aliquots of 1 µl of index PCR
145 products were assessed for quality with an Agilent TapeStation 4200 (Agilent Technologies), and
146 quantified with Qubit™ (Thermo Fisher Scientific). Libraries were prepared with a Nextera Flex DNA
147 Library prep kit (Illumina, Cambridge, UK), according to the manufacturers' instruction. Steps
148 included the introduction of indices with amplification of DNA and clean up, normalisation and pool.
149 All libraries were validated with an Agilent TapeStation 4200 (Agilent Technologies Ltd, Stockport,
150 UK) and final Concentration in nM was calculated based on the size of the library as determined by
151 Agilent TapeStation 4200. Concentrated libraries were diluted to 4 nM with 10 mM Tris pH 8.5.
152 Pooled DNA libraries were combined with 5 µl of 0.2 N NaOH, and incubated for 5 minutes at room
153 temperature to denature the DNA into single strands. Denatured DNA was then diluted to a final

154 concentration of 15 pM with Illumina Hybridization buffer (Illumina) and sequenced with the
155 Illumina MiSeq at CABI (Egham, UK) on an Illumina MiSeq V3 Cartridge (600 cycles) (Illumina).

156 *Metabarcoding data analysis.* Using DADA2 v1.16.0 (16,17), sequences were truncated and denoised
157 based on quality score using default values; maximum expected errors for forward and reverse reads
158 were 4 and 7 respectively and no truncation length was set for both ITS and 16S sequences. Chimera
159 removal and merge of reads were done with default parameters. 16S taxonomy was assigned on
160 representative ASVs against SILVA ref NR dataset v.138 (18). ITS taxonomy was assigned on
161 representative ASVs against UNITE fungal taxonomic reference v8.2 (19). Normalisation of reads,
162 richness and abundance analyses were undertaken with Phyloseq package v1.34.0 (20) and
163 visualised with ggplot2 (v3.3.3). Various helper functions were used for layout of ggplot2 objects
164 with cowplot, manipulating Phyloseq objects to work with other pipelines and calculating error bars.

165 *Statistical analyses.* All statistical analyses were performed within the computing environment R (v
166 3.5.0; R Core Development Team, 2005) and visualized using ggplot2 (v3.3.3). To visualize the overall
167 bacterial communities, normalised sequence counts from phyloseq were used to plot non-metric
168 multidimensional scaling (NMDS) on Bray-Curtis dissimilarity matrices to ordinate in two dimensions
169 the variance of beta diversity using the Vegan package (v2.5-7). NMDS analysis was performed
170 between samples grouped by cryopreservation treatments on ITS metabarcodes, and between
171 enrichment and cryopreservation treatments on 16S metabarcodes. These dissimilarity matrices
172 were also used to analyse the group effects between treatments and enrichment with pairwise
173 permutational analysis of variance (PERMANOVA) using the adonis() function in Vegan and
174 convenience wrapper functions in the pairwise.adonis package (v0.01). Pairwise PERMANOVA
175 analysis was performed between samples grouped by cryopreservation treatments on ITS
176 metabarcodes, and between enrichment and cryopreservation treatments on 16S metabarcodes
177 with 999 permutations. P values were false discovery rate corrected by the Benjamini-Hockberg
178 method (21). Heat tree plots were made using phyloseq objects converted to work with metacoder
179 (v0.3.4) (22).

180

181 **Results**

182 The impact of cryopreservation on the microbial communities was assessed by comparing the
183 microbiomes using culture-free approaches. Metabarcodes were generated for bacterial (16S rRNA)
184 and fungal (ITS rRNA) amplicons following different treatments of controlled rate cooling (CRC,
185 'rate'), and plunge cooling (PC, 'plunge'), compared to a non-treatment ambient control (control).

186

187 *Impact of cooling treatments on microbiome sequence content.* A total of 9,846,613 read counts
188 were obtained for 16S rRNA, and 8,493,298 for ITS rRNA (Table 2). In total, 4,915,778 sequences
189 were obtained for the 'control', 7,191,917 sequences for the 'plunge' and 6,232,216 sequences for
190 the 'rate' treatments across 44 replicates: five replicates for each treatment and enrichment pair.
191 One replicate of the 'control' was excluded from the data analysis due to sequence data corruption.

192 Enrichment produced PCR amplicon for the 16S rRNA PCR only with ITS rRNA PCR that did not
193 produce amplicons.

194 *Richness of the cryopreserved soil microbiome.* The overall richness of bacterial ASVs was statistically
195 comparable across the samples ($F=0.982$, $p=0.565$). Similarly, the richness of eukaryotic ASVs did not
196 produce significant differences ($F=1.392$, $p=0.163$). After enrichment, significant differences were
197 observed ($F=1.851$, $p=0.014$). Further analyses were undertaken to compare the treatments with
198 pairwise comparison of the relative abundance of ASVs, which was used to estimate the similarity of
199 the soil microbiome before and after cryopreservation (Table 3). The three treatments 'control',
200 'plunge' and 'rate' were statistically comparable. After enrichment, there were significant
201 differences in the composition of ASVs. When comparing the datasets after the enrichment step,
202 only 'rate' was statistically comparable to the control ($p=0.219$). Additional multivariate statistics
203 with non-metric multidimensional scaling supported the similarities of 'plunge' and 'rate' with the
204 'control' (Figure 1). With the ITS metabarcoding, 'plunge' and 'rate' had similar ASVs composition
205 compared to the 'control' ($p=0.358$) (Table 4).

206 *Taxonomic affiliation of the metabarcodes before and after cryopreservation.* When individual ASVs
207 were grouped by taxonomic affiliation, the similarity across the three treatments was supported
208 further (Figure 2). The ten most abundant taxa recovered from the three treatments were, on
209 average, 41.5% of the total number of DNA reads generated by 16S rRNA metabarcoding, and 58.3%
210 of ITS rRNA. In the 16S rRNA metabarcoding, several had little variability across the three
211 treatments: for example, Acidobacteriales were $5.79\% \pm 0.87$ in the 'control', $5.61\% \pm 0.43$ in
212 'plunge' and $5.56\% \pm 0.32$ in 'rate'. Tepidisphaerales had more variation and were more abundant in
213 'rate' ($9.88\% \pm 0.26$ vs $7.96\% \pm 1.04$ in 'control' and $7.69\% \pm 0.90$ in 'plunge') (Figure 2). In the ITS
214 rRNA metabarcoding, the ten most abundant genera were consistent in the three treatments (Figure
215 2). Relative abundances of the ten most abundant genera had minor variations and were consistent
216 overall. For example, the two most abundant genera in the three treatments were *Metarhizium* sp.
217 and *Mortierella* sp. Other less abundant genera were consistent in the three treatments.

218 *Microbial growth before and after cryopreservation.* With the enrichment method, some soil
219 bacterial taxa led to different trends when comparing 'control' vs 'plunge' and 'control' vs 'rate'.
220 After incubation of soil aliquots in enrichment broth, the ten most abundant taxa recovered from
221 the three treatments, which covered, on average, 93.1% of the total number of DNA reads, were
222 analysed further. When comparing 'control' vs 'plunge', the biggest variability was seen for
223 *Chryseobacterium* sp. (-13.9% after 'plunge') and Enterobacteriales ($+10.2\%$ after 'plunge'). When
224 comparing 'control' vs 'rate', the variability was smaller than 'plunge' with *Chryseobacterium* sp.
225 being the most affected genus (-8.4% after 'rate') and Bacillales ($+7.6\%$ after 'rate') (Figure 2).
226 Enterobacteriales after 'rate' were similar to those recovered from the 'control' (-0.3%). Overall,
227 'rate' was more similar to the 'control' than 'plunge'.

228

229 Discussion

230

231 We applied an optimised method for cryopreservation for which the genomic integrity of the
232 microbiome was conserved when compared with a 'control' treatment. The optimised
233 cryopreservation of soil samples is essential for the development of microbiome research in order to
234 retain stable, functionally intact microbiomes (7). Our results showed that metabarcoding of 16S and
235 ITS rRNA were useful methods to estimate successful cryopreservation. In addition, the microbiome
236 after enrichment delivered a similar response of cryopreserved soil and this was estimated with the
237 comparison of ten most abundant bacterial taxa accounting for, on average, 93.1% of DNA reads.
238 These findings support a successful process of cryopreservation and are promising for future use of
239 this technology. Previous success on cryopreservation and resuscitation of natural aquatic
240 prokaryotic communities has shown promising results also (2). To the best of our knowledge, this
241 study is the first report of cryopreservation of soil using a Stirling cycle cooling approach. Further, we
242 were able to cryopreserve and resuscitate a soil sample which maintained a representative
243 component of its genomic diversity.

244 Eukaryotes organisms had a similar response to that of the bacteria. 'Plunge' and 'rate' treatments
245 produced a consistent profile compared to the 'control', showing that these methods maintain the
246 genomic integrity of the eukaryotic portion of the microbiome similarly to the bacteria. This was
247 particularly evident with the comparison of the richness (Figure 1) and confirms previous findings for
248 which 'rate' is the optimal preservation approach for fungi (10). However, the method utilised has
249 yet to be optimised and more tests are required to improve the process. For example, the main aim
250 is to reduce ice formation which can be damaging for the cells in the samples. But mitigating the
251 damaging ice-effect will cause less damage to the organisms within the sample. Recent research by
252 McClure et al. (23), suggests that, under stress, microbial communities within soil samples may
253 produce trehalose, which has been well-characterised as a synthesised osmoprotectant in bacteria
254 (24). This is of significance as trehalose is a natural cryoprotectant and the presence of this would
255 reduce ice damage. Drying of samples to reduce residual moisture content in a controlled way that
256 promotes trehalose formation would therefore be worthy of further investigation.

257 Intact genomic content was recovered after cryopreservation. 16S rRNA metabarcodes were similar
258 in 'control', 'plunge' and 'rate' with 'plunge' being more similar to the 'control' than 'rate' without
259 enrichment. This trend could be explained by the 'plunge' method which rapidly stops metabolism
260 while 'rate' reaches similar output but gradually. Similarly, ITS rRNA metabarcodes were consistent
261 in 'control', 'plunge' and 'rate'. The overwhelming majority of microbes are essential for ecosystem
262 functioning and known for their interactions with other microorganisms as well as macroorganisms
263 (9), which supports the importance of maintaining the characteristics of a system so that species
264 interactions and communication remain untouched to provide sustainable population dynamics and
265 functional activities (25).

266 For the second hypothesis we compared bacterial growth after enrichment and forced the
267 microbiome in favour of bacterial development. The enrichment method was used as a measure of
268 the success of preservation to estimate bacterial growth before and after cryopreservation, and
269 measure live, metabolically active bacteria from the total gDNA component. This enrichment
270 method is non-selective; it is a rich medium that favours growth of fast-growing bacteria such as the
271 Enterobacterales. Under these conditions, 'plunge' and 'rate' had a different response. Overall,
272 similar microbial growth was obtained before and after cryopreservation, but 'rate' was similar to
273 the 'control' ($p=0.219$) as compared to 'plunge' ($p=0.044$). Our data indicated that cryopreservation

274 did not significantly impact community composition after the resuscitation process, and this was
275 particularly true for 'rate'. Specifically, *Chryseobacterium* sp. and Bacillales determined a better
276 recovery for 'rate' compared to 'plunge' (Figure 2). Aquatic microbiomes have shown similar trends
277 with no significant changes found after cryopreservation (2). However, these data reported on
278 genomic DNA isolated after microfiltration and do not provide indication of the preservation of
279 environmental factors and physical microstructures within the microbiome. More tests should be
280 undertaken to investigate the response of the microbiome after cryopreservation and explore the
281 resuscitation of organisms. The condition for the development of more complex eukaryotic
282 organisms remain more challenging.

283 With this experiment we did not prove that cryopreservation has kept physical space untouched. We
284 proved that the microbiome, under the same pressure generated by the enrichment with a liquid
285 medium, had an identical response and that 'rate' generated a bacterial profile statically comparable
286 to the 'control'. This proved that the microbiome of the selected soil sample reacted similarly to the
287 control, and that the bacterial fraction responded to the stress was statistically the same. The
288 improved integrity for bacteria suggests that the controlled rate cooling approach may greatly
289 improve the stringency of the cryopreservation preservation for prokaryotes. Although with only a
290 limited set of samples, the results provided a benchmark for further optimisation of process, but
291 importantly shows that cryopreservation of soil samples at ultra-low temperature is preferential for
292 conservation of key microbiome resources. This research provides a baseline for the further
293 optimisation of protocols, the development of standards and a long-term strategy for the
294 conservation of agricultural biodiversity.

295 In conclusion, this study provides evidence of a feasible method for the cryopreservation
296 encompassing controlled rate cooling techniques and resuscitation of a selected exemplar soil
297 microbiome. Similarly to other findings (2), the cryopreservation method developed and applied in
298 this study can be applied to other microbiomes. In particular, the enrichment method could be
299 particularly useful to investigate resuscitation of microbes, with further development that could
300 include different types of pressure beyond selective growth conditions as shown in this manuscript
301 with the enrichment method.

302 **Author statements**

303 **Authors and contributors**

304 GC designed and performed the experiment, analysed the data, and wrote first draft; JMB
305 performed the experiment, analysed the data; MR designed the experiment; IC provided soil type
306 information; all authors contributed to the final version of the manuscript.

307

308 **Conflicts of interest**

309 The authors declare that there are no conflicts of interest

310

311 **Funding information**

312 This work was undertaken through the Crop Microbiome Cryobank Project (Reference:
313 BB/T019700/1) supported by the BBSRC under the Bioinformatics and Biological Resources Fund.
314 CABI is an international intergovernmental organisation, and we gratefully acknowledge the core
315 financial support from our member countries (and lead agencies) including the United Kingdom
316 (Foreign, Commonwealth and Development Office), China (Chinese Ministry of Agriculture), Australia
317 (Australian Centre for International Agricultural Research), Canada (Agriculture and Agri-Food
318 Canada), Netherlands (Directorate-General for International Cooperation) and Switzerland (Swiss
319 Agency for Development and Cooperation). See [https://www.cabi.org/about-cabi/who-we-work-](https://www.cabi.org/about-cabi/who-we-work-with/key-donors/)
320 [with/key-donors/](https://www.cabi.org/about-cabi/who-we-work-with/key-donors/) for full details.

321

322 **Acknowledgements**

323 We thank Helen Stewart for technical support.

324

325 **References**

326

- 327 1. Faust K, Lahti L, Gonze D, de Vos WM, Raes J. Metagenomics meets time series analysis:
328 Unraveling microbial community dynamics. *Curr Opin Microbiol* [Internet]. 2015;25(May):56–
329 66. Available from: <http://dx.doi.org/10.1016/j.mib.2015.04.004>
- 330 2. Rain-Franco A, de Moraes GP, Beier S. Cryopreservation and Resuscitation of Natural Aquatic
331 Prokaryotic Communities. *Front Microbiol*. 2021;11(January):1–19.
- 332 3. Busby PE, Soman C, Wagner MR, Friesen ML, Kremer J, Bennett A, et al. Research priorities
333 for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol*. 2017;15(3):1–14.
- 334 4. Berg G, Rybakova D, Grube M, Köberl M, Price A. Europe PMC Funders Group The plant
335 microbiome explored : implications for experimental botany. 2017;67(4):995–1002.
- 336 5. Cavicchioli R, Ripple WJ, Timmis KN, Azam F, Bakken LR, Baylis M, et al. Scientists' warning to
337 humanity: microorganisms and climate change. *Nat Rev Microbiol* [Internet]. 2019;17(9):569–
338 86. Available from: <http://dx.doi.org/10.1038/s41579-019-0222-5>
- 339 6. Prosser JI, Bohannon BJM, Curtis TP, Ellis RJ, Firestone MK, Freckleton RP, et al. Prosser et al
340 2007_The role of ecological theory in microbial ecology. *Nat Rev Microbiol*.
341 2007;5(May):384–92.
- 342 7. Ryan MJ, Schloter M, Berg G, Kostic T, Kinkel LL, Eversole K, et al. Development of
343 Microbiome Biobanks – Challenges and Opportunities. *Trends Microbiol* [Internet].
344 2021;29(2):89–92. Available from: <https://doi.org/10.1016/j.tim.2020.06.009>

- 345 8. Gich F, Janys MA, König M, Overmann J. Enrichment of previously uncultured bacteria from
346 natural complex communities by adhesion to solid surfaces. *Environ Microbiol* [Internet].
347 2012 Nov 1;14(11):2984–97. Available from: [https://doi.org/10.1111/j.1462-](https://doi.org/10.1111/j.1462-2920.2012.02868.x)
348 2920.2012.02868.x
- 349 9. Berg G, Rybakova D, Fischer D, Cernava T, Vergès MCC, Charles T, et al. Microbiome definition
350 re-visited: old concepts and new challenges. *Microbiome*. 2020;8(1):1–22.
- 351 10. Ryan MJ, Kasulyte-Creasey D, Kermode A, San SP, Buddie AG. Controlled rate cooling of fungi
352 using a stirling cycle freezer. *Cryo-Letters*. 2014;35(1):63–9.
- 353 11. Yang F, Liu HM, Zhang R, Chen D Bin, Wang X, Li SP, et al. *Chryseobacterium shandongense*
354 sp. nov., isolated from soil. *Int J Syst Evol Microbiol*. 2015;65(6):1860–5.
- 355 12. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of general
356 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based
357 diversity studies. *Nucleic Acids Res*. 2013;41(1):1–11.
- 358 13. Schmidt P-A, Bálint M, Greshake B, Bandow C, Römbke J, Schmitt I. Illumina metabarcoding of
359 a soil fungal community. *Soil Biol Biochem* [Internet]. 2013;65:128–32. Available from:
360 <https://www.sciencedirect.com/science/article/pii/S0038071713001867>
- 361 14. White T, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA
362 genes for phylogenetics. New York, USA: Academic Press; 1990. 315–322 p.
- 363 15. Cafà G, Caggiano B, Reeve MA, Bhatti H, Honey SF, Bajwa B, et al. A polyphasic approach aids
364 early detection of potentially toxigenic aspergilli in soil. *Microorganisms*. 2019;7(9).
- 365 16. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
366 resolution sample inference from Illumina amplicon data. *Nat Methods* [Internet].
367 2016;13(7):581–3. Available from: <https://doi.org/10.1038/nmeth.3869>
- 368 17. Prodan A, Tremaroli V, Brolin H, Zwinderman AH, Nieuwdorp M, Levin E. Comparing
369 bioinformatic pipelines for microbial 16S rRNA amplicon sequencing. *PLoS One*. 2020;15(1):1–
370 19.
- 371 18. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA
372 gene database project: Improved data processing and web-based tools. *Nucleic Acids Res*.
373 2013;41(D1):590–6.
- 374 19. Abarenkov K, Henrik Nilsson R, Larsson K-H, Alexander IJ, Eberhardt U, Erland S, et al. The
375 UNITE database for molecular identification of fungi – recent updates and future
376 perspectives. *New Phytol* [Internet]. 2010 Apr 1;186(2):281–5. Available from:
377 <https://doi.org/10.1111/j.1469-8137.2009.03160.x>
- 378 20. McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and
379 Graphics of Microbiome Census Data. *PLoS One*. 2013;8(4).
- 380 21. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
381 Approach to Multiple Testing. *J R Stat Soc Ser B* [Internet]. 1995 May 4;57(1):289–300.

- 382 Available from: <http://www.jstor.org/stable/2346101>
- 383 22. Foster ZSL, Sharpton TJ, Grünwald NJ. Metacoder: An R package for visualization and
384 manipulation of community taxonomic diversity data. *PLoS Comput Biol*. 2017;13(2):1–15.
- 385 23. McClure RS, Lee JY, Chowdhury TR, Bottos EM, White RA, Kim YM, et al. Integrated network
386 modeling approach defines key metabolic responses of soil microbiomes to perturbations. *Sci*
387 *Rep* [Internet]. 2020;10(1):1–9. Available from: <https://doi.org/10.1038/s41598-020-67878-7>
- 388 24. Giaever HM, Styrvold OB, Kaasen I, Strøm AR. Biochemical and genetic characterization of
389 osmoregulatory trehalose synthesis in *Escherichia coli*. *J Bacteriol*. 1988;170(6):2841–9.
- 390 25. Bassler BL. Small talk: Cell-to-cell communication in bacteria. *Cell*. 2002;109(4):421–4.
- 391
- 392

393

394 **Figures and tables**

395

396 **Table 1.** Soil sample used in this study: site Stackyard, at the time of sampling land use was
397 “permanent Bare-Fallow”, Woburn Experimental Farm, Stackyard field, Bedfordshire, UK.

398

Collection Date	Soil_taxonomic classification	Geographic location				Texture		
		Depth	Elevation	Latitude	Longitude	Sand	Silt	Clay
16/06/2020	Sandy loam (Cottenham series)	0-20cm	100m	52.0004	0.614265	50%	27%	18%

399

400

401

402

403

404 **Table 2.** Raw statistics of metabarcodes obtained in this study

Treatment	Replicate	16S	16S_enrich	ITS
Control	1	246,069	267,068	127,049
	2	160,161	416,846	263,067
	3	435,272	576,261	77,441
	4	-	520,926	719,407
	5	194,165	301,919	610,127
Plunge	1	183,771	350,703	1,755,679
	2	446,374	163,302	447,575
	3	396,369	360,018	270,863
	4	200,425	386,624	604,740
	5	625,850	566,661	432,963
Rate	1	195,671	144,839	1,186,559
	2	260,284	333,558	389,127
	3	179,039	322,090	788,068
	4	578,534	398,167	453,732
	5	264,684	370,963	366,901

405

406

407 **Table 3.** Pairwise comparisons of corrected P values by PERMANOVA analysis based on Bray-Curtis
 408 dissimilarity matrix of normalised 16S sequence counts. P Value correction for false discovery rate
 409 preformed with Benjamini-Hochberg correction. C: No cryopreservation ('control'); P: Plunge cooled
 410 in LN ('plunge'); R: Controlled rate cooled ('rate'); CE: No cryopreservation ('control') after
 411 enrichment; PE: Plunged cooled in LN ('plunge') after enrichment; RE: Controlled rate cooled ('rate')
 412 after enrichment;

	C(16S)	P(16S)	R(16S)	CE(16S)	PE(16S)	RE(16S)
C(16S)	-	-	-	-	-	-
P(16S)	0.632	-	-	-	-	-
R(16S)	0.371	0.701	-	-	-	-
CE(16S)	0.017	0.018	0.017	-	-	-
PE(16S)	0.017	0.017	0.017	0.044	-	-
RE(16S)	0.017	0.017	0.017	0.219	0.043	-

413

414

415

416 **Table 4.** Pairwise comparisons of corrected P values by PERMANOVA analysis based on Bray-Curtis
417 dissimilarity matrix of normalised ITS sequence counts. P Value correction for false discovery rate
418 performed with Benjamini-Hochberg correction. C: No cryopreservation ('control'); P: Plunge cooled
419 in LN ('plunge'); and R: Controlled rate cooled ('rate').

420


	C(ITS)	P(ITS)	R(ITS)
C(ITS)	-	-	-
P(ITS)	0.358	-	-
R(ITS)	0.358	0.358	-

421

422

423


424



425

426 **Figure 1.** Richness of bacteria (A) and eukaryotes (B) of soil across different cryopreservation
427 treatments. Control: No cryopreservation; Plunge: Plunged cooled in LN; Rate: Controlled rate
428 cooled. Richness of non-enriched (dark bars) and enriched soil (light bars). Error bars are
429 bootstrapped 95% confidence intervals implemented in ggplot2; C-E) Non-metric multidimensional
430 scaling (NMDS) of 16S (C), Enriched 16S (D) and ITS (E) rRNA metabarcodes of soil across different
431 cryopreservation treatments. NMDS was derived from Bray-Curtis dissimilarity matrices calculated
432 from ASV sequence proportions among samples. Points are sample replicates and coloured ellipses
433 are calculated with standard deviation of sample centroids for each cryopreservation treatment
434 using 'ordiellipses' in the vegan package.

435



436

437 **Figure 2.** Relative abundance of dominant bacterial ASVs in control soil (left panels), enriched soil
 438 (central panels) and eukaryotic (right panels) across different cryopreservation treatments. Error
 439 bars are bootstrapped 95% confidence intervals implemented in ggplot2.

440