

## Iowa monarch conservation, pest management and crop production

Steven P. Bradbury, professor, Natural Resources Ecology and Management, Iowa State University; Tyler Grant, postdoctoral research associate, Natural Resources Ecology and Management, Iowa State University; Niranjana Krishnan, graduate student, Entomology, Iowa State University

### Introduction

The monarch butterfly population has experienced an 80% decline in North America over the past two decades (Brower et al., 2012; Pleasants and Oberhauser, 2013; Jepsen et al., 2015). The three to four hectares of occupied overwintering forest in 2016 and 2015 was well below a target of six hectares needed to support a resilient population and reduce the risk of quasi-extinction (loss of the North American migration) in the next 10 to 20 years (Semmens et al., 2016). In response to a petition to evaluate the status of the species, the U.S. Fish and Wildlife Service (USFWS) is evaluating listing the monarch as a threatened species under the Endangered Species Act (USFWS, 2014 a, b). Under a court-supervised schedule, the USFWS must propose a listing decision in June 2019, which underscores the urgency of establishing viable, voluntary, state-based monarch conservation programs to provide USFWS a credible rationale to not list the species. If the monarch butterfly is listed, it could lead to significant regulatory and management burdens for farmers and livestock producers.

Increasing summer reproductive success in the North Central United States has been identified as a high priority for monarch conservation (Oberhauser et al. 2017; Flockhart et al. 2015). Up to 58% of the monarchs overwintering in Mexico originate in Iowa and neighboring states in the North Central U.S. (Flockhart et al. 2017). Monarch butterflies oviposit only on milkweed species (mainly *Asclepias* spp.) and primarily on common milkweed (*Asclepias syriaca*) in the North Central states (Malcolm et al. 1993). To increase monarch populations to levels that would reduce the probability of quasi-extinction by 50% over 20 years, Thogmartin et al. (2017) estimated that 1.3 to 1.8 billion additional milkweed stems need to be added to the North Central U.S. landscape.

The current amount of common milkweed stems in the North Central U.S. is estimated to be approximately 1.3 billion, the majority of which is in publically owned grasslands, land enrolled in conservation programs, such as the United States Department of Agriculture's Conservation Reserve Program (CRP), crop field borders, road rights-of-way, urban and suburban land cover, and miscellaneous non-agricultural habitat (Pleasants and Oberhauser 2013; Thogmartin et al. 2017). Analyses by Thogmartin et al. (2017) indicate that adding 1.3 to 1.8 billion new milkweed stems in the North Central U.S. will require an 'all hands on deck' approach across all land cover/land use categories. Habitat conservation goals can only be achieved through significant adoption of habitat restoration in privately owned land in agricultural landscapes. Establishment of milkweed and forage plants in rural roadsides; marginal crop land; portions of existing CRP land, pastures and grassland; and grassy areas bordering crop fields, will create a patchwork of habitat within agricultural landscapes dominated by corn and soybean production.

### Overview of monarch conservation efforts in Iowa

In 2015, in response to the decline of the monarch butterfly, Iowa commodity and livestock organizations, pesticide technology providers, utility companies, city and county conservation organizations, Iowa colleges and universities, the Iowa Department of Agriculture and Land Stewardship, Iowa Department of Natural Resources and the College of Agriculture and Life Sciences at Iowa State University formed the Iowa Monarch Conservation Consortium (<https://monarch.ent.iastate.edu>). The consortium currently

includes 42 participating organizations, with USFWS and USDA serving as *ex officio* members. The mission of the Iowa Monarch Conservation Consortium is to enhance monarch butterfly reproduction and survival in Iowa through collaborative and coordinated efforts of farmers, private citizens and their organizations through research, education and direct action.

The consortium's outreach and extension efforts draw upon all the member organizations to ensure the broad delivery of practical, science-based information on monarch butterfly conservation practices for Iowa's landscapes. Habitat improvements in rural landscapes are being designed to target underutilized areas that do not conflict with agricultural production, are sufficient in scale to support improved monarch breeding success, and strive to complement other conservation programs. In this regard, monarch habitat will be beneficial for pollinators and other wildlife species and can be 'stacked' with other conservation practices designed to reduce soil erosion and minimize nutrient runoff. The consortium published Version 1.0 of Iowa Monarch Conservation Strategy in February 2017 (<https://monarch.ent.iastate.edu/files/file/iowa-monarch-conservation-strategy.pdf>). Version 2.0 of the strategy is under development and will include Iowa's habitat goals, which will be integrated with neighboring North Central States' habitat targets (see: [http://www.mafwa.org/?page\\_id=2347](http://www.mafwa.org/?page_id=2347)).

The research arm of the Iowa Monarch Conservation Consortium is based on the campus of Iowa State University and includes University and USDA scientists, technicians, graduate and undergraduate students. Below is a synopsis of ongoing demonstration and research activities.

### **Habitat establishment practices**

A seed mix has been developed that includes three milkweed species and several nectar-producing forbs that benefit butterflies and bees (Appelgate et al. 2017). Grass species included in the mix are mostly short to medium height in order to reduce competition with forbs. All species are native to Iowa and should thrive in well-drained and moderately-drained soils. This seed mix has been used to establish monarch habitat at 46 demonstration sites. Thirteen demonstration sites are associated with establishment of approved edge-of-field practices in the Iowa Nutrient Reduction Strategy, including six saturated buffers sites and seven bioreactors sites. The remaining sites are in underutilized grass areas (21 sites) and land near hog confinements (12 sites). Supporting experiments are ongoing to develop methods to establish milkweed into land dominated by smooth brome grass, *Bromus inermis*.

Research completed at Iowa State University (Pocius et al. 2017 a, b) indicate a diversity of milkweed species including: *Asclepias exaltata* (Poke Milkweed), *A. hirtella* (Tall Green Milkweed), *A. incarnata* (Swamp Milkweed), *A. speciosa* (Showy Milkweed), *A. sullivantii* (Prairie Milkweed), *A. syriaca* (Common Milkweed), *A. tuberosa* (Butterfly Milkweed), *A. verticillata* (Whorled Milkweed), and *Cynanchum laeve* (Honeyvine Milkweed) could benefit monarchs in their summer range. Ultimately, monarch seed mixes and planting guidelines will provide farmers and landowners practical and reliable methods to expand the benefits of existing groundcover to promote monarch caterpillar development and provide nectar sources for monarch butterfly adults. Pollinators and other wildlife species also will benefit from the increased diversity of plant species.

### **Monarch population modeling**

Research is needed to better understand how habitat patch size, composition, and spatial arrangements (Oberhauser et al. 2017; Pleasants and Oberhauser, 2013; Zalucki et al. 2001, 2002; Zalucki and Lammers 2010; Zalucki et al. 2016) influence monarch survival and breeding success. This understanding provides a foundation for evaluating how different habitat patch distributions influence monarch population growth rates, which in turn supports conservation planning. Modeling is a useful tool for investigating how monarch butterfly reproductive performance, larval development, and population growth respond to

habitat plant composition and quality, including milkweed stem density and spatial arrangement; habitat patch size; and habitat fragmentation patterns within a spatially explicit landscape. Previous modeling efforts in artificial landscapes have identified monarch perceptual range and habitat attractiveness, which influences the probability a butterfly will choose a given habitat patch, as important data inputs (Zalucki et al. 2016). Modeling efforts to date (Zalucki et al. 2016, Zalucki and Lammers, 2010) have also identified the important role individual, lone milkweed plants and small low-density patches of milkweed play in increasing the number of monarch eggs laid on a landscape. To further evaluate and advance understanding of how female monarchs utilize breeding habitat in Iowa, we have developed a model that uses spatially explicit landscapes that reflect current habitat conditions and spatial patterns (Grant et al., 2017a). The model can incorporate results from on-going studies that are refining estimates of monarch perceptual range and flight patterns (Fisher et al., 2017a, b).

## **Estimating risk of herbicides and insecticides**

Monarch conservation benefits and risks associated with restoring habitat near conventional crop fields, due to potential pesticide exposure, are uncertain. The National Resources Conservation Service (NRCS) *Monarch Butterfly Wildlife Habitat Evaluation Guide* (2016) discourages placement of monarch breeding habitat within 38m (125ft) of crop fields that are treated with herbicides or insecticides; a risk-based rationale for the buffer size is not available. Employing a buffer of this size would result in a significant area of land unavailable for the establishment of breeding habitat. In Story County, Iowa, a 38m buffer around corn and soybean fields would eliminate approximately 84% of rural roadside habitat and 38% of grassland, Conservation Reserve Program (CRP) land, pastures, railroad rights-of-way, riparian corridors and wetlands. Based on possible pesticide use patterns, do the potential risks from pesticide exposure reduce the benefits of establishing habitat in close proximity to crop fields?

### ***Herbicides***

The effect of nonlethal herbicide injury to milkweed on monarch oviposition and larval development has not been investigated. Utilization of forbs by pollinators was reduced by exposure to low concentrations of herbicides (Bohnenblust et al. 2016), largely due to fewer floral resources. The mechanism of toxicity for growth-regulator herbicides may be due to the stimulation of abscisic acid and ethylene production in affected plants (Grossman 2009). At sublethal doses, the growth-regulator herbicides cause responses such as epinasty and malformed leaves. Since a high percentage of milkweed occurs adjacent to crop fields, there is a need for an off-target risk assessment of herbicides on the value of milkweed to monarchs. Such an assessment is particularly necessary for evaluating the value of new habitat patches.

Common milkweed persists in crop fields due to perennial rootstocks that are resilient to most herbicides; post-emergence herbicides can cause significant damage to shoots present at the time of application. Hartzler and Lizotte-Hall (Department of Agronomy, ISU) are undertaking dose response studies to determine the effect of fomesafen, a herbicide commonly used on soybeans, on biomass production of common milkweed. Field experiments are determining how herbicide injury influences utilization of common milkweed by adult monarchs. Fomesafen is being used to evaluate effects of in-field herbicide use, whereas dicamba is being used to study effects of off-field exposure.

### ***Insecticides***

Insecticides for managing insect corn and soybean pests include organophosphate, pyrethroid, neonicotinoid, and anthranilic diamide (Hodgson and VanNostrand G 2016; University of Tennessee 2016; DuPont 2010). Monarch larvae in existing and newly established milkweed patches near crop fields could be exposed to insecticide spray drift following applications to manage soybean aphid (*Aphis glycines* Matsumura) between mid-July and mid-September (Iowa State University 2016), which coincides with estimated peak larval abundance of the 2nd and 3rd monarch generations in Iowa (Pleasants 2015; Prysby

and Oberhauser 2014; Nail et al. 2015). With expanding use of cover crops in Iowa, potential economic injury to corn from true armyworm (*Mythimna unipuncta* Haworth) has been observed, with foliar insecticide applications documented between mid-May and late June (Dunbar et al 2015), which overlaps with the 1st and 2nd generation larvae in Iowa. Monitoring studies designed to systematically document insecticide levels on milkweeds following foliar application are not available; however, models used by the U.S. EPA, such as AgDRIFT (USEPA 2017), indicate that spray drift exposure to milkweed up to 38m or more downwind is likely.

Monarch larvae could also be exposed to insecticides through ingestion of milkweed. Corn and soybean are typically planted with neonicotinoid-treated seed (Douglas and Tooker 2015), including 70% of soybean acres in Iowa (Hodgson et al. 2016). Chlorantraniliprole also is entering the market as a corn seed treatment option (Corn and Soybean Digest 2015). Imidacloprid, clothianidin, and thiamethoxam have moved into Iowa streams (Hladik et al 2014), presumably due to subsurface flow (Hladik et al 2017), which raises concerns that plants downslope of cropped fields could absorb neonicotinoids systemically. Several studies (Botias et al. 2016, 2015; David et al. 2016; Krupke et al. 2012; Long and Krupke 2016; Paola and Kaplan 2015; Pecenka and Lundgren 2015) indicate a variety of non-crop plants, including milkweed, in the margins of fields previously sowed with neonicotinoid-treated seeds can have detectable levels of imidacloprid, clothianidin, and thiamethoxam in leaves, pollen and nectar; although the frequency of detections and concentrations are highly variable.

There is a paucity of monarch toxicity data to interpret the significance of insecticide exposures due to spray drift or systemic uptake by milkweed. Consequently we are examining the toxicity of six representative insecticides: beta-cyfluthrin (pyrethroid, foliar uses); chlorpyrifos (organophosphate, foliar); imidacloprid (neonicotinoid, seed treatment and foliar); clothianidin (neonicotinoid, seed treatment); thiamethoxam (neonicotinoid, seed treatment and foliar); and chlorantraniliprole (anthranilic diamide, seed treatment and foliar) on monarch larvae, eggs, pupae and adults (Krishnan et al., 2017 a, b, c). Larvae, eggs and chrysalides are being treated topically to mimic exposure to a spray drift plume. Larvae and adults are also being exposed orally to mimic insecticide exposure from milkweed leaves (larvae - systemic uptake and spray drift deposition) and nectar of flowering plants (adults - systemic uptake), respectively. Dose response curves for mortality and growth/development are being compared to exposure levels obtained from spray drift models, on-going residue surveys, and existing ecological risk assessments and open literature studies. These analyses provide the means to assess risks at the habitat patch scale.

### ***Predicting landscape-scale risk of insecticide exposure to monarch populations***

Using instar-specific, 96-hour mortality dose-response curves for beta-cyfluthrin, chlorpyrifos, imidacloprid, thiamethoxam and chlorantraniliprole and estimates of exposure using EPA's AgDRIFT model, we have quantified spray drift risks at distances up to 38m downwind from treated fields following simulated aerial and ground applications (Krishnan et al 2017 a, b, c). In general, predicted mortality rates downwind were highest for chlorantraniliprole and beta-cyfluthrin and lowest for thiamethoxam and imidacloprid. To assess monarch population responses at a landscape scale we created a GIS layer that delineates areas within 38m of crop fields in Story County, Iowa. This layer was incorporated into the model described previously. A projection model is used to estimate survival from eggs to adults. Estimates of adult recruitment over a 10-year period are derived under three scenarios: a) no new milkweed planted within 38m of crop fields; b) milkweed patches placed within 38m of crop fields, with no insecticide exposure; and c) milkweed patches placed within 38m of crop fields with spray drift exposure. Different frequencies of economically-significant pest-pressure and associated foliar applications using typical wind direction and speed are used in the simulations (Bradbury et al. 2017; Grant et al 2017b). In future studies, the projection model will incorporate larval mortality rates based on systemic insecticide uptake in milkweed downslope of fields planted with treated corn or soybean seeds. Estimates of landscape-scale adult recruitment under varying spatial-temporal pest management scenarios provide the means to evaluate monarch conservation costs and benefits of establishing habitat in areas potentially exposed to insecticides.

## Summary

The USFWS is committed to make a decision regarding whether to list the monarch butterfly as a threatened species by June of 2019. On-going research is enhancing the scientific basis for supporting monarch conservation practices by elucidating relationships between landscape-level habitat patterns and monarch movement, survival, reproduction, and development in the context of pest management practices. Coordinated research, extension and outreach provides a foundation for advancing and implementing a proactive, science-based conservation program that will enhance North American recovery of the monarch while maintaining efficient agriculture and livestock production.

## Acknowledgments

Members of the Iowa State University monarch research team include: C. Abel, S. Appelgate, K. Bidne, R. Bitzer, T. Blader, S. Blodgett, D. Debinski, K. Fisher, T. Grant, R. Hartzler, R. Hellmich, N. Krishnan, S. Lizotte-Hall, J. Pleasants, V. Pocius, J. Pohl, D. Schweitzer, and N. Shryock. Special thanks to J. Appelhans, H. Arthur, J. Chavez-Martinez, K. Claus, W. Douglas, A. Euken, C. Haggard, , S. Hilby, B. Hoey, E. Kenyon, A. Kerker, N. Oppedal, J. Pfeiffer, R. Nylin and M. Van Loon, for technical help.

The research and demonstration efforts summarized here are supported by two NRCS Conservation Innovation Grants (69-6114-15-006; 69-3A75-16-006), grants from the Iowa Soybean Association, Iowa Pork Producers Association, and the Garden Club of America, as well as the Iowa Monarch Conservation Consortium and the College of Agriculture and Life Sciences, Iowa State University.

## References

- Appelgate, S., Blodgett, S., Bradbury, S., Debinski, D., Hartzler, R., Pleasants, J., Schweitzer, D. & Hellmich, R. (2017) Monarch Seed Mix: High Diversity. Iowa State University Extension and Outreach. ENT 0047.
- Bohnenblust E.W., Vaudo A.D., Egan J.F., et al (2016) Effects of the herbicide dicamba on nontarget plants and pollinator visitation. *Environmental Toxicology and Chemistry* 35:144–175.
- Botías C., David A., Hill E.M., Goulson D. (2016) Contamination of wild plants near neonicotinoid seed-treated crops, and implications for non-target insects. *Sci Total Environ* 566–567:269–278.
- Botías, C. D, A., Horwood J., Abdul-Sada A., et al (2015) Neonicotinoid residues in wildflowers, a potential route of chronic exposure for bees. *Environ Sci Technol* 49:12731–12740.
- Bradbury S.P., Grant, T.J., Krishnan, N. (2017) Landscape Scale Estimates of Monarch Butterfly (*Danaus plexippus*) Population Responses to Insecticide Exposure in an Iowa Agroecosystem. Society of Environmental Toxicology and Chemistry North America 38th Annual Meeting. November 12 -16, Minneapolis, MN.
- Brower L.P., Taylor O.R., Williams E.H., et al (2012) Decline of monarch butterflies overwintering in Mexico: Is the migratory phenomenn at risk? *Insect Conserv Divers* 5:95–100.
- Corn and Soybean Digest (2015) New seed treatment, high-tech corn coming to market.
- David A., Botías C., Abdul-Sada A., et al (2016) Widespread contamination of wildflower and bee-collected pollen with complex mixtures of neonicotinoids and fungicides commonly applied to crops. *Environ Int* 88:169–178.
- Douglas M.R., Tooker J.F. (2015) Large-scale deployment of seed treatments has driven rapid increase in use of neonicotinoid insecticides and preemptive pest management in U.S. field crops. *Environ Sci*



- Technol 49:5088–2097.
- Dunbar M.W., O'Neal M.E., Gassmann A.J. (2016) Increased risk of insect injury to corn following rye cover crop. *J Econ Entomol* 109:1697–1697.
- DuPont (2010) Supplemental labeling. Dupont™ CoragenR Insect Control. For foliar insect control on artichoke, asparagus, corn (field, pop, sweet), legume vegetables, okra, strawberry, sugarcane, tobacco & tuberous and corm vegetables. R-1070-02810 01-26-10. Du Pont Nemours Company, Crop Prot Wilmington, DE 4pp.
- Fisher, K.E., Adelman, J., and Bradbury, S.P. (2017a) Estimating perceptual range of the monarch butterfly (*Danaus plexippus*) with an automated radio telemetry system. Oral presentation. Annual meeting of the Entomological Society of America. November 5 – 8, 2017, Denver, CO.
- Fisher, K.E., Adelman, J., and Bradbury, S.P. (2017b) Testing methods for tracking monarch butterfly movement with radio telemetry. Poster presentation. Annual Meeting of the Ecological Society of America. August 6 – 11, 2017. Portland, OR.
- Flockhart D.T., Brower L.P., Ramirez M.I., et al (2017) Regional climate on the breeding grounds predicts variation in the natal origin of monarch butterflies overwintering in Mexico over 38 years. *Glob Chang Biol* 1–12. doi: 10.1111/gcb.13589
- Flockhart D.T.T., Pichancourt J.B., Norris D.R., Martin T.G. (2015) Unravelling the annual cycle in a migratory animal: Breeding-season habitat loss drives population declines of monarch butterflies. *J Anim Ecol* 84:155–165. doi: 10.1111/1365-2656.12253
- Grant T., Parry, H., R., Zalucki, M.J., and Bradbury, S.P. (2017a) Predicting monarch butterfly movement and egg laying with a spatially-explicit agent-based model: The role of monarch perceptual range and spatial memory. *Ecological Modelling* (Submitted).
- Grant, T.J., Krishnan, N. and Bradbury, S.P. (2017b) Spatially-explicit estimates of monarch butterfly (*Danaus plexippus*) population responses to insecticide spray drift exposure in north central agroecosystems. Poster presentation. Annual meeting of the Entomological Society of America. November 5 – 8, 2017, Denver, CO.
- Grossman K. (2009) Auxin herbicides: Current status of mechanism and mode of action. *Pest Manag Sci* 66:113–120.
- Hodgson E.W., Kemis M., Geisinger B. (2012) Assessment of Iowa soybean growers for insect pest management practices. *J Extension* 50:4RIB6. [http://www.joe.org/joe/2012august/pdf/JOE\\_v50\\_4rb6.pdf](http://www.joe.org/joe/2012august/pdf/JOE_v50_4rb6.pdf)
- Hodgson E.W., VanNostrand G. (2016) Soybean aphid efficacy screening program, 2015. *Arthropod Manag Tests* Section F:1–3.
- Iowa State University (2016) Soybean aphid efficacy evaluations 2016-2005. <http://www.ent.iastate.edu/soybeanresearch/content/extension>.
- Jepsen S., Schweitzer D.F., Young B., et al (2015) Conservation status and ecology of the monarch butterfly in the United States. [http://www.natureserve.org/sites/default/files/natureserve-xerces\\_monarchs\\_usfs-final.pdf](http://www.natureserve.org/sites/default/files/natureserve-xerces_monarchs_usfs-final.pdf).
- Krishnan, N., Bidne, K., Hellmich, R., Coats, J. and Bradbury, S.P. (2017a) Risk assessment of insecticides commonly used in corn and soybean production on monarch butterfly (*Danaus plexippus*) larvae. Society of Environmental Toxicology and Chemistry North America 38th Annual Meeting. November 12 -16, Minneapolis, MN.

- Krishnan, N., Bidne, K., Hellmich, R., Coats, J. and Bradbury, S.P. (2017b) Risk assessment of foliar insecticides commonly used in corn and soybean production on monarch butterfly (*Danaus plexippus*) larvae. Oral presentation. Annual meeting of the Entomological Society of America. November 5 – 8, 2017, Denver, CO.
- Krishnan, N., Bidne, K., Hellmich, R., Coats, J. and Bradbury, S.P. (2017c) Risk assessment of foliar insecticides commonly used in corn and soybean production on monarch butterfly (*Danaus plexippus*) larvae. Poster presentation. 254th American Chemical Society National Meeting. August 20-24, 2017. Washington, DC.
- Krupke C.H., Hunt G.J., Eitzer B.D., et al (2012) Multiple routes of pesticide exposure for honey bees living near agricultural fields. PLoS One. 7(1) doi: <http://doi.org/10.1371/journal.pone.0029268>.
- Long E.Y., Krupke C.H. (2016) Non-cultivated plants present a season-long route of pesticide exposure for honey bees. Nat Commun 7:11629.
- Malcolm, S.B., Cockrell, B.J., and Brower, L.P. 1993. Spring recolonization of eastern North America by the monarch butterfly: successive brood or single sweep migration? In S. B. Malcolm, & M. P. Zalucki (Eds.), *Biology and Conservation of the Monarch Butterfly* (pp. 253-267). Natural History Museum of Los Angeles County; Science Series, 38.
- Nail K.R., Stenoien C., Oberhauser K.S. (2015) Immature monarch survival: effects of site characteristics, density, and time. *Ann Entomol Soc Am* 108:680–690. doi: 10.1093/aesa/sav047.
- NRCS (2016) Monarch Butterfly Wildlife Habitat Evaluation Guide and Decision Support Tool: Midwest Edition 1.1. <https://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/plantsanimals/pollinate/?cid=nrcseprd402207>.
- Oberhauser K., Wiederholt R., Diffendorfer J.E., et al (2017) A trans-national monarch butterfly population model and implications for regional conservation priorities. *Ecol Entomol* 51–60. doi: 10.1111/een.12351.
- Paola O.A., Kaplan I. (2015) Non-target effects of agrochemicals on milkweed and monarch butterflies. 63rd Annu. Meet. Entomol. Soc. Am. Nov. 15-18, Minneapolis, MN.
- Pecenka J.R., Lundgren J.G. (2015) Non-target effects of clothianidin on monarch butterflies. *Sci Nat*. doi: 10.1007/s00114-015-1270-y
- Pleasants J.M., Oberhauser K.S. (2013) Milkweed loss in agricultural fields because of herbicide use: Effect on the monarch butterfly population. *Insect Conserv Divers* 6:135–144. doi: 10.1111/j.1752-4598.2012.00196.x
- Pleasants J.M. (2015) Monarch Butterflies and Agriculture. In: Oberhauser KS, Nail KR, Altizer S (eds) *Monarchs in a Changing World*. Cornell University Press, Ithaca, pp 169–178.
- Pocius, V.M., Debinski, D.M., Bidne, K.G., Hellmich, R.L., & Hunter, F.K. (2017a). Performance of early instar monarch butterflies (*Danaus plexippus* L.) on nine milkweed species native to Iowa. *The Journal of the Lepidopterists' Society*, 71(3), 153-161.
- Pocius, V.M., Debinski, D.M., Pleasants, J.M., Bidne, K.G., Hellmich, R.L., & Brower, L.P. (2017b). Milkweed matters: Monarch butterfly (Lepidoptera: Nymphalidae) survival and development on nine midwestern milkweed species. *Environmental Entomology*. doi.org/10.1093/ee/nvx137.
- Pryby M.D., Oberhauser K.S. (2004) Temporal and geographic variation in monarch densities: Citizen scientists document monarch population patterns. In: *Monarch butterfly Biology and Conservation*. pp 9–20.

- Semmens B.X., Semmens D.J., Thogmartin W.E., et al (2016) Quasi-extinction risk and population targets for the eastern, migratory population of monarch butterflies (*Danaus plexippus*). *Sci Rep* 6:1–7. doi: 10.1038/srep23265.
- Thogmartin W.E., López-Hoffman L., Rohweder J., et al (2017) Restoring monarch butterfly habitat in the Midwestern US: “all hands on deck.” *Environ Res Lett* 12:74005. doi: 10.1088/1748-9326/aa7637
- University of Tennessee Extension (2016) Insect control recommendations for field crops. <https://extension.tennessee.edu/publications/Pages/default.aspx>.
- USEPA (2017) Atmospheric Models. <https://www.epa.gov/pesticide-science-and-assessing-pesticide-risks/models-pesticide-risk-assessment#atmospheric>.
- USFWS (2014a) Monarch Butterfly Listing Petition. [http://www.biologicaldiversity.org/species/invertebrates/pdfs/Monarch\\_ESA\\_Petition.pdf](http://www.biologicaldiversity.org/species/invertebrates/pdfs/Monarch_ESA_Petition.pdf)
- USFWS (2014b) Monarch Butterfly, Status Review. <http://www.regulations.gov/#!documentDetail;D=FWS-R3-ES-2014-0056-0001>.
- Zalucki M.P., Parry H.R., Zalucki J.M. (2016) Movement and egg laying in Monarchs: To move or not to move, that is the equation. *Austral Ecol* 41:154–167. doi: 10.1111/aec.12285
- Zalucki M.P., Lammers J.H. (2010) Dispersal and egg shortfall in Monarch butterflies: What happens when the matrix is cleaned up? *Ecol Entomol* 35:84–91. doi: 10.1111/j.1365-2311.2009.01160.x
- Zalucki M.P., Clarke A.R., Malcolm S.B. (2002) Ecology and behavior of first instar larval Lepidoptera. *Annu Rev Entomol* 47:361–393.
- Zalucki M.P., Malcolm S.B., Paine T.D., et al (2001) It's the first bites that count: Survival of first-instar monarchs on milkweeds. *Austral Ecol* 26:547–555. doi: 10.1046/j.1442-9993.2001.01132.x