

Using opportunistic data to study the distribution and abundance of a warm water elasmobranch at the northern edge of its range

Hiddink, Jan Geert; Charles, Ryan; Moore, Alec

ICES Journal of Marine Science

DOI:

[10.1093/icesjms/fsad183](https://doi.org/10.1093/icesjms/fsad183)

Published: 05/12/2023

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Hiddink, J. G., Charles, R., & Moore, A. (2023). Using opportunistic data to study the distribution and abundance of a warm water elasmobranch at the northern edge of its range. *ICES Journal of Marine Science*. Advance online publication. <https://doi.org/10.1093/icesjms/fsad183>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Using opportunistic data to study the distribution and abundance of a warm water**
2 **elasmobranch at the northern edge of its range**

3

4 J.G. Hiddink, R. Charles & A. B. M. Moore

5

6 School of Ocean Sciences, Bangor University, Menai Bridge, LL59 5AB, United Kingdom,

7 j.hiddink@bangor.ac.uk, +44 1248 382864

8

9 **Abstract**

10 Detecting changes in the distribution and abundance of marine species that are cryptic or
11 occurring in very low abundances is difficult, but essential for assessing their status and
12 informing management. One way of quantifying these changes is through the collation of
13 opportunistic records. We reconstruct the population trajectory and distribution of the
14 common stingray *Dasyatis pastinaca* around Great Britain, using opportunistic records, mostly
15 obtained by recreational anglers. We tested if *D. pastinaca* declined in abundance and body
16 size in response to fishing and if their distribution has shifted northwards in response to
17 warming seas. We obtained 518 records covering the period 1838-2020. After correcting for
18 observation effort, *D. pastinaca* catches reported by anglers showed no long-term trend over
19 50 years, but with a decrease from 1970-1995 and an increase in abundance since 1995. While
20 records of species occurrence were found around much of Great Britain, nearly all were from
21 south of 54° latitude, and records have contracted southwards since 2000. No trend in
22 maximum size through time was detected. In conclusion, we did not find support for the
23 hypothesized declines in abundance and body size or a northward shift in distribution of *D.*
24 *pastinaca* and instead found a southward contraction.

25

26 Keywords: conservation, fisheries management, Dasyatidae, elasmobranch, historical
27 ecology, recreational angling, climate change

28 **Introduction**

29 Detecting changes in the distribution and abundance of uncommon marine species is difficult
30 (Pikitch, 2018), but essential for assessing their population status and identifying whether
31 management actions are required and effective. Many sharks and rays are threatened by
32 fisheries (e.g. Dulvy *et al.*, 2021; Pacoureau *et al.*, 2021). Their distribution and local
33 abundance may also shift in response to oceanographic and climate changes (e.g. Chin *et al.*,
34 2010; Osgood *et al.*, 2021; Hammerschlag *et al.*, 2022), with the latter likely to exacerbate
35 risks from fishing (Walker *et al.*, 2021). Yet, we have little idea of the conservation status of
36 hundreds of species of sharks and rays, including taxa that are infrequently recorded (Walls &
37 Dulvy, 2020, 2021).

38

39 Stingrays (Dasyatidae) are a diverse group of batoid elasmobranchs mainly found in tropical
40 and subtropical regions (Last *et al.*, 2016). The common stingray *Dasyatis pastinaca* (Linnaeus,
41 1758) occurs in the eastern Atlantic and Mediterranean Sea from the shore to about 200 m
42 depth but is more commonly recorded in shallow waters (< 50 m). While this species has quite
43 a large distribution including the Mediterranean, they are considered less common in
44 northern Europe, as this appears to be at the minimum thermal tolerance of its range
45 (Heessen *et al.*, 2016). The British Isles are on the northern edge of its distribution where it is
46 the only demersal stingray species frequently encountered (Last *et al.*, 2016; Ebert & Dando,
47 2020) and is not commercially fished. The species is suspected to have declined by at least
48 30% through European and north African waters over a three generation period (3 times 7.5
49 years), and is assessed as Vulnerable on the International Union for Conservation of Nature
50 (IUCN) Red List of Threatened Species (Jabado *et al.*, 2021), but the abundance data
51 underpinning this assessment are scarce. Because of their infrequent capture in systematic
52 fisheries-independent scientific demersal fish trawl surveys (which may relate to limited
53 overlap between surveys and the spatial distribution of the common stingray as well as their
54 small population size), such surveys are unable to provide reliable abundance trends (Martin
55 *et al.*, 2010; Heessen *et al.*, 2016; Rindorf *et al.*, 2020).

56

57 Common stingrays *Dasyatis pastinaca* have been documented by naturalists (under a wide
58 variety of names) as occurring around Great Britain for over 330 years, in Scotland, (Sibbald,
59 1684; Raye, 2018), England and Wales (Pennant, 1796; Couch, 1841; Herdman & Dawson,
60 1902). It was reported as most common on the English south coast (Yarrell, 1859) and
61 common in south Wales (Dillwyn, 1848), and was mostly recorded very close to shore (Couch,

62 1841; Yarrell, 1859). Over 100 years later Wheeler (1969) showed the distribution of *D.*
63 *pastinaca* all around Great Britain, noting the Thames estuary as a particularly important
64 habitat. Yet despite attracting this level of interest for many years, and being a warm water
65 species on the northern edge of its range in warming seas, there has been no research on how
66 the distribution or abundance of this species may have changed through time around Great
67 Britain. As a result of its larger size and distinctive nature (a relatively rare, venomous species
68 typically associated with warmer waters), the common stingray is popular in recreational sea
69 angling in the UK, where it can be caught on a wide range of baits fished on the seabed
70 (<https://britishseafishing.co.uk/common-stingray/#>, [https://hookpoint.co.uk/how-to-catch-](https://hookpoint.co.uk/how-to-catch-stingray/)
71 [stingray/](https://hookpoint.co.uk/how-to-catch-stingray/)). This means that there are records available that may provide insights, previously
72 overlooked, on contemporary population size and distribution. When these available
73 historical and contemporary records are considered together, there is an opportunity to
74 examine temporal changes to this population over the last two centuries. Historic data are
75 often being incomplete and patchy and subject to bias that can change through time. These
76 complications, however, do not lessen the value of historical records, and past studies have
77 shown that the detection and explanation of historical trends and variability are essential to
78 informed management (e.g. Swetnam *et al.*, 1999; Thurstan *et al.*, 2015).

79

80 We define ‘opportunistic records’ as those that are not the direct result of a scientific field
81 survey aimed at quantifying fish abundance. Opportunistic records therefore include those
82 from commercial fisheries, recreational fisheries, naturalists and the general public, and single
83 records in scientific papers, but not those from scientific trawl surveys. Opportunistic records
84 can be useful as an indicator of local population status or trends (e.g. Grant *et al.*, 2022),
85 although they can also be problematic due to a lack of time series consistency, and underlying
86 observation effort may be unclear (Swetnam *et al.*, 1999). The difficulty of monitoring
87 uncommon species that are seldom caught in scientific surveys is illustrated in a study on
88 angelsharks *Squatina squatina* in Wales, UK (Hiddink *et al.*, 2019). The northeast Atlantic is
89 one of the most intensively monitored oceans in the world, nevertheless only a handful of
90 angelshark were caught in >40 years of scientific fisheries-independent trawl surveys (>25,000
91 hauls) (Heessen *et al.*, 2016). In contrast to this paucity of records, 1,860 angelshark records
92 from a 50-year period were collated from the coastal waters of Wales using interviews,
93 charter-boat skipper log books and other opportunistic sources (Hiddink *et al.*, 2019). The
94 angelshark study estimated that there had been a 70% decline in abundance over 46 years,
95 with continued presence in Cardigan Bay. Although Hiddink *et al.* (2019) made corrections for

96 observation effort by taking account of the number and age distribution of the observers,
97 uncertainty remains about the reliability of opportunistic records for the estimation of
98 abundance and distribution trends. If trends estimated using opportunistic records are driven
99 mainly by changes in observation effort over time, it can be expected that different
100 uncommon species would show similar trends in abundance and distribution that are all
101 following the observation effort, when using the same type of opportunistic data for different
102 species. However, if the number of records for different species show divergent trends in
103 space and time, we can be more confident that opportunistic records provide an indication of
104 real population trends rather than changes in observation effort alone. For example, in the
105 Mediterranean Sea the number of opportunistic sightings of bluntnose sixgill shark *Hexanchus*
106 *griseus* have been decreasing while the number of opportunistic sightings of shortfin mako
107 *Isurus oxyrinchus* are increasing, meaning that it can be inferred that the population size
108 trend of shortfin mako is more positive than that for bluntnose sixgill shark (Bargnesi *et al.*,
109 2022).

110

111 We should also keep in mind that opportunistic recording and fishing activities may favour
112 different species at different times (Boersch-Supan *et al.*, 2019). For example, anglers may
113 target particular fish species, and this may guide exact fishing location, gear, bait, and other
114 factors (Lewin *et al.*, 2006). Collectively, opportunistic records can be useful for the study of
115 'rare' species, however, interpretations for how opportunistic records indicate changes at the
116 population level need to be carefully considered with respect to available information the
117 type of observations, on temporal changes and biases in observation effort (Swetnam *et al.*,
118 1999; Hiddink *et al.*, 2019).

119

120 Evidence from aquatic environments suggests that body size can decline as a result of climatic
121 warming at the level of the individual, population and community (Daufresne *et al.*, 2009;
122 Shackell *et al.*, 2010) as well as in response to exploitation pressure (e.g. Anderson *et al.*,
123 2008). Increasing sea bottom temperatures with climate change are likely to favour smaller
124 species and individuals (Atkinson, 1994; Hiddink & ter Hofstede, 2008). Climate change is also
125 likely to continue to result in a northward shift of the poleward edge of the distribution of
126 species with warm-water affinities (Perry *et al.*, 2005; Dulvy *et al.*, 2008). It has been shown
127 that fish species track temperatures under global warming by moving both latitudinally and
128 into deeper water (Dulvy *et al.*, 2008; Burrows *et al.*, 2019). Large shifts in the distribution and
129 abundance of elasmobranchs in the North Sea have been recorded (Sguotti *et al.*, 2016).

130 Species of highly mobile fish, such as sharks and rays, may be more responsive to temperature
131 change in time and space than analogous communities on land, potentially as a consequence
132 of living closer to their thermal limits (Burrows *et al.*, 2019). High fishing pressure has also
133 resulted in a decrease in the abundance and body size of large-bodied and slow-reproducing
134 fish species, including sharks and rays, over the last 100 years (Quero, 1998; Rogers & Ellis,
135 2000; Engelhard *et al.*, 2015).

136

137 This study aims to examine temporal occurrence records of *D. pastinaca* to determine if its
138 geographic range has changed over the last two centuries and whether maximum reported
139 sizes have reduced over time. Here we reconstruct the population trajectory and distribution
140 of the common stingray *Dasyatis pastinaca* over nearly two centuries around Great Britain
141 using opportunistic records, such as social media, newspapers, angling magazines and the
142 scientific literature (where records did not come from systematic fisheries surveys).
143 Recreational sea angling is popular in Great Britain (Hyder *et al.*, 2018) and provided most of
144 the records used in this study. We hypothesize that common stingrays, like other large
145 elasmobranchs (Walker & Hislop, 1998; Wolff, 2000; Rindorf *et al.*, 2020), have declined in
146 abundance and body size in response to fishing and may have shifted their distribution
147 northwards in response to climate change.

148

149 **Methods**

150 Opportunistic sightings of the common stingray *Dasyatis pastinaca* were compiled from
151 publicly accessible information sources. The study area is the island of Great Britain (England,
152 Scotland and Wales, including their component adjoining islands), and its surrounding seas.
153 All records included in this paper are considered as opportunistic records, which are defined
154 here as any records that were not the results of targeted survey effort in scientific trawl
155 campaigns (e.g. Heessen *et al.*, 2016), and can therefore include records from the scientific
156 literature..

157

158 A 'record' refers to an individual common stingray specimen encountered on one date,
159 whereas a 'report' refers to a unique common stingray encounter event: one report could
160 therefore comprise several common stingray records.

161 **Data sources**

162 ***Trophy-catch and magazine records***

163 This study updated past compilations of two major recreational sea angling data sources
164 which were identified and reviewed by Richardson *et al.* (2006); *SeaAngler* magazine (Kelsey
165 media) and the annual reports of the National Federation of Sea Anglers (NFSA). The most
166 popular recreational UK fishing magazine, *SeaAngler* produces 12-13 issues per year. Reader-
167 submitted UK common stingray catches are published in *SeaAngler* if specimens met or
168 exceeded the shore-based or vessel-based qualifying weight (9.1 kg and 11.3 kg, respectively),
169 but may be mentioned for other reasons too. The number of submitted records is likely to
170 depend on both the abundance of stingrays and the number of active anglers in a region.
171 Our search consisted of examining all pages of each *SeaAngler* issue, to find catch reports
172 and common stingray-specific articles. A total of 598 *SeaAngler* issues published from March
173 1972 to July 2021 were reviewed. Issues published onwards from 2014 were accessed through
174 a digitalised archive on go.readly.com, and hard copy issues published prior to 2014 were
175 reviewed at Kelsey Media, Kent. We also extracted all records of the angelshark *Squatina*
176 *squatina* from *SeaAngler* magazine using the same methodology for comparison of temporal
177 and spatial trends. The NFSA produced annual reports of trophy catches (catches above a
178 specified weight threshold) for a wide variety of species across the UK, including the species,
179 weight and location of capture, from 1976 to 2002, and had 40,000 members in 2006, but is
180 now defunct (Richardson *et al.*, 2006). These trophy records reports had a qualifying weight
181 threshold of 15 kg for common stingray.

182

183 ***Books, scientific papers and historical literature***

184 Where possible, Boolean operators (“and”, “or”) were used to ensure exact search terms were
185 incorporated; minimising time spent reviewing irrelevant material. All searches (across all
186 information sources) containing predefined terms (“Common stingray,” “stingray,” “*Dasyatis*
187 *pastinaca*” or “*pastinaca*” and any spelling variations that the searches returned) were then
188 examined for reports. If relevant data were not returned, the search string was adapted to
189 increase its sensitivity to the study question, by including locations (United Kingdom, Wales,
190 England, Scotland, Britain), and previous taxonomic classifications (*Trygon pastinaca*). The use
191 of historical local names for the common stingray (e.g. English local names “fire-flare”, “fiery-
192 flare”, or names in other languages (Welsh: “Morgath ddu”) did not yield any additional
193 records. Private browsing options were used to avoid previously cached terms. The identified
194 sources went through a manual four-stage screening process of search results to assess the
195 material suitability: title relevance, abstract relevance, figures/image thumbnail and full text.

196

197 Historical literature is accessible via the Biodiversity Heritage Library
198 (<https://www.biodiversitylibrary.org/>) and the websites archive.org and books.google.com.
199 Searches returned natural history literature, personal accounts from fishermen, museum
200 catalogues and zoological compilations of local fauna: collectively referred to here as “books”.
201 Peer-reviewed scientific literature (“scientific papers”) was systematically searched using the
202 search tools Google Scholar and ProQuest (Clarivate), and the publishers’ websites for Wiley
203 journals, Springer journals, ScienceDirect (Elsevier journals). Grey literature was also searched
204 to ensure the comprehensiveness of the study as suggested by Haddaway and Bayliss (2015).
205 This included other angling magazines separate from the trophy-catch dataset, and screening
206 bibliographies of relevant material identified other relevant literature.

207

208

209 ***Online searches: News articles, social media, forums, other sources***

210 Further common stingray records were found in 34 databases using a search conducted across
211 online angling forums (e.g. <https://norfolkfishing.com/>), social media platforms (e.g. Twitter,
212 Facebook) and ‘other’ sources: the latter refers to data portals, local environmental record
213 centres and museum catalogues (see the full list of sources in Supplementary material Table
214 S1). A ‘snowball’ sampling procedure searched forum responses, article comments and social
215 media post threads for common stingray mentions. This returned public discussion of a)
216 qualitative spatial-temporal distribution and b) the number of fish encountered. Despite
217 creating a non-random sample, this produced reports which would have otherwise gone
218 undocumented. We followed the guidelines for the ethics of using social media in fisheries
219 research in Monkman *et al.* (2018), and only included information that was accessible without
220 creating an account.

221

222 **Data extraction and handling**

223 For each record we collected where available the: number of individuals, size (disc-width,
224 weight and/or total length), sex, observation location, date-of-encounter (year and month)
225 and type of data source (e.g. book, trophy records, forum). Although *D. pastinaca* is the only
226 dasyatid stingray to occur regularly around British coasts, photographs were used where
227 available to confirm identification against the most recent comprehensive identification
228 source (Ebert & Dando, 2020), as some sources (e.g. social media, popular press) have
229 reported rajid skates as ‘stingrays’ (e.g.
230 <https://www.ayradvertiser.com/news/16051350.stingray-found-ballantrae-beach>). We also

231 acknowledged the possibility that the much less abundant pelagic stingray (*Pteroplatytrygon*
232 *violacea*) or vagrants (e.g. Tortonese's stingray *D. tortonesei*) could occur around Britain (Ellis,
233 2007; Ebert & Dando, 2020). When no photos were available, the identification of the original
234 source was accepted. Because anglers and other sources are more likely to report and
235 publicize large common stingrays (as 'trophy' or 'specimen' fish), the reported body sizes are
236 likely to be skewed towards larger sizes (although small specimens can also be reported,
237 especially in forums). Any changes in body size therefore need to be interpreted as changes
238 in the maximum size of common stingrays in the population, rather than changes in the mean.

239

240 The date-of-encounter was documented for each stingray rather than the date-of-publication.
241 The date-of-publication was used as a reference point if a past encounter was recalled. For
242 example, a report "seven years ago" was assigned a date seven years prior to the date of
243 publication. Positions of encounter locations were estimated based on given descriptions
244 when longitudes and latitudes were not stated. Commonly, the location name was given if the
245 encounter was shore-based. Less often, vessel-based encounters produced distance
246 estimations from a land-based reference point. Many reports did not report the size of the
247 stingray, but where the size was reported one of three measures could be given: disc width
248 (the widest point of the fish), total length or weight. Because weight was most commonly
249 reported, all measures were converted to weight from disc width (DW) and length,
250 respectively, using the following equations that were developed for *D. pastinaca*: weight (g) =
251 $0.0132 \text{ DW}^{3.06}$ (Froese & Pauly, 2004) where disc width was estimated from length using DW
252 (cm) = $0.60 \text{ total length (cm)} - 0.61$ (Heessen *et al.*, 2016) for 29% of records.

253

254 We checked for presumed duplicate records by evaluating if there were records from the
255 same location (0.1 degree precision) in the same year from different sources, and duplicates
256 were removed.

257

258 **Analysis**

259 Analysis was restricted up to the last complete year of data, 2020. For most types of data
260 collated here, it was not possible to correct for observation effort and calculate a catch per
261 unit effort. However, for the trophy and magazine catches only, we used the *SeaAngler*
262 magazine readership data as a proxy for sea angling effort to calculate the number of common
263 stingrays reported per 1000 readers per year. The number of readers was considered as likely
264 to broadly reflect angling effort and the number of catches that are reported (Richardson *et*

265 *al.*, 2006). Magazine readership was supplied by the publishers of *SeaAngler* as the combined
266 paper and online readership for the years 1974–2019. We did not find sources that could be
267 used as a proxy for observation effort for the other data sources.

268

269 Biogeographic distribution shifts were quantified by estimating the maximum, median and
270 minimum latitude of all records from a 5-year period. To reduce the influence of outliers when
271 assessing distributional shifts, the maximum and minimum latitude were estimated as the
272 0.95th and 0.05th quantile of the latitude of records per 5-year period. Results from this analysis
273 were only reported when there were >5 data points per 5-year period, resulting in estimates
274 of latitude for years after 1965 only. We tested if the fraction of records from southeast
275 England (east of 2° west longitude and south of 52° north latitude) changed over time. We
276 also tested if the months in which stingrays were recorded changed over time, as it can be
277 expected that they are seen earlier in the year with climate change (Schlaff *et al.*, 2014).

278

279 We also used the approach of McPherson and Myers (2009) to estimate the magnitude of any
280 change in the common stingray population and sensitivity of this to a range of observation
281 effort scenarios. This model builds on a different set of assumptions than the *SeaAngler*
282 analysis and a comparison between the outputs of the approaches therefore helps us in
283 assessing the robustness of our conclusions. The McPherson and Myers (2009) approach
284 extracts the relative magnitude of population change in the number of reported sightings by
285 fitting a series of generalized linear models the difference in the count data between any
286 reference date and the most recent point with data (2020), to provide multiple estimates of
287 declines under alternate scenarios of observation effort and explicitly address uncertainty
288 over variations in observation effort. This approach enables to simulate various scenarios for
289 proportional change in the observation effort. Values smaller than 0% suggest a declining
290 trend, while values equal to or larger than 0% suggest a stable or increasing population. For
291 more details about the analytical method see McPherson and Myers (2009).

292

293 **Results**

294 In total, we obtained 356 common stingray reports that contained 518 individual common
295 stingray records (Table 1), reflecting that most reports were of a single stingray. Records
296 covered the period from 1838 to 2020, but were very scarce and infrequent in the earlier years
297 (0-2 records per 5 years). Most records were catches reported by recreational anglers through
298 various sources, with around a third as catches reported in *SeaAngler* magazine. Forums, news

299 articles and social media together contributed 23% of records. Most of the remainder came
300 from online databases such as the NBN Gateway (<https://nbn.org.uk/>). There were several
301 records from commercial vessels that were reported in a series of papers on notable fish
302 captures (Wheeler & Blacker, 1969, 1972; Wheeler *et al.*, 1975).

303

304 The total number of common stingray records increased strongly over time, driven by an
305 increase in reports in *SeaAngler* magazine, forums, social media and regional newspapers that
306 have become available online since the year ~2000 (Figure 1A). *SeaAngler* contributed the
307 largest dataset where the observation effort was likely to be relatively constant (unlike for
308 example social media), and the temporal trend in the number of reports in this dataset was
309 seen to fluctuate with an increase in *SeaAngler* reports during a period of declining readership
310 since around the year 2000 (Figure 1B). Common stingray reports per 1,000 *SeaAngler* readers
311 show a U-shaped pattern, with a decrease from 1970 to 1995, followed by an increase to a
312 similar or even higher level than the start of the time-series by around 2015 (Figure 1C). The
313 dip is caused by a lower number of reports in a period with a high readership in the 1990s
314 (suggesting a lower abundance in that period), while the higher values at the start and the
315 end of the time-series relate to higher numbers of reports in periods with a lower readership
316 (suggesting a higher abundance). For comparison, angelshark records showed a very similar
317 decline in reports per 1,000 readers at the start of the observation period, while the uptick in
318 records and the sightings per 1,000 readers at the end of the observation period was very
319 limited compared to the common stingray (Figure 1C). For both species, the non-linear GAM
320 was a better fit to the data than a linear model (the AIC for the GAM was lower than for the
321 linear model). This divergence in the trend between the two species in recent years suggests
322 that the increase in stingray records since 1995 represents a real increase in abundance.

323

324 Records of common stingray were widely distributed around Great Britain, including all Welsh
325 and most English coasts and to the east of Scotland, but we did not find records from the north
326 and west of Scotland, few from northeast England, and only a single record from northwest
327 England (Figure 2A). Nearly all (97%) records were from southern Britain south of 54° latitude.
328 Although there were relatively few (n=8) records from Scottish waters, they occurred across
329 several time periods (Figure 2). The highest concentration of records was found in southeast
330 England: on the English Channel coast, centred around Hampshire and West Sussex and the
331 coast of the greater Thames estuary (>80% in recent decades, Figure 2A & 3C). While formerly
332 recorded around most of Britain, common stingray records are now only found in southern

333 Britain (Figure 2B-H). Despite an increase in the total number of stingray records over time in
334 Britain, the last common stingray recorded in the waters of Scotland and northern England
335 (>53°N) was 1991 and 1998 respectively. Most Welsh records were from the south (Bristol
336 Channel, bordering England) since 1975, and mid Wales (Cardigan Bay) in 1975–1999. Since
337 2000 we found few Welsh records (1 in north Wales in 2008, none in mid Wales, and 4 in south
338 Wales). Most records (76%) came from the coast (with a reconstructed position within 2 km
339 of land) rather than further offshore, reflecting the fact that most records were reported by
340 anglers fishing from the shore rather than reports by boat users. The spatial pattern for the
341 common stingray contrasts strongly with that for the much less commonly recorded
342 angelshark, for which *SeaAngler* records remain concentrated in Wales only where it is more
343 commonly recorded than the common stingray (Figure S1).

344

345 The maximum and median latitude at which common stingrays have been found in Britain has
346 significantly moved south by 150-350 km (Figure 3A, maximum latitude $R^2 = 0.47$, $F_{1,9} = 8.06$,
347 $p = 0.019$, median latitude $R^2 = 0.81$, $F_{1,9} = 39.4$, $p < 0.001$). Given that the southern boundary
348 of the distribution of the common stingray lies 1,000s of kms south of the study area, the
349 minimum latitude is effectively defined by the boundary of the study area and did not show a
350 change over time (minimum latitude $R^2 = 0.23$, $F_{1,9} = 2.76$, $p = 0.131$). The fraction of records
351 from southeast England (east of 2° west longitude and south of 52° north latitude) seems to
352 increase over time ($R^2 = 0.56$, $F_{1,4} = 5.19$, $p = 0.131$), with 88% of records from the period 2000–
353 2020 coming from this region (Figure 3C). This is driven by an increase over time in the number
354 of records from lower latitudes (Figure 3A, B&C, where stingray records were always more
355 numerous, while the number of records at higher latitudes decreased in recent years (Figure
356 3B). The months in which common stingrays were recorded did not change significantly over
357 time (linear regression, $F_{1,47} = 0.0021$, $p = 0.61$, Figure S2).

358

359 Estimates of the magnitude of change in stingray abundance from any given reference year to
360 2020, based on unstandardized reports from *SeaAngler* using the McPherson and Myers
361 method (2009) and assuming a linear change over time (Figure 4), indicates either no change
362 or an increase in common stingray abundance, if no change in observation effort is assumed
363 (0%). If observation effort is assumed to have halved (-50%), which Figure 1b suggests is
364 plausible, the model predicts that the common stingray has increased from the 1970s and
365 early 1980s to 2020 by a minimum of 10.2% (CIs -61.2 to +190.6%). If observation effort is
366 assumed to have doubled (100%), the model predicts that the common stingray has decreased

367 in abundance from the 1970s and early 1980s to 2020 by a maximum of 72.2%% (CIs 27.5 to
368 90.3%). This analysis therefore suggests that under the most plausible pattern of a decrease
369 or stable observation effort, stingray abundance has increased. The decrease in abundance to
370 1995 and the subsequent increase to 2020 is represented by the inverted U-shape of the
371 inferred changes in abundance in the figure. The low values on the left of the figure indicate
372 little change in abundance from the start to the end of the time-series, and the high values in
373 the middle of the figure indicate increases in abundance from around 1995 to the end of the
374 time-series.

375

376 Most of the recorded weights were provided for trophy catches, and therefore were
377 representative of the largest individuals in the population (15 to 20 kg) rather than an
378 indication of the body size of the population in general. No trend in the recorded weight over
379 time was detected (Figure 3D, $R^2 = 0.11$, $F_{1,3} = 0.39$, $p = 0.57$).

380

381 **Discussion**

382 Our study shows that changes in the distribution and abundance of an uncommon marine
383 species can be evaluated by a retrospective analysis of opportunistic records. Here, we
384 present evidence that, contrary to our expectations, the population of the common stingray
385 around Great Britain has shown no long-term trend over 50 years, but with a decrease from
386 1970 to 1995 and an increase in abundance since 1995. The spatial distribution of records has
387 become more concentrated in southern England, with very few recent Welsh records. This
388 southerly shift in apparent distribution is driven mainly by an increase in records from around
389 the Isle of Wight (possibly resulting from an increase in abundance or of targeted angling
390 efforts for stingrays). Unfortunately, the spatio-temporal distribution of angling has not been
391 mapped and it is therefore difficult to make detailed corrections for observation effort.
392 Common stingray sightings have always been sporadic in Scotland, and none have been
393 recorded there in the last 20 years. Because they have likely never been abundant in Scotland,
394 records will have been more newsworthy, which may have lead to a higher likelihood of
395 reporting. As such, an absence of records is likely to reflect a current rarity or absence of
396 stingray in Scottish waters. As observation effort and reporting through online sources has
397 been increasing substantially over time, it seems likely that this lack of records in Scotland and
398 most of Wales represents a lower abundance of common stingrays there from 1975 onwards.
399 Unsurprisingly, the number of records obtained from social media, forums and other internet-
400 based media increased strongly after 2000, and these increases will reflect increases in the

401 availability and use of these types of sources rather than an actual increase in common
402 stingray abundance.

403

404 As for a previous study involving angelsharks (Hiddink et al., 2019), our findings from rich
405 opportunistic data for stingrays provide a valuable complement to scientific trawl surveys:
406 common stingrays were only caught in small numbers in an extensive program of trawling
407 across the NW European shelf during the period 1977-2013 (Heessen et al., 2016), in contrast
408 to the 356 stingray records we obtained. Additionally, both the trawling and opportunistic
409 datasets appear to corroborate each other in terms of spatial occurrence, with most records
410 around southeastern England/the Eastern English Channel (Heessen et al., 2016).

411

412 The ultimate challenge in the interpretation of opportunistic records is separating true
413 population trends from changes in the observation effort (e.g. McPherson & Myers, 2009),
414 and we could only estimate observation effort for one of the used data sources, the trophy
415 fish catches. The interest in reporting of seldomly encountered and unusual fish may have
416 varied over time (e.g. there was a set of papers describing 'rare and little-known' catches
417 spanning 1966 to 1971, Wheeler & Blacker, 1969, 1972; Wheeler *et al.*, 1975). Only *SeaAngler*
418 records yielded a suitable number of records for a quantitative analysis, limiting this most
419 rigorous analysis to the years 1974-2019. The analysis of this dataset showed that common
420 stingray abundance fluctuated with no long-term trend over 50 years, but with a decrease
421 from the 1970s to 1995 and increased again since 1995. The comparison of the temporal
422 and spatial distribution of common stingrays with patterns of angelsharks recorded using the
423 same method (*SeaAngler* reports) is informative (reported here, and in Hiddink *et al.*, 2019,
424 although scale of the angelshark work there was more extensive because it also include
425 interviews with fishers): while the number of common stingray reports was fairly stable from
426 1970, angelshark records declined strongly since the 1990s and has not recovered. While
427 there have been very few records of common stingrays from north and mid Wales in recent
428 decades, angelsharks maintain a stronghold in mid Wales, and Hiddink *et al.* (2019) report
429 hundreds of records there since 2000. The divergent trends between these two species
430 suggest that the observed patterns are unlikely to be solely driven by changes in observation
431 and reporting effort, and instead reflect actual changes in stingray populations. They could
432 nevertheless be partly driven by observation and reporting effort given that angling catches,
433 especially for seldomly encountered species, can be highly reliant on a handful of anglers or

434 charter boats, and if these change, this can make a large difference to the numbers of local
435 records.

436

437 This study adds to a growing body of research using opportunistic records to reconstruct
438 population and distribution trends of uncommon and cryptic marine animals (e.g. McPherson
439 & Myers, 2009; Curtis et al., 2014; Barbini et al., 2015; Olson et al., 2018; Hiddink et al., 2019).
440 For each of these studies, it seems likely that the probability of reporting an observation has
441 been increasing over time, and the key to robust conclusions in each of these studies has been
442 to separate the trends in abundance from the trends in observation and/or reporting effort,
443 and different studies have taken different approaches. Olson et al. (2018) implicitly assumed
444 that observation and reporting effort of killer whales *Orcinus orca* was constant as they did
445 not correct for potentially changing effort levels. Barbini et al. (2015) used the number of
446 classified advertisements offering fishing guide services published per year in a magazine as a
447 measure of observation effort. Several studies have used the approach developed by
448 McPherson and Myers (2009) that simulates various scenarios of change in the observation
449 effort to evaluate the sensitivity of the inferred magnitude of decline to observation effort
450 (Curtis et al., 2014; Hiddink et al., 2019), while Hiddink et al. (2019) also inferred observation
451 effort from the age distribution and number of respondents. Despite the similarity in the
452 methods and analyses among the studies mentioned above, the inferred relative abundance
453 patterns vary widely among those studies and in comparison to the current results. As it seems
454 likely that the probability of reporting an observation has increased over time for all studies,
455 it is reassuring to see that these trends in the number of records and inferred abundance are
456 different between stingrays (strong increase in recent years) and angelsharks (no or weak
457 increase in recent years), as this gives confidence that we have been evaluating real trends in
458 abundance and distribution rather than artefacts of the recording effort only, and it shows
459 how opportunistic datasets can be valuable tools for illuminating spatial and temporal trends.
460 The sensitivity analysis of our dataset using the McPherson and Myers (2009) method suggests
461 that the long-term trends in common stingray abundance are not particularly sensitive to the
462 halving in observation effort that the *SeaAngler* readership data suggest (the confidence
463 intervals overlap with zero). A doubling in observation effort could have masked substantial
464 declines in their abundance, but we have no evidence to suggest that observation effort is
465 likely to have substantially increased.

466

467 The records presented here are largely from shallow coastal areas, because most of them
468 were obtained from anglers fishing from the shore or small inshore boats. In other studies,
469 very different habitat preferences of common stingray were inferred. For example, Martin *et*
470 *al.* (2010) reported that this species was found offshore in deep waters where tidal currents
471 are moderately intense, based on trawl survey data. This suggests that the mode of data
472 collection has a major effect on the inferred pattern of habitat use. Because our records were
473 mostly from coastal waters, we can only draw conclusions about changes in the distribution
474 in coastal waters. Size may also play a role in habitat preference: based on limited common
475 stingray records from trawl surveys, Heessen *et al.*, (2016) reported smaller individuals in
476 shallower (mode = 40 cm DW at <50m water depth) water, with larger individuals in deeper
477 water (mode = 70 cm DW).

478

479 Our hypothesis that common stingrays declined in abundance and maximum body size in
480 response to fishing was not supported by the results, as no long-term decline in either
481 parameter was obvious. Several other large species of rays have in fact increased in
482 abundance in the Celtic Seas and displayed a fairly stable abundance in the North Sea in the
483 last few decades (Engelhard *et al.*, 2015; Heessen *et al.*, 2016). Similarly, a shorter-term study
484 using only scientific trawl records showed no significant change in common stingray
485 abundance from 1995 to 2015 in the North Sea (Rindorf *et al.*, 2020). The observed maximum
486 body sizes in our study (130 cm disc width (DW), 137 cm length which converts to 82 cm DW,
487 36 kg which converts to 126 cm DW) are very large compared to reported maximum body
488 sizes of common stingray of 60–68 cm DW in other sources (Heessen *et al.*, 2016; Last *et al.*,
489 2016). Although these conversions from length and weight to DW depend on the accuracy of
490 the relationship being used and are highly uncertain, it seems plausible that common stingray
491 grows much larger than 60–68 cm DW. Last *et al.* (2016) and Ebert and Dando (2020) note
492 reports of up to 140 cm DW for common stingray are dubious, but lengths of up to 164 cm
493 were reported in Heessen *et al.*, (2016). Furthermore, the subtly different Tortonese's stingray
494 *D. tortonesei* (not yet known to occur in British waters) is reported as attaining a greater DW
495 than common stingray (84 cm), but the two species have only fairly recently been separated
496 due to their morphological similarity, and thus their biological parameters are less well
497 defined and possibly confounded. It is likely that the larger specimens we recorded represent
498 expanded maximum sizes for common stingray, although we do not discount the possibility
499 that individual Tortonese's stingray could occur as vagrants in British waters. It may be

500 possible that this species attains larger sizes in these higher latitudes, given that this is a
501 common pattern (Atkinson, 1994).

502

503 Our hypothesis that common stingrays would have shifted their distribution northwards to
504 track changes in sea bottom temperature in response to climate change (Hiddink *et al.*, 2015)
505 was not supported and we instead found an opposite pattern, with a contraction in common
506 stingray record distribution towards the south-east coast of England. Our finding of a
507 southward contraction is consistent with apparent disappearance from the south-eastern
508 North Sea since the mid-1960s (Wolff, 2000). The distribution of angling effort alone cannot
509 explain this, as areas of high angling effort are found throughout England and Wales and some
510 parts of Scotland, and there is no reason to assume that rays are more favoured as a target
511 species in the southern and southern-eastern English coasts than in the southwest or
512 northeast (Monkman *et al.*, 2018; MMO, 2020). The southward shift of the distribution of
513 common stingray records could be related to the divergent spatial patterns in human
514 population growth in the UK, with greater growth in the south, and therefore the readership
515 of the *SeaAngler* could have become more concentrated in the south. However, the
516 geographic centre of human population density in Great Britain only shifted ~34 km to the
517 southeast between 1970 and 2020 (https://en.wikipedia.org/wiki/Center_of_population),
518 while the centre of stingray records shifted ~155 km south over that period, and the human
519 population shift can therefore only explain a minor amount of this southward shift.
520 Nevertheless, angling effort is not uniformly distributed and undergoes spatial and temporal
521 changes in distribution, and a loss of effort (e.g. charter boats) for whatever reason might
522 partly explain observed changes such as decreases in numbers observed in certain areas. Such
523 a pattern of southward contraction of geographic range has only been observed (based on
524 long-term scientific surveys) for a few fish species (Dover sole *Solea solea* and the Norway
525 pout *Trisopterus esmarkii*) and about 10% of species of benthic invertebrates in the North Sea,
526 in contrast to the majority of species which expanded north (Perry *et al.*, 2005; Hiddink *et al.*,
527 2015)(Burrows *et al.*, 2019). It is hard to explain this contraction of the distribution of the
528 common stingray, in particular when the total number of records was highest in the later
529 periods because of the increasing use of social media and online forums. The most plausible
530 explanation is that parameters other than mean temperature, such as fishing pressure, food
531 availability or availability of spawning habitat, are driving their range-shifts and -contractions
532 (VanDerWal *et al.*, 2013). Commercial fishing continues around all of Great Britain without
533 great shifts in effort distribution and is therefore not expected to lead to a spatial shift in

534 records (Jennings *et al.*, 1999). Nevertheless, if populations in northern Britain are at their
535 lower thermal tolerance limit, which is likely even though thermal limits have not been
536 independently estimated, it could be expected that population density has always been very
537 low. Therefore, it is possible that their naturally low abundance in the north of Britain has
538 been reduced even further by factors such as commercial fishing, while denser populations in
539 the south of Britain have tolerated fishing pressure better.

540

541 Future work aiming to use historical data, angling catches and other opportunistic to better
542 understand population trends of uncommon marine fauna would benefit from approaches
543 that avoid, or correct for, the biases that may be introduced by uneven observation and
544 reporting efforts. For current angling effort, it should be possible to map the spatio-temporal
545 distribution of angling effort through approaches such as field surveys, interviews and angling
546 shop expenditure, but correcting fully for observation effort in historical data is unlikely to be
547 possible. Comparison of the inferred trends in abundance using opportunistic sources with
548 commercial landings and scientific trawl surveys might help us to understand which of these
549 sources of data are most suitable for reconstructing abundance trends.

550

551 In conclusion, we did not find support for the hypothesized declines in abundance and body
552 size or for northward shifts for the common stingray, and instead found a contraction of
553 distribution records towards southern England. Divergence between the temporal and spatial
554 distribution patterns of common stingrays and angelsharks using a similar methodology show
555 that the observed patterns are not only driven by patterns in observation effort for large
556 elasmobranchs in British waters. Therefore, opportunistic records can be suitable to better
557 understand population trends of seldomly encountered, large and charismatic marine species.
558 We did not find evidence of long-term decline in the population size of the common stingray
559 around Great Britain, and abundance seems to have increased since 1995. The trend in
560 records of common stingray around Great Britain therefore may not match the overall trend
561 for the NE Atlantic as a whole that led to it being classified as Vulnerable by the IUCN based
562 on scarce data in 2021 (Jabado *et al.*, 2021).

563

564 **Acknowledgements**

565 We thank Kelsey Media for giving us access to the *SeaAngler* magazine back catalogue.

566

567 **Contributions**

568 JGH and AM generated the idea. RC collated the records. JGH and RC analysed the data. All
569 authors wrote and reviewed the manuscript.

570

571 **Data Availability statement**

572 On publication, the collated records will be made available as supplementary material.

573

574

References

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594 Anderson, C.N.K., Hsieh, C.-h., Sandin, S.A., Hewitt, R., Hollowed, A., Beddington, J., May,
595 R.M. & Sugihara, G. (2008) Why fishing magnifies fluctuations in fish abundance.
596 *Nature*, **452**, 835-839.

597 Atkinson, D. (1994) Temperature and organism size: a biological law for ectotherms?
598 *Advances in ecological research*, **25**, 1-1.

599 Bargnesi, F., Moro, S., Leone, A., Giovos, I. & Ferretti, F. (2022) New technologies can
600 support data collection on endangered shark species in the Mediterranean Sea.
601 *Marine Ecology Progress Series*, **689**, 57-76.

602 Boersch-Supan, P.H., Trask, A.E. & Baillie, S.R. (2019) Robustness of simple avian population
603 trend models for semi-structured citizen science data is species-dependent.
604 *Biological Conservation*, **240**, 108286.

605 Burrows, M.T., Bates, A.E., Costello, M.J., Edwards, M., Edgar, G.J., Fox, C.J., Halpern, B.S.,
606 Hiddink, J.G., Pinsky, M.L., Batt, R.D., García Molinos, J., Payne, B.L., Schoeman, D.S.,
607 Stuart-Smith, R.D. & Poloczanska, E.S. (2019) Ocean community warming responses
608 explained by thermal affinities and temperature gradients. *Nature Climate Change*,
609 **9**, 959–963

610 Chin, A., Kyne, P.M., Walker, T.I. & McAuley, R.B. (2010) An integrated risk assessment for
611 climate change: analysing the vulnerability of sharks and rays on Australia's Great
612 Barrier Reef. *Global Change Biology*, **16**, 1936-1953.

613 Couch, J. (1841) *A history of the fishes of the British Islands. Volume 1*. J. Van Voorst, London.

- 614 Daufresne, M., Lengfellner, K. & Sommer, U. (2009) Global warming benefits the small in
615 aquatic ecosystems. *Proceedings of the National Academy of Sciences*, **106**, 12788–
616 12793.
- 617 Dillwyn, L.W. (1848) *Materials for a fauna and flora of Swansea and the neighbourhood*. D.
618 Rees.
- 619 Dulvy, N.K., Rogers, S.I., Jennings, S., Stelzenmuller, V., Dye, S.R. & Skjoldal, H.R. (2008)
620 Climate change and deepening of the North Sea fish assemblage: a biotic indicator of
621 warming seas. *Journal of Applied Ecology*, **45**, 1029-1039.
- 622 Dulvy, N.K., Pacoureau, N., Rigby, C.L., Pollom, R.A., Jabado, R.W., Ebert, D.A., Finucci, B.,
623 Pollock, C.M., Cheok, J. & Derrick, D.H. (2021) Overfishing drives over one-third of all
624 sharks and rays toward a global extinction crisis. *Current Biology*, **31**, 4773-4787. e8.
- 625 Ebert, D.A. & Dando, M. (2020) *Field Guide to Sharks, Rays & Chimaeras of Europe and the*
626 *Mediterranean*. Princeton University Press.
- 627 Ellis, J. (2007) Occurrence of pelagic stingray *Pteroplatytrygon violacea* (Bonaparte, 1832) in
628 the North Sea. *Journal of Fish Biology*, **71**, 933-937.
- 629 Engelhard, G.H., Lynam, C.P., Garcia-Carreras, B., Dolder, P.J. & Mackinson, S. (2015) Effort
630 reduction and the large fish indicator: spatial trends reveal positive impacts of
631 recent European fleet reduction schemes. *Environmental Conservation*, **42**, 227-236.
- 632 Froese, R. & Pauly, D. (2004) *FishBase. World Wide Web electronic publication*.
633 www.fishbase.org, version (09/2004).
- 634 Grant, M.I., Bicknell, A.W.J., Htut, T., Maung, A., Maung, T., Myo Myo, K., Rein, T., San, M.K.,
635 White, W.T., Ya, K.Z. & Mizrahi, M. (2022) Market surveys and social media provide
636 confirmation of the endangered giant freshwater whipray *Urogyrnus polylepsis* in
637 Myanmar. *J Fish Biol*, **101**, 302-307.
- 638 Haddaway, N.R. & Bayliss, H.R. (2015) Shades of grey: two forms of grey literature important
639 for reviews in conservation. *Biological Conservation*, **191**, 827-829.
- 640 Hammerschlag, N., McDonnell, L.H., Rider, M.J., Street, G.M., Hazen, E.L., Natanson, L.J.,
641 McCandless, C.T., Boudreau, M.R., Gallagher, A.J. & Pinsky, M.L. (2022) Ocean
642 warming alters the distributional range, migratory timing, and spatial protections of
643 an apex predator, the tiger shark (*Galeocerdo cuvier*). *Global Change Biology*, **28**,
644 1990-2005.
- 645 Heessen, H.J.L., Daan, N. & Ellis, J.R. (2016) *Fish atlas of the Celtic Sea, North Sea, and Baltic*
646 *Sea: Based on international research-vessel surveys*. KNNV.
- 647 Herdman, W. & Dawson, R. (1902) Fish and fisheries of the Irish Sea. In. London
- 648 Hiddink, J.G. & ter Hofstede, R. (2008) Climate induced increases in species richness of
649 marine fishes. *Global Change Biology*, **14**, 453–460.
- 650 Hiddink, J.G., Burrows, M.T. & Molinos, J.G. (2015) Temperature tracking by North Sea
651 benthic invertebrates in response to climate change. *Global Change Biology*, **21**,
652 117-129.
- 653 Hiddink, J.G., Shepperson, J., Bater, R., Goonesekera, D. & Dulvy, N.K. (2019) Near
654 disappearance of the Angelshark *Squatina squatina* over half a century of
655 observations. *Conservation Science and Practice*, **1**, e97.
- 656 Hyder, K., Weltersbach, M.S., Armstrong, M., Ferter, K., Townhill, B., Ahvonen, A.,
657 Arlinghaus, R., Baikov, A., Bellanger, M. & Birzaks, J. (2018) Recreational sea fishing
658 in Europe in a global context—participation rates, fishing effort, expenditure, and
659 implications for monitoring and assessment. *Fish and Fisheries*, **19**, 225-243.
- 660 Jabado, R.W., Chartrain, E., De Bruyne, G., Derrick, D., Dia, M., Diop, M., Doherty, P., Leurs,
661 G.H.L., Metcalfe, K., Pacoureau, N., Pires, J.D., Ratão, S., Seidu, I., Serena, F., Soares,
662 A.-L., Tamo, A., VanderWright, W.J. & Williams, A.B. (2021) *Dasyatis pastinaca*. The
663 IUCN Red List of Threatened Species 2021: e.T161453A124488102.

664 <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T161453A124488102.en>. Accessed
665 on 20 October 2023.

666 Jennings, S., Alvsvag, J., Cotter, A.J.R., Ehrich, S., Greenstreet, S.P.R., Jarre-Teichmann, A.,
667 Mergardt, N., Rijnsdorp, A.D. & Smedstad, O. (1999) Fishing effects in northeast
668 Atlantic shelf seas: patterns in fishing effort, diversity and community structure. III.
669 International trawling effort in the North Sea: an analysis of spatial and temporal
670 trends. *Fisheries Research*, **40**, 125-134.

671 Last, P., Naylor, G., Séret, B., White, W., de Carvalho, M. & Stehmann, M. (2016) *Rays of the*
672 *World*. CSIRO publishing.

673 Lewin, W.-C., Arlinghaus, R. & Mehner, T. (2006) Documented and potential biological
674 impacts of recreational fishing: insights for management and conservation. *Reviews*
675 *in Fisheries Science*, **14**, 305-367.

676 Martin, C., Vaz, S., Ellis, J., Coppin, F., Le Roy, D. & Carpentier, A. (2010) Spatio-temporal
677 patterns in demersal elasmobranchs from trawl surveys in the eastern English
678 Channel (1988–2008). *Marine Ecology Progress Series*, **417**, 211-228.

679 McPherson, J.M. & Myers, R.A. (2009) How to infer population trends in sparse data:
680 examples with opportunistic sighting records for great white sharks. *Diversity and*
681 *Distributions*, **15**, 880-890.

682 MMO (2020) Mapping recreational sea anglers in English waters. A report produced
683 for the Marine Management Organisation, MMO Project No: 1163, February 2020,. In, p.
684 129

685 Monkman, G.G., Kaiser, M.J. & Hyder, K. (2018) Text and data mining of social media to map
686 wildlife recreation activity. *Biological conservation*, **228**, 89-99.

687 Osgood, G.J., White, E.R. & Baum, J.K. (2021) Effects of climate-change-driven gradual and
688 acute temperature changes on shark and ray species. *Journal of Animal Ecology*,

689 Pacoureaux, N., Rigby, C.L., Kyne, P.M., Sherley, R.B., Winker, H., Carlson, J.K., Fordham, S.V.,
690 Barreto, R., Fernando, D., Francis, M.P., Jabado, R.W., Herman, K.B., Liu, K.M.,
691 Marshall, A.D., Pollom, R.A., Romanov, E.V., Simpfendorfer, C.A., Yin, J.S., Kindsvater,
692 H.K. & Dulvy, N.K. (2021) Half a century of global decline in oceanic sharks and rays.
693 *Nature*, **589**, 567-571.

694 Pennant, T. (1796) *The history of the parishes of Whiteford and Holywell*. B. and J. White.

695 Perry, A.L., Low, P.J., Ellis, J.R. & Reynolds, J.D. (2005) Climate Change and Distribution Shifts
696 in Marine Fishes. *Science*, **308**, 1912-1915.

697 Pritchard, E.K. (2018) A tool for finding rare marine species. *Science*, **360**, 1180-1182.

698 Quero, J.C. (1998) Changes in the Euro-Atlantic fish species composition resulting from
699 fishing and ocean warming. *Italian Journal of Zoology*, **65**, 493-499.

700 Raye, L. (2018) Robert Sibbald's Scotia Illustrata (1684): A faunal baseline for Britain. *Notes*
701 *and Records: the Royal Society Journal of the History of Science*, **72**, 383-405.

702 Richardson, E.A., Kaiser, M.J., Edwards-Jones, G. & Ramsay, K. (2006) Trends in sea anglers'
703 catches of trophy fish in relation to stock size. *Fisheries Research*, **82**, 253-262.

704 Rindorf, A., Gislason, H., Burns, F., Ellis, J.R. & Reid, D. (2020) Are fish sensitive to trawling
705 recovering in the Northeast Atlantic? *Journal of Applied Ecology*, **57**, 1936-1947.

706 Rogers, S. & Ellis, J. (2000) Changes in the demersal fish assemblages of British coastal
707 waters during the 20th century. *ICES Journal of Marine Science*, **57**, 866-881.

708 Schlaff, A.M., Heupel, M.R. & Simpfendorfer, C.A. (2014) Influence of environmental factors
709 on shark and ray movement, behaviour and habitat use: a review. *Reviews in Fish*
710 *Biology and Fisheries*, **24**, 1089-1103.

711 Serena, F., Mancusi, C., Morey, G. & Ellis, J.R. (2015) *Dasyatis pastinaca*. The IUCN Red List of
712 Threatened Species 2015: e.T161453A48933979.

713 Sguotti, C., Lynam, C.P., García-Carreras, B., Ellis, J.R. & Engelhard, G.H. (2016) Distribution of
714 skates and sharks in the North Sea: 112 years of change. *Global change biology*, **22**,
715 2729-2743.

716 Shackell, N.L., Frank, K.T., Fisher, J.A.D., Petrie, B. & Leggett, W.C. (2010) Decline in top
717 predator body size and changing climate alter trophic structure in an oceanic
718 ecosystem. *Proceedings of the Royal Society B: Biological Sciences*, **277**, 1353.

719 Sibbald, R. (1684) *Scotia Illustrata. Tomus secundus de animalibus Scotiae tam feris quam*
720 *domesticis et de mineralibus metallis et marinis Scotiae*. Ex Officinâ Typographicâ
721 Jacobi Kniblio, Josuae Solingensis & Johannis Colmarii, Sumptibus Auctoris,
722 Edinburgh.

723 Swetnam, T.W., Allen, C.D. & Betancourt, J.L. (1999) Applied historical ecology: using the
724 past to manage for the future. *Ecological applications*, **9**, 1189-1206.

725 Thurstan, R., McClenachan, L., Crowder, L., Drew, J., Kittinger, J., Levin, P., Roberts, C. &
726 Pandolfi, J. (2015) Filling historical data gaps to foster solutions in marine
727 conservation. *Ocean & Coastal Management*, **115**, 31-40.

728 VanDerWal, J., Murphy, H.T., Kutt, A.S., Perkins, G.C., Bateman, B.L., Perry, J.J. & Reside, A.E.
729 (2013) Focus on poleward shifts in species' distribution underestimates the
730 fingerprint of climate change. *Nature Climate Change*, **3**, 239-243.

731 Walker, P.A. & Hislop, J.R.G. (1998) Sensitive skates or resilient rays? Spatial and temporal
732 shifts in ray species composition in the central and north-western North Sea
733 between 1930 and the present day. *ICES Journal of Marine Science*, **55**, 392-402.

734 Walker, T.I., Day, R.W., Awruch, C.A., Bell, J.D., Braccini, J.M., Dapp, D.R., Finotto, L., Frick,
735 L.H., Garcés-García, K.C. & Guida, L. (2021) Ecological vulnerability of the
736 chondrichthyan fauna of southern Australia to the stressors of climate change,
737 fishing and other anthropogenic hazards. *Fish and Fisheries*,

738 Walls, R.H. & Dulvy, N.K. (2020) Eliminating the dark matter of Data Deficiency by predicting
739 the conservation status of Northeast Atlantic and Mediterranean Sea sharks and
740 rays. *Biological Conservation*, **246**, 108459.

741 Walls, R.H. & Dulvy, N.K. (2021) Tracking the rising extinction risk of sharks and rays in the
742 Northeast Atlantic Ocean and Mediterranean Sea. *Scientific reports*, **11**, 1-15.

743 Wheeler, A. & Blacker, R. (1969) Rare and little-known fishes in British seas in 1966 and
744 1967. *Journal of Fish Biology*, **1**, 311-331.

745 Wheeler, A. & Blacker, R. (1972) Rare and little-known fishes in British seas in 1968 and
746 1969. *Journal of Fish Biology*, **4**, 141-170.

747 Wheeler, A., Blacker, R.W. & Pirie, S.F. (1975) Rare and little-known fishes in British seas in
748 1970 and 1971. *Journal of Fish Biology*, **7**, 183-201.

749 Wheeler, A.C. (1969) *The fishes of the British Isles and north-west Europe*. Macmillan.

750 Wolff, W.J. (2000) The south-eastern North Sea: losses of vertebrate fauna during the past
751 2000 years. *Biological Conservation*, **95**, 209-217.

752 Yarrell, W. (1859) *A history of British fishes*. John Van Voorst.


753

754 Table 1. Summary statistics of the eight sources of common stingray record used, their
 755 temporal range, the number of reports (of one or more stingrays at a particular time and
 756 location) and records (total number of individuals reported) in Great Britain

Type	First record	Last record	Reports	Recorded numbers
SeaAngler magazine	1952	2020	155	155
Online forum	1905	2020	75	79
Other	1857	2015	36	161
Book	1838	1987	30	50
News article	1980	2020	26	26
Social media	2011	2020	17	17
Scientific paper	1989	2003	9	10
NFSA	1970	1989	8	20
Total			356	518

757


758



759

760 Figure 1. A) Temporal trends in the number of common stingray reports (one or more stingrays
 761 at one time and place) around Great Britain obtained from all data sources as total records
 762 per 5-year period from 1830 to 2020 (see Table 1). B) *SeaAngler* magazine readership (online
 763 and paper subscribers, left axis, line) and the number of common stingray (black points) and
 764 angelshark (blue points) reports in *SeaAngler* (right axis, per year) over the period from 1974
 765 to 2020. C) The number of common stingray (black) and angelshark (blue) reports per 1,000
 766 *SeaAngler* readers. The lines are GAMs fitted through the data (black: common stingray,
 767 effective df=2.91, F=5.236, p=0.0033, deviance explained = 29.9%, Δ AIC with linear model = -
 768 7.87, light blue: angelshark, effective df=2.79, F=14.32 p<0.0001, Δ AIC with linear model = -
 769 7.87, deviance explained = 49.8%). Note that the x-axes for A) covers a much longer period.


770



771

772 Figure 2. (A) The distribution of common stingray records around Great Britain from all sources
 773 based on reconstructed positions, showing degrees of latitude and longitude. All records are
 774 plotted in transparent grey scale. This results in the most persistent observation locations
 775 being represented by more intense shades and individual isolated locations being represented
 776 by transparent points. (B–H) Distribution by 25-year period of common stingray, over nearly
 777 two centuries. The 2000 onwards observation period is inevitably truncated and comprised of
 778 20 years of records to 2020. The black crosses indicates the 95% confidence intervals around
 779 the mean latitude and longitude (centre of the cross) of all records for each period.

780



781

782 Figure 3. Temporal trends in the common stingray distribution and size around Great Britain.

783 (A) The maximum (blue, 0.95th quantile), median (black) and minimum (red, 0.05th quantile)

784 latitude of common stingray records per 5-year period. The minimum latitude of the study

785 area is ~50°N. (B) Number of common stingray records by 4-degree latitude band per 25-year

786 period. Red = 48 to 52°, black = 52 to 56°, blue = 56 to 60°. (C) The fraction of the total number


787 of records from SE England. (D) The median reconstructed wet weight of stingrays (± 95%

788 confidence intervals) per 10-year period. Points are only plotted for periods where the

789 number of records n > 5. The number of observations, N, is given below or above the data

790 points. The indicated years are the years at the start of a time period. X-axes differ between plots.


791



792
 793
 794
 795
 796
 797

Figure 4. Estimates of the magnitude of changes in abundance of the common stingray in the UK using opportunistic data from multiple data sources, with 95% confidence bounds, between any chosen reference year and 2020, based on all reported observations. Different lines represent different assumed changes in observation effort.

798 **Supplementary material**




799

800 Figure S1. Angelshark distribution from magazine records only, extracted using the same
801 methodology as for the common stingray. All records are plotted in transparent blue grey
802 scale. This results in the most persistent observation locations being represented by more
803 intense shades and individual isolated locations being represented by transparent
804 dots. (B–C) Distribution by 25-year period of angelshark. The 2000 onwards
805 observation period is inevitably truncated and comprised of 20 years of records to 2020. The
806 black crosses indicates the 95% confidence intervals around the mean latitude and longitude
807 (centre of the cross) of all records for each period.

808

809



810

811 Figure S2. Month in which common stingray was recorded from 1970 onwards. The line
812 indicates a binomial regression (because values are bounded between 1 and 12, converted
813 to 0 to 1 for model fitting) with confidence intervals (GLM, $z_{1,265} = -1.318$, $p = 0.187$). There
814 was no significant change in the length of the sighting period over time ($t_{47}=0.505$, $p=0.616$).

815

816 Table S1. Information sources from which stingray records were compiled.

Source	Link	Source category
Chelmsford Weekly News	www.chelmsfordweeklynews.co.uk	News article
Daily Echo	www.dailyecho.co.uk	News article
East Anglian Daily Times	www.eadt.co.uk	News article
Express	www.express.co.uk	News article
Gazette News	www.gazette-news.co.uk	News article
ITV	www.itv.com	News article
Kent Online	www.kentonline.co.uk	News article
Planet Sea Fishing	www.planetseafishing.com	Forum
The News - Portsmouth	www.portsmouth.co.uk	News article
Press Reader	www.pressreader.com	News article
Veals Mail Order	www.veals.co.uk	News article
Wales Online	www.walesonline.co.uk	News article
West Sussex Today	www.westsussextoday.co.uk	News article
Archive	https://archive.org	Other
Biodiversity Library	www.biodiversitylibrary.org	Other
Angling Addicts	www.anglingaddicts.co.uk	Forum
Bristol Channel Federation of Sea Anglers (BCFSA)	www.bristolchannelfishing.com	Other
Charter Boats UK	www.charterboats-uk.co.uk	Other
British Marine Life Study Society	www.glaucus.org.uk	Other
World Sea Fishing	www.worldseafishing.com	Forum
Anchorman Charters	www.anchormancharters.co.uk	Other
The Database of Trawl Surveys	https://datras.ices.dk	Other
Fish UK	www.fish-uk.com	Forum
Freshwater Habitats	https://freshwaterhabitats.org.uk	Forum
National Biodiversity Network (NBN)	https://records.nbnatlas.org	Other

Natural History Museum (NHM)	https://data.nhm.ac.uk	Other
Norfolk Fishing	https://norfolkfishing.com	Forum
South-West Federation of Sea Angling (SWFSA)	https://swfsa.co.uk	Other
Underwater Fishing	www.underwaterfishing.co.uk	Forum
Wales Federation of Sea Anglers (WFSA)	https://www.wfsa.org.uk/	Other
Facebook (i.e., Natur Dyfi, BluePlanetSoc)	www.facebook.com	Social media
Tumblr	www.tumblr.com	Social media
Youtube	www.youtube.com	Social media
Instagram accounts (i.e., @southwalesfishing)	www.instagram.com	Social media
First Nature	www.first-nature.com	Forum
The Archive for Marine Species and Habitats Data (DASSH)	https://www.dassh.ac.uk/	Other

817

818

819