




*Foras na Mara*  
*Marine Institute*



## **Porcupine Bank *Nephrops* Grounds (FU16) 2023 UWTV Survey Report and catch scenarios for 2024**

**Mikel Aristegui<sup>1</sup>, Gráinne Ryan<sup>1</sup>, James Fahy<sup>1</sup>, Laurence Manning<sup>1</sup>, Neve McCann<sup>1</sup>, Kirsty Woodcock<sup>2</sup>, Grant Course<sup>3</sup>, Rita Meireles de Castro<sup>4</sup>, Natasha Murphy<sup>1</sup>**

<sup>1</sup> Fisheries Ecosystems Advisory Services, Marine Institute, Rinville, Oranmore, Galway, Ireland.

<sup>2</sup> Joint Nature Conservation Committee, Monkstone House, Peterborough, UK.

<sup>3</sup> SeaScope Fisheries Research, Powfoot, Annan, Dumfries & Galloway, Scotland, UK.

<sup>4</sup> Marine and Freshwater Research Centre, Atlantic Technological University, Galway, H91 T8NW, Ireland.



© Marine Institute 2023



Ireland's European Structural and  
Investment Funds Programmes  
2014-2020

Co-funded by the Irish Government  
and the European Union



**An Roinn Talmhaíochta,  
Bia agus Mara**  
Department of Agriculture,  
Food and the Marine



EUROPEAN MARITIME AND  
FISHERIES FUND

**Suggested citation:**

Aristegui, M., Ryan, G., Fahy, J., Manning, L., McCann, N., Woodcock, K., Course, G., Meireles de Castro, R. and Murphy, N. 2023. Porcupine Bank *Nephrops* Grounds (FU16) 2023 UWTV Survey Report and catch scenarios for 2024. Marine Institute UWTV Survey report.

**Acknowledgments:**

We would like to express our sincere thanks and gratitude to the Master and crew of the RV *Tom Crean*. Thanks also to the P&O Maritime IT & Instrumentation Technician Anthony English who maintained the UWTV system throughout the survey. Thanks to Aodhán Fitzgerald, and Frank Armstrong (RVOPs) and Dave Tully and John Enright (FEAS) at the Marine Institute for organising survey logistics. Thanks to Gordon Furey, Damian Crean, Barry Kavanagh, John Barry (P&O Maritime) for shore side support. Thanks to Chris Allsop (OCIS/FEAS) for supporting the UWTV database management.

**Photography credit:**

Photograph used on the cover has been provided by © Mikel Aristegui Ezquibela.

**Disclaimer:**

Although every effort has been made to ensure the accuracy of the material contained in this publication, complete accuracy cannot be guaranteed. Neither the Marine Institute nor the author accepts any responsibility whatsoever for loss or damage occasioned, or claimed to have been occasioned, in part or in full as a consequence of any person acting or refraining from acting, as a result of a matter contained in this publication. All or part of this publication may be reproduced without further permission, provided the source is acknowledged.

## Abstract

This report provides the results of the eleventh underwater television on the 'Porcupine Bank *Nephrops* grounds' ICES assessment area; Functional Unit 16. The survey was multi-disciplinary in nature collecting UWTV and other ecosystem data. In total 71 UWTV stations were successfully completed (100% of the planned stations) in a randomised 6 nautical mile isometric grid covering the full spatial extent of the stock. The mean burrow density observed in 2023, adjusted for edge effect, was 0.27 burrows/m<sup>2</sup>. The final krigged abundance estimate was 2002 million burrows with a CV of 3% and an estimated stock area of 7,130 km<sup>2</sup>. The 2023 abundance estimate was 47% higher than in 2022. Using the 2023 estimate of abundance and updated stock data imply that catches in 2024 should be between 3677 and 4560 tonnes, according to the EU MAP and ICES MSY approach (assuming that all catch is landed). Four species of sea-pen (*Virgularia mirabilis*, *Funiculina quadrangularis*, *Pennatula phosphorea* and the deepwater sea-pen *Kophobelemnion stelliferum*) were observed during the survey. Trawl marks were also observed on 20% of the stations surveyed.

Key words: *Nephrops norvegicus*, Porcupine Bank, stock assessment, geostatistics, underwater television (UWTV), sea-pens, benthos.

## 1. Introduction

*Nephrops norvegicus* are common around the Irish coast occurring in geographically distinct sandy/muddy areas where the sediment is suitable for them to construct their burrows. The *Nephrops* fishery in ICES Subarea 7 is extremely valuable with Irish landings in 2022 worth around €69 million at first sale. The *Nephrops* fishery on the Porcupine Bank takes place on a large area approximately 7130 km<sup>2</sup> of complex muddy habitat between depths of 330 to 580m. The fishery typically yields very large individual *Nephrops* that attain very high market prices relative to other fisheries around Ireland. International landings from the fishery peaked in the early 1980s around 4000 tonnes, declining since then with some fluctuations (ICES, 2022). The total estimated Irish landings in 2022 were 1649 tonnes.

In the past sustainability of the Porcupine Bank *Nephrops* stock has been a major concern. Consequently, a spatio-temporal closed area was developed and proposed by the North Western Waters Advisory Council (NWWAC) and implemented between 1<sup>st</sup> June and 31<sup>st</sup> July in 2010-2012. Since 2013 Irish fishing has closed for one month from 1<sup>st</sup> to 31<sup>st</sup> of May. Since 2011 a functional unit catch limit (actually landings) has also been in place as part of the TAC regulation (ICES, 2014). These measures were introduced due to negative trends in the various indicators used to assess the stock and ICES advised for a closure of the fishery in 2009 and 2010. The stock situation is known to have improved since 2010 following good recruitment. Scientific information for this area improved with the introduction of a dedicated Irish fisheries-science partnership trawl survey between 2010 and 2012 and the provision of commercial grade data by the Irish fishing industry since 2010 (ICES, 2014).

*Nephrops* spend a great deal of time in their burrows and their emergence behaviour is influenced by many factors; time of year, light intensity and tidal strength. Assessment methodologies, based on underwater television surveys, have been developed by ICES to provide a fishery independent estimate of stock size, exploitation status and catch advice (ICES, 2009 & 2013). Since 2012, UWTV surveys have been used to assess and provide catch scenarios and advice for this stock (ICES, 2013).

This was the eleventh UWTV survey of the Porcupine Bank *Nephrops* grounds (FU16). The 2023 survey was multi-disciplinary in nature. The specific objectives are listed below:

1. To obtain 2023 quality assured estimates of *Nephrops* burrow densities from a randomised isometric grid of UWTV stations at 6 nautical mile spacing over the known spatial and bathymetric distribution of the stock (Figure 1).
2. To collect ancillary information from the UWTV footage collected at each station such as the presence of sea-pens, other macrobenthos and fish species and trawl marks on the sea bed.
3. To collect relative abundance and distribution of marine mammals in the area by a Marine Mammal Observer (MMO).

This report details the final UWTV results of the 2023 survey and also documents other data collected during the survey. Operational survey details are available in the form of a survey narrative from the scientist in charge (MAE). The 2023 abundance data are used to generate catch options for 2024 in line with the recommendations and procedures outlined at the

2013 ICES benchmark (ICES, 2013) and in the stock annex (ICES, 2013) and using the  $F_{MSY}$  reference points proposed by FMSYREF4 (ICES, 2016).

## 2. Material and methods

A randomised isometric grid of stations at 6 nautical mile or 11.1km intervals was planned for the area. The boundary used to delineate the edge of the ground was based on VMS (Vessel Monitoring System) data of fishing activity between 2006-2011 targeting *Nephrops* (Lordan *et al.* 2012). The grid spacing was determined based on a time constraint of getting the survey completed within a time window of around 5-6 days. This resulted in 71 planned stations and were generated using the “spsamp1” function in the “sp” package (Pebesma & Bivand, 2005) of “R” (R Core Team, 2022). Data on bathymetry and backscatter were also available from the Irish National Seabed Survey and INFOMAR project (<http://www.infomar.ie/>). The stations ranged from 350m to 582m in depth with an average depth of 459m (Figure 1). Survey timing has generally been standardised to June each year. In 2015 the national research vessel broke down prior to the survey and the survey was not carried out despite several attempts to get to this ground. In 2023, the FU16 survey was carried out between the 30<sup>th</sup> of May and the 10<sup>th</sup> of June on the RV. *Tom Crean* during the TC23012 Marine Institute survey.

In 2023 all the stations were surveyed using the High Definition (HD) camera system. These stations recorded at 12 frames per second with a camera angle of 75°. The digital images were stored on a server and were reviewed during the survey through an in-house developed Image Annotation R Shiny app (Aristegui, 2020). This app allows each reviewer to annotate burrows for each randomly assigned station in an efficient manner.

The operational protocols used were those reviewed by the Workshop on the use of UWTV surveys for determining abundance in *Nephrops* stocks throughout European waters (ICES, 2007) and follow internationally agreed standards as recommended in the Manual for the *Nephrops* Underwater TV Surveys (TIMES) (Dobby *et al.*, 2021). These protocols are employed on other UWTV surveys in Irish waters and can be summarised as follows: at each station the UWTV sledge was deployed. Once stable on the seabed a 10 minute tow was recorded. Time referenced high definition image data were collected with a field of view (FOV) of 1.02 metre. Vessel position (DGPS) and position of sledge (using a USBL transponder) were recorded every 2 seconds. The navigational data were quality controlled using an “R” script developed by the Marine Institute (ICES, 2009). The USBL Sonardyne navigational data were used to calculate distance over ground for 96% all of stations. Due to technical problems however, corrected ship navigation data had to be used for three stations.

In line with recommendations of the Workshop on *Nephrops* Burrow Counting (WKNEPS), all scientists were trained/re-familiarised using HD image data for training material and a reference set (ICES, 2018). All counts were conducted by four trained scientists independent of each other during the survey. The numbers of *Nephrops* burrow systems were counted, where multiple burrows in close proximity which appeared to be part of a single system were counted as one. *Nephrops* activity in and out of burrows were counted and recorded for each station.

Following the recommendation of ICES Study Group on *Nephrops* Surveys (SGNEPS) the time for verified recounts was seven minutes (ICES, 2009) but this was increased to nine minutes for the Porcupine until 2022. This was because at the lower densities observed the relative scale of variation between minutes was higher than typical compared to *other Nephrops* grounds. Therefore, counting more minutes resulted in a more stable mean density estimates for each station. Due to a burrow density increase at the Porcupine in the recent years, in 2023 seven minutes were counted in each station, following the SGNEPS recommendation.

Presence / absence for the occurrence of trawl marks, fish and other species were also recorded at each station. Presence / absence of sea-pen species were recorded to fulfil an OSPAR Special Request (ICES 2011).

Finally, if there was any time during each minute where counting was not possible due to sediment clouds or other reasons, this time was recorded and removed from the distance over ground calculations. The “R” quality control tool allowed for the data quality of navigation, speed, visual clarity and consistency in counts to be checked (an example is given in Figure 2).

In 2023 survey count data were screened to check for any unusual discrepancies using Lin’s Concordance Correlation Coefficient (CCC) with a threshold of 0.6. Lin’s CCC (Lin, 1989) measures the ability of counters to reproduce each other’s counts on a scale of 1 to -1, where 1 is perfect concordance (i.e. a pairwise plot will have all points lying along the 1:1 line; a value of -1 would be generated by all points lying on the -1:1 line and a value of 0 indicates no correspondence at all). Lin’s CCC quality control plots of survey count data for stations 50 to 52 are shown in Figure 3. When the count data fell below the threshold of 0.6 a third review was carried out. The paired count data that passed the Lin’s CCC threshold was used in the analysis. When the paired counts did not pass the threshold an average of the three reviewers was deemed appropriate to use in the analysis.

Mean density was calculated by dividing the total number of burrow systems by the survey area observed. In 2023 the field of view of the camera at the bottom of the screen was estimated by extrapolation at 1.02m assuming that the sledge was flat on the seabed (i.e. no sinking).

From 2012 to 2014 to account for the spatial co-variance and other spatial structuring a geo-statistical analysis of the mean and variance was carried out using SURFER Version 10.7.972 and the krigged estimation variance or coefficient of variance (CV) was carried out using the EVA: Estimation VAriance software (Petitgas and Lafont, 1997).

Since 2016, the geostatistical analysis has been carried out using “RGeostats” package Version 11.1.1 (Renard D., *et al*, 2015) in “R” and is available as a separate R markdown document. The same basic steps were carried out as in previous years; construction of experimental variogram, and a model variogram ( $h$ ) produced with an exponential model, a krigged grid file was created using all data points as neighbours, the same boundary used to

estimate the domain area, the mean density, total burrow abundance and survey precision calculated.

For each UWTV station a CTD was logged for the duration of each tow using a sled mounted and calibrated RBR*brevio* (rated to 750m depth). This data will be processed at a later stage.

For the second time in an UWTV survey carried out by the Marine Institute in FU16, a marine mammal survey was conducted at the same time. This survey was carried out by a dedicated MMO on board who used a standard single platform line-transect methodology to collect data of relative abundance and distribution of marine mammals in the Porcupine Bank. This work was funded through the Marine Biodiversity Scheme, carried out under Ireland's Operational Programme, co-funded by the European Maritime and Fisheries Fund (EMFF) and by the Irish Government. Data have been processed by the MMO and fully documented in a separate report (Meireles de Castro, 2023) available through the Marine Institute Biodiversity portal.

### 3. Results

In 2023, 71 stations were completed successfully on the Porcupine Bank (Figure 1). Figure 4 shows bubble plots of the variability between minutes for each station. At the lower densities observed the relative scale of variation between minutes was higher than typical in other areas, reflecting the patchy and low density.

A combined violin and box plot of the observed burrow densities is presented in Figure 5. This shows that median, mean burrow densities and inter-quartile ranges were similar in most years, but have increased since 2021. The mean burrow density observed in 2023, adjusted<sup>1</sup> for edge effect, was 0.27 burrows/m<sup>2</sup>.

The final modelled density surfaces from 2012 to 2014 and 2016 to 2023 are shown as a heat map in Figure 6. The 2023 burrow surface shows a general increase in burrow densities in all the ground. The abundance estimate derived from the kriged burrow surfaces (and adjusted for edge effect) increased by 47% from 1363 million burrows in 2022 to 2002 million in 2023 (Figure 7 and Table 1) with an estimated area of the ground or domain area of 7,130 km<sup>2</sup>. The estimation CV on the abundance was around 3.1% in 2023.

Trawl marks were observed at 14 out of the 71 surveyed stations (20%). The distribution of the various sea-pen species observed on the UWTV footage is shown in Figure 8. Three sea-pen species occur in the deep mud habitats around the coastal British Isles and are usually observed in the UWTV surveys: *Virgularia mirabilis*, *Pennatula phosphorea* and *Funiculina quadrangularis* (Hughes, 1998). The distribution over time of a fourth sea-pen, *Kophobelemnion stelliferum*, is shown in Figure 9 as part of a species review in the UWTV database. This species has been recorded at the Porcupine Seabight in depths to 1600 m (Rice *et al.*, 1992).

---

<sup>1</sup> Note the “adjusted” density estimates in this report are adjusted by dividing by 1.26 to take account of edge effect over estimation of area viewed during UWTV transects (see Campbell *et al.*, 2009).

The UWTV abundance data together with data from the fishery; landings, removals in number, and mean weight in the landings are shown (Table 2). The basis to the catch options table is given in Table 3. The catch and landings options calculated using the 2023 survey abundance estimate and updated stock data, imply that catches in 2024 should be between 3677 and 4560 tonnes, according to ICES MSY approach (assuming that all catch is landed) (Table 4).

During the TC23012 UWTV survey (which covered FU16, FU17 and part of FU19), the MMO recorded a total of 70 common dolphins (17 sightings), 15 bottlenose dolphins (one sighting), five unidentified dolphins and one unidentified cetacean (Meireles de Castro, 2023).

#### 4. Discussion

This was the eleventh systematic UWTV *Nephrops* survey of the Porcupine Bank. The distance from shore (~ 120 nautical miles), exposed nature of the area, the significant water depths involved (350m to 582m) and relatively large size of the area (>7100km<sup>2</sup>) presents significant logistical, technical and survey design challenges. The Marine Institutes carries out UWTV surveys in three pre-planned survey legs (of 12, 11 and 10 days). Priority was given to the Porcupine Bank which was successfully completed on the first leg in 2023, together with FU17. The visibility and footage quality was normally excellent, burrow morphology and size were similar to other areas and burrow identification was relatively easy.

Due to the large increase in stock abundance in 2023 (>40% increase compared to 2022), a review process was implemented according to the ICES UWTV Survey Protocols (Dobby *et al.*, 2021). 14 stations were randomly selected (20% of the total number of stations) and reviewed independently by trained burrow counters. The review of these stations corroborated the accuracy of the original counts and confirmed the increase in stock abundance compared to 2022.

The survey design, with a randomised 6 nautical mile isometric grid and fixed ground boundary, was the same as that used previously (Lordan, *et al.*, 2012). The total abundance estimate has increased. Catch scenarios for 2024 have been calculated using updated landings mean weight data from WGCSE (ICES, 2023). These data are usually estimated using the graded landings information for the fishing industry and sampling data carried out at sea by scientific observer. However, since 2020, due to low sampling levels caused by Covid-19 restrictions, sampling was adapted and mean weights estimated from catch samples. An average over the last three years (2020-2022) is used to account for the mean weight in the landings estimated in the recent past. The resulting catch advice for 2024 fishing at the  $F_{MSY}$  point is an increase of 20%), due to the higher abundance estimate compared to last year. Carrying out annual UWTV surveys to generate catch advice, while challenging, should be continued annually given the limited number of UWTV observations to date and evolving knowledge base on the spatial and temporal dynamics of this stock.

In addition to estimating *Nephrops* stock abundance UWTV surveys can be used to monitor the presence of certain benthic fauna (ICES, 2011). Sea-pens and burrowing megafauna



communities have been included in the OSPAR list of threatened and/or declining species and habitats (OSPAR, 2010). As previously observed, all three sea-pen species which occur on mud habitat around Ireland are found on the Porcupine Bank, and also the deepwater sea-pen *Kophobelemnion stelliferum*. Data of the four sea-pen species from FU16 are submitted as part of an annual ICES data call to Working Group on Deep-water Ecology (WGDEC). This dataset provides information regarding the impact of fisheries on components of the ecosystem. The occurrence of *F. quadrangularis* in particular is significant owing to its particularly vulnerability to trawl mortality. *F. quadrangularis* is largely absent from other *Nephrops* grounds around Ireland, although there are catches on groundfish surveys in areas where *Nephrops* are not commercially fished (Power and Lordan, 2012). The majority of the Porcupine Bank is fished at least once annually based on the methods described in Gerritsen, *et al.* (2013). The observation that 20% of stations showed some trawl marks is lower than previous years.

## 5. References

- Aristegui, M. 2020. Image annotation R Shiny app. Marine Institute. <http://doi.org/d24n>
- Campbell, N., Dobby, H., and Bailey, N. 2009. Investigating and mitigating uncertainties in the assessment of Scottish *Nephrops norvegicus* populations using simulated underwater television data. ICES Journal of Marine Science 66: 646–655. doi: 10.1093/icesjms/fsp046.
- Dobby, H., Doyle, J., Jónasson, J., Jonsson, P., Leocádio, A., Lordan, C., Weetman, A., and Wieland, K. 2021. ICES Survey Protocols – Manual for *Nephrops* underwater TV surveys, coordinated under ICES Working Group on *Nephrops* Surveys (WGNEPS). ICES Techniques in Marine Environmental Sciences Vol. 65. 44 pp. <https://doi.org/10.17895/ices.pub.8014>.
- EU. 2019. Regulation (EU) 2019/472 of the European Parliament and of the Council of 19 March 2019 establishing a multiannual plan for stocks fished in the Western Waters and adjacent waters, and for fisheries exploiting those stocks, amending Regulations (EU) 2016/1139 and (EU) 2018/973, and repealing Council Regulations (EC) No 811/2004, (EC) No 2166/2005, (EC) No 388/2006, (EC) No 509/2007 and (EC) No 1300/2008. Official Journal of the European Union, L 83: 117. <http://data.europa.eu/eli/reg/2019/472/oj>.
- Gerritsen, H.D., Minto, C., and Lordan, C. 2013. How much of the seabed is impacted by mobile fishing gear? Absolute estimates from Vessel Monitoring System (VMS) point data. ICES Journal of Marine Science. doi: 0.1093/icesjms/fst017.
- Hughes, D. J. 1998. Sea Pens and Burrowing Megafauna. An overview of dynamics and sensitivity characteristics for conservation management of marine SACs. Scottish Association for Marine Science.
- ICES. 2007. Report of the Workshop on the use of UWTV surveys for determining abundance in *Nephrops* stocks throughout European waters (WKNEPHTV). ICES CM: 2007/ACFM: 14 Ref: LRC, PGCCDBS.
- ICES. 2009. Report of the Study Group on *Nephrops* Surveys (SGNEPS). ICES CM 2009/LRC: 15. Ref: TGISUR.
- ICES. 2011. Protocols for assessing the status of sea-pen and burrowing megafauna communities. ICES Advice 2011, Book 1, 2011.
- ICES. 2013. Report of the Benchmark Workshop on *Nephrops* Stocks (WKNEPH), 25 February–1 March 2013, Lysekil, Sweden. ICES CM 2013/ACOM:45. 230 pp.
- ICES. 2014. Report of the Working Group for Celtic Seas Ecoregion (WGCSE). ICES CM: 2014/ACOM:12.
- ICES. 2016. EU request to ICES to provide FMSY ranges for selected stocks in ICES subareas 5 to 10. In Report of the ICES Advisory Committee, 2016. ICES Advice 2016, Book 5, Section 5.2.3.1.
- ICES. 2018. Report of the Workshop on *Nephrops* burrow counting (WKNEPS). 2-5 October 2018, Aberdeen, UK. ICES CM 2018/ EOSG:25.47 pp.
- ICES. 2023. Report of the Working Group for Celtic Seas Ecoregion (WGCSE). ICES (in draft).

- Lin, L. I. 1989. A Concordance Correlation Coefficient to Evaluate Reproducibility. *Biometrics* 45, 255-268.
- Lordan, C., Doyle, J., Dobby, H., Hehir, I. Fee, D., Allsop, C. and O'Neil, R. 2012. Porcupine Bank *Nephrops* Grounds (FU16) 2012 UWTV Survey Report and catch options for 2013. Marine Institute UWTV Survey report. <http://hdl.handle.net/10793/832>
- Meireles de Castro, R. 2023. Underwater Television Survey; Marine Mammal Observer Report.
- OSPAR. 2010. Background Document for Seapen and Burrowing megafauna communities. OSPAR Commission 2010, London. Publication number: 481/2010
- Petitgas, P., and Lafont, T. 1997. EVA (Estimation VAriance). A geostatistical software on IBM-PC for structure characterization and variance computation. Version 2.
- Power, J., and Lordan, C. 2012. A review of the effects of bottom trawling on soft sediments; sea pens and burrowing megafauna biotope complexes. Marine Institute internal report Version 1, 30 July 2012.
- Renard D., Bez N., Desassis N., Beucher H., Ors F. and Laporte F. 2015. RGeostats: The Geostatistical package [version:11.0.3]. MINES ParisTech. Free download from: <http://cg.ensmp.fr/rgeostats>
- Rice A.L., Tyler P.A. and Paterson G.J.L. 1992. The Pennatulid *Kophobelemnon stelliferum* (Cnidaria:Octocorallia) in the Porcupine Seabight (north-east Atlantic Ocean). *Journal of the Marine Biological Association of the UK* 72(02):417 - 434 · May 1992.
- R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>




Figure 1: FU16 UWTV 2023. UWTV map of station positions and bathymetry (source: INFOMAR\*). Red line indicates survey ground polygon.

\*INFOMAR is the Department of Communications, Climate Action and Environment (DCCAE) funded national seabed mapping programme, jointly managed and delivered by Geological Survey Ireland and Marine Institute.




Figure 2: FU16 UWTV 2023. UWTV example quality control plot for the navigational and recount data.




Figure 3: FU16 UWTV 2023. Lin's CCC quality control plot of count data for stations 50 to 52 from the 2023 survey.




Figure 4: FU16 UWTV 2023. UWTV quality control plot showing variability between minutes for each UWTV station




Figure 5: FU16 UWTV 2023. Violin and box plot of adjusted burrow density distributions by year from 2012-2023. The blue line indicates the mean density over time. The horizontal black line represents the median, white box is the inter quartile range, the black vertical line

is the range and the black dots are outliers. No UWTV survey in 2015.




Figure 6: FU16 UWTV 2023. Heat map of *Nephrops* burrow density observations from 2012 to 2023.




Figure 7: FU16 UWTV 2023. Time series of total abundance estimates for FU16 and 95% confidence intervals. No UWTV survey in 2015.






Figure 8: FU16 UWTV 2023. Stations where sea-pens *Virgularia mirabilis* (VAM), *Funiculina quadrangularis* (FAQ), *Pennatula phosphorea* (PNP) and *Kophobelemnion stelliferum* (KOP) were identified from video footage. Coloured circles denote presence in the TV station and empty circles denote TV stations with no sea-pen observations.




Figure 9: FU16 UWTV 2023. The presence/absence distribution of the deep water sea-pen species *Kophobelemnion stelliferum* (KOP) observed on the video footage from 2012 to 2023 (no survey in 2015). Coloured circles denote presence in the TV station and empty circles denote absence.

Table 1: FU16 UWTV 2023. Summary of univariate statistics and geostatistics for the burrow density estimates (bias corrected) on the Porcupine Bank UWTV survey in 2012-2023. No TV survey in 2015.

Year	Univariate Statistics					Geostatistics		
	Number of Observations	Min	Max	Mean	Standard Deviation	Adjusted abundance estimate (millions)	Domain area (km <sup>2</sup> )	Coef. Of Variation
2012	47	0.014	0.358	0.151	0.063	787	7108	0.049
2013	68	0.012	0.233	0.106	0.051	768	7108	0.044
2014	67	0	0.226	0.099	0.049	722	7108	0.025
2015	0							
2016	65	0.011	0.325	0.132	0.0055	958	7108	0.036
2017	63	0.037	0.496	0.118	0.0082	850	7134	0.054
2018	69	0.095	0.524	0.156	0.011	1117	7130	0.042
2019	65	0.012	0.440	0.139	0.011	1010	7131	0.051
2020	65	0.0097	0.364	0.174	0.0097	1264	7133	0.038
2021	71	0	0.450	0.141	0.0096	1018	7129	0.046
2022	58	0.0050	0.428	0.190	0.0131	1363	7124	0.034
2023	71	0.047	0.598	0.274	0.0224	2002	7130	0.031

Table 2: FU16 UWTV 2023. Assessment summary.

Year	UWTV abundance estimate	High 95% confidence interval	Low 95% confidence interval	Landings in number	Total discards in number*	Removals in number	Harvest rate (by number)**	Landings	Total discards*	Discard rate (by number)	Dead discard rate (by number)	Mean weight in landings	Mean weight in discards
	millions						%	tonnes	%		grammes		
2012	787	866	708	25	0	25	3.2	1258	0	0	0	50.36	n/a
2013	768	829	707	20	0	20	2.6	1141	0	0	0	57.54	n/a
2014	722	757	687	17.3	0	17.3	2.4	1189	0	0	0	68.54	n/a
2015	No survey			27	0	27	3.3***	1394	0	0	0	50.86	NA
2016	958	1026	890	53	n/a	53	5.6	2154	n/a	n/a	n/a	40.29	n/a
2017	850	939	760	85	n/a	85	10.0	2632	n/a	n/a	n/a	31.01	n/a
2018	1117	1208	1025	66	n/a	66	5.9	2751	n/a	n/a	n/a	41.55	n/a
2019	1010	1111	910	42	n/a	42	4.1	2229	n/a	n/a	n/a	53.38	n/a
2020	1264	1358	1170	50	n/a	50	3.9	1899	n/a	n/a	n/a	38.26	n/a
2021	1018	1110	927	58	n/a	58	5.7	2476	n/a	n/a	n/a	42.82	n/a
2022	1363	1454	1271	98	n/a	98	7.2	2846	n/a	n/a	n/a	29.13	n/a
2023	2002	2124	1880										

\* Discarding up to 2015 was considered to be negligible. Discard estimates are not available since 2016 and are therefore not included in the assessment.

\*\* Values since 2016 onwards may be underestimated owing to insufficient discard data.

\*\*\* The harvest rate is estimated based on a linear interpolation of abundance, as no survey was carried out in this year.

n/a = not available.

Table 3: FU16 UWTV 2023. The basis for the catch advice and scenarios.

Variable	Value	Notes
Stock abundance (2024)	2002	UWTV survey 2023; individuals in millions
Mean weight in projected landings (2024)	36.74	Average 2020–2022; in grammes
Mean weight in projected discards (2024)	-	Unknown
Projected discards rate (2024)	-	Unknown
Discards survival rate (2024)	-	Not applicable

Table 4: FU16 UWTV 2023. Annual catch advice and scenarios.

Basis	Total catch	Projected landings	Projected discards	Harvest rate * %	% advice change **
	PL + PD	PL	PD	For PL + PD	
ICES advice basis					
EU MAP <sup>^</sup> : F <sub>MSY</sub>	4560	4560	0	6.2	20
EU MAP <sup>^</sup> : F <sub>MSY lower</sub>	3677	3677	0	5.0	20
EU MAP <sup>^</sup> : F <sub>MSY upper</sub> ***	4560	4560	0	6.2	20
Other scenarios					
MSY approach	4560	4560	0	6.2	20
F <sub>2022</sub>	5272	5272	0	7.2	39

\* By number.

\*\* Advice values for 2024 relative to the corresponding 2023 values (MAP advice of 3787, 3054, and 3787 tonnes, respectively); other option values are relative to 3787 tonnes.

\*\*\* F<sub>MSY upper</sub> = F<sub>MSY</sub> for this stock.

<sup>^</sup> EU multiannual plan (MAP) for the Western Waters (EU, 2019).