

12-20-2023

## End-Stage Acute Thoracic Aortic Care Patients' Interventions and Two-Year Survival: the New York State Experience

Joshua Helali

*Scripps Clinic/Scripps Green Hospital, helali.joshua@scrippshealth.org*

Annet Kuruvilla

*State University of New York at Stony Brook, Annet.Kuruvilla@stonybrookmedicine.edu*

Ashutosh R. Yaligar

*State University of New York at Stony Brook, ashutosh.yaligar1@gmail.com*

Daniel H. Wolbrom

*State University of New York at Stony Brook*

Sohaib Agha

*New York Institute of Technology*

*See next page for additional authors*

Follow this and additional works at: <https://commons.library.stonybrook.edu/dos-articles>



Part of the [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), and the [Surgery Commons](#)

---

### Recommended Citation

Helali, Joshua; Kuruvilla, Annet; Yaligar, Ashutosh; Wolbrom, Daniel; Agha, Sohaib; Yang, Jie; Pryor, Aurora; Price, Jonathan; Tannous, Henry; Bilfinger, Thomas; McLarty, Allison; Shroyer, Annie-Laurie; "End-Stage Acute Thoracic Aortic Care Patients' Interventions and Two-Year Survival: the New York State Experience" (2023). Department of Surgery Faculty Publications.

This Report is brought to you for free and open access by the Department of Surgery at Academic Commons. It has been accepted for inclusion in Department of Surgery Faculty Publications by an authorized administrator of Academic Commons. For more information, please contact [mona.ramonetti@stonybrook.edu](mailto:mona.ramonetti@stonybrook.edu), [hu.wang.2@stonybrook.edu](mailto:hu.wang.2@stonybrook.edu).

---

## Authors

Joshua Helali, Annet Kuruvilla, Ashutosh R. Yaligar, Daniel H. Wolbrom, Sohaib Agha, Jie Yang, Aurora D. Pryor, Jonathan Price, Henry J. Tannous, Thomas J. Bilfinger, Allison J. McLarty, and Annie-Laurie W. Shroyer

**Title:** End-Stage Acute Thoracic Aortic Care Patients' Interventions and Two-Year Survival: the New York State Experience

**Principal Investigator:** Dr. Allison J. McLarty, MD

**Co-Principal Investigators:** Dr. A. Laurie Shroyer, PhD, MSHA, Dr. Thomas Bilfinger, MD, ScD, Dr. Henry J. Tannous, MD, and Dr. Jie Yang, PhD

**Residents:** Joshua Helali, MD, Daniel Wolbrom, MD, Annet Kuruvilla, MD

**Medical Students:** Sohaib Agha

**Undergraduate Students:** Ashutosh Yaligar

**Biostatistician:** Jie Yang, PhD, Junying Wang

**Research Question:** What are the risk factors, treatments, and outcomes for patients *initially presenting to a hospital* with a *ruptured and/or dissected thoracic aortic aneurysm (TAA)*?

**PURPOSE AND SPECIFIC AIMS:**

Using the de-identified New York Statewide Planning and Research Cooperative System (SPARCS) comprehensive data system, this descriptive study seeks to analyze data for patients presenting initially with either a ruptured and/or dissected thoracic aortic aneurysm (TAA) from 2005-2018:

- a. Patients with an initial diagnosis billing code for TAA rupture/dissection and/or an initial procedure billing code for TAA-related surgery between 2005-2018 will be included. Examples of exclusions may include:
  - a. Patients with unknown UPID, unknown gender, age < 18, non-NY residents, and duplicate records will be excluded.
  - b. Patients at the treating hospital will be included in the study. For example, exclusions include patients transferred out from external facilities.
  - c. Patients with confounding conditions such as those with a concurrent diagnosis code for bicuspid aortic valve, a TAA-predisposing genetic disease, and cancer patients will be excluded. Cancer patients were defined based on Elixhauser/Charlson comorbidities, which include any hematologic or solid tumor malignancy, except malignant neoplasm of the skin.
  - d. Patients without any follow-up records will be excluded. Patients with an initial TAA rupture/dissection in 2017 or 2018 will be excluded to allow for consistent assessment of 2-year follow up.
  - e. As an example, patients may be stratified by TAA dissection-only, TAA rupture-only, and concurrent TAA rupture and dissection.
- b. Data on baseline demographics, prior comorbidities, admission acuity, and acuity and/or timing of surgery received will be extracted.

- c. Across all of these categories, examples of outcomes range from risk-adjusted 30-outcomes to 2-year composite outcomes. 30-day operative death will be defined as death within the same hospital admission as the initial diagnosis even if it was after 30 days, or 30-day mortality in general. Composite outcomes include death, rupture/dissection after the index hospitalization, or emergent/urgent surgery after the index hospitalization.

The following hypotheses will be tested:

H(0): For TAA dissected-only, TAA rupture-only, and concurrent TAA rupture and dissection patients, there will be no differences in risk-adjusted outcomes across baseline demographics, prior comorbidities, admission type, and acuity and/or timing of surgery received, holding other factors constant.

- As an example, patients may be evaluated in three subgroups: index intervention, subsequent intervention, no intervention received (e.g. type B dissection medical management).

H(0): There will be no differences in baseline demographics, prior comorbidities, admission acuity, and acuity and/or timing of surgery received when comparing patients who initially presented with TAA dissection-only, TAA rupture-only, and concurrent TAA rupture and dissection.

Given this TAA project is research anticipated to advance the frontier of knowledge, ad hoc exploratory analyses (e.g., addressing other endpoints or expanding the time frame for follow-up) may be required to provide details to explain these hypothesis-based findings and/or to identify additional topics warranting future research (i.e., generate pilot data to initiate a new research project). Additional analyses may be performed to explore opportunities to develop future research projects.

## **BACKGROUND AND SIGNIFICANCE:**

### **Ruptured/Dissected Thoracic Aortic Aneurysms**

Thoracic aortic aneurysms (TAAs) are overall rare but when present are a major cause of morbidity and mortality in the United States [Kent, 2014]. TAAs have been mostly found in the ascending aorta but can also present in the aortic arch or descending thoracic aorta. In most cases, TAAs are diagnosed incidentally and are often referred to as a “silent killer.” Previous studies have determined certain traditional risk factors that may contribute to the development of TAAs and subsequent complications of rupture or dissection, which include conditions that predispose to cystic medial degeneration of the aortic wall [Goldfinger et al., 2014].

Some of the major factors that typically lead to such degeneration include aging, specific inflammatory conditions, familial conditions, increases in aortic wall stress secondary to hypertension, and rapid aneurysm expansion. Autosomal dominant diseases that can predispose patients to TAAs include connective tissue diseases such as Marfan syndrome, Ehlers–Danlos syndrome, Loeys–Dietz syndrome, and Turner syndrome, but only make up 5% of TAAs. There also appears to be a difference in gender and TAA outcomes, where females are deemed to have

more adverse events with TAA with an increased likelihood of rupture. Anatomic location of the TAA has been associated with a variability in outcomes as well [Saeyeldin et al., 2019].

TAAAs are usually asymptomatic until a certain threshold size is reached, at which point there is an increasing risk of dissection or rupture. The complications of rupture or dissection or both are considered to be an emergency with a high morbidity rate [Goldfinger et al., 2014]. According to current guidelines, the decision to pursue surgical intervention depends on the size and location of the TAA and whether or not there is a concurrent rupture/dissection. In most cases, surgical intervention involves resection and replacement or stenting of at least an aortic segment. There is currently a gap in the literature in regard to short-term and long-term outcomes of ruptured/dissected TAAAs with and without surgical intervention.

### **Rationale of Study**

Current guidelines state that symptomatic TAAAs including ruptured TAAAs and Type A aortic dissections should be operated on surgically [Hiratzka et al., 2010]. In the acute setting, Type B dissections are conventionally treated medically, but in practice they are often chronically treated with surgery after acute medical management [Alfson et al, 2017]. Contrary to what might be expected based on these guidelines, data from our patient population suggests that in practice, most ruptured and dissected TAA survivors were found to have no surgical intervention. Our study therefore assesses the status-quo of management trends of ruptured and dissected TAAAs.

In addition, since ruptured and dissected TAAAs carry such a high mortality, management in the acute setting is more focused on salvaging survival than consideration of longer-term outcomes [Geisbüsch et al, 2010]. We therefore have limited knowledge of how these ruptured and dissected TAA patients do down the road. This study therefore aims to assess the risk-factors associated with risk-adjusted outcomes of ruptured and dissected TAA patients. Furthermore, for this very high-risk TAA patient cohort, a screening algorithm may be developed from our findings to help guide future clinical care and optimize patient outcomes.

### **RESEARCH DESIGN AND METHODS:**

Using the SPARCS database (i.e. IRB approval previously received by Dr. Allison J. McLarty), a retrospective observational cohort study will be performed using the SPARCS Health Facts dataset. Multivariable regression analysis using NYS records in the SPARCS dataset ranging from 2005- 2018 will be performed.

With the assistance of the SBU SOM Bioinformatics Department and Biostatistics Core Lab, the SPARCS database will be matched and merged to the enclosed coding listings to generate a study-specific de-identified thoracic aortic aneurysm database. Bioinformatics and Biostatistics team members will be responsible for providing the descriptive statistics listed, as well as providing a study-database for future analyses. SAS version 9.4 will be used to complete all necessary statistical analysis for this study. Given the concern for multiple comparisons performed, multivariable significance level will be set at 0.05, with all p-values reported for separate interpretation by readers. All secondary and tertiary analyses, as well as additional exploratory analyses, will use a p-value of <0.01.

*Please note, the SPARCS database de-identified reports will be used, and a non-human subjects research (NHSR).*

**SAMPLE TABLES:**

Table 1: Descriptive table for patient characteristics and risk factors by disease type

| <b>Variable</b>                    | <b>Total</b> | <b>TAA dissection only (N= )</b> | <b>TAA ruptured only (N= )</b> | <b>TAA ruptured and dissection (N= )</b> | <b>P-value</b> |
|------------------------------------|--------------|----------------------------------|--------------------------------|--|----------------|
| <b>Patient Characteristics</b>     |              |                                  |                                |  |                |
| <b>Gender</b>                      |              |                                  |                                |  |                |
| Female                             |              |                                  |                                |  |                |
| Male                               |              |                                  |                                |  |                |
| <b>Age group</b>                   |              |                                  |                                |  |                |
| <80                                |              |                                  |                                |  |                |
| >=80                               |              |                                  |                                |  |                |
| <b>Race</b>                        |              |                                  |                                |  |                |
| Black                              |              |                                  |                                |  |                |
| Non Black                          |              |                                  |                                |  |                |
| <b>Insurance type</b>              |              |                                  |                                |  |                |
| Commercial                         |              |                                  |                                |  |                |
| Other                              |              |                                  |                                |  |                |
| <b>Year group</b>                  |              |                                  |                                |  |                |
| <2014                              |              |                                  |                                |  |                |
| >=2014                             |              |                                  |                                |  |                |
| <b>Risk Factors</b>                |              |                                  |                                |  |                |
| <b>Admission type</b>              |              |                                  |                                |  |                |
| Elective                           |              |                                  |                                |  |                |
| Urgent/Emergent                    |              |                                  |                                |  |                |
| Carotid Disease                    |              |                                  |                                |  |                |
| Coronary Artery Disease            |              |                                  |                                |  |                |
| Acute Visceral/Mesenteric Ischemia |              |                                  |                                |  |                |
| Hypotension                        |              |                                  |                                |  |                |
| Myocardial Infarction              |              |                                  |                                |  |                |
| Arrhythmia                         |              |                                  |                                |  |                |
| Resuscitation                      |              |                                  |                                |  |                |
| Aortic Valve Replacement           |              |                                  |                                |  |                |

|               |  |  |  |  |  |
|---------------|--|--|--|--|--|
| Elderly Women |  |  |  |  |  |
|---------------|--|--|--|--|--|

Table 2: Multivariate model results for 30-day operative death

| Variable | Odds ratio (95% CI) | P-value* |
|----------|---------------------|----------|
|          |                     |          |

Table 3: Multivariate model results for 2-year composite outcomes

| Variable | Odds ratio (95% CI) | P-value* |
|----------|---------------------|----------|
|          |                     |          |

Table 4: Univariate analysis of surgery categories stratified by disease type – 30-day operative death



| <b>Variable</b>                          | <b>Total</b> | <b>Without 30-day<br/>operative death</b> | <b>With 30-day<br/>operative death</b> | <b>P-value</b> |
|--|--------------|---|--|----------------|
| <b>All Patients (N= )</b>                |              |   |  |                |
| <b>Any Surgery</b>                       |              |   |  |                |
| <b>Surgery Type</b>                      |              |   |  |                |
| No Surgery                               |              |   |  |                |
| Elective Surgery                         |              |   |  |                |
| Urgent Surgery                           |              |   |  |                |
| Emergent Surgery                         |              |   |  |                |
| <b>TAA Dissection Only (N= )</b>         |              |   |  |                |
| <b>Any Surgery</b>                       |              |   |  |                |
| <b>Surgery Type</b>                      |              |   |  |                |
| No Surgery                               |              |   |  |                |
| Elective Surgery                         |              |   |  |                |
| Urgent Surgery                           |              |   |  |                |
| Emergent Surgery                         |              |   |  |                |
| <b>TAA Ruptured Only (N= )</b>           |              |   |  |                |
| <b>Any Surgery</b>                       |              |   |  |                |
| <b>Surgery Type</b>                      |              |   |  |                |
| No Surgery                               |              |   |  |                |
| Elective Surgery                         |              |   |  |                |
| Urgent Surgery                           |              |   |  |                |
| Emergent Surgery                         |              |   |  |                |
| <b>TAA Ruptured and Dissection (N= )</b> |              |   |  |                |
| <b>Any Surgery</b>                       |              |   |  |                |
| <b>Surgery Type</b>                      |              |   |  |                |
| No Surgery                               |              |   |  |                |
| Emergent Surgery                         |              |   |  |                |

Table 5: Univariate analysis of surgery categories stratified by disease type – 2-year composite outcome.

| Variable                                 | Total | Without 2-year Composite Outcome | With 2-year Composite Outcome | P-value |
|--|-------|----------------------------------|-------------------------------|---------|
| <b>All Patients (N= )</b>                |       |                                  |                               |         |
| <b>Any Surgery</b>                       |       |                                  |                               |         |
| <b>Surgery Type</b>                      |       |                                  |                               |         |
| No Surgery                               |       |                                  |                               |         |
| Elective Surgery                         |       |                                  |                               |         |
| Urgent Surgery                           |       |                                  |                               |         |
| Emergent Surgery                         |       |                                  |                               |         |
| <b>TAA Dissection Only (N= )</b>         |       |                                  |                               |         |
| <b>Any Surgery</b>                       |       |                                  |                               |         |
| <b>Surgery Type</b>                      |       |                                  |                               |         |
| No Surgery                               |       |                                  |                               |         |
| Elective Surgery                         |       |                                  |                               |         |
| Urgent Surgery                           |       |                                  |                               |         |
| Emergent Surgery                         |       |                                  |                               |         |
| <b>TAA Ruptured Only (N= )</b>           |       |                                  |                               |         |
| <b>Any Surgery</b>                       |       |                                  |                               |         |
| <b>Surgery Type</b>                      |       |                                  |                               |         |
| No Surgery                               |       |                                  |                               |         |
| Elective Surgery                         |       |                                  |                               |         |
| Urgent Surgery                           |       |                                  |                               |         |
| Emergent Surgery                         |       |                                  |                               |         |
| <b>TAA Ruptured and Dissection (N= )</b> |       |                                  |                               |         |
| <b>Any Surgery</b>                       |       |                                  |                               |         |
| <b>Surgery Type</b>                      |       |                                  |                               |         |
| No Surgery                               |       |                                  |                               |         |
| Emergent Surgery                         |       |                                  |                               |         |

Table 6: O/E ratio for NYS region for 30-day operative death

| Variable | Level       | O/E Ratio (95% CI) |
|----------|-------------|--------------------|
| Region   | Long Island |                    |
|          | Mid/North   |                    |

| Variable | Level    | O/E Ratio (95% CI) |
|----------|----------|--------------------|
|          | NYC Area |                    |
|          | West     |                    |

Table 7: Average annual facility volume by region

| Variable | Level       | Average annual facility volume |      |    |     |        |     |     |        |
|----------|-------------|--------------------------------|------|----|-----|--------|-----|-----|--------|
|          |             | N                              | Mean | SD | Min | Median | Max | IQR | 95% CI |
| Region   | Long Island |                                |      |    |     |        |     |     |        |
|          | Mid/North   |                                |      |    |     |        |     |     |        |
|          | NYC Area    |                                |      |    |     |        |     |     |        |
|          | West        |                                |      |    |     |        |     |     |        |
| Total    |             |                                |      |    |     |        |     |     |        |

## DIAGNOSIS AND PROCEDURE CODES

### **TAA Diagnosis Codes**

| Risk Factor                             | ICD-10   | ICD-9  | CPT |
|---|--|--|-----|
| <b>Atherosclerotic Disease of Aorta</b> | I70.0  | 440.0  |     |
| <b>Carotid Disease</b>                  | I77.71, I65.21, I65.22, I65.23, I65.29, G45.1, I65.1 I65.01 I65.02 I65.03 I65.09 | 433.0-433.3, 435.8, 443.21   |     |
| <b>Coronary Artery Disease</b>          | I25-I25.4, I25.6-I25.9   | 414-414.9  |     |
| <b>Congestive Heart Failure</b>         | I50-I50.9, I09.9 I11.0 I13.0 I13.2 I25.5 I42.0 I42.5 - I42.9 I43 P29.0           | 428.0-428.9, 398.91 402.01 402.11 402.91 404.01 404.03 404.11 404.13 404.91 404.93 |     |

|                              |   |  |  |
|------------------------------|---|--|--|
|                              |   | 425.4 - 425.99,<br>414.8   |  |
| <b>Hypertension</b>          | I10, I11.0, I11.9,<br>I12.0, I12.9, I13.0,<br>I13.1, I13.10,<br>I13.11, I13.2, I15,<br>I15.0, I15.1, I15.2,<br>I15.8, I15.9 I16,<br>I16.0, I16.1, I16.9 | 401.0, 401.1,<br>401.9, 402.01,<br>402.11, 402.91,<br>402.00, 402.10,<br>402.90, 403.01,<br>403.11, 403.91,<br>403.00, 403.10,<br>403.90, 404.01,<br>404.11, 404.91,<br>404.00, 404.10,<br>404.90, 404.02,<br>404.12, 404.92,<br>404.03, 404.13,<br>404.93, 405.01,<br>405.11, 405.91,<br>405.91, 405.99,<br>405.09, 405.19,<br>405.99 |  |
| <b>Myocardial Infarction</b> | I25.2, I21-I21.9,<br>I21.A1, I21.A9,<br>I22.0-I22.9   | 412, 410.00-<br>410.92   |  |
| <b>Aortic Valve Disease</b>  | I06.0, I06.2, I08.0,<br>I08.2, I08.3, I35.0,<br>I35.2, I06.1, I35.1,<br>I35.8, I35.9  | 395.0, 395.2,<br>424.1, 396.3,<br>396.1  |  |
| <b>Aortic Coarctation</b>    | Q25.1   | 747.1  |  |
| <b>Diabetes mellitus</b>     | E08.00-E13.9  | 249.00-249.91,<br>250-250.03,<br>250.1-250.13,<br>250.2-250.23,<br>250.3-250.33,<br>250.4-250.43,<br>250.5-250.53,<br>250.6-250.63,<br>250.7-250.73,<br>250.8-250.83,<br>250.9,250.93  |  |

|   |  |   |  |
|---|--|---|--|
| <b>Chronic Obstructive Pulmonary Disease</b>    | J41.0, J41.1, J41.8, J42, J43.0, J43.1, J43.2, J43.8, J43.9, J44.0, J44.1, J44.9<br>*Asthma and Bronchiectasis were not counted as a chronic obstructive pulmonary disease | 491.0, 491.1, 491.20, 491.21, 491.22, 491.8, 491.9, 492.0, 492.8, 496<br>*Asthma and Bronchiectasis were not counted as a chronic obstructive pulmonary disease |  |
| <b>Tobacco/Smoking</b>                          | Z72.0, F17.21-F17.299, Z87.891   | V15.82, 305.1   |  |
| <b>Cerebrovascular Disease</b>                  | I60-I69.998, Z86.73, G46.0-G46.8, G45.0-G45.9  | 430-438.9, V12.54   |  |
| <b>Peripheral Vascular Disease</b>              | I73.00, I73.01, I73.1, I73.81, I73.89, I73.9, I70.20-I70.25, I70.8, I70.92   | 443.0, 443.1, 443.21, 443.22, 443.23, 443.24, 443.29, 443.81, 443.82, 443.89, 443.9   |  |
| <b>Prior Percutaneous Coronary Intervention</b> | Z98.61   | V45.82  |  |
| <b>Dialysis</b>                                 | Z99.2  | V45.11  |  |
| <b>Hyperlipidemia</b>                           | E78.00, E78.01, E78.1, E78.2, E78.3, E78.41, E78.49, E78.5   | 272.0, 272.1, 272.2, 272.3, 272.4,  |  |
| <b>Dyslipidemia- Literature Codes</b>           | E78.0-E78.9  | 272.0-272.5, 272.8, 272.9   |  |
| <b>Dyslipidemia- Dr. Bilfinger Codes</b>        | E78.00, E78.01, E78.5, E78.79, E78.9   | 272.0, 272.4, 272.8, 272.9  |  |
| BMI:<br>< 19.9<br>20-29<br>30-39<br>³ 40.0      | Z68.1<br>Z68.20-Z68.29<br>Z68.30-Z68.39<br>Z68.41-Z68.45   | < 19: V85.0<br>19-24: V85.1<br>25.0-29.9:<br>V85.21-V85.25<br>30.0-39.9:<br>V85.30-V85.39<br>³ 40.0: V85.41-V85.44  |  |
| <b>Acute Renal Failure</b>                      | N17.0-N17.9  | 584.5-584.9   |  |

|                                 |   |   |  |
|---------------------------------|---|---|--|
| Chronic kidney disease          |   |   |  |
| Stage I                         | N18.1   | 585.1   |  |
| Stage II                        | N18.2   | 585.2   |  |
| Stage III                       | N18.3   | 585.3   |  |
| Stage IV                        | N18.4   | 585.4   |  |
| Stage V                         | N18.5   | 585.5   |  |
| ESRD                            | N18.6   | 585.6   |  |
| CKD, with dialysis              | Z99.2   | V45.11  |  |
| CKD, without dialysis           | N18.1-N18.9   | 585.1-585.9   |  |
| CKD + Hypertension              | I12.0, I12.9, I13.0,<br>I13.1, I13.10,<br>I13.11, I13.2   | 403.00-403.91,<br>404.00-404.93   |  |
| <b>Obesity</b>                  | E66-E66.9   | 278-278.3   |  |
| <b>Resuscitation</b>            | 5A12012, 5A19054  | 93.93, 99.60  | 92950  |
| <b>Arrhythmia</b>               | R00.0-R00.1,<br>R00.8-R00.9<br>I44-I44.7, I45-I45.9,<br>I47.0-I47.9, I48-<br>I48.92, I49-I49.9,   | 426.0-426.9,<br>427.0-427.9,<br>785.0   |  |
| <b>Hypovolemic Shock</b>        | R57.1   | 785.59  |  |
| <b>Cardiogenic Shock</b>        | R57.0   | 785.51  |  |
| <b>Hypotension</b>              | I95.0-I95.3, I95.89,<br>I95.9   | 458.0-458.1,<br>458.29-458.9  |  |
| <b>Aortic Valve Replacement</b> | 02RF07Z, 02RF0JZ,<br>02RF08Z,<br>02RF0KZ,<br>02RF37H,<br>02RF38H,<br>02RF3JH,<br>02RF3KH,<br>02RF37Z,<br>02RF38Z, 02RF3JZ,<br>02RF3KZ,<br>02RF47Z,<br>02RF48Z, 02RF4JZ,<br>02RF4KZ, Z95.2-<br>Z95.4 | 35.21, 35.22,<br>35.05, 35.06,<br>V43.3, V42.2  | 33405<br>33406<br>33410<br>33361<br>33362<br>33363<br>33364<br>33365<br>33366<br>33367<br>33368<br>33369 |
| <b>Chest Pain</b>               | I20.0-I20.9   | 413.1, 413.9,<br>411.1  |  |
| <b>Liver dysfunction</b>        | K76.0-K76.9,<br>K70-K70.9, K71.0-<br>K71.9, K72.0-<br>K74.9, B18.0-B18.9  | 570, 571.0-<br>571.9, 572.2-<br>572.8, 573.3-<br>573.9, 070.22,<br>070.23, 070.32,<br>070.33, 070.44, |  |

|  |   |                                 |  |
|--|---|---------------------------------|--|
|  |   | 070.54, 070.59,<br>070.6, 070.9 |  |
| <b>Immunosuppression</b>                             | Z79.51-Z79.52<br>D80.0-D89.9  | V58.65,<br>279.00-279.9         |  |
| <b>Cardiopulmonary Bypass Time</b>                   | 5A1221Z   | 39.61, 39.66                    | 33367, 33368, 33369,<br>33390, 33391, 33405,<br>33406, 33410, 33858,<br>33859, 33863, 33864,<br>33871, 33870 |
| <b>Neurological Deficit (hemiplegia, paraplegia)</b> | G81.00-G81.94,<br>G82.2-G82.22  | 342.00-342.92,<br>344.1         |  |
| <b>Atrial Fibrillation</b>                           | Paroxysmal: I48.0<br>Persistent: I48.11-<br>I48.19<br>Chronic: I48.20-<br>I48.21<br>Unspecified: I48.91 | 427.31                          |  |
| <b>Atrial Flutter</b>                                | Typical: I48.3<br>Atypical: I48.4<br>Unspecified: I48.92  | 427.32                          |  |
| <b>Use of Anticoagulants</b>                         | Z79.01-Z79.02   | V58.61, V58.63                  |  |
| <b>Use of Systemic Steroids</b>                      | Z79.51-Z79.52   | V58.65                          |  |
| <b>Acute Visceral/Mesenteric Ischemia</b>            | K55.0-K55.069   | 557.0                           |  |
| <b>Coma at Admission</b>                             | R40.20  | 780.01                          |  |
| <b>Vasopressor Use</b>                               | 3E030XZ,<br>3E033XZ,<br>3E040XZ, 3E043XZ  | 0.17                            |  |

|   |         |
|---|---------|
| <b>POST-OPERATIVE OUTCOME CODE TO BE USED IN CONJUNCTION WITH OUTCOME:</b>                            |         |
| Other post procedural cardiac functional disturbances following cardiac surgery                       | I97.190 |
| Other post procedural cardiac functional disturbances following other surgery                         | I97.191 |
| Other post procedural complications and disorders of the circulatory system, not elsewhere classified | I97.89  |

| Disease                                    |                                   | ICD10  | ICD9                                    |
|--|-----------------------------------|--|---|
| Thoracic Aortic Aneurysm (TAA)-non rupture |                                   | I71.2  | 441.2                                   |
| TAA-ruptured                               |                                   | I71.1  | 441.1                                   |
| TAA-dissection                             |                                   | I71.01   | 441.01                                  |
| Bicuspid Aortic Valve Dx                   |                                   | Q23.1  | 746.4                                   |
| Other                                      | Marfan Syndrome                   | Q87.40   | 759.82                                  |
|  | Marfan Syndrome-CV manifestations | Q87.41, Q87.410, Q87.418   | NA                                      |
|  | Marfan Syndrome-Aortic Dilation   | Q87.418  | NA                                      |
|  | Ehler-Danlos Syndrome             | Q79.6 (Q79.60, Q79.61, Q79.62, Q79.63, Q79.69 were used instead) | 756.83                                  |
|  | Turner Syndrome                   | Q96.0, Q96.9   | 758.6 (includes > than Turner syndrome) |
|  | Ehler-Danlos Syndrome-Vascular    | Q79.63   | NA                                      |
|  |                                   |  |   |

**TAA Procedure Codes**

| Open Approach   |         |  |       |  |
|---|---------|--|-------|--|
| ICD-10  |         | ICD-9  |       |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Autologous Tissue Substitute, Open Approach    | 02RX07Z | Resection of vessel with anastomosis, other thoracic vessels | 38.35 |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Zooplastic Tissue, Open Approach               | 02RX08Z | Resection of vessel with replacement, thoracic vessels       | 38.45 |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Synthetic Substitute, Open Approach            | 02RX0JZ |  |       |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Nonautologous Tissue Substitute, Open Approach | 02RX0KZ |  |       |  |



|   |             |  |  |  |
|---|-------------|--|--|--|
| Replacement of Thoracic Aorta, Descending with Autologous Tissue Substitute, Open Approach    | 02RW<br>07Z |  |  |  |
| Replacement of Thoracic Aorta, Descending with Zooplastic Tissue, Open Approach               | 02RW<br>08Z |  |  |  |
| Replacement of Thoracic Aorta, Descending with Synthetic Substitute, Open Approach            | 02RW<br>0JZ |  |  |  |
| Replacement of Thoracic Aorta, Descending with Nonautologous Tissue Substitute, Open Approach | 02RW<br>0KZ |  |  |  |
| Supplement Thoracic Aorta, Descending with Autologous Tissue Substitute, Open Approach        | 02UW<br>07Z |  |  |  |
| Supplement Thoracic Aorta, Descending with Zooplastic Tissue, Open Approach                   | 02UW<br>08Z |  |  |  |
| Supplement Thoracic Aorta, Descending with Synthetic Substitute, Open Approach                | 02UW<br>0JZ |  |  |  |
| Supplement Thoracic Aorta, Descending with Nonautologous Tissue Substitute, Open Approach     | 02UW<br>0KZ |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Autologous Tissue Substitute, Open Approach    | 02UX0<br>7Z |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Zooplastic Tissue, Open Approach               | 02UX0<br>8Z |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Synthetic Substitute, Open Approach            | 02UX0<br>JZ |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Nonautologous Tissue Substitute, Open Approach | 02UX0<br>KZ |  |  |  |
| Restriction of Thoracic Aorta, Descending with Intraluminal Device, Open Approach             | 02VW<br>0DZ |  |  |  |
| Restriction of Thoracic Aorta, Ascending/Arch with Intraluminal Device, Open Approach         | 02VX0<br>DZ |  |  |  |
| Repair Thoracic Aorta, Descending, Open Approach  | 02QW<br>0ZZ |  |  |  |

|  |             |  |       |  |
|--|-------------|--|-------|--|
| Repair Thoracic Aorta, Ascending/Arch, Open Approach   | 02QX0<br>ZZ |  |       |  |
|  |             |  |       |  |
| <b>Percutaneous/Hybrid Approach</b>  |             |  |       |  |
| <b>ICD-10</b>  |             | <b>ICD-9</b>   |       |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Autologous Tissue Substitute, Percutaneous Endoscopic Approach    | 02RX4<br>7Z | Endovascular implantation of graft in thoracic aorta | 39.73 |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Zooplastic Tissue, Percutaneous Endoscopic Approach               | 02RX4<br>8Z |  |       |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Synthetic Substitute, Percutaneous Endoscopic Approach            | 02RX4<br>JZ |  |       |  |
| Replacement of Thoracic Aorta, Ascending/Arch with Nonautologous Tissue Substitute, Percutaneous Endoscopic Approach | 02RX4<br>KZ |  |       |  |
| Replacement of Thoracic Aorta, Descending with Autologous Tissue Substitute, Percutaneous Endoscopic Approach        | 02RW<br>47Z |  |       |  |
| Replacement of Thoracic Aorta, Descending with Zooplastic Tissue, Percutaneous Endoscopic Approach                   | 02RW<br>48Z |  |       |  |
| Replacement of Thoracic Aorta, Descending with Synthetic Substitute, Percutaneous Endoscopic Approach                | 02RW<br>4JZ |  |       |  |
| Replacement of Thoracic Aorta, Descending with Nonautologous Tissue Substitute, Percutaneous Endoscopic Approach     | 02RW<br>4KZ |  |       |  |
| Supplement Thoracic Aorta, Descending with Autologous Tissue Substitute, Percutaneous Approach                       | 02UW<br>37Z |  |       |  |

|   |             |  |  |  |
|---|-------------|--|--|--|
| Supplement Thoracic Aorta, Descending with Zooplastic Tissue, Percutaneous Approach                           | 02UW<br>38Z |  |  |  |
| Supplement Thoracic Aorta, Descending with Synthetic Substitute, Percutaneous Approach                        | 02UW<br>3JZ |  |  |  |
| Supplement Thoracic Aorta, Descending with Nonautologous Tissue Substitute, Percutaneous Approach             | 02UW<br>3KZ |  |  |  |
| Supplement Thoracic Aorta, Descending with Autologous Tissue Substitute, Percutaneous Endoscopic Approach     | 02UW<br>47Z |  |  |  |
| Supplement Thoracic Aorta, Descending with Zooplastic Tissue, Percutaneous Endoscopic Approach                | 02UW<br>48Z |  |  |  |
| Supplement Thoracic Aorta, Descending with Synthetic Substitute, Percutaneous Endoscopic Approach             | 02UW<br>4JZ |  |  |  |
| Supplement Thoracic Aorta, Descending with Nonautologous Tissue Substitute, Percutaneous Endoscopic Approach  | 02UW<br>4KZ |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Autologous Tissue Substitute, Percutaneous Approach            | 02UX3<br>7Z |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Zooplastic Tissue, Percutaneous Approach                       | 02UX3<br>8Z |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Synthetic Substitute, Percutaneous Approach                    | 02UX3<br>JZ |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Nonautologous Tissue Substitute, Percutaneous Approach         | 02UX3<br>KZ |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Autologous Tissue Substitute, Percutaneous Endoscopic Approach | 02UX4<br>7Z |  |  |  |

|  |             |  |  |  |
|--|-------------|--|--|--|
| Supplement Thoracic Aorta, Ascending/Arch with Zooplastic Tissue, Percutaneous Endoscopic Approach               | 02UX4<br>8Z |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Synthetic Substitute, Percutaneous Endoscopic Approach            | 02UX4<br>JZ |  |  |  |
| Supplement Thoracic Aorta, Ascending/Arch with Nonautologous Tissue Substitute, Percutaneous Endoscopic Approach | 02UX4<br>KZ |  |  |  |
| Restriction of Thoracic Aorta, Ascending/Arch with Intraluminal Device, Percutaneous Approach                    | 02VX3<br>DZ |  |  |  |
| Restriction of Thoracic Aorta, Ascending/Arch with Intraluminal Device, Percutaneous Endoscopic Approach         | 02VX4<br>DZ |  |  |  |
| Repair Thoracic Aorta, Ascending/Arch, Percutaneous Approach   | 02QX3<br>ZZ |  |  |  |
| Repair Thoracic Aorta, Ascending/Arch, Percutaneous Endoscopic Approach  | 02QX4<br>ZZ |  |  |  |
| Restriction of Thoracic Aorta, Descending with Intraluminal Device, Percutaneous Approach                        | 02VW<br>3DZ |  |  |  |
| Restriction of Thoracic Aorta, Descending with Intraluminal Device, Percutaneous Endoscopic Approach             | 02VW<br>4DZ |  |  |  |
| Repair Thoracic Aorta, Descending, Percutaneous Approach   | 02QW<br>3ZZ |  |  |  |
| Repair Thoracic Aorta, Descending, Percutaneous Endoscopic Approach  | 02QW<br>4ZZ |  |  |  |
|  |             |  |  |  |

| <b>CPT Description</b>  | <b>CPT Code</b>  |
|---|--|
| TAA repair-sternotomy   | 33859, 33860, 33863, 33864, 33866, 33870, 33871, 33875 |
| Ascending aorta graft, with cardiopulmonary bypass, includes valve suspension, when performed; for aortic dissection  | 33858  |
| Ascending aorta graft, with cardiopulmonary bypass, includes valve suspension, when performed; for aortic disease other than dissection (eg, aneurysm)  | 33859  |
| Ascending aorta graft, with cardiopulmonary bypass, includes valve suspension, when performed   | 33860  |
| Ascending aorta graft, with cardiopulmonary bypass, with aortic root replacement using valved conduit and coronary reconstruction (eg, Bentall)   | 33863  |
| Ascending aorta graft, with cardiopulmonary bypass with valve suspension, with coronary reconstruction and valve-sparing aortic root remodeling (eg, David Procedure, Yacoub Procedure)   | 33864  |
| Aortic hemiarch graft including isolation and control of the arch vessels, beveled open distal aortic anastomosis extending under one or more of the arch vessels, and total circulatory arrest or isolated cerebral perfusion (List separately in addition to code for primary procedure)  | 33866  |
| Transverse arch graft, with cardiopulmonary bypass  | 33870  |
| Repair Procedures for Thoracic Aortic Aneurysm  | 33871  |
| Descending thoracic aorta graft, with or without bypass   | 33875  |
| TEVAR- Ascending & Arch-Not present   | NO CODE  |
| TEVAR-Descending  | 33880, 33881, 33883, 33884, 33886, 33889, 33891        |
| Endovascular repair of descending thoracic aorta (eg, aneurysm, pseudoaneurysm, dissection, penetrating ulcer, intramural hematoma, or traumatic disruption); involving coverage of left subclavian artery origin, initial endoprosthesis plus descending thoracic aortic extension(s), if required, to level of celiac artery origin | 33880  |

|   |       |
|---|-------|
| Endovascular repair of descending thoracic aorta (eg, aneurysm, pseudoaneurysm, dissection, penetrating ulcer, intramural hematoma, or traumatic disruption); not involving coverage of left subclavian artery origin, initial endoprosthesis plus descending thoracic aortic extension(s), if required, to level of celiac artery origin | 33881 |
| Placement of proximal extension prosthesis for endovascular repair of descending thoracic aorta (eg, aneurysm, pseudoaneurysm, dissection, penetrating ulcer, intramural hematoma, or traumatic disruption); initial extension  | 33883 |
| Placement of proximal extension prosthesis for endovascular repair of descending thoracic aorta (eg, aneurysm, pseudoaneurysm, dissection, penetrating ulcer, intramural hematoma, or traumatic disruption); each additional proximal extension   | 33884 |
| Placement of distal extension prosthesis(s) delayed after endovascular repair of descending thoracic aorta  | 33886 |
| Open subclavian to carotid artery transposition performed in conjunction with endovascular repair of descending thoracic aorta, by neck incision, unilateral  | 33889 |
| Bypass graft, with other than vein, transcervical retropharyngeal carotidcarotid, performed in conjunction with endovascular repair of descending thoracic aorta, by neck incision  | 33891 |

| Procedure                       | ICD-10  | ICD-9               | CPT                 |                                |
|---------------------------------|---|---------------------|---------------------|--------------------------------|
| Aortic Valve Replacement (Open) | 02RF07Z, 02RF0JZ, 02RF08Z, 02RF0KZ  | 35.21, 35.22, 35.11 | 33405, 33406, 33410 |                                |
| Aortic Valve Open               | 024F07J, 024F08J, 024F0JJ, 025F0ZZ, 027F04Z, 027F0DZ, 027F0ZZ, 02BF0ZX, 02BF0ZZ, 02CF0ZZ, 02NF0ZZ, 02QF0ZJ, 02QF0ZZ, 02UF07J, 02UF07Z, 02UF08J, 02UF08Z, 02UF0JJ, 02UF0JZ, 02UF0KJ, 02UF0KZ, 02WF07Z, 02WF08Z, 02WF0JZ, 02WF0KZ | 35.21, 35.22, 35.12 | 33405, 33406, 33411 | *use these codes for exclusion |

|  |   |                           |   |                                      |
|--|---|---------------------------|---|--------------------------------------|
| CABG   | 0210083, 0210088, 0210089, 021008C,<br>021008F, 021008W, 0210093, 0210098,<br>0210099, 021009C, 021009F, 021009W,<br>02100A3, 02100A8, 02100A9, 02100AC,<br>02100AF, 02100AW, 02100J3, 02100J8,<br>02100J9, 02100JC, 02100JF, 02100JW,<br>02100K3, 02100K8, 02100K9, 02100KC,<br>02100KF, 02100KW, 02100Z3, 02100Z8,<br>02100Z9, 02100ZC, 02100ZF, 0211083,<br>0211088, 0211089, 021108C, 021108F,<br>021108W, 0211093, 0211098, 0211099,<br>021109C, 021109F, 021109W, 02110A3,<br>02110A8, 02110A9, 02110AC, 02110AF,<br>02110AW, 02110J3, 02110J8, 02110J9,<br>02110JC, 02110JF, 02110JW, 02110K3,<br>02110K8, 02110K9, 02110KC, 02110KF,<br>02110KW, 02110Z3, 02110Z8, 02110Z9,<br>02110ZC, 02110ZF, 0212083, 0212088,<br>0212089, 021208C, 021208F, 021208W,<br>0212093, 0212098, 0212099, 021209C,<br>021209F, 021209W, 02120A3, 02120A8,<br>02120A9, 02120AC, 02120AF,<br>02120AW, 02120J3, 02120J8, 02120J9,<br>02120JC, 02120JF, 02120JW, 02120K3,<br>02120K8, 02120K9, 02120KC, 02120KF,<br>02120KW, 02120Z3, 02120Z8, 02120Z9,<br>02120ZC, 02120ZF, 0213083, 0213088,<br>0213089, 021308C, 021308F, 021308W,<br>0213093, 0213098, 0213099, 021309C,<br>021309F, 021309W, 02130A3, 02130A8,<br>02130A9, 02130AC, 02130AF, 02130AW,<br>02130J3, 02130J8, 02130J9, 02130JC,<br>02130JF, 02130JW, 02130K3, 02130K8,<br>02130K9, 02130KC, 02130KF, 02130KW,<br>02130Z3, 02130Z8, 02130Z9, 02130ZC,<br>02130ZF | 36.10-<br>36.19           | 33510,<br>33511,<br>33512,<br>33513,<br>33514,<br>33516,<br>33517,<br>33518,<br>33519,<br>33521,<br>33522,<br>33523,<br>33530,<br>33533,<br>33534,<br>33535,<br>33536 | *use these<br>codes for<br>exclusion |
| Mitral<br>Valve<br>Replace<br>ment<br>(Open) | 02QG0ZZ, 02QG0ZE, 02RG07Z,<br>02RG08Z, 02RG0JZ, 02RG0KZ   | 35.12,<br>35.24,<br>35.23 | 33430   |                                      |
| Mitral<br>Valve<br>Open                      | 02NG0ZZ, 02QG0ZZ, 02UG08Z,<br>025G0ZZ, 027G0ZZ, 02BG0ZX,<br>02BG0ZZ, 02UG07Z, 02UG0JZ,<br>02UG0KZ, 02VG0ZZ, 02RG08Z,<br>02QG0ZE, 027G04Z, 027G0DZ,  | 35.12,<br>35.24,<br>35.23 | 33422,<br>33425,<br>33426,<br>33427,<br>33430   | *use these<br>codes for<br>exclusion |

|                                    |   |                           |   |                                |
|------------------------------------|---|---------------------------|---|--------------------------------|
|                                    | 02RG07Z, 02RG0JZ, 02RG0KZ,<br>024G082, 02UG08E, 024G072, 024G0J2,<br>024G0K2, 02UG07E, 02UG0JE,<br>02UG0KE, 02CG0ZZ, 02WG07Z,<br>02WG08Z, 02WG0JZ, 02WG0KZ  |                           |   |                                |
| Tricuspid Valve Replacement (Open) | 02QJ0ZZ, 02QJ0ZG, 02RJ07Z, 02RJ0JZ,<br>02RJ08Z, 02RJ0KZ   | 35.14,<br>35.27,<br>35.28 | 33465   |                                |
| Tricuspid Valve Open               | 02NJ0ZZ, 02QJ0ZZ, 02UJ08Z, 025J0ZZ,<br>027J0ZZ, 02BJ0ZX, 02BJ0ZZ, 02UJ07Z,<br>02UJ0JZ, 02UJ0KZ, 02QJ0ZG, 027J0DZ,<br>02RJ07Z, 02RJ0JZ, 02CJ0ZZ, 02RJ08Z,<br>027J04Z, 02RJ0KZ, 02WJ08Z, 02UJ08G,<br>024J082, 02UJ07G, 02UJ0JG, 02UJ0KG,<br>02WJ0JZ, 024J072, 024J0J2, 024J0K2,<br>02WJ07Z, 02WJ0KZ | 35.14,<br>35.27,<br>35.28 | 33460,<br>33463,<br>33464,<br>33465,<br>33468   | *use these codes for exclusion |
| Pulmonary Valve Replacement (Open) | 02QH0ZZ, 02RH07Z, 02RH0JZ,<br>02RH08Z, 02RH0KZ  | 35.13,<br>35.26,<br>35.25 | 33475   |                                |
| Pulmonary Valve Open               | 02NH0ZZ, 02QH0ZZ, 02UH08Z,<br>025H0ZZ, 027H0ZZ, 02BH0ZX,<br>02BH0ZZ, 02TH0ZZ, 02UH07Z,<br>02UH0JZ, 02UH0KZ, 02LH0ZZ,<br>027H0DZ, 02RH07Z, 02RH0JZ,<br>02CH0ZZ, 02RH08Z, 027H04Z,<br>02RH0KZ, 02LH0CZ,<br>02LH0DZ, 02WH08Z, 02WH0JZ,<br>02WH07Z, 02WH0KZ   | 35.13,<br>35.26,<br>35.25 | 33474,<br>33475,<br>33476,<br>33478   | *use these codes for exclusion |
| Maze                               | 02560ZZ, 02570ZZ, 025K0ZZ, 025L0ZZ,<br>02B60ZZ, 02B70ZZ, 02BK0ZZ,<br>02BL0ZZ, 02T80ZZ   | 37.33                     | 33254,<br>33255,<br>33256   | *use these codes for exclusion |
| Sternotomy                         | 0P800ZZ   | 77.31                     | 33202,<br>33365,<br>33955,<br>33956,<br>33963,<br>33964,<br>33985,<br>33986,<br>33988,<br>33989 |                                |



|             |  |   |   |  |
|-------------|--|---|---|--|
| Thoracotomy | 02JA0ZZ, 0WJC0ZZ, 0P810ZZ, 0P820ZZ, 0P850ZZ, 0P860ZZ, 0P890ZZ, 0P8B0ZZ | 34.02   | 33366, 33140, 33202, 33203, 33236, 33238, 33243, 33955, 33956, 33963, 33964, 33985, 33986, 33988, 33989 |  |
| Other       | Z95.1-Z95.818, Z95.9   | V45.81, V43.3, V42.1, V42.2, V43.21, V43.22, V45.09, V45.00 | .   | *use these codes for exclusion: Note, this should be accompanied with either sternotomy or thoracotomy codes |

**REFERENCES:**

Kent, KC (2014). Abdominal aortic aneurysms. *New England Journal of Medicine*, 371(22), 2101-2108.

Saeyeldin, A. A., Velasquez, C. A., Mahmood, S., Brownstein, A. J., Zafar, M. A., Ziganshin, B. A., & Elefteriades, J. A. (2019). Thoracic aortic aneurysm: unlocking the "silent killer" secrets. *General thoracic and cardiovascular surgery*, 67(1), 1–11.

Goldfinger, J. Z., Halperin, J. L., Marin, M. L., Stewart, A. S., Eagle, K. A., & Fuster, V. (2014). Thoracic aortic aneurysm and dissection. *Journal of the American College of Cardiology*, 64(16), 1725-1739.

Alfson, D. B., & Ham, S. W. (2017). Type B Aortic Dissections: Current Guidelines for Treatment. *Cardiology clinics*, 35(3), 387–410.

Geisbüsch, P., Kotelis, D., Weber, T. F., Hyhlik-Dürr, A., & Böckler, D. (2010). Endovascular repair of ruptured thoracic aortic aneurysms is associated with high perioperative mortality and morbidity. *Journal of vascular surgery*, 51(2), 299–304.

Hiratzka LF, Bakris GL, Beckman JA, Bersin RM, Carr VF, Casey DE Jr, Eagle KA, Hermann LK, Isselbacher EM, Kazerooni EA, Kouchoukos NT, Lytle BW, Milewicz DM, Reich DL, Sen S, Shinn JA, Svensson LG, Williams DM; American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines; American Association for Thoracic Surgery; American College of Radiology; American Stroke Association; Society of Cardiovascular Anesthesiologists; Society for Cardiovascular Angiography and Interventions; Society of Interventional Radiology; Society of Thoracic Surgeons; Society for Vascular Medicine. 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM guidelines for the diagnosis and management of patients with Thoracic Aortic Disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, American Association for Thoracic Surgery, American College of Radiology, American Stroke Association, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of Thoracic Surgeons, and Society for Vascular Medicine. *Circulation*. 2010 Apr 6;121(13):e266-369. doi: 10.1161/CIR.0b013e3181d4739e.