

## Article

---

"Effect of media and temperature on sporulation of *Septoria apiicola*, and of inoculum density on septoria blight severity in celery"

I.W. Mudita et A.C. Kushalappa

*Phytoprotection*, vol. 72, n° 3, 1991, p. 97-103.

Pour citer cet article, utiliser l'information suivante :

URI: <http://id.erudit.org/iderudit/706009ar>

DOI: 10.7202/706009ar

Note : les règles d'écriture des références bibliographiques peuvent varier selon les différents domaines du savoir.

---

Ce document est protégé par la loi sur le droit d'auteur. L'utilisation des services d'Érudit (y compris la reproduction) est assujettie à sa politique d'utilisation que vous pouvez consulter à l'URI <https://apropos.erudit.org/fr/usagers/politique-dutilisation/>

---

Érudit est un consortium interuniversitaire sans but lucratif composé de l'Université de Montréal, l'Université Laval et l'Université du Québec à Montréal. Il a pour mission la promotion et la valorisation de la recherche. Érudit offre des services d'édition numérique de documents scientifiques depuis 1998.

Pour communiquer avec les responsables d'Érudit : [info@erudit.org](mailto:info@erudit.org)

## Effect of media and temperature on sporulation of *Septoria apiicola*, and of inoculum density on septoria blight severity in celery

I. W. Mudita and A.C. Kushalappa

Department of Plant Science, Macdonald Campus of McGill University,  
21 111 Lakeshore Rd., Sainte-Anne-de-Bellevue, Québec, Canada H9X 1C0

(Received 1991-05-07; accepted 1991-09-23)

The number of conidia produced by *Septoria apiicola* was quantified on seven media incubated at 20°C. Conidia were produced abundantly on celery agar (CA) and celery potato dextrose agar (CPDA) after 12 days of incubation. In general, media containing celery decoction, including cleared V8 juice agar (CVA), produced more conidia than those without celery decoction. The number of conidia and the colony area produced per plate on CA increased with increase in temperature up to an optimum temperature and then decreased markedly. A second-order linear regression equation explained 99% of the variation in the number of conidia produced on CA plates incubated at temperatures ranging from 15 to 30°C, with the predicted optimum temperature of 22.4°C. Ninety eight percent of the variations in the colony area were explained by a linearized BETE-equation with the predicted optimal temperature of 22.7°C. Blight severity in celery (*Apium graveolens* var. *dulce*) increased with increase in inoculum density. For inoculum density ranging from 5-140 conidia/cm<sup>2</sup> of leaf surface, 92% of the variations in the number of lesions per leaf, pooled from two experiments, were explained by a second-order linear regression equation. The model was less reliable for inoculum densities less than 17 conidia/cm<sup>2</sup> of leaf surface.

**Mudita, I.W., and A.C. Kushalappa. 1991. Effect of media and temperature on sporulation of *Septoria apiicola*, and of inoculum density on septoria blight severity in celery. PHYTOPROTECTION 72: 97-103.**

Nous avons étudié la production de conidies du *Septoria apiicola* sur sept milieux gélosés à 20°C. Après 12 jours d'incubation, la production de conidies fut abondante sur les géloses d'infusion de feuilles de céleri seules (CA) ou additionnées de pomme de terre dextrosée (CPDA). En général, les géloses contenant des extraits de la plante hôte, incluant le jus V-8 (CVA), ont produit davantage de conidies que les autres géloses. Le nombre de conidies par boîte et la surface des colonies ont augmenté au fur et à mesure que la température augmentait jusqu'à un maximum puis a diminué rapidement. Un polynôme du deuxième degré a expliqué 99% de la variation du nombre de conidies par boîte. La température optimale prédite par le modèle fut de 22,4°C. Un modèle BETE linéarisé a expliqué 98% de la variation de la surface des colonies. La température optimale prédite par le modèle fut de 22,7°C. La gravité de l'infection sur des feuilles de céleri (*Apium graveolens* var. *dulce*) a augmenté au fur et à mesure que la concentration de l'inoculum augmentait. Un polynôme du deuxième degré a expliqué 92% de la variation du nombre de lésions par feuilles pour des concentrations de 5-140 conidies/cm<sup>2</sup> de feuille. Néanmoins, les prévisions du modèle furent moins fiables pour des concentrations inférieures à 17 conidies/cm<sup>2</sup> de feuille.

### Introduction

Celery [*Apium graveolens* L. var. *dulce* (Mill.) Pers.] is an economically important vegetable crop in Québec. Grown mainly in the muck-soil region of the province, it was valued at \$ 5.4 million for a total production area of 406 ha in 1989 (Statistics Canada 1990). *Septoria* blight, induced by *Septoria apiicola* Speg., threatens the crop in years of abundant rainfall or where overhead sprinkler irrigation is used, conditions that provide long durations of wetness required for infection (Berger 1970). The pathogen produces chlorotic spots on leaf laminae and later on petioles. The lesions enlarge rapidly and turn necrotic (Dixon 1981). When the disease is severe, the entire crop may be lost (Sherf and MacNab 1986). To reduce this risk, early and frequent fungicide applications are recommended to manage the disease (CPVQ

1987), but an efficient management strategy is not presently available. In order to develop a more efficient and environmentally sound management practice for septoria blight, better understanding of the various biological aspects of the pathogen and the disease is necessary.

The information available on factors influencing the production of conidia of *S. apiicola* is scarce. Sheridan (1968) produced conidia on potato dextrose agar but Wolcan (1987) obtained abundant conidia using Fries modified agar and V8 juice agar. Similarly, the relationship between inoculum density and blight severity is also not clearly understood. Berger (1970) reported that the number of lesions produced by *S. apiicola* increased with increase in inoculum concentration but at very high inoculum concentrations the number of lesions was reduced. More information on production of conidia of *S. apiicola* and the effect of inoculum density on blight severity is re-

quired to study specific environmental factors critical for blight development.

The objectives of this study were to select a suitable medium and temperature level for inoculum production, and to model the effect of inoculum density on the number of lesions produced by *S. apiicola*.

## Materials and methods

**Production of inoculum.** *S. apiicola* was isolated from a diseased celery plant collected from a commercial field in Sainte-Clotilde, Québec, in 1989. The isolate was used to inoculate plants grown in the greenhouse from which one diseased leaf was selected for further isolation. The leaf was cut into discs and the discs were surface disinfected by immersing in 1% NaOCl for 2 min. After being rinsed, the discs were suspended in sterile water for 10-15 min to release the conidia. The suspension of conidia was diluted to approximately 1 000 conidia/mL and 5 mL of the suspension was dispersed on water-agar medium in a Petri plate. Following overnight incubation at 20°C, one germinating conidium was transferred into each of five plates of celery-agar medium and incubated at 20°C and a 16-h photoperiod in a growth chamber (Forma Scientific Model 3740) for 12 days. Only the conidia cultured in this medium were used in all the experiments, with no subsequent sub-culturing.

Inoculum for all experiments was obtained by flooding 12-day-old cultures with 2 mL sterile distilled water. After 10-15 min, the resulting suspensions of conidia were filtered through two layers of sterile muslin cloth and the concentration of the suspensions was determined using a hemacytometer.

**Preparation of media.** Media tested were potato dextrose agar (PDA), potato dextrose peptone agar (PDPA), cleared V8 juice agar (CVA), celery agar (CA), celery dextrose agar (CDA), celery peptone agar (CPA), and celery potato dextrose agar (CPDA). PDA and CVA were prepared as recommended by Tuite (1969) and PDPA by following the method of Louis and Cooke (1985). CA was prepared by adding 320 mL of celery decoction into 680 mL of water containing 12 g bacto-agar (Difco). Celery decoction was prepared from 50 g of shredded celery leaves steamed in 500 mL distilled water for 1 h. CDA and CPA were prepared by adding 5 g dextrose and 2.5 g peptone, respec-

tively, into 250 mL CA. CPDA was prepared from CA by replacing bacto-agar with dehydrated bacto-PDA (Difco). All media were autoclaved for 20 min at 100 kPa and then were adjusted to a pH of 5.0 with 25% lactic acid.

**Inoculation, incubation and quantification of conidia.** Five 10- $\mu$ L droplets of conidium suspension were dispensed onto each plate (100mm x 15 mm) using a digital micropipette (Eppendorf). Inoculum concentrations used for media and temperature experiments were 200 000 and 150 000 conidia/mL, respectively. For the media experiment, plates were incubated at 20°C in the growth chamber (Forma Scientific Model 3740) for 12 and 18 days whereas for the temperature experiment, plates were incubated at 15°, 20°, 25° and 30°C for 12 days. Two 15-W cool black fluorescent tubes positioned at 0.3 m above the cultures provided the lighting for a photoperiod of 16 h. There were four and three replicates for the media and the temperature experiment, respectively.

After incubation, 15 mL of 1% formaldehyde with 0.01% Tween-80 in sterile distilled water was poured into each plate. Fifteen minutes later, the resulting suspension of conidia was then filtered using two layers of muslin cloth. The number of conidia in each suspension was counted, four counts per suspension, with the aid of a hemacytometer. The shape of the conidia was also noted. Colony diameter was measured at five random directions per colony and the colony area was calculated based on the average value.

**Preparation and inoculation of plants.** Transplants of celery cv. Florida 683 were produced by placing one seed on the surface of each cell of a seedling tray previously filled with Profi-Mix B type VPW 30 (a mixture of peat moss, dolomitic lime stone, perlite, and vermiculite) and then lightly covered with a layer of fine sand. The seedling tray was then placed in a growth chamber (Conviron Model E15) adjusted to a 22°C day/18°C night temperature and a 14-h photoperiod. After six weeks, seedlings were transplanted into plastic pots (12-cm deep, 12-cm upper-internal diameter) filled with a mixture of 2/3 part (v/v) soil containing 20% organic matter, 1/6 part peat, and 1/6 part perlite. Potted plants were maintained in a growth chamber and were watered daily. A 15-15-18 fertilizer (Plant-Prod) prepared at 0.3 g/L was applied daily along with the

irrigation water.  $\text{CaCO}_3$  was applied twice a week at 8 g/L to prevent black-heart disease.

At the sixth-leaf stage, three fully expanded leaves per plant were tagged for blight monitoring. Plants were inoculated using an automatic spray chamber (Incom Int.). The canopy level was placed at 1.3 m below the solid-cone-type spray nozzle which moved horizontally. Prior to spraying each suspension, at a constant pressure of 172 kPa, the speed of nozzle movement was adjusted to deposit 140, 70, 35.5, 17.5, 8.8, and 4.4 conidia/cm<sup>2</sup> from inoculum suspensions of 80 000, 40 000, 20 000, 10 000, 5 000, and 2500, respectively. The density of deposition of conidia was determined by placing three water-agar plates (100 mm x 15 mm) at the canopy level and the counts were made on four 1-cm<sup>2</sup> circles per plate. The actual number of conidia deposited from each suspension was 140.04, 69.98, 35.02, 17.51, 8.76, and 4.80, respectively. After spraying the inoculum, each plant was enclosed in a transparent plastic bag and placed in the growth chamber (Incom Int.) maintained at 20°C for 72 h. The bags were then removed and the plants were transferred into the greenhouse and arranged according to a randomized complete block design on a bench. Lesions were counted daily on the three-tagged leaves after their first appearance until they began to coalesce. The experiment was conducted twice.

**Data analyses.** Analysis of variance was performed on the number of conidia and the colony area from the media experiment. The Duncan's new multiple range test was used to

separate treatment means. Regression analysis was used to model the effect of temperature on the number of conidia and on the colony area and the effect of inoculum density on the number of lesions. A second model considered appropriate for the temperature experiment is the BETE-function (Analytis 1977). The linearized form of this function was fitted to the proportion of maximum number of conidia and to the proportion of maximum colony area. For the number of lesions, the analysis was carried out on the data from the individual experiment and also on the pooled data. All analyses were conducted using SAS/STAT™ software for IBM-PC (SAS Institute Inc. 1987).

## Results

**Effect of media on sporulation.** Conidia were produced most abundantly on CA and on CPDA after 12 days of incubation at 20°C (Table 1). In general, media containing celery decoction produced more conidia than those without celery decoction. Two exceptions were noted, CDA after 12 days of incubation and CPA after 18 days of incubation. Dextrose or peptone combined with celery decoction enhanced colony development. Large colony area on CPDA after 12 days of incubation was followed by marked increase in the production of conidia after 18 days of incubation. Marked increase in the number of conidia was also observed on CVA, CDA, and CPA.

Differences in shape were observed among conidia produced on different media. Conidia produced on PDA and PDPA were slightly

**Table 1.** Effect of various media on number of conidia and colony area of *S. apiicola* at 20°C

Media <sup>§</sup>	Number of conidia per plate (x 10 <sup>6</sup> ) <sup>†</sup>		Colony area (mm <sup>2</sup> ) <sup>†</sup>	
	12 days	18 days	12 days	18 days
CA	5.44 a	7.00 a	91 a	123 ac
CPDA	5.39 a	7.03 a	135 b	146 b
CPA	5.03 b	6.30 b	122 c	147 b
CVA	2.74 c	7.00 a	76 d	128 a
PDA	0.81 d	1.16 d	86 ad	109 c
PDPA	0.52 d	1.27 d	87 ad	121 ac
CDA	0.51 d	2.32 c	112 c	117 c

§ CA=celery agar; CPDA=celery potato dextrose agar; CPA=celery peptone agar; CVA=cleared V8 juice agar; PDA=potato dextrose agar; PDPA=potato dextrose peptone agar; CDA=celery dextrose agar.

† Average of four replicates. Numbers in a column followed by the same letter are not significantly different ( $P > 0.05$ ) according to the Duncan's new multiple range test.

atypical while those produced on other media were similar to those found in lesions.


**Effect of temperature on sporulation.** The number of conidia produced per plate increased with increase in temperature up to a maximum level and then decreased markedly (Fig. 1A). Colony area also followed the same pattern (Fig. 1B). A second-degree linear polynomial regression equation explained significantly

( $P \leq 0.01$ ) the effects of temperature on the number of conidia per plate (Eq. 1):

$$CONIDIA = -48.06 + 4.84 T - 0.11 T^2, \quad [1]$$

$$R^2 = 0.99$$

where *CONIDIA* is the predicted number of conidia produced per plate ( $\times 10^6$ ) and *T* is the temperature ( $^{\circ}\text{C}$ ). The optimum temperature for




**Figure 1.** Effect of temperature (*T*) on **A**) number of conidia ( $\times 10^6$ ) and **B**) colony area ( $\text{mm}^2$ ). The equation to predict production of conidia is  $CONIDIA = -48.06 + 4.84 T - 0.11 T^2$ ,  $R^2 = 0.99$  and the equation to predict colony development is  $PMCA = \exp(3.58 + 3.29 \ln T' + 2.16 \ln [1 - T'])$ ,  $R^2 = 0.98$  where *CONIDIA* is the predicted number of conidia per plate, *PMCA* is the proportion of maximum colony area (colony area in  $\text{mm}^2$ ), *T* is the temperature, *T'* is  $(T - T_{min}) / (T_{max} - T_{min})$ , and  $R^2$  is the coefficient of determination.

production of conidia predicted using Eq. 1 was 22.4°C. For the colony area, the polynomial model underestimated the colony area at 25°C since the overall responses were skewed to the lower temperatures. A good fit to the proportion of maximum colony area was obtained from use of the linearized BETE-function. The back-

transformed BETE-equation for the proportion of the maximum colony area (the observed maximum =  $1.97 \times 10^2 \text{ mm}^2$ ) is:

$$PMCA = \exp(3.58 + 3.29 \ln T' + 2.16 \ln [1 - T']) \quad [2]$$

$$R^2 = 0.98$$



**Figure 2.** Effect of inoculum density on disease severity induced by *S. apicola* for A) experiment 1 and experiment 2 and B) pooled data. Equations to predict number of lesions are:  $NLES_1 = 6.02 + 2.89INODEN - 0.01INODEN^2$ ,  $R^2 = 0.98$ ;  $NLES_2 = 15.79 + 3.35INODEN - 0.01INODEN^2$ ,  $R^2 = 0.91$ ; and  $NLES_P = 10.99 + 3.13INODEN - 0.01INODEN^2$ ,  $R^2 = 0.91$ , respectively, where  $NLES$  is the predicted number of lesions per leaf, subscripts 1, 2, and P indicate the number of experiment and pooled data, respectively,  $INODEN$  is the inoculum density (conidia/cm<sup>2</sup> of leaf surface), and  $R^2$  is the coefficient of determination.

where *PMCA* is the predicted proportion of maximum colony area (colony area in mm<sup>2</sup>) and  $T' = (T - T_{min}) / (T_{max} - T_{min})$  in which *T* is the temperature (°C) and subscripts *min* and *max* indicate the minimum and the maximum temperature for activity, respectively.  $T_{min}$  and  $T_{max}$  were selected iteratively until the highest coefficient of determination (*R*<sup>2</sup>) was obtained. Equation 2 was obtained at the  $T_{min}$  and  $T_{max}$  value of 11.0 and 30.5°C, respectively. Based on this equation, the predicted optimum temperature for colony development was 22.78°C.

**Effect of inoculum densities on disease severity.** Lesions were large and non-coalescing on plants inoculated with 4.8 conidia/cm<sup>2</sup> and were small and readily coalesced on plants inoculated with 70 and 140 conidia/cm<sup>2</sup>. Although the lesions tended to appear earlier on older leaves, the final number of lesions depended more on the leaf size than on the leaf position. The maximum number of lesions was observed 18 days after inoculation and this maximum was used for further analyses. The number of lesions increased with the increase in inoculum density (Fig. 2). Second-degree linear regression equations explained significantly (*P* ≤ 0.01) the variation in the number of lesions from the individual experiment. Since the *F*-test carried out to examine the homogeneity of experimental error from both experiments indicated no significant difference (*P* > 0.01), a regression model was also fitted to the pooled data. Equations to predict the number of lesions based on the data from experiment 1, experiment 2, and pooled data are:

$$NLES_1 = -6.02 + 2.89 INODEN - 0.01 INODEN^2, \quad [3]$$

$$R^2 = 0.98$$

$$NLES_2 = -15.79 + 3.35 INODEN - 0.01 INODEN^2, \quad [4]$$

$$R^2 = 0.91$$

$$NLES_p = -10.99 + 3.13 INODEN - 0.01 INODEN^2, \quad [5]$$

$$R^2 = 0.91$$

where *NLES* is the predicted number of lesions per leaf, subscripts 1, 2 and *P* indicate experiment 1, experiment 2, and the pooled data, respectively, and *INODEN* is the inoculum density (conidia/cm<sup>2</sup>). Models based on data from individual experiments and pooled data

accounted for up to 91% of the variation in the number of lesions observed. The model based on experiment 1 had a higher coefficient of determination (*R*<sup>2</sup>) and a lower mean square error (MSE) but a less desirable pattern of residuals as compared to the model based on experiment 2 or to that based on pooled data. Models based on data from individual experiments gave poorer predictions than that based on pooled data at the lower inoculum densities. All models had negative intercepts which were not significantly different from zero (*P* > 0.01).

## Discussion

Production of sufficient amount of inoculum is a primary concern for quantitative epidemiological studies of many diseases including septoria blight. Abundant conidia were obtained when celery decoction was added into media. The magnitude of the increment, however, was influenced by the basal medium used. When celery decoction was added into media containing dextrose or peptone, more conidia were produced after 18 days than after 12 days of incubation. Larger colony area as a result of addition of peptone or dextrose to media containing celery decoction accounted for more sporulation.

CA and CPDA readily produced abundant conidia, but preparation of these media required a supply of celery leaves. When celery leaves are not readily available, however, CVA which contains celery could be considered as an alternative. Wolcan (1987) reported that V8 juice agar was one of the best media for production of secondary conidia of *S. apiicola*. Here, CVA provided abundant conidia but significantly less than those produced on CA, CPDA, and CPA after 12 days.

Media not containing celery decoction produced slightly atypical conidia. The shape of conidia varied more in conidia produced on media containing no celery decoction than in those produced on media containing celery decoction. However, according to Gabrielson and Grogan (1964) this variation was common in *S. apiicola* and was not likely attributed to a particular isolate.

Temperature may have affected sporulation directly by influencing production of conidia or indirectly by influencing colony development which later provided a larger colony area for further production of conidia. As a result, both

production of conidia and colony area had similar response curves. However, the optimum temperature for the development of colonies was closer to the maximum than that for the production of conidia. Consequently, use of second-order polynomial to model the effect of temperature on the colony area appeared inadequate. The linearized BETE-functions gave a better fit to the colony area. For fitting BETE-functions to the colony area,  $T_{min}$  and  $T_{max}$  were selected merely based on the coefficients of determination. Experimental proof of  $T_{min}$  and  $T_{max}$  would improve the model.

The variations in the pooled number of lesions per leaf obtained from different inoculum densities were explained by a second-order linear regression equation with a non-significant negative intercept. The equation implied that, within the range of inoculum densities tested, the ratio of disease to inoculum decreased with the increase in inoculum density and that the minimum inoculum threshold for the pathogen was not critical. A decrease in the ratio of disease to inoculum may occur when there is competition among conidia for susceptible infection sites (Van der Plank 1975).

The model based on the pooled data was less reliable for inoculum densities lower than 17.5 conidia/cm<sup>2</sup>. Densities above 70 conidia/cm<sup>2</sup> resulted in more crowded and easily coalesced lesions, making reading more difficult. Therefore, for quantitative studies involving plant inoculations and lesion counting, an inoculum density from 17-35 conidia/cm<sup>2</sup> of leaf surface is considered the most appropriate.

This research was funded by the Conseil des recherches en pêche et agroalimentaire du Québec. The scholarship provided by the government of Indonesia to the first author is gratefully acknowledged. We thank O. Carisse for French translation of the abstract and Dr. T.C. Paulitz for his useful suggestions.

- Analytis, S. 1977.** Über die Relation zwischen biologischer Entwicklung und Temperatur bei phytopathogenen Pilzen. *Phytopathol. Z.* 90: 64-76.
- Berger, R.D. 1970.** Epiphytology of celery late blight. *Proc. Fla. State Hort. Soc.* 83: 208-212.
- CPVQ (Conseil des productions végétales du Québec). 1987.** Légumes protection 1983-1984. Agdex 250/605. Ministère de l'Agriculture, des Pêcheries et de l'Alimentation du Québec. 86 pp.
- Dixon, G.R. 1981.** Vegetable crop diseases. Macmillan Publ., London. 404 pp.
- Gabrielson, R.L., and R.G. Grogan. 1964.** The celery late blight organism *Septoria apiicola*. *Phytopathology* 54: 1251-1257.
- Louis, I., and R. C. Cooke. 1985.** Conidial matrix and spore germination in some plant pathogens. *Trans. Br. Mycol. Soc.* 84: 661-667.
- SAS Institute Inc. 1987.** SAS/STAT Guide for personal computers. Version 6 edition. SAS Institute Inc., Cary, NC. 1028 pp.
- Sherf, A.F., and A.A. MacNab. 1986.** Vegetable diseases and their control. 2nd ed. John Wiley & Sons, New York. 728 pp.
- Sheridan, J.E. 1968.** Condition for germination of pycnidiospores of *Septoria apiicola* Speg. *N.Z. J. Bot.* 6: 315-322.
- Statistic Canada. 1990.** Fruit and vegetable production. Catalogue 22-0033 Seasonal. Ministry of Supply and Service Canada, Ottawa. 29 pp.
- Tuite, J. 1969.** Plant pathological methods, fungi and bacteria. Burgess Publ., Minneapolis, MN. 239 pp.
- Van der Plank, J.E. 1975.** Principles of plant infection. Academic Press, New York. 216 pp.
- Wolcan, S. 1987.** Conidios secundarios de *Septoria apiicola* Speg. I. Comparación con las picnidiosporas. Medios de cultivo que estimulan la producción de los dos tipos de esporas. *Turrialba* 37: 239-244.