

# *ALGEBRAIC COMBINATORICS*

Tan N. Tran & Shuhei Tsujie

**MAT-free graphic arrangements and a characterization of strongly chordal graphs by edge-labeling**

Volume 6, issue 6 (2023), p. 1447-1467.

<https://doi.org/10.5802/alco.319>


© The author(s), 2023.

 This article is licensed under the  
CREATIVE COMMONS ATTRIBUTION (CC-BY) 4.0 LICENSE.  
<http://creativecommons.org/licenses/by/4.0/>



*Algebraic Combinatorics is published by The Combinatorics Consortium  
and is a member of the Centre Mersenne for Open Scientific Publishing*  
[www.tccpublishing.org](http://www.tccpublishing.org)    [www.centre-mersenne.org](http://www.centre-mersenne.org)  
e-ISSN: 2589-5486





# MAT-free graphic arrangements and a characterization of strongly chordal graphs by edge-labeling

Tan N. Tran & Shuhei Tsujie

**ABSTRACT** Ideal subarrangements of a Weyl arrangement are proved to be free by the multiple addition theorem (MAT) due to Abe–Barakat–Cuntz–Hoge–Terao (2016). They form a significant class among Weyl subarrangements that are known to be free so far. The concept of MAT-free arrangements was introduced recently by Cuntz–Mücksch (2020) to capture a core of the MAT, which enlarges the ideal subarrangements from the perspective of freeness. The aim of this paper is to give a precise characterization of the MAT-freeness in the case of type  $A$  Weyl subarrangements (or graphic arrangements). It is known that the ideal and free graphic arrangements correspond to the unit interval and chordal graphs, respectively. We prove that a graphic arrangement is MAT-free if and only if the underlying graph is strongly chordal. In particular, it affirmatively answers a question of Cuntz–Mücksch that MAT-freeness is closed under taking localization in the case of graphic arrangements.

## 1. INTRODUCTION

A hyperplane arrangement is said to be *free* if its logarithmic derivation module is a free module [17, 12] (cf. Definition 2.1). An important class of free arrangements is that of Weyl arrangements defined by the positive roots of irreducible root systems in Euclidean spaces. There has been considerable interest in finding and characterizing free subarrangements of a Weyl arrangement by combinatorial structures. In the case of type  $A$ , Weyl subarrangements are completely determined by *graphic arrangements* whose freeness is fully characterized by *chordal* (or *supersolvable*) *graphs* [15, 5] (cf. Theorem 2.3). Several certain cases of type  $B$  were studied in the connection with signed graphs [5, 16, 19] yet no complete characterization is known. Arguably, the most fundamental and significant class of Weyl subarrangements known to be free is that of *ideal subarrangements* derived by the multiple addition theorem (MAT) due to Abe–Barakat–Cuntz–Hoge–Terao [1] (cf. Theorem 2.7). In fact, the MAT applies in a more general setting; based on that Cuntz–Mücksch [3] introduced a new class of free arrangements, the so called *MAT-free arrangements* (cf. Definition 2.4). MAT-free arrangements are new examples of arrangements having *combinatorial freeness* [3, Lemma 18] that has received increasing attention in recent years (e.g. [2, 10]).

---

*Manuscript received 29th June 2022, revised 1st June 2023, accepted 5th June 2023.*

**KEYWORDS.** Hyperplane arrangement, free arrangement, MAT-free arrangement, ideal subarrangement, graphic arrangement, strongly chordal graph, edge-labeling of graph.

**ACKNOWLEDGEMENTS.** The first author was supported by JSPS Research Fellowship for Young Scientists Grant Number 19J12024 at Hokkaido University and a postdoctoral fellowship of the Alexander von Humboldt Foundation at Ruhr-Universität Bochum.

We are especially interested in characterizations of freeness-related properties of type  $A$  Weyl subarrangements (or graphic arrangements in  $\mathbb{R}^\ell$ ) in which considerable power and development of graph theory would be brought to bear. It is known that the ideal graphic arrangements are parametrized by *unit interval graphs* e.g. [20] (cf. Theorem 2.8). Our main result is a complete characterization of MAT-free graphic arrangements over arbitrary fields by means of *strongly chordal graphs* (cf. Definition 2.9). We summarize the results in Table 1.

We find it interesting that the concept of MAT-freeness which was recently introduced is captured by the notion of strongly chordal graphs which appeared much earlier in literature. This can also be regarded as an analogue of the classical theory of freeness and chordality. We will see in §5.1 that many important concepts in the classical theory such as simplicial vertices and perfect elimination orderings of chordal graphs have their analogous MAT- versions.

Graph class	Weyl subarrangement class	Location
chordal	free	[15], [5, Theorem 3.3]
<b>strongly chordal</b>	<b>MAT-free</b>	<b>Theorem 2.10</b>
unit interval	ideal	e.g. [20, Theorem 16]

TABLE 1. Interplay between graphs and Weyl arrangements in type  $A$ . The third row (in bold) indicates the main result of the paper.

Our main result has a number of applications. From the viewpoint of arrangement theory, it gives an affirmative answer to a question of Cuntz–Mücksch in the case of graphic arrangements that MAT-freeness is closed under taking localization (cf. Corollary 2.11). Thanks to the relation  $\{\text{unit interval graphs}\} \subsetneq \{\text{strongly chordal graphs}\}$ , it also gives a different and graphical proof that the ideal graphic arrangements are MAT-free (cf. §6(A)). From the viewpoint of graph theory, our main result contributes a new characterization of strongly chordal graphs via a special type of edge-labelings, which we shall call *MAT-labelings* (cf. Definition 4.2).

A key ingredient in our proof that strong chordality implies MAT-freeness (the harder part) is a characterization of strongly chordal graphs by their *clique intersection posets* due to Nevries–Rosenke [11] (cf. Theorem 3.4). Our strategy is to construct an MAT-labeling for a given strong chordal graph which building blocks are complete induced subgraphs of the graph. The clique intersection poset of a chordal graph consists of all intersections of maximal cliques of the graph which serves as essential machinery in the construction. We believe that it is worth pursuing further the notion of clique intersection poset for MAT-freeness of larger class of arrangements (see §6(F) for more details).

The remainder of the paper is organized as follows. In §2.1, we recall the definitions and basic facts of free arrangements and chordal graphs. In §2.2, we recall the definitions of MAT-free arrangements and strongly chordal graphs, and give the statement of our main result. In §3, we recall some other useful characterizations of (strongly) chordal graphs. In §4.1, we introduce the notion of MAT-labeling of graphs. In §4.2, we prove the “only if” part of our main Theorem 2.10 (MAT-freeness implies strong chordality). In §5.1, we introduce the notions of MAT-simplicial vertices and MAT-perfect elimination orderings. In §5.2, we prove the “if” part of our main Theorem 2.10 (strong chordality implies MAT-freeness). The proof the main theorem is

also included with an example in the end of this section. Finally, in §6, we address some further remarks and suggest some problems for future research.

2. ARRANGEMENTS, GRAPHS AND STATEMENT OF THE MAIN RESULT

2.1. FREE ARRANGEMENTS. We first review some basic concepts and preliminary results on free arrangements. Our standard reference is [12]. Let  $\mathbb{K}$  be a field,  $\ell$  be a positive integer and  $V = \mathbb{K}^\ell$  be the  $\ell$ -dimensional vector space over  $\mathbb{K}$ . A **hyperplane** in  $V$  is a *linear* subspace of codimension one of  $V$ . An **arrangement** is a finite set of hyperplanes in  $V$ . Let  $\mathcal{A}$  be an arrangement in  $V$ . Define the **intersection lattice**  $L(\mathcal{A})$  (of flats) of  $\mathcal{A}$  by

$$L(\mathcal{A}) := \left\{ \bigcap_{H \in \mathcal{B}} H \mid \mathcal{B} \subseteq \mathcal{A} \right\},$$

with the partial order given by reverse inclusion:  $X \leq Y \Leftrightarrow Y \subseteq X$  for  $X, Y \in L(\mathcal{A})$ . We agree that  $V$  is the unique minimal element in  $L(\mathcal{A})$  as the intersection over the empty set. Thus  $L(\mathcal{A})$  is a geometric lattice which can be equipped with rank function  $r(X) := \text{codim}(X)$  for  $X \in L(\mathcal{A})$ . We also define  $\text{rank}(\mathcal{A})$  as the rank of the maximal element  $\bigcap_{H \in \mathcal{A}} H$  of  $L(\mathcal{A})$ .

The **characteristic polynomial**  $\chi_{\mathcal{A}}(t) \in \mathbb{Z}[t]$  of  $\mathcal{A}$  is defined by

$$\chi_{\mathcal{A}}(t) := \sum_{X \in L(\mathcal{A})} \mu(X)t^{\dim X},$$

where  $\mu$  denotes the **Möbius function**  $\mu: L(\mathcal{A}) \rightarrow \mathbb{Z}$  defined recursively by

$$\mu(V) = 1 \quad \text{and} \quad \mu(X) = - \sum_{\substack{Y \in L(\mathcal{A}) \\ X \subsetneq Y}} \mu(Y).$$

Let  $\{x_1, \dots, x_\ell\}$  be a basis for the dual space  $V^*$  and let  $\mathbf{S} = \mathbb{K}[x_1, \dots, x_\ell]$ . The **defining polynomial**  $Q(\mathcal{A})$  of  $\mathcal{A}$  is given by

$$Q = Q(\mathcal{A}) := \prod_{H \in \mathcal{A}} \alpha_H \in \mathbf{S},$$

where  $\alpha_H = a_1x_1 + \dots + a_\ell x_\ell$  ( $a_i \in \mathbb{K}$ ) satisfies  $H = \ker \alpha_H$ . The **module**  $D(\mathcal{A})$  of **logarithmic derivations** is defined by

$$D(\mathcal{A}) := \{ \theta \in \text{Der}(\mathbf{S}) \mid \theta(Q) \in Q\mathbf{S} \},$$

where  $\text{Der}(\mathbf{S}) = \{ \varphi : \mathbf{S} \rightarrow \mathbf{S} \mid \varphi \text{ is } \mathbb{K}\text{-linear, } \varphi(fg) = f\varphi(g) + g\varphi(f) \text{ for any } f, g \in \mathbf{S} \}$  is the set of all derivations of  $\mathbf{S}$  over  $\mathbb{K}$ . Note that  $\text{Der}(\mathbf{S})$  is a free  $\mathbf{S}$ -module with basis  $\{ \partial/\partial x_1, \dots, \partial/\partial x_\ell \}$  consisting of the usual partial derivatives. A non-zero element  $\varphi = f_1 \cdot \partial/\partial x_1 + \dots + f_\ell \cdot \partial/\partial x_\ell \in \text{Der}(\mathbf{S})$  is **homogeneous of degree**  $b$  written  $\text{deg } \varphi = b$  if each non-zero polynomial  $f_i \in \mathbf{S}$  for  $1 \leq i \leq \ell$  is homogeneous of degree  $b$ .

DEFINITION 2.1 (Free arrangements and their exponents [17, 12]). *An arrangement  $\mathcal{A}$  is called **free** with the multiset  $\text{exp}(\mathcal{A}) = \{d_1, \dots, d_\ell\}$  of **exponents** if  $D(\mathcal{A})$  is a free  $\mathbf{S}$ -module with a homogeneous basis  $\{\theta_1, \dots, \theta_\ell\}$  such that  $\text{deg } \theta_i = d_i$  for each  $i$ .*

Remarkably, when an arrangement is free, the exponents turn out to be the roots of the characteristic polynomial by the following result due to Terao.

THEOREM 2.2 (Factorization theorem [18], [12, Theorem 4.137]). *If  $\mathcal{A}$  is free with  $\text{exp}(\mathcal{A}) = \{d_1, \dots, d_\ell\}$ , then*

$$\chi_{\mathcal{A}}(t) = \prod_{i=1}^{\ell} (t - d_i).$$

In general it is very hard to characterize the freeness of an arrangement by combinatorial data. It is only possible in some very special classes of arrangements. One of those is that of graphic arrangements which we shall recall shortly. Let  $G$  be a simple graph (i.e. no loops and no multiple edges) with vertex set  $V_G = \{v_1, \dots, v_\ell\}$  and edge set  $E_G$ . The **graphic arrangement**  $\mathcal{A}_G$  in  $\mathbb{K}^\ell$  is defined by

$$\mathcal{A}_G := \{x_i - x_j = 0 \mid \{v_i, v_j\} \in E_G\}.$$

A simple graph is **chordal** if it does not contain an induced cycle of length greater than three, or  $C_n$ -free<sup>(1)</sup> for all  $n > 3$  in shorthand notation. The freeness of graphic arrangements is completely characterized by chordality.

**THEOREM 2.3** (Freeness and chordality [15], [5, Theorem 3.3]). *Let  $G$  be a simple graph. The graphic arrangement  $\mathcal{A}_G$  is free if and only if  $G$  is chordal.*

**2.2. MAT-FREE ARRANGEMENTS AND THE MAIN RESULT.** Now we recall the concept of MAT-free arrangements following [3]. For  $X \in L(\mathcal{A})$ , we define the **localization** of  $\mathcal{A}$  on  $X$  by

$$\mathcal{A}_X := \{K \in \mathcal{A} \mid X \subseteq K\},$$

and define the **restriction**  $\mathcal{A}^X$  of  $\mathcal{A}$  to  $X$  by

$$\mathcal{A}^X := \{K \cap X \mid K \in \mathcal{A} \setminus \mathcal{A}_X\}.$$

For a positive integer  $n$ , denote  $[n] := \{1, 2, \dots, n\}$ .

**DEFINITION 2.4** (MAT-partition and MAT-free arrangements [3, Lemma 19 and Definition 20]). *Let  $\mathcal{A}$  be a nonempty arrangement. A partition (disjoint union of nonempty subsets)  $\pi = (\pi_1, \dots, \pi_n)$  of  $\mathcal{A}$  is called an **MAT-partition** if the following three conditions hold for every  $k \in [n]$ .*

(MP1)  $\text{rank}(\pi_k) = |\pi_k|$ .

(MP2) *There does not exist  $H' \in \mathcal{A}_{k-1}$  such that  $\bigcap_{H \in \pi_k} H \subseteq H'$ , where  $\mathcal{A}_{k-1} := \pi_1 \sqcup \dots \sqcup \pi_{k-1}$  (disjoint union) and  $\mathcal{A}_0 := \emptyset_\ell$  is the  $\ell$ -empty arrangement.*

(MP3) *For each  $H \in \pi_k$ ,  $|\mathcal{A}_{k-1}| - |(\mathcal{A}_{k-1} \cup \{H\})^H| = k - 1$ .*

*An arrangement is called **MAT-free** if it is empty or admits an MAT-partition.*

All irreducible complex reflection arrangements that are MAT-free were characterized in [3]. The name “MAT-free arrangement” is made by inspiration of the multiple addition theorem due to Abe-Barakat-Cuntz-Hoge-Terao.

**THEOREM 2.5** (Multiple addition theorem (MAT) [1, Theorem 3.1]). *Let  $\mathcal{A}'$  be a free arrangement with  $\text{exp}(\mathcal{A}') = \{d_1, \dots, d_\ell\}_\leq$  (this notation means  $d_1 \leq \dots \leq d_\ell$ ), and  $p \in [\ell]$  the number of maximal exponents. Let  $H_1, \dots, H_q \notin \mathcal{A}'$  be hyperplanes. Define  $\mathcal{A} := \mathcal{A}' \cup \{H_1, \dots, H_q\}$  and  $\mathcal{A}''_j := (\mathcal{A}' \cup \{H_j\})^{H_j}$ . Assume that the following three conditions are satisfied:*

- (1)  $X := H_1 \cap \dots \cap H_q$  is  $q$ -codimensional.
- (2)  $X \not\subseteq \bigcup_{H \in \mathcal{A}'} H$ .
- (3)  $|\mathcal{A}'| - |\mathcal{A}''_j| = d$  ( $j \in [q]$ ).

*Then  $q \leq p$  and  $\mathcal{A}$  is free with  $\text{exp}(\mathcal{A}) = \{d_1, \dots, d_{\ell-p}, d^{p-q}, (d+1)^q\}$ . Here  $d^e$  means  $d$  appears  $e \geq 0$  times in the multiset of exponents.*

The addition of  $\{H_1, \dots, H_q\}$  to the arrangement  $\mathcal{A}'$  resulting in  $\mathcal{A} = \mathcal{A}' \cup \{H_1, \dots, H_q\}$  in Theorem 2.5 is sometimes called an **MAT-step** [10, Definition 2.11]. Thus any MAT-free arrangement is a free arrangement which can be constructed

<sup>(1)</sup>In general, a graph is called  $H$ -free if it does not contain  $H$  as an induced subgraph. It is not to be confused with “MAT-free arrangement.”

inductively from the empty arrangement by MAT-steps. For later use, we state the following corollary (see also [10, Remark 2.15]).

**COROLLARY 2.6.** *Suppose that  $\mathcal{A}$  is MAT-free with MAT-partition  $\pi = (\pi_1, \dots, \pi_n)$ . Then the following statements hold.*

- (1) *For each  $k \in [n]$ ,  $\mathcal{A}_k$  is MAT-free with MAT-partition  $(\pi_1, \dots, \pi_k)$ .*
- (2)  *$\mathcal{A}$  is free with  $\exp(\mathcal{A}) = \{d_1, \dots, d_\ell\}_\leq$  given by the block sizes of the dual partition of  $\pi$ , that is,  $d_i = |\{k \mid |\pi_k| \geq \ell - i + 1\}|$  for all  $i \in [n]$ .*

A remarkable application of the multiple addition theorem (MAT) is an affirmative answer for a conjecture of Sommers–Tymoczko [14] on the freeness of ideal subarrangements of Weyl arrangements.

Let us recall it. Let  $\mathbb{K} = \mathbb{R}$  and  $V = \mathbb{R}^\ell$  with the standard inner product  $(\cdot, \cdot)$ . Let  $\Phi$  be an irreducible (crystallographic) root system in  $V$ , with a fixed positive system  $\Phi^+ \subseteq \Phi$  and the associated set of simple roots  $\Delta := \{\alpha_1, \dots, \alpha_\ell\}$ . For  $\alpha \in \Phi$ , define  $H_\alpha := \{x \in V \mid (\alpha, x) = 0\}$ . For  $\Psi \subseteq \Phi^+$ , the **Weyl subarrangement**  $\mathcal{A}_\Psi$  is defined by  $\mathcal{A}_\Psi := \{H_\alpha \mid \alpha \in \Psi\}$ . In particular,  $\mathcal{A}_{\Phi^+}$  is called the **Weyl arrangement**.

Define the partial order  $\geq$  on  $\Phi^+$  as follows:  $\beta_1 \geq \beta_2$  if  $\beta_1 - \beta_2 \in \sum_{i=1}^\ell \mathbb{Z}_{\geq 0} \alpha_i$ . A subset  $I \subseteq \Phi^+$  is an **ideal** of  $\Phi^+$  if for  $\beta_1, \beta_2 \in \Phi^+$ ,  $\beta_1 \geq \beta_2, \beta_1 \in I$  implies  $\beta_2 \in I$ . For an ideal  $I \subseteq \Phi^+$ , the corresponding Weyl subarrangement  $\mathcal{A}_I$  is called the **ideal subarrangement**.

**THEOREM 2.7** (Ideal MAT-free theorem [1, Theorem 1.1]). *Any ideal subarrangement  $\mathcal{A}_I$  is MAT-free, hence free.*

In this paper we are mainly interested in the graphic arrangements hence root systems of type  $A$ . We will use the following construction of type  $A$  root systems. Let  $\{\epsilon_1, \dots, \epsilon_\ell\}$  be an orthonormal basis for  $V$ , and define  $U := \{\sum_{i=1}^\ell r_i \epsilon_i \in V \mid \sum_{i=1}^\ell r_i = 0\} \simeq \mathbb{R}^{\ell-1}$ . The set  $\Phi(A_{\ell-1}) = \{\pm(\epsilon_i - \epsilon_j) \mid 1 \leq i < j \leq \ell\}$  is a root system of type  $A_{\ell-1}$  in  $U$ , with a positive system  $\Phi^+(A_{\ell-1}) = \{\epsilon_i - \epsilon_j \mid 1 \leq i < j \leq \ell\}$  and the associated set of simple roots  $\Delta(A_{\ell-1}) = \{\alpha_i := \epsilon_i - \epsilon_{i+1} \mid 1 \leq i \leq \ell - 1\}$ . Thus one can see that there is a one-to-one correspondence between the graphic arrangements in  $\mathbb{R}^\ell$  and type  $A_{\ell-1}$  Weyl subarrangements.

In the case of type  $A$ , the ideal subarrangements can be parametrized by unit interval graphs. Recall that a simple graph is a **unit interval graph** if it is chordal and (claw, net, 3-sun)-free (see Figure 1).

**THEOREM 2.8** (Ideals and unit interval graphs e.g. [20, Theorem 16]). *Let  $G$  be a simple graph with  $\ell$  vertices. There exists a vertex-labeling of  $G$  using elements from  $[\ell]$  so that the graphic arrangement  $\mathcal{A}_G$  is an ideal subarrangement of the Weyl arrangement  $\mathcal{A}_{\Phi^+(A_{\ell-1})}$  if and only if  $G$  is a unit interval graph.*

In the study of interplay between arrangements and graphs it is thus natural to ask which graph class corresponds to the MAT-free graphic arrangements? An answer to this question concerns strongly chordal graphs, a class squeezed between the classes of unit interval and chordal graphs.

**DEFINITION 2.9** (Strongly chordal graphs e.g. [6]). *An  $n$ -sun (or trampoline)  $S_n$  ( $n \geq 3$ ) is a (chordal) graph with vertex set  $V_{S_n} = \{u_1, \dots, u_n\} \cup \{v_1, \dots, v_n\}$  and edge set*

$$E_{S_n} = \{\{u_i, u_j\} \mid 1 \leq i < j \leq n\} \cup \{\{v_i, u_j\} \mid 1 \leq i \leq n, j \in \{i, i + 1\}\},$$

where we let  $u_{n+1} = u_1$ . A simple graph is **strongly chordal** if it is chordal and  $n$ -sun-free for  $n \geq 3$ . See Figure 1 for the  $n$ -suns with  $n = 3, 4$ .




FIGURE 1. Some obstructions to unit interval and strongly chordal graphs.

We are ready to state the main result of the paper (whose proof will be presented in the end of §5.2).

**THEOREM 2.10** (MAT-freeness and strong chordality). *Let  $G$  be a simple graph and  $\mathbb{K}$  be a field. The graphic arrangement  $\mathcal{A}_G$  in  $\mathbb{K}^\ell$  is MAT-free if and only if  $G$  is strongly chordal.*

Cuntz–Mücksch [3, Problem 48] asked if the class of MAT-free arrangements is closed under taking localization. An important consequence of our Theorem 2.10 is an affirmative answer for this question in the case of graphic arrangements.

**COROLLARY 2.11.** *MAT-freeness of graphic arrangements is closed under taking localization.*

### 3. MORE ON (STRONGLY) CHORDAL GRAPHS

In this section, we recall some other characterizations of (strongly) chordal graphs that will be useful for our discussion later.

First we collect terminology and notation from graph theory. Let  $G = (V_G, E_G)$  be a simple graph. For  $S \subseteq V_G$ , denote the **(vertex-)induced subgraph** of  $S$  by  $G[S] = (S, E_{G[S]})$ , where  $E_{G[S]} = \{\{u, v\} \in E_G \mid u, v \in S\}$ . If  $v$  is a vertex of  $G$  (sometimes  $v \in G$  is used) then by  $G \setminus v$  we mean the induced subgraph  $G[V_G \setminus \{v\}]$ . For  $F \subseteq E_G$ , define the subgraphs  $G_F := (V_G, F)$  and  $G \setminus F := (V_G, E_G \setminus F)$ . If  $e$  is an edge of  $G$  (sometimes  $e \in G$  is used) then by  $G \setminus e$  we mean the subgraph  $G \setminus \{e\}$ .

An  $n$ -**cycle**  $C_n$  ( $n \geq 3$ ) is a graph with vertex set  $\{v_1, v_2, \dots, v_n\}$  and edge set  $\{\{v_i, v_{i+1}\} \mid 1 \leq i \leq n\}$  where  $v_{n+1} = v_1$ . The 3-cycle is also called a **triangle**. The **length** of a cycle is its number of edges. A **chord** of  $C$  is an edge not in the edge set of  $C$  whose endvertices are in the vertex set.

A **clique** of  $G$  is a subset of  $V_G$  such that every two distinct vertices are adjacent. For each  $v \in V_G$ , its **neighborhood** in  $G$  is  $N_G(v) = \{u \in V_G \mid \{u, v\} \in E_G\}$ . A vertex  $v \in V_G$  is called **simplicial** if its neighborhood is a clique. An ordering  $(v_1, \dots, v_\ell)$  of  $G$  (a linear order on  $V_G$ ) is called a **perfect elimination ordering (PEO)** if  $v_i$  is simplicial in the induced subgraph  $G[\{v_1, \dots, v_i\}]$  for each  $i \in [\ell]$ . The following characterization of chordal graphs is useful to determine the exponents of the corresponding graphic arrangement (e.g. [5, Lemma 3.4]).

**THEOREM 3.1** (Chordality and PEO [7]). *A simple graph is chordal if and only if it has a perfect elimination ordering.*

Let  $a, b \in V_G$  be two distinct vertices which belong to the same connected component of  $G$ . A subset  $S \subseteq V_G$  is called an  $(a, b)$ -**separator** if  $a$  and  $b$  belong to different connected components of  $G[V_G \setminus S]$ . An  $(a, b)$ -separator is **minimal** if it does not

properly contain any  $(a, b)$ -separator. A **minimal vertex separator** is a minimal  $(a, b)$ -separator for some  $a, b \in V_G$ . The following characterization of chordality will also be useful for some inductive arguments later.

**THEOREM 3.2** ([4, Theorem 1]). *A simple graph is chordal if and only if every minimal vertex separator is a clique.*

A **maximal clique** is a clique that it is not a subset of any other clique. A **largest (or maximum) clique** is a clique that has the largest possible number of vertices. Denote by  $\mathcal{K}(G)$  the set of all maximal cliques of  $G$ .

Let  $G$  be a chordal graph. Let  $\mathcal{P}_G$  be the poset consisting of all *possibly-empty* intersections of maximal cliques of  $G$ , i.e.,

$$\mathcal{P}_G = \left\{ \bigcap_{C \in \mathcal{B}} C \mid \emptyset \neq \mathcal{B} \subseteq \mathcal{K}(G) \right\},$$

where the partial order is given by inclusion  $X_1 \leq X_2 \Leftrightarrow X_1 \subseteq X_2$  for  $X_1, X_2 \in \mathcal{P}_G$ . We call  $\mathcal{P}_G$  the **clique intersection poset**<sup>(2)</sup> of  $G$ . Note that  $\mathcal{P}_G$  is a meet-semilattice (not necessarily graded) whose maximal elements are the maximal cliques of  $G$ , and minimal element  $\hat{0} := \bigcap_{C \in \mathcal{K}(G)} C \in \mathcal{P}_G$  is the clique consisting of the dominating vertices<sup>(3)</sup> of  $G$ . We call an element of  $\mathcal{P}_G$  a **node**.

**REMARK 3.3.** Ho-Lee [8, Lemma 2.1] showed that a nonempty subset  $S \subseteq V_G$  is a minimal vertex separator if and only if  $S = C \cap C'$  for distinct maximal cliques  $C$  and  $C'$  forming an edge in some clique tree of  $G$ . Therefore every minimal vertex separator of  $G$  belongs to  $\mathcal{P}_G$ .

A  **$k$ -crown**<sup>(4)</sup> ( $k \geq 1$ ) is a poset on  $\{x_1, \dots, x_k, y_1, \dots, y_k\}$  with relations  $x_i < y_i$  and  $x_i < y_{i+1}$  for all  $1 \leq i \leq k$  (counted modulo  $k$ ) and there are no other relations. See Figure 2 for the  $k$ -crowns with  $k = 3, 4$ . A poset  $P$  is called  **$k$ -crown-free** if there exists no induced subposet<sup>(5)</sup> of  $P$  isomorphic to the  $k$ -crown. The following characterization of strongly chordal graphs will play a crucial role in the proof of the “if” part of our main Theorem 2.10 (strong chordality implies MAT-freeness §5.2).

**THEOREM 3.4** (Strong chordality and clique intersection poset [11, Theorem 1]). *A chordal graph  $G$  is strongly chordal if and only if its clique intersection poset  $\mathcal{P}_G$  is  $k$ -crown-free for all  $k \geq 3$ .*<sup>(6)</sup>

## 4. MAT-FREENESS IMPLIES STRONG CHORDALITY

**4.1. MAT-LABELING OF GRAPHS.** In this subsection, we show that MAT-free graphic arrangements (Definition 2.4) can be completely determined by a special edge-labeling

<sup>(2)</sup>The poset  $\mathcal{P}_G \setminus \{\emptyset\}$  where  $\emptyset$  is the empty set was first defined in [11] where its Hasse diagram is called **clique arrangement**. It is not to be confused with “hyperplane arrangement.”

<sup>(3)</sup>A dominating vertex is a vertex that is adjacent to all other vertices of the graph. The presence of minimum element  $\hat{0}$  (possibly the empty set) is helpful for us, e.g. to define the rank of nodes in Lemma 5.10.

<sup>(4)</sup>The  $k$ -crown here plays a role of **bad  $k$ -cycle** in [11]. More precisely, there exists an induced  $k$ -crown of the clique intersection poset if and only if there exists an induced bad  $k$ -cycle of the clique arrangement. The bottom and top elements of a  $k$ -crown are the starters and terminals of the corresponding bad  $k$ -cycle respectively. Also, it is not to be confused with the  **$k$ -crown graph** which is a graph on  $\{u_1, \dots, u_n\} \cup \{v_1, \dots, v_n\}$  and with an edge from  $u_i$  to  $v_j$  whenever  $i \neq j$ .

<sup>(5)</sup>A poset  $(Q, \leq_Q)$  is an **induced subposet** of a poset  $(P, \leq_P)$  if  $Q \subseteq P$  and for any  $a, b \in Q$  it holds that  $a \leq_Q b$  if and only if  $a \leq_P b$ .

<sup>(6)</sup>It is not hard to see that  $\mathcal{P}_G$  is  $k$ -crown-free for all  $k \geq 3$  if and only if  $\mathcal{P}_G \setminus \{\emptyset\}$  is  $k$ -crown-free for all  $k \geq 3$ .




FIGURE 2. Hasse diagrams of crowns.

of graphs. First we show that the condition of being admitted a “partition” of a (nonempty) MAT-free arrangement is actually implied by the three conditions (MP1), (MP2), and (MP3).

PROPOSITION 4.1. *An arrangement  $\mathcal{A}$  is MAT-free if and only if  $\mathcal{A}$  can be decomposed into a disjoint union of possibly-empty subsets  $\pi_1, \dots, \pi_n$  of  $\mathcal{A}$  satisfying (MP1), (MP2), and (MP3).*

*Proof.* If  $\mathcal{A} = \emptyset$ , then the statement is clear. Suppose  $\mathcal{A} \neq \emptyset$ . We only need to show the “if” part, namely, the existence of a disjoint union of possibly-empty subsets  $\pi_1, \dots, \pi_n$  of  $\mathcal{A}$  satisfying (MP1), (MP2), and (MP3) implies the existence of an MAT-partition of  $\mathcal{A}$ .

First we show that  $\pi_1 \neq \emptyset$ . Suppose to the contrary that  $\pi_1 = \emptyset$ . Thus  $n \geq 2$ . Let  $k \geq 2$  be the minimal integer such that  $\pi_k \neq \emptyset$ . Then (MP3) yields  $0 = |\mathcal{A}_{k-1}| \geq k - 1 \geq 1$ , a contradiction. Thus  $\pi_1 \neq \emptyset$ .

Let  $p := \max\{1 \leq i \leq n \mid \pi_i \neq \emptyset\}$ . We will show that  $\pi_k \neq \emptyset$  for all  $k \in [p]$  which in turn implies that  $(\pi_1, \dots, \pi_p)$  is an MAT-partition of  $\mathcal{A}$ . Suppose to the contrary that there exists  $2 \leq k < p$  such that  $\pi_k = \emptyset$  and choose minimal such  $k$ . Set  $\mathcal{B}' := \pi_1 \cup \dots \cup \pi_{k-1}$ . By definition,  $\mathcal{B}'$  is MAT-free with MAT-partition  $(\pi_1, \dots, \pi_{k-1})$ . Also, the maximal exponents of  $\mathcal{B}'$  are equal to  $k - 1$ . Let  $q := \min\{k < i \leq p \mid \pi_i \neq \emptyset\}$ . Since  $\pi_q \neq \emptyset$ , we can take  $H \in \pi_q$  and write  $\mathcal{B} := \mathcal{B}' \cup \{H\}$ . It is a known fact<sup>(7)</sup> in the theory of free arrangements that  $|\mathcal{B}'| - |\mathcal{B}^H| \leq k - 1$ . However, (MP3) implies  $|\mathcal{B}'| - |\mathcal{B}^H| = q - 1 > k - 1$ , a contradiction. This completes the proof.  $\square$

An **edge-labeled graph** is pair  $(G, \lambda)$  where  $G$  is a simple graph and  $\lambda: E_G \rightarrow \mathbb{Z}_{>0}$  is a map, called **(edge-)labeling**. Now we define a labeling of graphs which characterizes the MAT-freeness of graphic arrangements.

DEFINITION 4.2 (MAT-labelings). *Let  $(G, \lambda)$  be an edge-labeled graph. Let  $\pi_k := \lambda^{-1}(k) \subseteq E_G$  and  $E_k := \pi_1 \sqcup \dots \sqcup \pi_k$  for every  $k \in \mathbb{Z}_{>0}$  and  $E_0 := \emptyset$ . We say that  $\lambda$  is an **MAT-labeling** if the following conditions hold for every  $k \in \mathbb{Z}_{>0}$ .*

- (ML1)  $\pi_k$  is a forest.
- (ML2)  $\text{cl}(\pi_k) \cap E_{k-1} = \emptyset$ . Here  $\text{cl}(F)$  for  $F \subseteq E_G$  denotes the **closure** of  $F$  in the matroid sense. Namely, an edge  $e \in E_G$  is in  $\text{cl}(F)$  when the two endvertices of  $e$  are connected by edges in  $F$ .
- (ML3) Every  $e \in \pi_k$  forms exactly  $k - 1$  triangles (3-cycles) with edges in  $E_{k-1}$ .

<sup>(7)</sup>Here is the precise statement: “Let  $\mathcal{A}$  be an arrangement and let  $H \in \mathcal{A}$ . If  $\mathcal{A}' := \mathcal{A} \setminus \{H\}$  is free with maximal exponent  $m$ , then  $|\mathcal{A}'| - |\mathcal{A}^H| \leq m$ .” We believe that this fact is well known among experts, but we give here a short proof for the sake of completeness. There exists a polynomial  $B$  such that  $\deg B = |\mathcal{A}'| - |\mathcal{A}^H|$  and  $D(\mathcal{A}')\alpha_H$  is contained in the ideal  $(\alpha_H, B) \subseteq S$  [12, Lemma 4.39 and Proposition 4.41]. If  $\deg B > m$ , then  $D(\mathcal{A}')\alpha_H \subseteq (\alpha_H)$ , hence  $D(\mathcal{A}') = D(\mathcal{A})$ . Therefore  $\mathcal{A}$  is free and  $\exp(\mathcal{A}) = \exp(\mathcal{A}')$ . This implies  $|\mathcal{A}'| = |\mathcal{A}|$ , a contradiction. Thus  $|\mathcal{A}'| - |\mathcal{A}^H| = \deg B \leq m$ .

PROPOSITION 4.3. *Let  $G$  be a simple graph and  $\mathbb{K}$  be a field. The graphic arrangement  $\mathcal{A}_G$  in  $\mathbb{K}^\ell$  is MAT-free if and only if  $G$  admits an MAT-labeling.*

*Proof.* This is a translation of Definition 2.4 into graphical terms with the use of Proposition 4.1. □

Thus characterizing the MAT-freeness of graphic arrangements amounts to characterizing the graphs having MAT-labelings. Here are first and simple facts on MAT-labelings. Denote by  $K_\ell$  the complete graph on  $\ell$  vertices.

PROPOSITION 4.4. *If  $\lambda$  is an MAT-labeling of  $K_\ell$ , then  $|\pi_k| = \ell - k$  for all  $k \in [\ell - 1]$ .*

*Proof.* The graphic arrangement  $\mathcal{A}_{K_\ell}$  (in  $\mathbb{R}^\ell$ ) is precisely the Weyl arrangement of type  $A_{\ell-1}$  (also known as the **braid arrangement**) which has exponents  $\{0, 1, 2, \dots, \ell - 1\}$ . Corollary 2.6 completes the proof. □

REMARK 4.5. In fact,  $K_\ell$  always has an MAT-labeling  $\lambda$ . We can see this from Theorem 2.7 as  $K_\ell$  corresponds to a positive system (in particular, an ideal) of a root system of type  $A$ . More precisely,  $\lambda: E_{K_\ell} \rightarrow \mathbb{Z}_{>0}$  is given by  $\lambda(\{v_i, v_j\}) = j - i$  for  $1 \leq i < j \leq \ell$  according to the height<sup>(8)</sup> of positive roots. Also, one can check directly that this labeling satisfies (ML1), (ML2) and (ML3). We will see a different<sup>(9)</sup> (or nonisomorphic) labeling of  $K_\ell$  in Lemma 5.6(2) (see also Figure 7).

It is important to know whether or not a restriction of an MAT-labeling is also an MAT-labeling. The proposition below states that it is enough to check the third condition.

PROPOSITION 4.6. *Let  $\lambda$  be an MAT-labeling of a simple graph  $G$  and  $F \subseteq E_G$ . Then the restriction  $\lambda|_F$  is an MAT-labeling of the subgraph  $G_F = (V_G, F)$  if and only if  $\lambda|_F$  satisfies (ML3).*

*Proof.* Since  $(\lambda|_F)^{-1}(k) = \pi_k \cap F \subseteq \pi_k$  and  $\pi_k$  is a forest,  $(\lambda|_F)^{-1}(k)$  is also a forest. Moreover,  $\text{cl}_{G_F}(\pi_k \cap F) \cap (E_{k-1} \cap F) \subseteq \text{cl}_G(\pi_k) \cap E_{k-1} = \emptyset$ . Thus (ML1) and (ML2) are automatically satisfied. □

LEMMA 4.7. *Let  $\lambda$  be an MAT-labeling of a simple graph  $G$  and  $F_1, F_2 \subseteq E_G$ . If  $\lambda|_{F_1}$  and  $\lambda|_{F_2}$  are MAT-labelings, then  $\lambda|_{F_1 \cup F_2}$  is an MAT-labeling.*

*Proof.* For every subset  $F \subseteq E_G$  and  $k \in \mathbb{Z}_{>0}$ , let  $\pi_k^F := \lambda|_F^{-1}(k) = \pi_k \cap F$ . By Proposition 4.6, it suffices to prove that  $\lambda|_{F_1 \cup F_2}$  satisfies (ML3).

Let  $e \in \pi_k^{F_1 \cup F_2} = \pi_k^{F_1} \cup \pi_k^{F_2}$ . Without loss of generality, we may assume  $e \in \pi_k^{F_1}$ . Since  $\lambda|_{F_1}$  is an MAT-labeling,  $e$  forms at least  $k - 1$  triangles with edges in  $E_{k-1} \cap (F_1 \cup F_2)$ . Moreover, since  $\lambda$  is an MAT-labeling,  $e$  forms at most  $k - 1$  triangles with edges in  $E_{k-1} \cap (F_1 \cup F_2)$ . Therefore  $e$  forms exactly  $k - 1$  triangles with edges in  $E_{k-1} \cap (F_1 \cup F_2)$ . □

4.2. PROOF OF THE IMPLICATION “MAT-FREE  $\Rightarrow$  STRONGLY CHORDAL”. In this subsection we prove the “only if” part of our main Theorem 2.10 (MAT-freeness implies strong chordality).

First we need a few preliminary results. Let  $G = (V_G, E_G)$  be a simple graph. Let  $\chi_G(t)$  be the **chromatic polynomial** of  $G$  (the polynomial that counts the number of proper vertex colorings of  $G$ ). It is known that  $\chi_G(t) = \chi_{\mathcal{A}_G}(t)$  (e.g. [12,

<sup>(8)</sup>The **height** of a positive root  $\beta = \sum_{\alpha \in \Delta} c_\alpha \alpha \in \Phi^+$  is defined by  $\sum_{\alpha \in \Delta} c_\alpha$ .

<sup>(9)</sup>Two edge-labeled graphs  $(G_1, \lambda_1)$  and  $(G_2, \lambda_2)$  are **isomorphic** if there exists a bijection  $\sigma: V_{G_1} \rightarrow V_{G_2}$  such that  $\{v_i, v_j\} \in E_{G_1}$  if and only if  $\{\sigma(v_i), \sigma(v_j)\} \in E_{G_2}$  with  $\lambda_1(\{v_i, v_j\}) = \lambda_2(\{\sigma(v_i), \sigma(v_j)\})$ . If  $G_1 = G_2 = G$ , then we say that two labelings  $\lambda_1$  and  $\lambda_2$  are the same (or isomorphic) if  $(G, \lambda_1)$  and  $(G, \lambda_2)$  are isomorphic.

Theorem 2.88]). Let  $\omega(G)$  denote the **clique number**, the number of vertices in a largest clique of  $G$ . Recall that  $\mathcal{K}(G)$  denotes the set of all maximal cliques of  $G$ .

PROPOSITION 4.8. *Let  $G$  be a chordal graph. Then the following statements hold.*

- (1) *The maximal exponents of  $\mathcal{A}_G$  are equal to  $\omega(G) - 1$ . In addition, the number of maximal exponents of  $\mathcal{A}_G$  equals the number of largest cliques of  $G$ .*
- (2) *If  $\lambda$  is an MAT-labeling of  $G$ , then the endvertices of each  $e \in \pi_n$  where  $n = \omega(G) - 1$  are contained in a unique maximal clique of  $G$ . Furthermore, the map  $\phi : \pi_n \rightarrow \mathcal{K}(G)$  defined by  $\phi(e) =$  the maximal clique containing the endvertices of  $e$  induces a bijection  $\pi_n \simeq \phi(\pi_n)$  and  $\phi(\pi_n) = \{\text{all largest cliques of } G\}$ .*

*Proof.* First we prove (1). When  $G = K_\ell$ ,  $\omega(G) = \ell$  and  $\chi_G(t) = t(t-1) \cdots (t-\ell+1)$ . The assertions clearly hold.

We may assume that  $\ell \geq 3$  and  $G$  is not complete. We proceed by induction on  $\ell = |V_G|$ . Since  $G$  is not complete, there exist two nonadjacent vertices  $a, b \in V_G$ . Let  $S \subseteq V_G$  be a minimal  $(a, b)$ -separator. Then  $V_G$  is decomposed as  $V_G = A \sqcup S \sqcup B$ , where  $a \in A$  and  $b \in B$ . Let  $G_1 := G[A \sqcup S]$  and  $G_2 := G[S \sqcup B]$ . Note that these  $G_1$  and  $G_2$  are chordal. Moreover,  $G[S]$  is complete (Theorem 3.2) and  $\chi_G(t) = \chi_{G_1}(t)\chi_{G_2}(t)/\chi_{G[S]}(t)$  (e.g. [13, Theorem 3]).

Let  $m_1$  and  $m_2$  denote the numbers of cliques consisting of  $\omega(G)$  many vertices in  $G_1$  and  $G_2$ , respectively. Since there is no clique of  $G$  containing both vertices in  $A$  and in  $B$ , the number of largest cliques of  $G$  equals  $m_1 + m_2$ . By the induction hypothesis, the chromatic polynomials of  $G_1$  and  $G_2$  can be expressed as  $\chi_{G_1}(t) = (t - \omega(G) + 1)^{m_1} f(t)$  and  $\chi_{G_2}(t) = (t - \omega(G) + 1)^{m_2} g(t)$ , where  $f(t), g(t) \in \mathbb{Z}[t]$  are the products of some linear factors with roots strictly smaller than  $\omega(G) - 1$ . Therefore  $\chi_G(t) = (t - \omega(G) + 1)^{m_1+m_2} f(t)g(t)/\chi_{G[S]}(t)$ . Since  $\chi_{G[S]}(t) = t(t-1) \cdots (t-|S|+1)$  and  $|S| < \omega(G)$  (Remark 3.3), the maximal exponents of  $G$  are equal to  $\omega(G) - 1$  and the number of maximal exponents of  $G$  is  $m_1 + m_2$ .

Now we prove part (2). If  $n = 0$ , the assertions hold trivially. If  $n = 1$ , then  $G$  is a forest and  $\lambda$  is a constant labeling whose value is 1. Hence the assertions also hold.

Now suppose  $n \geq 2$ . Let  $e \in \pi_n$ . Clearly, there exists  $C \in \mathcal{K}(G)$  such that  $e \in G[C]$ . Suppose that there exist two distinct  $C_1, C_2 \in \mathcal{K}(G)$  such that  $e \in G[C_1 \cap C_2]$ . Since  $\lambda_{E_{G \setminus e}}$  is an MAT-labeling,  $\mathcal{A}_{G \setminus e}$  is free and hence  $G \setminus e$  is chordal. By the maximality of  $C_1, C_2$ , there exist  $u \in C_1 \setminus C_2$  and  $v \in C_2 \setminus C_1$  such that  $\{u, v\} \notin E_G$ . Then  $u, v$  and the endvertices of  $e$  form a 4-cycle which is chordless in  $G \setminus e$ , a contradiction. Thus the endvertices of each edge in  $\pi_n$  are contained in exactly one maximal clique of  $G$ . Therefore the map  $\phi$  is well-defined. Moreover,  $|\phi(\pi_n)| \leq |\pi_n|$ .

Let  $G' := G \setminus \pi_n$ . The restriction  $\lambda|_{E_{G'}}$  is an MAT-labeling of  $G'$  hence  $\mathcal{A}_{G'}$  is MAT-free. Therefore the maximal exponents of  $\mathcal{A}_{G'}$  are equal to  $n - 1$ . By part (1),  $\omega(G') = \omega(G) - 1$ . Thus every largest clique of  $G$  contains the endvertices of at least one edge in  $\pi_n$ . Hence every largest clique of  $G$  belongs to  $\phi(\pi_n)$  and  $|\phi(\pi_n)| \geq |\pi_n|$ . Thus  $\phi(\pi_n)$  is precisely the set consisting of the largest cliques. This completes the proof.  $\square$

In general, a restriction of an MAT-labeling is not an MAT-labeling (e.g. when we restrict to any edge in  $\pi_k$  with  $k > 1$ ). We show below that it is the case for restriction to certain subset. Recall that  $\mathcal{P}_G$  denotes the clique intersection poset of a chordal graph  $G$  (§3).

LEMMA 4.9. *If  $\lambda$  is an MAT-labeling of a chordal graph  $G$ , then the restriction  $\lambda|_{E_{G[X]}}$  is an MAT-labeling of the subgraph  $(V_G, E_{G[X]})$  for any node  $X \in \mathcal{P}_G$ .*

*Proof.* We proceed by induction on  $n = \omega(G) - 1$ . Again it is easily seen that the assertion holds true for  $n = 0, 1$ . Now suppose  $n \geq 2$ . First we consider the case  $X = C$  where  $C$  is a largest clique of  $G$ . Note that by Proposition 4.6, it suffices to prove that  $\lambda|_{E_{G[C]}}$  satisfies (ML3).

Let  $e = \{u, v\} \in \pi_n$  be the unique edge in  $\pi_n$  whose endvertices are contained in  $C$  (Proposition 4.8). The clique  $C$  is the union of two largest cliques  $C' = C \setminus \{v\}$  and  $C'' = C \setminus \{u\}$  of  $G' := G \setminus \pi_n$ . For  $W \in \{C, C', C''\}$ , define  $\pi_k^W := \lambda|_{E_{G[W]}}^{-1}(k) = \pi_k \cap E_{G[W]}$  for  $k \in [n]$ . Note that  $\pi_n^C = \{e\}$  and  $\pi_k^C = \pi_k^{C'} \cup \pi_k^{C''}$  for  $k \in [n-1]$  since the restrictions  $\lambda|_{E_{G[C'()]}}$  and  $\lambda|_{E_{G[C''()]}}$  are MAT-labelings by the induction hypothesis.

To show (ML3) of  $\lambda|_{E_{G[C]}}$ , it suffices to prove that every edge in  $\pi_k^C$  forms at least  $k-1$  triangles with edges in  $E_{k-1} \cap E_{G[C]}$  because every edge in  $\pi_k^C$  forms at most  $k-1$  triangles with edges in  $E_{k-1} \cap E_{G[C]}$  by (ML3) of  $\lambda$ . When  $k = n$ , the edge  $e \in \pi_n^C$  forms  $n-1$  triangles with edges in  $E_{G[C]} \setminus \{e\} \subseteq E_{n-1} \cap E_{G[C]}$ . We are left with  $k \in [n-1]$ . Let  $f \in \pi_k^C$ , then  $f \in \pi_k^{C'} \cup \pi_k^{C''}$ . Without loss of generality, we may assume  $f \in \pi_k^{C'}$ . Then  $f$  forms  $k-1$  triangles with edges in  $E_{k-1} \cap E_{G[C']} \subseteq E_{k-1} \cap E_{G[C]}$ . Therefore  $\lambda|_{E_{G[C]}}$  satisfies (ML3) and hence  $\lambda|_{E_{G[C]}}$  is an MAT-labeling.

Now we treat the case  $X \in \mathcal{P}_G$  is not a largest clique of  $G$ . First we claim that  $E_{G[X]} \cap \pi_n = \emptyset$ . If not, we can find an edge  $e \in E_{G[X]} \cap \pi_n$ . Let  $C$  denote the largest clique such that  $e \in E_{G[C]}$ . By the definition of  $X$ , there exists a maximal clique  $D$  of  $G$  such that  $X \subseteq D \neq C$ . Thus  $e \in E_{G[C \cap D]}$ . Take a vertex  $c \in C \setminus D$ . By the maximality of  $D$ , there exists  $d \in D$  such that  $\{c, d\} \notin E_G$ . Then we obtain a chordless 4-cycle in  $G' = G \setminus \pi_n$  formed by  $c, d$  and the endvertices of  $e$ , which contradicts to the chordality of  $G'$ . Thus  $E_{G[X]} \cap \pi_n = \emptyset$ .

Observe that any maximal but not a largest clique of  $G$  is a maximal clique of  $G'$ . If  $X$  is the intersection of non-largest maximal cliques in  $G$ , then  $X \in \mathcal{P}_{G'}$ . By the induction hypothesis,  $\lambda|_{E_{G[X]}} = (\lambda|_{E_{G'}})|_{E_{G'[X]}}$  is an MAT-labeling. Otherwise,  $X$  is contained in a largest clique of  $G$ , say  $C$ . Let  $e = \{u, v\} \in \pi_n$  be the unique edge in  $\pi_n$  whose endvertices are contained in  $C$ . Then  $C = C' \cup C''$  where  $C' = C \setminus \{v\}$  and  $C'' = C \setminus \{u\}$  are largest cliques of  $G'$ . By the preceding paragraph,  $e \notin E_{G[X]}$  hence either  $u \notin X$  or  $v \notin X$ . Thus either  $X \subseteq C''$  or  $X \subseteq C'$ . Hence we may replace  $C$  by  $C'$  or  $C''$  in the definition of  $X$ . Therefore the node  $X$  is the intersection of some maximal cliques of  $G'$ . Again the induction hypothesis applies.  $\square$

We are ready to prove the main result of this subsection.

**THEOREM 4.10** (MAT-freeness implies strong chordality). *If a simple graph  $G$  admits an MAT-labeling, then  $G$  is strongly chordal.*

*Proof.* Suppose that  $G$  is not strongly chordal. Note that  $G$  is chordal by Proposition 4.3. Then  $G$  contains an  $n$ -sun  $S_n$  (Definition 2.9) as an induced subgraph for some  $n \geq 3$ .

Let  $Z := \{u_1, \dots, u_n\}$  be the central clique of  $S_n$ ,  $T_i := \{u_i, v_i, u_{i+1}\}$  the vertex set of the triangle around  $Z$ , and  $C_i$  a maximal clique of  $G$  containing  $T_i$  for  $i \in [n]$ . Let  $G_0$  be the subgraph of  $G$  with vertex set  $V_{G_0} := C_1 \cup \dots \cup C_n$  and edge set  $E_{G_0} := E_{G[C_1]} \cup \dots \cup E_{G[C_n]}$ . By Lemmas 4.9 and 4.7,  $G_0$  admits an MAT-labeling.

Suppose  $n \geq 4$ . If  $\{u_i, u_j\}$  is an edge of  $G_0$ , then  $\{u_i, u_j\} \in E_{G[C_k]}$  for some  $k \in [n]$ . Therefore both  $u_i$  and  $u_j$  are adjacent to  $v_k$ . This implies  $\{u_i, u_j, v_k\} = T_k$  and hence  $j = i \pm 1$ . Thus the cycle in  $G_0$  consisting of edges  $\{u_1, u_2\}, \dots, \{u_{n-1}, u_n\}, \{u_n, u_1\}$  has no chords and its length is four or more. Therefore  $G_0$  is not chordal, which is a contradiction.

Now we consider  $n = 3$ . Then  $Z = \{u_1, u_2, u_3\}$  is a clique of  $G_0$ . Let  $K$  be a maximal clique of  $G_0$  containing  $Z$ . Since  $u_i \notin C_{i+1}$  for each  $i \in \{1, 2, 3\}$  ( $C_4 = C_1$ ),

the clique  $K$  is neither  $C_1, C_2$ , nor  $C_3$ . Let  $X_i := C_i \cap K \subsetneq K$  for each  $i \in \{1, 2, 3\}$ . Then the restrictions  $\lambda|_{E_{G_0[K]}}$  and  $\lambda|_{E_{G_0[X_i]}}$  are MAT-labelings by Lemma 4.9, where  $\lambda$  denotes an MAT-labeling of  $G_0$ . Since  $E_{G_0} = E_{G[C_1]} \cup E_{G[C_2]} \cup E_{G[C_3]}$ , we have  $E_{G_0[K]} = E_{G_0[X_1]} \cup E_{G_0[X_2]} \cup E_{G_0[X_3]}$ . Therefore

$$\max \left\{ \lambda|_{E_{G_0[K]}}(e) \mid e \in E_{G_0[K]} \right\} = \max \left\{ \lambda|_{E_{G_0[X_i]}}(e) \mid i \in \{1, 2, 3\}, e \in E_{G_0[X_i]} \right\}.$$

Hence  $|K| = |X_i|$  for some  $i \in \{1, 2, 3\}$ , which is a contradiction. □

COROLLARY 4.11. *The  $n$ -sun  $S_n$  admits no MAT-labelings.*

### 5. STRONG CHORDALITY IMPLIES MAT-FREENESS

5.1. MAT-SIMPLICIAL VERTICES AND MAT-PERFECT ELIMINATION ORDERINGS. The proof of the “if” part of our main Theorem 2.10 (strong chordality implies MAT-freeness) requires more effort. We need a deeper understanding of the structure of graphs having MAT-labelings. In this subsection, we develop a fundamental study on such graphs analogous to the theory of (strongly) chordal graphs by introducing MAT- versions of simplicial vertex and perfect elimination ordering.

DEFINITION 5.1 (MAT-simplicial vertices). *Given an edge-labeled graph  $(G, \lambda)$ , a vertex  $v \in V_G$  is said to be **MAT-simplicial** if the following conditions hold.*

- (MS1)  *$v$  is a simplicial vertex of  $G$ , that is, its neighborhood  $N_G(v)$  is a clique of  $G$ .*
- (MS2)  $\{ \lambda(\{u, v\}) \in \mathbb{Z}_{>0} \mid u \in N_G(v) \} = \{1, 2, \dots, \deg_G(v)\}$ , where  $\deg_G(v) = |N_G(v)|$  denotes the degree of  $v$  in  $G$ .
- (MS3) *For any distinct  $u_1, u_2 \in N_G(v)$ ,  $\lambda(\{u_1, u_2\}) < \max\{\lambda(\{u_1, v\}), \lambda(\{u_2, v\})\}$ .*

Next we show the existence of MAT-simplicial vertices in the graphs having MAT-labelings.

LEMMA 5.2. *Let  $(G, \lambda)$  be an edge-labeled graph such that  $|V_G| \geq 2$  and  $\lambda$  is an MAT-labeling of  $G$ .*

- (1) *If  $G = K_\ell$  is a complete graph, then the endvertices of the edge with maximal label are MAT-simplicial.*
- (2) *If  $G$  is noncomplete, then  $(G, \lambda)$  has two nonadjacent MAT-simplicial vertices.*

*Proof.* First we prove part (1). Let  $e_0 = \{u_0, v_0\} \in E_G$  be the edge with maximal label. It suffices to prove that  $v_0$  is MAT-simplicial. First, (MS1) is clear. Next we show (MS2). Note that  $\lambda|_{E_{G \setminus e_0}}$  is an MAT-labeling. Then by Lemma 4.9, the labelings  $\lambda|_{E_{G[C]}}$  and  $\lambda|_{E_{G[X]}}$  are MAT-labelings, where  $C := V_G \setminus \{u_0\}$  and  $X := V_G \setminus \{u_0, v_0\}$ . Comparing the exponents of  $\mathcal{A}_{G[C]}$  and  $\mathcal{A}_{G[X]}$ , we have  $\{ \lambda(\{v, v_0\}) \in \mathbb{Z}_{>0} \mid v \in X \} = [\ell - 2]$ . Since  $\lambda(e_0) = \ell - 1$ ,  $\{ \lambda(\{v, v_0\}) \in \mathbb{Z}_{>0} \mid v \in N_G(v_0) \} = [\ell - 1]$ . Thus (MS2) is satisfied.

Lastly, we show (MS3). Let  $u, v \in X$  and write  $\lambda(\{u, v\}) = k$ . We want to show  $k < \max\{\lambda(\{u, v_0\}), \lambda(\{v, v_0\})\}$ . Since both  $\lambda|_{E_{G[C]}}$  and  $\lambda|_{E_{G[X]}}$  are MAT-labelings, (ML3) implies  $k \leq \max\{\lambda(\{u, v_0\}), \lambda(\{v, v_0\})\}$ . If the equality happens, then it contradicts to (ML2) of  $\lambda|_{E_{G[C]}}$ . Therefore  $k < \max\{\lambda(\{u, v_0\}), \lambda(\{v, v_0\})\}$ . Moreover,  $\max \{ \lambda(\{v, u_0\}) \mid v \in X \} = \ell - 2 < \ell - 1 = \lambda(e_0)$ , since  $\lambda|_{E_{G \setminus v_0}}$  is also an MAT-labeling by Lemma 4.9. Therefore (MS3) holds. Thus  $v_0$  is MAT-simplicial.

Now we prove part (2). We proceed induction on  $\ell = |V_G|$ . If  $\ell = 2$ , then the assertion holds trivially. Suppose  $\ell \geq 3$ . We may assume that  $G$  is connected. So let  $a, b \in V_G$  be nonadjacent vertices and  $S$  a minimal  $(a, b)$ -separator. Then  $S$  is a clique by Theorem 3.2 and  $S \in \mathcal{P}_G$  by Remark 3.3. Hence  $\lambda|_{E_{G[S]}}$  is an MAT-labeling of  $G[S]$  by Lemma 4.9.

Let  $A$  be the vertex set of the connected component containing  $a$  of  $G \setminus S$  and  $B := V_G \setminus (A \cup S)$ . We will show that  $\lambda|_{E_{G[A \cup S]}}$  and  $\lambda|_{E_{G[B \cup S]}}$  are MAT-labelings. By Proposition 4.6, it suffices to show (ML3). Let  $e \in \pi_k \cap E_{G[A \cup S]}$ . Then  $e$  forms at most  $k - 1$  triangles with edges in  $E_{k-1} \cap E_{G[A \cup S]}$  since  $\lambda$  is an MAT-labeling. When  $e \in E_{G[S]}$ ,  $e$  forms exactly  $k - 1$  triangles with edges in  $E_{k-1} \cap E_{G[S]}$ . Therefore  $e$  forms exactly  $k - 1$  triangles with edges in  $E_{k-1} \cap E_{G[A \cup S]}$ . Suppose that at least one endvertex of  $e$  belongs to  $A$ . Then  $e$  cannot form a triangle with a vertex in  $B$ . Hence  $e$  forms exactly  $k - 1$  triangles with edges in  $E_{k-1} \cap E_{G[A \cup S]}$ . Thus  $\lambda|_{E_{G[A \cup S]}}$  is an MAT-labeling. By a similar way, one can prove that  $\lambda|_{E_{G[B \cup S]}}$  is an MAT-labeling.

Next we show that  $A \setminus S$  contains an MAT-simplicial vertex of  $G$ . Note that if  $v \in A \setminus S$  is MAT-simplicial in  $G[A \cup S]$ , then  $v$  is also MAT-simplicial in  $G$  since  $N_G(v) \subseteq A \cup S$ . If  $G[A \cup S]$  is a complete graph, then the endvertices of the edge  $e_0$  in  $G[A \cup S]$  with maximal label is MAT-simplicial in  $G[A \cup S]$  by part (1). Since  $\lambda|_{E_{G[S]}}$  is an MAT-labeling, at least one endvertex of  $e_0$  belongs to  $A \setminus S$  by Proposition 4.4, which is a desired MAT-simplicial vertex. If  $G[A \cup S]$  is not a complete graph, then by the induction hypothesis  $G[A \cup S]$  has two nonadjacent MAT-simplicial vertices. At least one of them belongs to  $A \setminus S$  since  $S$  is a clique. Thus  $A \setminus S$  contains an MAT-simplicial vertex of  $G$ .

Similarly,  $B \setminus S$  contains an MAT-simplicial vertex of  $G$ . Therefore  $G$  has two nonadjacent MAT-simplicial vertices.  $\square$

The following is a first important property of MAT-simplicial vertices.

**PROPOSITION 5.3.** *Let  $(G, \lambda)$  be an edge-labeled graph with  $|V_G| \geq 2$ . Suppose that  $v \in V_G$  is an MAT-simplicial vertex of  $(G, \lambda)$ . The following are equivalent.*

- (1)  $\lambda$  is an MAT-labeling of  $G$ .
- (2)  $\lambda|_{E_{G \setminus v}}$  is an MAT-labeling of  $G \setminus v$ .

*Proof.* First we prove (1)  $\Rightarrow$  (2). Let  $\pi'_k := (\lambda|_{E_{G \setminus v}})^{-1}(k) = \pi_k \cap E_{G \setminus v}$  and  $E'_{k-1} := \pi'_1 \sqcup \dots \sqcup \pi'_{k-1}$  for  $k \in \mathbb{Z}_{>0}$ . By Proposition 4.6, we only need to prove (ML3) of  $\lambda|_{E_{G \setminus v}}$ .

Let  $e \in \pi'_k$ . Since  $e \in \pi'_k \subseteq \pi_k$ ,  $e$  forms exactly  $k - 1$  triangles with edges in  $E_{k-1}$ . These triangles do not contain the vertex  $v$  because by (MS3) the number of edges incident to  $v$  with label less than  $k$  is at most 1. Therefore  $e$  forms exactly  $k - 1$  triangles with edges in  $E'_{k-1}$ . Thus  $\lambda|_{E_{G \setminus v}}$  is an MAT-labeling of  $G \setminus v$ .

Next we prove (2)  $\Rightarrow$  (1). Let  $k \in \mathbb{Z}_{>0}$ . By (MS2),  $v$  is a leaf or an isolated vertex of  $\pi_k$ . Moreover, since  $\pi'_k$  is a forest,  $\pi_k$  is a forest. This shows (ML1).

To show (ML2), suppose  $\text{cl}_G(\pi_k) \cap E_{k-1} \neq \emptyset$  and take  $e \in \text{cl}_G(\pi_k) \cap E_{k-1}$ . Then there exists a cycle  $C$  in  $G$  such that  $e \in C$  and  $C \setminus e \subseteq \pi_k$ . If  $e$  is not incident to  $v$  (in particular,  $v$  is not a vertex of  $C$ ), then  $e \in E'_{k-1}$  and  $C \setminus e \subseteq \pi'_k$ . Therefore  $e \in \text{cl}_{G \setminus v}(\pi'_k) \cap E'_{k-1} = \emptyset$ , a contradiction. Hence  $e$  is incident to  $v$ , and  $C$  contains an edge  $\{v, w\}$  with  $\lambda(\{v, w\}) = k$ . Write  $e = \{u, v\}$ . Then  $\{u, w\} \in E_G$  by (MS1) and  $\lambda(\{u, w\}) < \max\{\lambda(\{u, v\}), \lambda(\{v, w\})\} = k$  by (MS3). Hence  $\{u, w\} \in E'_{k-1}$  (in particular,  $C$  has length at least 4). Moreover,  $\{u, w\}$  and  $C \setminus \{\{v, w\}, e\} \subseteq \pi'_k$  form a cycle and hence  $\{u, w\} \in \text{cl}_{G \setminus v}(\pi'_k) \cap E'_{k-1} = \emptyset$ , a contradiction.

Finally, we prove (ML3). Let  $e \in \pi_k$ . If  $e \in \pi'_k$  (i.e.  $e$  is not incident to  $v$ ), then  $e$  forms exactly  $k - 1$  triangles with some edges in  $E'_{k-1}$ . If one endvertex of  $e$  is not adjacent to  $v$ , then  $e$  and  $v$  cannot form a triangle. If both endvertices of  $e$  are adjacent to  $v$ , then at least one edge of the triangle containing  $e$  and  $v$  has label greater than  $k$  by (MS3). In either case,  $e$  forms exactly  $k - 1$  triangles with some edges in  $E_{k-1}$ . Now consider  $e \in \pi_k \setminus \pi'_k$  (i.e.  $e$  is incident to  $v$ ). By (MS2), we can index the elements of  $N_G(v)$  as  $\{u_1, \dots, u_d\}$  where  $k \leq d = \deg_G(v)$  so that  $e = \{u_k, v\}$  and  $\lambda(\{u_i, v\}) = i$  for every  $i \in [d]$ . Hence  $e$  forms exactly  $k - 1$  triangles given by  $\{u_i, u_k, v\}$  for  $i \in [k - 1]$  with some edges in  $E_{k-1}$ . Thus  $\lambda$  is an MAT-labeling.  $\square$

DEFINITION 5.4 (MAT-PEO). *Given an edge-labeled graph  $(G, \lambda)$ , an ordering  $(v_1, \dots, v_\ell)$  of  $G$  is said to be a **MAT-perfect elimination ordering (MAT-PEO)** of  $(G, \lambda)$  if  $v_i$  is MAT-simplicial in  $(G_i, \lambda_i)$  for each  $i \in [\ell]$ , where  $G_i := G[\{v_1, \dots, v_i\}]$  and  $\lambda_i := \lambda|_{E_{G_i}}$ .*

In particular, any MAT-PEO is a PEO. The theorem below exhibits a strong connection between MAT-labelings and MAT-PEOs which can be seen as an analogue of Theorem 3.1.

THEOREM 5.5 (MAT-labelings and MAT-PEOs). *Given an edge-labeled graph  $(G, \lambda)$ , the following are equivalent.*

- (1)  $\lambda$  is an MAT-labeling of  $G$ .
- (2) There exists an MAT-PEO of  $(G, \lambda)$ .

*Proof.* Both implications can be proved by induction on  $\ell = |V_G|$ . The implication (2)  $\Rightarrow$  (1) is easy thanks to Proposition 5.3. Let us show the converse (which is also not hard). When  $\ell = 1$ , the assertion is true. Suppose  $\ell \geq 2$ . By Lemma 5.2, there exists an MAT-simplicial vertex  $v_\ell$  of  $(G, \lambda)$ . Then  $\lambda|_{E_{G \setminus v_\ell}}$  is an MAT-labeling of  $G \setminus v_\ell$  by Proposition 5.3. By the induction hypothesis,  $(G \setminus v_\ell, \lambda|_{E_{G \setminus v_\ell}})$  has an MAT-PEO  $(v_1, \dots, v_{\ell-1})$ . Thus  $(v_1, \dots, v_\ell)$  is an MAT-PEO of  $(G, \lambda)$ .  $\square$

We complete this subsection by giving two lemmas on (extensions of) MAT-labelings and MAT-PEOs of complete graphs. MAT-labelings of complete graphs will play a crucial role in the next subsection.

LEMMA 5.6. *Let  $G = K_\ell$  be a complete graph and  $W \subseteq V_G$ .*

- (1) *Let  $\lambda$  be an MAT-labeling of  $G$ . If  $(v_1, \dots, v_r)$  is an MAT-PEO of  $(G[W], \lambda|_{E_{G[W]}})$ , then  $(v_1, \dots, v_r)$  can be extended to an MAT-PEO  $(v_1, \dots, v_r, \dots, v_\ell)$  of  $(G, \lambda)$ .*
- (2) *If  $\lambda_W$  is an MAT-labeling of  $G[W]$ , then  $\lambda_W$  can be extended to an MAT-labeling of  $G$ .*

*As a consequence, a complete graph always has an MAT-labeling (and an MAT-PEO) which can be constructed inductively from any vertex of the graph.*

*Proof.* (1) We proceed by induction on  $\ell = |V_G|$ . When  $\ell = 1$ , we have nothing to prove. Now suppose  $\ell \geq 2$ . If  $W = V_G$ , the assertion holds trivially. Suppose  $W \subsetneq V_G$ . Let  $e_0 \in E_G$  be the edge with maximal label. Since  $\max \{ \lambda(e) \in \mathbb{Z}_{>0} \mid e \in E_{G[W]} \} < \lambda(e_0) = \ell - 1$ , at least one endvertex of  $e_0$ , say  $v_\ell$  does not belong to  $W$ . By Lemma 5.2(1),  $v_\ell$  is MAT-simplicial in  $G$ . By the induction hypothesis, there exists an MAT-PEO  $(v_1, \dots, v_r, \dots, v_{\ell-1})$  of  $G \setminus v_\ell$ . Hence  $(v_1, \dots, v_\ell)$  is a desired ordering.

(2) Without loss of generality, we may assume  $\ell \geq 2$  and  $|W| = \ell - 1$ . By Theorem 5.5, there exists an MAT-PEO  $(v_1, \dots, v_{\ell-1})$  of  $(G[W], \lambda_W)$ . Let  $v_\ell$  denote the vertex in  $V_G \setminus W$ . We define a labeling  $\lambda$  of  $G$  by

$$\lambda(e) := \begin{cases} \lambda_W(e) & \text{if } e \in E_{G[W]}; \\ i & \text{if } e = \{v_i, v_\ell\} \text{ for } i \in [\ell - 1]. \end{cases}$$

We will show that  $v_\ell$  is MAT-simplicial in  $(G, \lambda)$ . Firstly, (MS1) is clear since  $G$  is complete. Secondly, we have  $\{ \lambda(\{v_i, v_\ell\}) \mid i \in [\ell - 1] \} = [\ell - 1]$  and hence (MS2) holds. Thirdly, for  $1 \leq i < j < \ell$ , we have  $\lambda(\{v_i, v_j\}) \leq j - 1 < j = \lambda(\{v_j, v_\ell\})$ , which shows (MS3). Therefore  $v_\ell$  is MAT-simplicial in  $(G, \lambda)$ . Thus  $(v_1, \dots, v_\ell)$  is an MAT-PEO in  $(G, \lambda)$  and  $\lambda$  is an MAT-labeling by Proposition 5.3.  $\square$

LEMMA 5.7. *Let  $G = K_\ell$ . Suppose that  $V_G = A \cup B$  and there exist MAT-labelings  $\lambda_A, \lambda_B, \lambda_{A \cap B}$  of  $G[A], G[B], G[A \cap B]$ , respectively such that  $\lambda_A|_{E_{G[A \cap B]}} =$*

$\lambda_B|_{E_{G[A \cap B]}} = \lambda_{A \cap B}$ . Then there exists an MAT-labeling  $\lambda$  of  $G$  such that  $\lambda|_{E_{G[A]}} = \lambda_A$  and  $\lambda|_{E_{G[B]}} = \lambda_B$ .

*Proof.* By Lemma 5.6(1), there exists an MAT-PEO  $(a_1, \dots, a_p)$  of  $G[A \cap B]$  and its extensions  $(a_1, \dots, a_p, a_{p+1}, \dots, a_{p+q})$  of  $G[A]$  and  $(a_1, \dots, a_p, b_1, \dots, b_r)$  of  $G[B]$  (where  $p + q + r = \ell$ ). Define a labeling  $\lambda: E_G \rightarrow \mathbb{Z}_{>0}$  by

$$\lambda(e) := \begin{cases} \lambda_A(e) & \text{if } e \in E_{G[A]}; \\ \lambda_B(e) & \text{if } e \in E_{G[B]}; \\ p + i + j - 1 & \text{if } e = \{a_{p+i}, b_j\} \quad (i \in [q], j \in [r]). \end{cases}$$

We claim that  $\lambda$  is a desired MAT-labeling of  $G$  by induction on  $r$ . If  $r = 0$ , then  $\lambda = \lambda_A$  and hence the claim holds. Suppose  $r \geq 1$ . We will prove that  $b_r$  is MAT-simplicial in  $(G, \lambda)$ . The condition (MS1) is clear. Since  $b_r$  is MAT-simplicial in  $(G[B], \lambda_B)$ ,

$$\{\lambda(\{a_i, b_r\}) \mid i \in [p]\} \cup \{\lambda(\{b_j, b_r\}) \mid j \in [r-1]\} = [p + r - 1].$$

By the definition of  $\lambda$  we have  $\lambda(\{a_{p+i}, b_r\}) = p + i + r - 1$  ( $i \in [q]$ ). Therefore  $\{\lambda(\{v, b_r\}) \mid v \in N_G(b_r)\} = [p + r - 1]$  and hence (MS2) holds.

Next we show (MS3), i.e.  $\lambda(\{u, v\}) < \max\{\lambda(\{u, b_r\}), \lambda(\{v, b_r\})\}$  for any distinct vertices  $u, v \in N_G(b_r)$ . It is clear when  $u, v \in B$  since  $b_r$  is MAT-simplicial in  $(G[B], \lambda_B)$ . Consider the case  $u = a_{p+i} \in A \setminus B, v = a_j \in A \cap B$  for some  $i, j$  with  $p + i > j$ . Then  $\lambda(\{a_{p+i}, a_j\}) \leq p + i - 1 < p + i + r - 1 = \lambda(\{a_{p+i}, b_r\})$  since  $a_{p+i}$  is MAT-simplicial in  $(G, \lambda|_{E_{G[\{a_1, \dots, a_{p+i}\}]}})$ . Now consider  $u = a_{p+i} \in A \setminus B, v = b_j \in B \setminus A$  for some  $i, j$  with  $1 \leq i \leq q$  and  $1 \leq j < r$ . Then  $\lambda(\{a_{p+i}, b_j\}) = p + i + j - 1 < p + i + r - 1 = \lambda(\{a_{p+i}, b_r\})$ . Thus (MS3) holds and  $b_r$  is an MAT-simplicial vertex of  $(G, \lambda)$ .

By the induction hypothesis,  $\lambda|_{E_{G \setminus b_r}}$  is an MAT-labeling. Using Proposition 5.3, we conclude that  $\lambda$  is an MAT-labeling.  $\square$

5.2. PROOF OF THE IMPLICATION “STRONGLY CHORDAL  $\Rightarrow$  MAT-FREE”. In this subsection we prove the “if” part of our main Theorem 2.10 (strong chordality implies MAT-freeness). To find an MAT-labeling for a given strongly chordal graph, our strategy is to find compatible MAT-labelings of the subgraphs induced by all maximal cliques, then combine the constructions by the following “gluing trick.”

**THEOREM 5.8** (“Gluing trick”). *Let  $G$  be a simple graph and suppose that  $V_G = A \cup B, E_G = E_{G[A]} \cup E_{G[B]}$ , and  $A \cap B$  is a clique. Assume that there exist MAT-labelings  $\lambda_A, \lambda_B, \lambda_{A \cap B}$  of  $G[A], G[B], G[A \cap B]$ , respectively such that  $\lambda_A|_{E_{G[A \cap B]}} = \lambda_B|_{E_{G[A \cap B]}} = \lambda_{A \cap B}$ . Define  $\lambda := \lambda_A \cup \lambda_B: E_G \rightarrow \mathbb{Z}_{>0}$  by  $\lambda|_A = \lambda_A, \lambda|_B = \lambda_B$ , i.e.*

$$\lambda(e) := \begin{cases} \lambda_A(e), & \text{if } e \in E_{G[A]}, \\ \lambda_B(e), & \text{if } e \in E_{G[B]}. \end{cases}$$

Then  $\lambda$  is an MAT-labeling of  $G$ .

*Proof.* We proceed by induction on  $\ell = |V_G|$ . When  $\ell \leq 2$  the assertion is trivial. Suppose  $\ell \geq 3$ . We may assume  $A \setminus B \neq \emptyset$ .

We claim that  $G[A]$  has an MAT-simplicial vertex in  $A \setminus B$ . First consider the case  $A$  is a clique. Then any MAT-PEO of  $(G[A \cap B], \lambda_{A \cap B})$  is extended to an MAT-PEO of  $(G[A], \lambda_A)$  by Lemma 5.6(1), which shows the claim. Next suppose that  $A$  is not a clique. Then  $G[A]$  has two nonadjacent MAT-simplicial vertices by Lemma 5.2(2). At least one of them belongs to  $A \setminus B$  since  $A \cap B$  is a clique. Thus, in either case,  $G[A]$  has an MAT-simplicial vertex, say  $v_\ell$  in  $A \setminus B$ . Note that  $v_\ell$  is MAT-simplicial also in  $(G, \lambda)$  since  $N_G(v_\ell) \subseteq E_{G[A]}$ .


By Lemma 5.3,  $\lambda_A|_{E_{G[A] \setminus v_\ell}}$  is an MAT-labeling. Consider the graph  $G \setminus v_\ell$  with the decomposition  $V_{G \setminus v_\ell} = (A \setminus \{v_\ell\}) \cup B$ . By the induction hypothesis, we have that  $\lambda|_{E_{G \setminus v_\ell}}$  is an MAT-labeling. Using Lemma 5.3 again, we conclude that  $\lambda$  is an MAT-labeling of  $G$ .  $\square$

The lemma below describes an important property (of antichains) of the clique intersection poset  $\mathcal{P}_G$ .

LEMMA 5.9. *Let  $G$  be a strongly chordal graph and let  $T \subseteq \mathcal{P}_G$  be an antichain with  $|T| \geq 2$ . Then there exist distinct  $X_0, Y_0 \in T$  such that  $X_0 \cap Y_0 \supseteq X_0 \cap Y$  for all  $Y \in T \setminus \{X_0\}$ .*

*Proof.* Let  $Q$  be the subposet of  $\mathcal{P}_G$  induced by  $\{X \cap Y \in \mathcal{P}_G \mid X, Y \in T\} \supseteq T$ . We will show that there exists a node  $X_0 \in T$  such that  $X_0$  is a leaf of the Hasse diagram  $\mathcal{H}(Q)$  of  $Q$ .


Consider induced subposets of  $Q$  whose Hasse diagrams have the following form:



where  $m \geq 0$  and  $X_i \in T$  for all  $i \in \{0, \dots, m\}$ . Let  $F = \{X_0, Z_1, \dots, Z_m, X_m\} \subseteq Q$  be a poset of the form above such that  $m$  is maximum. Since the Hasse diagram does not change when we replace  $Z_i$  by  $X_{i-1} \cap X_i$  for each  $i \in \{1, \dots, m\}$ , we may assume  $Z_i = X_{i-1} \cap X_i$ .

If  $X_0$  is not a leaf of  $\mathcal{H}(Q)$ , then there exists a node  $Z' \in Q$  such that in  $\mathcal{H}(Q)$ ,  $Z'$  is covered by  $X_0$  and the pair  $\{Z', Z_1\}$  is incomparable. Since every element in  $Q \setminus T$  is the intersection of some elements in  $T$ , there exists a node  $X' \in T$  such that  $X' \supseteq Z'$  and  $X' \neq X_0$ . Similarly, we may assume  $Z' = X' \cap X_0$ .

Since  $m$  is maximum, either  $X'$  contains some  $Z_j \in F$ , or  $Z'$  is contained in some  $X_j \in F \setminus \{X_0\}$ . In either case, we obtain an induced subposet of  $Q$  whose Hasse diagram has the form:



where  $X'_j$  denotes  $X_j$  or  $X'$ .

Since  $G$  is strongly chordal,  $\mathcal{P}_G$  is  $k$ -crown-free for all  $k \geq 3$  by Theorem 3.4. Hence  $j = 1$ . Thus this leads to one of the following induced subposets:



If the first case occurs, then  $Z_1 \subseteq X_0 \cap X' = Z'$ , which contradicts the incomparability of  $Z_1$  and  $Z'$ . When the second case occurs,  $Z' \subseteq X_0 \cap X_1 = Z_1$ , a contradiction again. In summary,  $X_0$  is a leaf of  $\mathcal{H}(Q)$ .

Now let  $Y_0 \cap Y_1 \in Q$  for  $Y_0, Y_1 \in T$  denote a unique node in  $Q$  that is covered by  $X_0$ . We may assume that one of  $Y_0, Y_1$  is not  $X_0$ , say  $Y_0 \neq X_0$ . Then  $X_0 \supseteq X_0 \cap Y_0 \supseteq X_0 \cap Y_0 \cap Y_1 = Y_0 \cap Y_1$ . Hence  $X_0 \cap Y_0 = Y_0 \cap Y_1$ .

Finally let  $Y \in T \setminus \{X_0\}$ . Then  $X_0 \cap Y \in Q$  and  $X_0 \cap Y \subsetneq X_0$ . Since every path from  $X_0 \cap Y$  to  $X_0$  passes through  $Y_0 \cap Y_1$  in  $\mathcal{H}(Q)$ , we must have  $X_0 \cap Y \subseteq X_0 \cap Y_0$ .  $\square$

The lemma below shows the existence of constituent MAT-labelings compatible with the “gluing trick.”

LEMMA 5.10. *Let  $G$  be a strongly chordal graph. Then there exists a family  $\mathcal{F}(\mathcal{P}_G) = \{\lambda_X\}_{X \in \mathcal{P}_G}$  consisting of MAT-labelings  $\lambda_X$  of  $G[X]$  such that  $\mathcal{F}(\mathcal{P}_G)$  is closed under restriction in the sense that  $\lambda_X|_{E_{G[Y]}} = \lambda_Y$  whenever  $X \supseteq Y$ .*

*Proof.* We define the rank of a node  $X \in \mathcal{P}_G$  as the length of a maximum chain connecting  $X$  and  $\hat{0}$ . Let  $\mathcal{P}_G^r$  denote the set consisting of the nodes of rank at most  $r$ . We will show by induction on  $r$  that there exists a family  $\mathcal{F}(\mathcal{P}_G^r) = \{\lambda_X\}_{X \in \mathcal{P}_G^r}$  consisting of MAT-labelings  $\lambda_X$  of  $G[X]$  such that  $\lambda_X|_{E_{G[Y]}} = \lambda_Y$  whenever  $X \supseteq Y$ .

When  $r = 0$ ,  $\mathcal{P}_G^0 = \{\hat{0}\}$ . Since  $G[\hat{0}]$  is a complete graph (or null graph), there exists an MAT-labeling of  $G[\hat{0}]$  by Lemma 5.6.

Now suppose  $r > 0$ . Then by the induction hypothesis there exists a family  $\mathcal{F}(\mathcal{P}_G^{r-1}) = \{\lambda_Y\}_{Y \in \mathcal{P}_G^{r-1}}$  consisting of MAT-labelings  $\lambda_Y$  of  $G[Y]$  such that  $\lambda_{Y_1}|_{E_{G[Y_2]}} = \lambda_{Y_2}$  whenever  $Y_1 \supseteq Y_2$ . We prove the following claim.

CLAIM 5.11. *Let  $X \in \mathcal{P}_G^r \setminus \mathcal{P}_G^{r-1}$  and  $T \subseteq \mathcal{P}_G^{r-1}$  a set consisting of some nodes covered by  $X$ . Then there exists an MAT-labeling  $\lambda_T$  of  $G[\cup_{Y \in T} Y]$  satisfying  $\lambda_T|_{E_{G[Y]}} = \lambda_Y$  for any  $Y \in T$ .*

*Proof of Claim 5.11.* We prove by induction on  $|T|$ . If  $|T| = 1$ , then it is clear. Suppose  $|T| \geq 2$ . By Lemma 5.9, there exist distinct  $X_0, Y_0 \in T$  such that  $X_0 \cap Y_0 \supseteq X_0 \cap Y$  for all  $Y \in T \setminus \{X_0\}$ . By the induction hypothesis on  $|T|$ , there exists an MAT-labeling  $\lambda'$  of  $G[\cup_{Y \in T \setminus \{X_0\}} Y]$  such that  $\lambda'|_{E_{G[Y]}} = \lambda_Y$  for any  $Y \in T \setminus \{X_0\}$ . Note that

$$X_0 \cap \left( \bigcup_{Y \in T \setminus \{X_0\}} Y \right) = \bigcup_{Y \in T \setminus \{X_0\}} (X_0 \cap Y) = X_0 \cap Y_0.$$

By Lemma 5.7, there exists an MAT-labeling  $\lambda_T$  of  $G[\cup_{Y \in T} Y]$  such that  $\lambda_T|_{E_{G[X_0]}} = \lambda_{X_0}$  and  $\lambda_T|_{E_{G[\cup_{Y \in T \setminus \{X_0\}} Y]}} = \lambda'$ . Therefore  $\lambda_T|_{E_{G[Y]}} = \lambda_Y$  for any  $Y \in T$ .  $\square$

Now we return to the proof of Lemma 5.10. Let  $T$  be the set consisting of all nodes covered by  $X$ . Then use Lemma 5.6(2) to extend  $\lambda_T$  to  $\lambda_X$  of  $G[X]$ .  $\square$

We are ready to prove the main result of this subsection.

THEOREM 5.12 (Strong chordality implies MAT-freeness). *If  $G$  is a strongly chordal graph, then  $G$  admits an MAT-labeling.*

*Proof.* By Lemma 5.10, there exists a family  $\{\lambda_X\}_{X \in \mathcal{P}_G}$  consisting of MAT-labelings  $\lambda_X$  of  $G[X]$  such that  $\lambda_X|_{E_{G[Y]}} = \lambda_Y$  whenever  $X \supseteq Y$ . Considering the antichain  $\mathcal{K}(G)$  of  $\mathcal{P}_G$  consisting of the maximal cliques of  $G$ , we can construct an MAT-labeling  $\lambda$  of  $G$  by using Lemma 5.9 and the “gluing trick” (Theorem 5.8). More precisely, we show that for any  $T \subseteq \mathcal{K}(G)$ , there exists an MAT-labeling  $\lambda_T := \cup_{Y \in T} \lambda_Y$  of  $G[\cup_{Y \in T} Y]$  satisfying  $\lambda_T|_{E_{G[Y]}} = \lambda_Y$  for every  $Y \in T$ . This can be done by induction on  $|T|$  very similar to the proof of Claim 5.11. Then take  $T = \mathcal{K}(G)$ .  $\square$

Finally we present the proofs of the main result of the paper and its corollary.

*Proof of Theorem 2.10.* It follows from Theorems 4.10, 5.12 and Proposition 4.3.  $\square$

*Proof of Corollary 2.11.* Taking localization on a flat of a graphic arrangement is equivalent to taking an induced subgraph of the underlying graph. The proof follows from Theorem 2.10 and a simple fact that the class of strongly chordal graphs is closed under taking induced subgraphs.  $\square$

We close this section by giving an example to illustrate the construction in Theorem 5.12.

EXAMPLE 5.13. Let  $G$  be a unit interval graph in Figure 3. Its clique intersection poset  $\mathcal{P}_G$  is given in Figure 4. First we need to find a family  $\mathcal{F}(\mathcal{P}_G) = \{\lambda_X\}_{X \in \mathcal{P}_G}$  consisting of MAT-labelings one for each  $G[X]$  such that  $\mathcal{F}(\mathcal{P}_G)$  is closed under inclusion mentioned in Lemma 5.10. This can be done inductively from the bottom to top starting from the minimum element  $\hat{0}$ . For example, to find a desired MAT-labeling  $\lambda_3 \in \mathcal{F}(\mathcal{P}_G)$  of  $G[X]$  where  $X = \{v_2, v_3, v_4, v_5\}$  provided that the compatible MAT-labelings of  $G[Y]$  for all  $Y$ 's covered by  $X$  (in this case  $\{v_4, v_5\}$  and  $\{v_2, v_3, v_4\}$ ) were given, we use Lemma 5.7 (and Lemma 5.6(2) if  $\cup Y \subsetneq X$ ). Combining the resulting MAT-labelings  $\lambda_i \in \mathcal{F}(\mathcal{P}_G)$  ( $1 \leq i \leq 4$ ) of the maximal cliques by the “gluing trick” (Theorem 5.8) yields an MAT-labeling of  $G$ . Figure 5 shows a gluing  $((\lambda_1 \cup \lambda_2) \cup \lambda_3) \cup \lambda_4$  and how the exponents change in each inductive step, which we call an “exponent growth process.” Note that although MAT-labeling of  $G$  is uniquely determined by  $\lambda_i$ 's, gluing order is not necessarily unique. For example, the gluing  $\lambda_1 \cup (\lambda_2 \cup (\lambda_3 \cup \lambda_4))$  derived from the same method gives the same output but different exponent growth process:  $\{0, 1, 2, 3\} \rightarrow \{0, 1, 2, 3, 3\} \rightarrow \{0, 1, 2, 2, 3, 3\} \rightarrow \{0, 1, 2, 2, 2, 3, 3\}$ .




FIGURE 3. A unit interval (hence strongly chordal) graph  $G$  on 7 vertices with an MAT-labeling constructed by using Theorem 5.12. The corresponding graphic arrangement  $\mathcal{A}_G$  is free with exponents  $\{0, 1, 2, 2, 2, 3, 3\}$ .

### 6. FURTHER REMARKS AND OPEN PROBLEMS

In this section we address some remarks and suggest problems for future research.

- (A) As noted in Introduction, our Theorem 2.10 gives an alternative proof that the ideal graphic arrangements are MAT-free (type A of Theorem 2.7). We give here two examples to illustrate the difference between two methods. The original proof of the ideal MAT-free theorem is inductive on the height of ideals [1, §5], and in each inductive MAT-step only some of maximal exponents get increased by 1. This yields a rigorous exponent growth process hence differs from our construction in Theorem 5.12. For example, the unit interval graph  $G$  in Figure 3 with the given vertex-labeling has its corresponding graphic arrangement  $\mathcal{A}_G$  an ideal subarrangement of the Weyl arrangement  $\mathcal{A}_{\Phi^+(A_6)}$ . The exponent growth process following the ideal MAT-free theorem is given in Figure 6 which differs from that in Figure 5. Our construction




FIGURE 4. The clique intersection poset of the graph in Figure 3 with MAT-labelings  $\lambda_i$  ( $1 \leq i \leq 4$ ) of the maximal cliques constructed by using Lemma 5.10.




FIGURE 5. An exponent growth process for the graph in Figure 3 following the “gluing trick” in Theorem 5.12.

applies also to strongly chordal graphs that are not unit interval graphs. Another way to see the difference between two methods is to consider MAT-labelings of complete graphs, see Remark 4.5, Lemma 5.6(2) and Figure 7.




FIGURE 6. Exponent growth process for the graph in Figure 3 following the ideal MAT-free theorem.




FIGURE 7. Two nonisomorphic MAT-labelings of  $K_4$  constructed by using Lemma 5.6(2) (left) and the ideal MAT-free theorem (right).

- (B) Cuntz–Mücksch [3, Example 22] showed that MAT-freeness is in general not closed under taking restriction. Their example is a non-MAT-free restriction to a hyperplane of the Weyl arrangement of type  $E_6$ . We give here a different example (with a smaller number of hyperplanes) thanks to the fact that the class of strongly chordal graph is not closed under taking edge-contraction. Consider the rising sun (which is a strongly chordal graph) with its edge  $e$  displayed in Figure 8. Taking the contraction of  $e$  results in the 3-sun which is not strongly chordal.




FIGURE 8. The rising sun.

- (C) Strongly chordal graphs are the intersection graphs of unit balls in  $\mathbb{R}$ -trees [9]. Therefore they can be considered as generalization of unit interval graphs in the perspective of intersection graphs.
- (D) Strongly chordal graphs are also known as the graphs having a **strong perfect elimination ordering (SPEO)** [6], i.e. a PEO  $(v_1, \dots, v_\ell)$  with the property that for all  $i < j, k < q$  if  $\{v_i, v_k\}, \{v_i, v_q\}, \{v_j, v_k\}$  are edges, then  $\{v_j, v_q\}$  is an edge. It would be interesting to find a (more direct) connection between SPEO and MAT-PEO.
- (E) If an arrangement  $\mathcal{A}$  is MAT-free, then  $\mathcal{A}$  is *accurate* [10, Theorem 1.2] i.e.  $\mathcal{A}$  is free with  $\exp(\mathcal{A}) = \{d_1, \dots, d_\ell\}_{\leq}$  and there exists for each  $0 \leq p < \ell$  a  $p$ -codimensional flat  $X \in L(\mathcal{A})$  such that  $\mathcal{A}^X$  is free with  $\exp(\mathcal{A}^X) = \{d_1, \dots, d_{\ell-p}\}_{\leq}$ . Characterize the accuracy of graphic arrangements. We are able to show that if  $G$  is an  $n$ -sun, then  $\mathcal{A}_G$  is accurate (but not MAT-free).
- (F) From Theorems 2.10 and 3.4, we now know that the MAT-freeness of graphic arrangements can be characterized by a poset structure, the clique intersection poset of chordal graphs. Define a “clique intersection poset” of an arbitrary (supersolvable) arrangement and characterize the MAT-freeness of the arrangement by the poset. It is related to another question of Cuntz–Mücksch [3, Problem 47] which asked if the MAT-freeness can be characterized by a partial order on the hyperplanes, generalizing the classical partial order (§2.2) on the positive roots of an irreducible root system.

*Acknowledgements.* We thank Professor Takuro Abe for posing a question that whether or not strongly chordal graphs are closed under taking contraction which

guides us to the rising sun example in §6(B). We thank Professor Gerhard Röhrle for suggesting the problem of characterizing accurate graphic arrangements in §6(E). We thank the reviewers for the valuable comments which helped to improve the manuscript.

## REFERENCES

- [1] T. Abe, M. Barakat, M. Cuntz, T. Hoge, and H. Terao, *The freeness of ideal subarrangements of Weyl arrangements*, J. Eur. Math. Soc. **18** (2016), 1339–1348.
- [2] T. Abe and H. Terao, *Multiple addition, deletion and restriction theorems for hyperplane arrangements*, Proc. Amer. Math. Soc. **147** (2019), no. 11, 4835–4845.
- [3] M. Cuntz and P. Mücksch, *MAT-free reflection arrangements*, Electron. J. Combin. **27** (2020), no. 1, article no. 1.28 (28 pages).
- [4] G. A. Dirac, *On rigid circuit graphs*, Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg, vol. 25, Springer, 1961, pp. 71–76.
- [5] P. H. Edelman and V. Reiner, *Free hyperplane arrangements between  $A_{n-1}$  and  $B_n$* , Math. Z. **215** (1994), 347–365.
- [6] M. Farber, *Characterizations of strongly chordal graphs*, Discrete Math. **43** (1983), no. 2–3, 173–189.
- [7] D. Fulkerson and O. Gross, *Incidence matrices and interval graphs*, Pacific J. Math. **15** (1965), no. 3, 835–855.
- [8] C. Ho and R. C. T. Lee, *Counting clique trees and computing perfect elimination schemes in parallel*, Inform. Process. Lett. **31** (1989), no. 2, 61–68.
- [9] M. Kuroda and S. Tsujie, *Unit ball graphs on geodesic spaces*, Graphs Combin. **37** (2021), no. 1, 111–125.
- [10] P. Mücksch and G. Röhrle, *Accurate arrangements*, Adv. in Math. **383** (2021), article no. 107702 (30 pages).
- [11] R. Nevries and C. Rosenke, *Characterizing and computing the structure of clique intersections in strongly chordal graphs*, Discrete Appl. Math. **181** (2015), 221–234.
- [12] P. Orlik and H. Terao, *Arrangements of hyperplanes*, Grundlehren der Mathematischen Wissenschaften 300, Springer-Verlag, Berlin, 1992.
- [13] R. C. Read, *An introduction to chromatic polynomials*, J. Combin. Theory **4** (1968), 52–71.
- [14] E. Sommers and J. Tymoczko, *Exponents for  $B$ -stable ideals*, Trans. Amer. Math. Soc. **358** (2006), 3493–3509.
- [15] R. P. Stanley, *Supersolvable lattices*, Algebra Univers. **2** (1972), 197–217.
- [16] D. Suyama, M. Torielli, and S. Tsujie, *Signed graphs and the freeness of the Weyl subarrangements of type  $B_\ell$* , Discrete Math. **342** (2019), no. 1, 233–249.
- [17] H. Terao, *Arrangements of hyperplanes and their freeness I, II*, J. Fac. Sci. Univ. Tokyo **27** (1980), 293–320.
- [18] ———, *Generalized exponents of a free arrangement of hyperplanes and Shepherd-Todd-Brieskorn formula*, Invent. Math. **63** (1981), no. 1, 159–179.
- [19] M. Torielli and S. Tsujie, *Freeness of hyperplane arrangements between boolean arrangements and Weyl arrangements of type  $B_\ell$* , Electron. J. Combin. **27** (2020), no. 3, article no. 3.10 (15 pages).
- [20] T. N. Tran and A. Tsuchiya, *Worpitzky-compatible subarrangements of braid arrangements and cocomparability graphs*, C. R. Math. Acad. Sci. Paris **359** (2021), no. 6, 665–674.

TAN N. TRAN, Leibniz Universität Hannover, Fakultät für Mathematik und Physik, Institut für Algebra, Zahlentheorie und Diskrete Mathematik, Welfengarten 1, D-30167, Hannover, Germany  
*E-mail* : [tan.tran@math.uni-hannover.de](mailto:tan.tran@math.uni-hannover.de)

SHUHEI TSUJIE, Hokkaido University of Education, Department of Mathematics, Asahikawa, Hokkaido 070-8621, Japan  
*E-mail* : [tsujie.shuhei@hokkyodai.ac.jp](mailto:tsujie.shuhei@hokkyodai.ac.jp)