

Modelling basin-wide variations in Amazon forest productivity – Part 1: Model calibration, evaluation and upscaling functions for canopy photosynthesis

L. M. Mercado^{1,2}, J. Lloyd³, A. J. Dolman⁴, S. Sitch⁵, and S. Patiño^{2,3,6}

¹Centre for Ecology and Hydrology, Wallingford, Oxon, OX10 8BB, UK

²Max Planck Institute for Biogeochemistry, 07745 Jena, Germany

³School of Geography, University of Leeds, LS2 9JT, UK

⁴Free University of Amsterdam, 1081 HV Amsterdam, The Netherlands

⁵Met Office Hadley Centre, JCHMR, Wallingford, Oxon, OX10 8BB, UK

⁶Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Diagonal 27 No. 15-09, Bogotá D.C, Colombia

Received: 22 December 2008 – Published in Biogeosciences Discuss.: 11 March 2009

Revised: 12 June 2009 – Accepted: 26 June 2009 – Published: 28 July 2009

Abstract. Given the importance of Amazon rainforest in the global carbon and hydrological cycles, there is a need to parameterize and validate ecosystem gas exchange and vegetation models for this region in order to adequately simulate present and future carbon and water balances. In this study, a sun and shade canopy gas exchange model is calibrated and evaluated at five rainforest sites using eddy correlation measurements of carbon and energy fluxes.

Results from the model-data evaluation suggest that with adequate parameterisation, photosynthesis models taking into account the separation of diffuse and direct irradiance and the dynamics of sunlit and shaded leaves can accurately represent photosynthesis in these forests. Also, stomatal conductance formulations that only take into account atmospheric demand fail to correctly simulate moisture and CO₂ fluxes in forests with a pronounced dry season, particularly during afternoon conditions. Nevertheless, it is also the case that large uncertainties are associated not only with the eddy correlation data, but also with the estimates of ecosystem respiration required for model validation. To accurately simulate Gross Primary Productivity (GPP) and energy partitioning the most critical parameters and model processes are the quantum yield of photosynthetic uptake, the maximum carboxylation capacity of Rubisco, and simulation of stomatal conductance.

Using this model-data synergy, we developed scaling functions to provide estimates of canopy photosynthetic parameters for a range of diverse forests across the Amazon region, utilising the best fitted parameter for maximum carboxylation capacity of Rubisco, and foliar nutrients (N and P) for all sites.

1 Introduction

The Amazon forest is the largest example of tropical rain forest in the world and it is estimated to be among the most productive ecosystems in the world (Grace et al., 2001). It has received considerable attention due to its potential capacity to act as a source or sink of atmospheric carbon dioxide (Grace et al., 1995a; Lloyd and Farquhar, 1996, 2008; Phillips et al., 1998; Prentice and Lloyd, 1998; Tian et al., 1998; Chambers and Silver, 2004; Lewis et al., 2004). Results from global carbon cycle models have suggested a potential significant effect of Amazonian forests on estimates of the global carbon budget (Prentice and Lloyd, 1998; Tian et al., 1998; Cox et al., 2000). Indeed, the study of Cox et al. (2000), predicts large-scale forest dieback across Amazonia in the 21st century, with associated emissions contributing to an amplification of human-induced climate change. Such results need to be refined using calibrated models that are better constrained by data from the Amazon region. These should lead to improved present and future carbon balances predictions for Amazonia. This is of vital importance



Correspondence to: L. M. Mercado
(lmme@ceh.ac.uk)

for the accuracy of global climate-carbon cycle models and their projections of future atmospheric carbon dioxide concentration and climate.

Although land surface schemes within Global Circulation Models (GCMs) currently represent vegetation in terms of a small set of plant functional types (PFTs) (Sitch et al., 2003), progress is now being made in this respect, associated with a greater appreciation of the physiological diversity of the many different species and the wide diversity of soil types in the region (Fyllas et al., 2009). Thus although the entire Amazon rainforest is taken as one physiologically uniform entity, it is now appropriate for models of Amazon forest gas exchange to incorporate variations in plant physiological characteristics and soil properties.

The increased availability of eddy correlation flux data has been very useful for calibrating and evaluating model ability to simulate measurements and ecosystem responses to different environmental variables. Nevertheless it is important to note some limitations of the method, such as a failure to close the energy balance (Massman and Lee, 2002; Wilson et al., 2002; Baldocchi, 2003; Finnigan et al., 2003) and that the method only estimates the net ecosystem carbon exchange. This comprises two large and opposing fluxes associated with processes that often occur simultaneously: viz. photosynthesis by leaves, and whole-ecosystem respiration, i.e. the combination of plant and heterotrophic respiration. Separating the net flux into its components is non-trivial, and is usually achieved by relating respiration to the net ecosystem exchange measured during night-time. There are two drawbacks in estimating ecosystem respiration using this method: first, daytime ecosystem respiration differs from night-time due to temperature differences, especially at the canopy level. Second, and most importantly, there is a frequent failure of the system to measure night-time respiration fluxes at low wind speed conditions (Aubinet et al., 2002; Massman and Lee, 2002; Pattey et al., 2002; Saleska et al., 2003).

Carbon isotope measurements of plant tissues ($\delta^{13}\text{C}$) place a constraint on models of isotopic discrimination during photosynthesis (Aranibar et al., 2006). These measurements for C_3 plants give an indication of the average intercellular CO_2 concentration (C_i) during photosynthetic periods (Farquhar et al., 1982). This metric has been used in modelling studies to constrain the stomatal conductance parameters (Aranibar et al., 2006).

Under the assumption of nitrogen (N) limitation, leaf photosynthesis is usually modelled based on the measured linearity between photosynthetic capacity and N content per unit leaf area (Hirose and Werger, 1987; Evans, 1993). Leaf photosynthesis is scaled up to canopy level based on the hypothesis that N partitioning within canopies varies with irradiance in such a way as to maximize whole-canopy photosynthesis (Evans, 1993). This optimal approach to N partitioning is frequently used in the modelling community. However, for tropical ecosystems leaf phosphorous (P) rather than

leaf N may be the key nutrient limiting productivity of low-land rainforests, where soils are highly weathered, phosphorous availability is low and nitrogen is relatively abundant (Vitousek and Sanford, 1986). Aside from the importance of many sugar-phosphates in photosynthesis and respiration, phosphorous plays an essential role in energy metabolism because of its presence in important molecules that store energy which are essential to the Calvin cycle. Therefore, deficiencies in phosphorous can limit the rates of RuBP regeneration (RuBP is the CO_2 acceptor molecule in the Calvin cycle) and consequently carbon assimilation (Campbell and Sage, 2006). It is likely the low P concentrations in tropical forest leaves may constrain photosynthetic rate, at least for some tropical forests (Lloyd et al., 2001). Lloyd et al. (2009) have also shown that the “optimal” gradient of N and P within plant canopies is actually less steep than has often been considered the case, especially when overall nutrient concentrations are low.

The motivation behind this study is thus to refine ecosystem gas exchange models in order to better represent the gross carbon uptake of forests in the Amazon Basin, utilising available eddy correlation data and other field observations, for model calibration and evaluation. Then, we can derive upscaling relationships of canopy photosynthesis appropriate for basin-wide simulations. We first assess model performance of simulated carbon uptake and energy partition at five rainforest sites in the Amazon basin using eddy correlation flux data and identify the main limitations of such a model-data evaluation exercise. We take the sun/shade model for photosynthesis (de Pury and Farquhar, 1997) coupled to a stomatal conductance model (Cowan and Farquhar, 1977), previously calibrated at one rainforest site, as described in Mercado et al. (2006). This coupled model is applied at 5 eddy correlation sites across Amazonia. It is calibrated and then evaluated using fluxes of carbon and energy derived from eddy correlation data, and measurements of foliar carbon isotope fractionation (Lloyd et al., 2009; Fyllas et al., 2009). Then, we derive a canopy scaling function to scale GPP to the basin level. Here we relate canopy maximum carboxylation capacity of Rubisco activity (V_{max}) at each site to foliar N. Additionally, given the possibility of phosphorous deficiency in tropical leaves being an important constraint on photosynthesis (Lloyd et al., 2001), relationships between V_{max} and foliar phosphorous are also explored.

2 Methods

2.1 Data

2.1.1 Tower sites

Eddy correlation measurements made above five primary rainforest sites in the Brazilian Amazon (averaged over hourly time steps) with associated meteorological variables,

Table 1. Information about eddy correlation data used from five sites in the Brazilian Amazon.

	Man C14	Man K34	Jaru	Tapajos	Caxiuana
Period used	10/1995–07/1996	10/1999–05/2000 10/2000–12/2000	All 1999	All 2002	1999
Storage Flux	modelled	measured	1999 measured 2000 modelled	measured	Measured few small gaps modelled
Energy balance closure	94%	70%	70%	87% including heat storage from ground and vegetation	70%
Correction applied to data used	Low frequency contributions	no corrections applied to data set used at the time simulations were performed	no corrections applied to data set used at the time simulations were performed	Night time u*	Low frequency contributions

Table 2. General characteristics of five rainforest sites in the Brazilian Amazon.

	Man C14	Man K34	Jaru	Tapajos km 67	Caxiuana
Geographical coordinates	2°35'21.08" S 60°06'53.63" W	2°36'32.67" S 60°12'33.48" W	10°4.706' S 61°56.03' W	03°03' S 54°56' W	01.42° S 51.32° W
Location	Cuieiras Reserve Previously known as ZF2 Manaus, Amazonas	10 km away from the C14 site Manaus, Amazonas	Reserva Jaru 100 km north of Ji-Parana, Rondonia	Tapajos National forest 70 km south of Santarem, Para	1 km north of field station floresta Nacional de Caxiuana, Para
Tower height (m)	41.5	52	62	65	51.5
Mean elevation (m)	100–150	100±50	150–200	90	15
Landscape	Undulating: Plateau and valleys	Undulating: Plateau and valleys	Gently sloping plain	Flat plateau	Flat plateau
Forest type	Terra firme	Terra firme	Terra firme	Terra firme	Terra firme
Canopy height (m)	30–35	30–35	35 up to 45	40 emergent up to 55	~35
LAI (m ² m ⁻²)	5.6	4.4	4	6.7	5.4
Type of soil	Geric carric Ferrasol (Alumic Hyperdystric, Clayic) ^a	Geric Ferrasol (Alumirc Hyperdystric, Clayic) ^a	Red-yellow Acrisol ^b	Oxisol Geric Ferrasol (Alumic, Hyperdystric, Clayic, Xanthic) ^a	Geric, Acric, Ferrasol (Alumic Hyperdystric, Clayic) ^a
Mean temperature (°C)	26.7	26.7	25–27	max [24–32] & min [20–25]	27
Mean precipitation (mm year ⁻¹)	1900–2300	1900–2300	1900	1920	2300
Dry season length (months with rainfall < 100 mm)	Jun–Sep	Jun–Sep	May/June–Sep	Jun–Dec	Jun–Aug
Mean aboveground biomass (Tonne ha ⁻¹)	300–350 ^c	300–350 ^c	220 ^d	349.1 ^c	371.7 ^c
Reference to site and eddy correlation data	Malhi et al. (1998); Malhi et al. (2002)	Araújo et al. (2002); Chambers et al. (2004)	von Randow et al. (2004)	Hutyra et al. (2008); Saleska et al. (2003), From a neighbouring site (km 83): Goulden et al. (2004); Miller et al. (2004); Da Rocha et al. (2004)	Carswell et al. (2002); Iwata et al. (2005)

^a Quesada et al. (2009); ^b von Randow et al. (2004); ^c Baker et al. (2004); ^d Meir et al. (2001)

were used to calibrate and evaluate the model used in this study. A summary of data used and site characteristics is given in Tables 1 and 2, respectively, and site locations are shown in Fig. 1. Information on instruments used for measurements and methods for flux calculations at each site can be found in the original references for these measurements as given in Table 2.

2.1.2 Meteorology

The meteorological data used for model input comes from automatic weather stations located at the top of the towers. Global solar radiation, wind speed and air temperature were available at all towers. To determine atmospheric water vapour content, methodologies differed between sites –



Fig. 1. Locations of rainforest sites with eddy correlation systems used in this study.

values are derived from wet bulb temperatures at Manaus C14 and Caxiuna, from dew point temperatures at Tapajos and from relative humidity measurements at Jaru and Man K34.

2.1.3 Carbon fluxes

Fluxes of carbon dioxide correspond to net ecosystem exchange rates (N_E). In order to determine the net canopy assimilation rate (i.e. total photosynthesis minus daytime canopy leaf respiration R_C), G_P or gross canopy assimilation rate (i.e. total photosynthesis, which includes daytime leaf respiration) (G_{P^*}), it is necessary to consider the non-leaf ecosystem respiration rate R_{E^*} or the total ecosystem respiration rate R_E , respectively, and carbon dioxide accumulated inside the canopy:

$$G_P = -N_E + R_{E^*} \quad (1)$$

$$G_{P^*} = G_P + R_C = -N_E + R_E \quad (2)$$

$$N_E = F_c + \int_0^{h_t} \frac{\partial C_a}{\partial t} dz \quad (3)$$

where F_c is the flux of CO_2 measured by eddy correlation ($\mu\text{mol m}^{-2} \text{s}^{-1}$) and the integrand, often referred to as “the change in canopy storage flux” ($\mu\text{mol m}^{-2} \text{s}^{-1}$), represents the rate of change in the CO_2 concentration (C_a) within the canopy between the forest floor and the eddy correlation measurement height, h_t . Canopy CO_2 storage flux (S_T) is estimated from measurements of within canopy CO_2 concentrations and it is usually measured along with eddy correlation flux data.

Of the five sites, the canopy storage flux was determined from measurements at Manaus K34, Tapajos, Jaru and Caxiuna. Data from Manaus C14 only includes storage mea-

surements for the period October–November 1995. Lacking measurements at Manaus C14 for the remaining period of study and at Jaru, simulated storage fluxes (provided by collaborators) were used.

To calculate photosynthesis from measured eddy correlation N_E , it was necessary to use either G_P or G_{P^*} depending upon the ecosystem respiration available at each individual site. At sites where it was possible to estimate the separate contributions to ecosystem respiration, i.e. from stems and branches, soil and coarse litter (Man K34, Jaru and Man C14), the net assimilation rate, G_P , was used. This term accounts for the balance of the parallel processes of photosynthesis and leaf respiration. At Man K34, Jaru and Man C14 the “observed” G_P was calculated as:

$$G_P = -N_E + R_{E^*} = -N_E + R_S + R_W + R_{CS} \quad (4)$$

R_{E_NL} is the sum of all respiratory contributions from soil (autotrophic and heterotrophic on a ground area basis), coarse litter, stems and branches. R_W represents the respiration contribution from stems and branches, R_S from soil (which includes root and fine litter decomposition at the soil surface) and R_{CS} accounts for coarse litter respiration. At sites where ecosystem respiration was given as a single flux (Caxiuna and Tapajos), Gross photosynthesis, G_{P^*} , is used. At Tapajos and Caxiuna, total ecosystem respiration was available and G_{P^*} was calculated as:

$$G_{P^*} = -N_E + R_E \quad (5)$$

where R_E is defined as R_{E^*} plus respiratory contributions from daytime leaf canopy respiration.

To derive G_P or G_{P^*} at the five tower sites, different methodologies for estimating ecosystem respiration had to be used, these again dependent upon the techniques employed by the original investigators.

Ecosystem respiration for both Manaus sites, and Jaru, was calculated using measurements from its single components (Eq. 4) and for Tapajos and Caxiuna, ecosystem respiration was derived from night time measurements of N_E . A summary of how ecosystem respiration was calculated for each site is included in Table 3.

2.1.4 Foliar carbon isotopes, N and P content and leaf area index

Measurements of foliar carbon isotopic composition as described in Lloyd et al. (2009) and Fyllas et al. (2009) were used to test model predictions of C_i with the exception of the Jaru site where we take data from the study of Ometto et al. (2006).

Foliar N, P (on a dry weight (DW) basis) and leaf mass per unit area for leaves taken from the top of the canopy are as in Fyllas et al. (2009) with nitrogen and phosphorus data from Jaru kindly provided by Martinelli and Nardoto (unpublished data).

Table 3. Calculation of ecosystem respiration for the different sites. All respiration terms are in $\mu\text{mol m}^{-2} \text{s}^{-1}$ (ground area basis). The range of R_E represents seasonal variation.

Site	Method of calculation	Source	R_E
Man C14	Sum of individual components of R_E		
	R_S at soil T	min–max [2.1–4.10]	Chambers et al. (2004)
	R_{CS} (at 25°C)	0.5	Chambers et al. (2000)
	R_w (at 25°C)	1.1	Chambers et al. (2004)
	$R_{C,25}$	3.5	modelled
Man K34	Sum of individual components of R_E		
	R_S at soil T	min–max [2.1–4.10]	Chambers et al. (2004)
	R_{CS} (at 25°C)	0.5	Chambers et al. (2000)
	R_w (at 25°C)	1.1	Chambers et al. (2004)
	$R_{C,25}$	2.5	modelled
Jaru	Sum of individual components of R_E		avg 9.3
	R_S (at soil T)	5.22	Meir et al. (1996)
	R_{CS} (at 25°C)	0.5	Chambers et al. (2000)
	R_w (at 25°C)	0.75	Meir and Grace (2002)
	$R_{C,25}$	2.66	modelled
Tapajos	Inferred from night time fluxes by filtering out data below particular u^* thresholds. Hourly values are calculated as the average of valid nighttime N_E within a centered 5-day wide window, assuming at least 24 valid hours of nighttime N_E were available.	Restrepo-Coupe (2009) Hutyra et al. (2008)	min–max [9–10.6]
Caxiuaana	Inferred from night time fluxes by filtering out data below particular u^* thresholds, and applying the Michaelis Menten equation with a fixed Q_{10} of 2 to soil temperature data to estimate daytime respiration rates	Iwata et al. (2005)	avg 8.6
		modelled	
	R_E (at 25°C)	8.6	
	$R_{C,25}$	2.7	

Leaf area index (LAI) in m^2 of leaves per m^2 of ground was derived from hemispherical photographs using the method of Keeling and Phillips (2007) taken at 4 of the 5 sites during the end of the rainy season (Patiño, unpublished data). LAI at the remaining site, Jaru was taken from the study of Meir et al. (2000).

2.2 Model description

2.2.1 Sun and shade model

The canopy gas exchange sun and shade model used in this study (de Pury and Farquhar, 1997) simulates canopy photosynthesis and energy balance by dividing the canopy into a sunlit and shaded component. This division of sunlit and shaded foliage changes with both the portion of the incoming irradiance that is diffuse, as well as the solar elevation angle, and therefore canopy photosynthetic capacity of the sunlit and shaded portions is dynamic. Sunlit leaves are modelled to receive direct and diffuse radiation while shaded

leaves receive only diffuse radiation. Details about the model and equations for canopy photosynthesis are included in Appendix A.

2.2.2 Radiation absorption and energy balance

Using formulations developed by Goudriaan (1977) as presented in Wang and Leuning (1998) and de Pury and Farquhar (1997), absorption of Photosynthetically Active Radiation (PAR) and near infrared radiation (NIR) was estimated for the sun/shade model. The net energy available to the sunlit and shaded leaf was calculated as the sum of net absorbed PAR, net absorbed NIR and net absorbed long wave by each leaf.

Since leaf temperature is needed to calculate long wave radiation, net isothermal radiation is calculated for the sun and shaded leaves, which are partitioned between the latent and sensible heat fluxes following the isothermal form of the Penman-Monteith equation (Jones, 1992), assuming ground evaporation and soil heat flux to be negligible. This

Table 4. Leaf P and Leaf N in area and dry weight and SLA (Fyllas et al., 2009).

Site	Leaf P (mg g ⁻¹)	Leaf P (g m ⁻²)	Leaf N (mg g ⁻¹)	Leaf N (g m ⁻²)	SLA (cm ² g ⁻¹)
Manaus C14	0.69	0.07	19.93	1.98	102.87
Manaus K34	0.64	0.06	23.10	2.36	96.82
Jaru	0.89 ^a	0.084	27.37 ^a	2.6	105.2 ^b
Caxiuna	0.60	0.05	19.83	1.75	111.17
Tapajos	0.76	0.07	23.48	2.16	103.06

^a Martinelli and Nardoto (unpublished data)

^b Meir et al. (2001)

assumption is based on the observation that typically only 1% of the radiation received at the top of the canopy reaches the forest floor (Shuttleworth, 1989). For example, soil heat fluxes represent only 2% of the daytime net radiation at Tapajos (da Rocha et al., 2004). Since energy stored in biomass and in the canopy air and chemical energy used for CO₂ exchange are small in quantity, these terms were also neglected.

2.2.3 Leaf biochemistry and stomatal conductance

The leaf level photosynthesis model used here includes the biochemistry of C₃ photosynthesis from Farquhar and von Caemmerer (1982) (Eqs. A3 to A6) and stomatal conductance is calculated with the equation from Cowan and Farquhar (1977), called the “Lambda” model (Eq. A7). Description of both models is included in Appendix A.

2.2.4 Isotopic composition of leaves δ¹³C

Isotopic composition of leaves δ¹³C (‰) was calculated for the canopy following Farquhar et al. (1982) as:

$$\delta^{13}\text{C} = \delta_{\text{atm}} - a - (b - a)C_i / C_a \quad (6)$$

where a corresponds to the maximum fractionation due to diffusion of CO₂ in air (4.4‰) and b is the maximum fractionation in the carboxylation reaction (30‰) (Farquhar et al., 1982). δ_{atm} is δ¹³C of the air, taken as −8.0(‰), the late twentieth century mean atmospheric background value (Ehleringer et al., 1987). The ratio C_i / C_a corresponds to simulated daytime hourly values from the whole canopy. The simulated average integral of photosynthetic discrimination during the different seasons analyzed ($\bar{\delta}^{13}\text{C}$) is calculated with Eq. (6) using the photosynthetic flux weighted average (C_i / C_a):

$$\bar{\delta}^{13}\text{C} = -12.4 - 25.6 \frac{\sum_{i=1}^n G_{Pi} * (C_i / C_a)}{\sum_{i=1}^n G_{Pi}} \quad (7)$$

Equation 6 does not include re-fixation of soil and plant respired CO₂. However, measured values of δ¹³C are from top of the canopy leaves and as shown by Lloyd et al. (1996), recycling at this level in the canopy is minimal (<5%).

2.3 Model parameterisation

Canopy level maximum carboxylation activity of Rubisco (V_{max}) was estimated using Eqs. (A1) and (A2). This assumes an exponential decrease in V_{max} with height (or more precisely with cumulative leaf area) as was observed for the profile of V_{max} derived from gas exchange measurements at the Manaus C14 site (Carswell et al., 2000) and parameterised in Mercado et al. (2006) for the sun and shade model for the Manaus C14 site. From the Mercado et al. (2006) parameterisation, V_{max} for the top and bottom 50% of canopy leaves averages 65% and 35% of the V_{max} of the uppermost leaves, respectively. Having defined this distribution of V_{max} with cumulative leaf area, total canopy V_{max} was then calculated at each site as the integral of leaf level V_{max} over cumulative leaf area index, as described in Eq. (A1). Lloyd et al. (2009) discuss gradients in nutrients within Amazon forests in general and conclude that for trees growing on such relatively infertile soils, such shallow within canopy gradients should, indeed, exist.

Canopy level respiration (expressed on a ground area basis) was simulated for all sites using the parameterisation for the sun and shade model from Mercado et al. (2006) as derived from Carswell et al. (2000), and was assumed to decrease exponentially with cumulative leaf area index. Respiration at the top of the canopy was taken as a constant fraction of V_{max} from top leaves (i.e. $0.022 \times V_{\text{max}}$) for all sites (de Pury and Farquhar, 1997). Total canopy respiration R_C was subsequently calculated using the same nitrogen allocation coefficient (k_n) obtained for the exponential regression of V_{max} against cumulative leaf area index (Eq. A2), with a temperature dependence of canopy respiration given by Lloyd and Taylor (1994).

The model was calibrated using a simplex procedure to minimize the error sum of squares of the modelled photosynthesis minus the sum of net ecosystem exchange as measured by eddy correlation plus the estimated ecosystem respiration (Table 3) from a selected subset of data from each site (Nelder and Mead, 1965). Criteria of selection followed the same conditions as in Lloyd et al. (1995) and in Grace et al. (1995b). To avoid the CO₂ flush or so-called morning peak, data before 09:00 was not used for model parameterisation. Storage flux values larger than $10 \mu\text{mol m}^{-2} \text{s}^{-1}$ and smaller than $-10 \mu\text{mol m}^{-2} \text{s}^{-1}$ were also neglected as were data points collected during and after rainfall. Measurements where radiation fluctuated as a result of a moving cloud (i.e. abrupt changes in solar radiation from hour to hour) were also filtered together with aerodynamic conductances lower than $0.1 \mu\text{mol m}^{-2} \text{s}^{-1}$.

The following parameters (from Eqs. A3, A5 and A6) were estimated from the fitting procedure: top of the canopy V_{max} , the ratio $J_{\text{max}} / V_{\text{max}}$ and the canopy level quantum yield (Φ). We decided to fit quantum yield to the observations because a comparison from the light response curves of N_E derived from eddy correlation measurements for the

various sites (Fig. B1), indicates variations in slope of the linear part of the curves at the different sites. The remaining parameters for the sun/shade model, viz. the curvature factor of the light response curve, θ (Eq. A5), and the temperature sensitivity parameters of the electron transport rate (Eq. A8 S_J and H_J) and Rubisco (Eq. A7) and leaf respiration temperature sensitivities (Eq. A10) were taken from Mercado et al. (2006) (i.e. θ and H_J were taken as parameterised in the study of Carswell et al. (2000) at the leaf level and S_J was taken as fitted for the sun/shade model in Mercado et al., 2006). The model was calibrated using eddy correlation data from end of the dry season for all sites. The parameterization for Manaus C14 was taken from Mercado et al. (2006).

Leaf reflectance and transmittance were assumed identical at all sites. Values were taken from the study of Poorter et al. (1995) where measurements were made along a vertical gradient in canopies in a tropical rainforest in Costa Rica.

As in Katul et al. (2000) and Aranibar et al. (2006), we use foliar carbon isotopes measurements from the different sites to help constrain the parameterisation of lambda (λ) model of stomatal conductance (Eq. A11). Initial values of λ were calculated using the Farquhar and Lloyd (1994) version of the model that allows λ to be estimated from \bar{C}_i/\bar{C}_a , the derived from the $\delta^{13}\text{C}$ measurements and the long-term vapour pressure deficit (\bar{D}):

$$\lambda \approx \frac{1.6\bar{D}(C_a - T^*)}{C_c^2 \left[1 - \frac{\bar{C}_i}{C_a}\right]^2} \quad (8)$$

Subsequently, following Aranibar et al. (2006), values of λ that produced similar simulated discrimination as foliar isotope ratios measured at the different sites were chosen.

2.4 Model evaluation

Simulated net carbon uptake and energy partition were evaluated using eddy correlation data for all sites. Because eddy correlation measurements above forest ecosystems often fail energy balance closure tests (Massman and Lee, 2002; Wilson et al., 2002; Baldocchi, 2003; Finnigan et al., 2003), it is more practical to test model performance of energy partition using the evaporative fraction (defined as the ratio between the latent heat flux and the sum of the latent and sensible heat fluxes). There is considerable variability in the hourly calculated evaporative fraction, therefore model performance is evaluated using the mean diurnal cycles of measured and simulated evaporative fraction. In addition, simulated $\delta^{13}\text{C}$ values are compared to measurements of foliar isotopic fractionation as described above. Having an accurate simulated $\delta^{13}\text{C}$ helps to substantiate the results for simulated photosynthetic uptake and at the same time $\delta^{13}\text{C}$ fixes the range within which λ in Eq. (A11) can vary. By increasing or decreasing λ , the partition of energy into sensible and latent heat fluxes is changed and the model can easily fit the measured energy balance, but at the same time the lambda parameter

affects the photosynthetic rate through C_i/C_a and thus the simulated $\delta^{13}\text{C}$. Therefore a realistically simulated $\delta^{13}\text{C}$ is a valuable diagnostic for testing the validity of λ , applied in a fully linked carbon/water/energy exchange model as used here.

Simulations of carbon uptake are compared with the observations using RMSE and regression coefficient (R^2) and slope of the linear regression line of observed and simulated data pairs. Model performance is also evaluated by comparing simulated light response of photosynthesis, diurnal cycle of evaporative fraction and canopy $\delta^{13}\text{C}$ with observations.

3 Results

3.1 Data-model calibration

Best fitting values for top of the canopy V_{\max} , ratio of J_{\max}/V_{\max} and canopy level quantum yield, and non fitted parameters used for all simulations are presented in Table 5.

3.2 Model performance: carbon uptake and energy partition

Statistics of model performance for carbon uptake are included in Table 6 for all sites and seasons studied. The mean light response and diurnal cycles of hourly observations and simulations of G_P (for both Manaus sites and Jaru) and G_{P^*} for Caxiuana and Tapajos are presented in Figs. 2 and 3, respectively. Mean diurnal cycle of observed and simulated evaporative fraction is shown in Fig. 4 and comparison of observed and simulated $\delta^{13}\text{C}$ is shown in Table 7. Results for each study site are discussed separately.

3.2.1 Manaus C14

The sun/shade model simulates reasonably well the three seasons tested within the error bars of the observations at this site (see Table 6 for statistics). However, G_P estimated from eddy correlation measurements during the end of dry season of 1995 and the dry season of 1996 is overestimated by the model by 8% and 10%, respectively. Such overestimation seems to occur mainly during the morning during both seasons (Fig. 3). In 1995 this occurs during high radiation values (PAR higher than $1000 \mu\text{mol m}^{-2} \text{s}^{-1}$) and in 1996 at all radiation levels (Fig. 2, top row). This is associated with the storage fluxes used for calculation of N_E (Eq. 3). For this site, the storage fluxes used were model output. Comparison of simulated storage fluxes against observations (for the available period) show a tendency to under predict the measurements in the morning, implying low estimates of measured N_E , respectively (Fig. B2). This may be one reason for the model to show a tendency to overestimate morning net carbon uptake fluxes.

Simulated energy partition (Fig. 4) shows good agreement with observations at all studied seasons for this site.




Fig. 2. Light response of observed (black) and simulated (grey) values of G_p during the seasons tested.




Fig. 3. Diurnal cycle of observed (black) and simulated (grey) G_p .




Fig. 4. Mean diurnal cycle of observed (black) and simulated (grey) evaporative fraction.

Table 5. Parameters for sun/shade simulations at each site.

	Man C14	Man K34	Jaru	Tapajos	Caxiuana
V_{\max} top canopy ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	52.3	40.	51.89	47.49	32.11
J_{\max}/V_{\max}	1.9	1.9	1.82	2.96	1.9
Quantum yield, Φ	0.4	0.35	0.4	0.16	0.5
Theta, θ^* curvature factor	0.7	0.7	0.7	0.7	0.7
S_J ($\text{J mol}^{-1} \text{K}^{-1}$)	693.124	693.124	693.124	693.124	693.124
H_J (J mol^{-1})	220 000	220 000	220 000	220 000	220 000
Lambda, λ (mol mol^{-1})	1200	2000	2000	5000	1800
LAI ($\text{m}^2 \text{m}^{-2}$)	5.63	4.4	4.0	6.5	5.43
Canopy V_{\max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	185.1	127.8	118.5	100.2	109.7

Table 6. Statistics of model-data comparison for G_p^* during the seasons tested. The regression model is modelled flux = $a \times$ measured (or estimated from eddy correlation) flux.

Site	Season	a	R^2	RMSE*
Manaus C14	End of dry 1995	1.08	0.94	5.29
Manaus C14	Wet 1996	0.96	0.94	5.20
Manaus C14	Dry 1996	1.1	0.95	5.29
Manaus K34	End of dry 1999	1.03	0.96	4.36
Manaus K34	End of dry 2000	1.08	0.96	4.47
Manaus K34	Rain 2000	0.97	0.95	4.36
Jaru	Rain 2000 (all data)	0.96	0.96	4.00
Jaru	Dry 2000 (all data)	1.13	0.96	4.35
Jaru	End of dry 2000 (all data)	1.04	0.95	5.00
Caxiuana	Rain 1999 (all data)	0.96	0.96	4.58
Caxiuana	Dry 1999 (all data)	0.96	0.97	4.24
Caxiuana	End of dry 1999 (all data)	0.95	0.97	4.24
Tapajos	Rain 2002 (all data)	0.90	0.92	5.29
Tapajos	Dry 2002 (all data)	0.95	0.93	5.29

However, under the current parameterisation simulated $\delta^{13}\text{C}$ over predicts the observed $\delta^{13}\text{C}$ (i.e. simulated values are less negative than observations) (Table 7), meaning that the modelled C_i/C_a is lower than what is suggested by the observations. Measured values of $\delta^{13}\text{C}$ could only be explained by the model using a higher λ than $1200 \text{ mol mol}^{-1}$. However, with a higher λ , the model predicts a higher evaporation rate, consequently increasing the simulated evaporative fraction, which leads to an over prediction of observed energy partition (not shown).

3.2.2 Man K34

In terms of carbon uptake, there is good agreement between observations and simulations (see Table 6 for statistics), with only an 8% model overestimation during the end of the dry season in 2000. From the comparison of observed and simulated mean diurnal cycle of net photosynthetic uptake

Table 7. Observed (at Jaru from Ometto et al., 2006 and from Fyllas et al., 2009 for the remaining sites) and simulated $\delta^{13}\text{C}$.

Site		$\delta^{13}\text{C}$ (‰)
Manaus C14	Measured	32.05±2.0
	Simulated with $\lambda=1200 \text{ mol mol}^{-1}$	
	End of dry season 1995	-26.95
Manaus K34	Rainy season 1996	-25.49
	Dry season 1996	-24.56
	Measured	-31.6±1.6
	Simulated with $\lambda=2000 \text{ mol mol}^{-1}$	
Jaru	End of dry season 1999	-29.03
	End of dry season 2000	-29.15
	Rainy season 2000	-30.99
	Measured	-32.3±2.0
	Simulated with $\lambda=2000 \text{ mol mol}^{-1}$	
Caxiuana	Rainy season 2000	-30.47
	Dry season 2000	-28.92
	End of dry season 2000	-29.7
	Measured	-32.5±1.7
	Simulated with $\lambda=1800 \text{ mol mol}^{-1}$	
Tapajos	Rainy season 1999	-29.84
	Dry season 1999	-29.01
	End of dry season 1999	-29.32
	Measured	-31.8±1.3
	Simulated with $\lambda=5000 \text{ mol mol}^{-1}$	
	Rain 2002	-28.95
	Dry 2002	-30.18

(Fig. 3), we do, however, find a bias in the model to simulate higher G_p during afternoons compared with estimates from eddy correlation, which coincides with higher radiation values in the light response curve (Fig. 2). Closer inspection reveals that this is linked to high vapour pressure deficit (D_C) and air temperatures usually after 11:00 local time. This problem could be linked to the stomatal conductance model used not being sufficiently sensitive to such conditions. The sensitivity of the model was tested using different λ , values (results not shown). Nevertheless, the bias towards high D_C and high temperatures persisted.

According to Lloyd et al. (2009) and Fyllas et al. (2009), $\delta^{13}\text{C}$ at the top of the canopy varies within a range of (-30 and -34%) with a mean value of -32.05 ± 2.0 (‰) from leaves collected at all heights, implying the sun and shade simulations are probably satisfactory in terms of modelled carbon exchange during all seasons (Table 7).

Energy partition was reasonably well simulated in the dry and end of dry season with the exception of the rainy season of 2000 where the model under-estimated the measured evaporative fraction (Fig. 4). Note that during the season when the best model data fit was obtained for carbon uptake (i.e. rainy season 2000), with closest simulated $\delta^{13}\text{C}$ to observations, the worst fit for energy partition was obtained.

A sensitivity test showed that the model fit to observed evaporative fraction during the rainy season of 2000 can be improved by artificially increasing the λ parameter from 2000 to 3000 mol mol^{-1} . However, increasing λ to 3000 mol mol^{-1} changes the partition between latent and sensible heat (increasing evaporation rates), improving the model comparison to evaporative fraction but at the cost of a decrease in the simulated $\delta^{13}\text{C}$ (more negative) which leads to model underestimation of $\delta^{13}\text{C}$ with respect to the measurements (results not shown).

3.2.3 Jaru

Simulated carbon uptake was best during the rainy and end of the dry seasons. However, the model overestimated observed carbon uptake by 13% during the dry season (Figs. 2–4). As for Manaus K34, the model has a bias towards overestimating net carbon uptake after midday, which is linked to high D_C and high air temperature conditions. Nevertheless, the model obtains a good comparison to leaf carbon isotope measurements (Table 7). In terms of energy partition, the model agrees with the observations during the dry and end of dry seasons but underestimates evaporation rates during the rainy season. Again, as for Manaus K34, the best model fit to observations for photosynthesis is obtained at the cost of the worst model fit to observations of energy partition.

As the model was calibrated on data from the end of the dry season at all sites, a tendency to overestimate net carbon uptake fluxes during the dry season was not totally unexpected and could be explained as follows by a variety of phenomena, perhaps acting in concert: (1) decreases in canopy conductance due to higher D_C during the dry season, (2) decreases in canopy conductance due to low soil water availability and (3) underestimation of the prescribed soil respiration used to calculate G_P from eddy correlation and (4) reductions of canopy LAI due to leaf fall. Hereafter, each possibility is explored.

(1) Decreases in canopy conductance due to high D_C during the dry season

Measured values of D_C during the dry season at this site in year 2000 were higher than during the rainy and end of dry season. The model over prediction of carbon uptake during

this time could be linked to an insufficient stomatal closure at high D_C . Aiming to improve the model fit to carbon uptake, λ , was therefore artificially decreased from 2000 mol mol^{-1} to 1000 mol mol^{-1} and a very good fit to carbon uptake observations was then obtained. However, this resulted in a reduced simulated evaporative fraction, with values lower than observed (not shown). Under this parameterisation, simulated $\delta^{13}\text{C}$ (-24.65) is much less negative than observed (-32.3 ± 2.0).

(2) Decreases in canopy conductance due to low soil water availability

According to von Randow et al. (2004), during 2000 no apparent direct effects of soil moisture deficit on canopy gas-exchange were detectable. Therefore, it seems unlikely that modelled fluxes overestimation during 2000 are a consequence of lower canopy conductances due to soil moisture deficits. In another modelling approach, Lloyd et al. (1995) used eddy correlation flux data for the same site during dry and wet seasons of 1992 and 1993 and also found no indication of soil moisture deficits on canopy carbon uptake once the D_C effects had been taken into account. Indeed, von Randow et al. (2004) measured the soil moisture profile down to 3.4 m for the period February 1999–September 2002 and found indications of ongoing water extraction from depth, especially during the dry season, this being attributed to continued root water uptake to satisfy transpirational demands, although some lateral drainage was not excluded.

(3) Low values of prescribed soil respiration rate used to calculate G_P from eddy correlation

Ecosystem respiration at this site was estimated using soil respiration that was assumed diurnally and seasonally invariant, as there is little indication of seasonal differences in efflux rates from rainforest soil at Jaru and near constant soil temperatures as already discussed by Meir et al. (1996) and Lloyd et al. (1995). However, it has been shown that for other sites (Davidson et al., 2000; Sotta et al., 2004) soil water may play an important role in regulating the CO_2 efflux from the soil especially in dry periods during and soon after precipitation events. Davidson et al. (2000) reports optimum conditions for soil respiration at intermediate water contents with decreases in respiration at water contents both above and below that value. Therefore, soil respiration could have been higher than assumed during the dry season. However, Meir et al. (2008) conclude that short term moisture limitation can lead to significant reductions of total soil respiration and recognize the lack of a mechanistically and spatially accurately description of the response of soil respiration to variations in soil moisture across the Amazon basin.

(4) Possible reductions of canopy LAI due to leaf fall

A large increase in litterfall at the onset of the dry season has been well documented for the Manaus region (Klinge, 1968; Luizao and Schubart, 1987). If this increase in litterfall gives rise to a decrease in LAI, forest photosynthetic carbon uptake during the dry season would naturally be expected to decrease. To test this possibility, model runs were performed

for the dry season assuming a 10% and 20% decrease in total LAI, respectively (Nepstad et al., 2002; Asner et al., 2004). But even in these cases, simulated G_P overestimates the observations (results not shown).

3.2.4 Caxiuana

For this site a good model fit to observed mean diurnal cycle and light response of net carbon uptake for all seasons was obtained (Figs. 2–4). Simulated $\delta^{13}\text{C}$ values are in within the higher range of the reported values from this site (Table 7). For this site, no D_C bias to carbon uptake was obtained, although there were fewer data points available here than for other sites (Man K34 and Jaru). However, in terms of energy partition, the model overestimates afternoon measurements during all seasons, which could be linked to an insensitive stomatal conductance model to higher afternoon D_C and air temperatures.

3.2.5 Tapajos

The sun/shade model provided a good fit to the mean observed diurnal cycle and light response of gross carbon uptake for this site, (Figs. 2–4) but with a 10% underestimation in observed net carbon uptake during the rainy season. Notice the linear response of simulated photosynthesis at all light levels. This is driven by the obtained parameterisation for this site using the optimization procedure, (i.e. a high J_{\max}/V_{\max} ratio of 2.92, a very low quantum yield of 0.16 and high λ , compared to that obtained at the other sites). A combination of a very high J_{\max}/V_{\max} ratio and low quantum yield of photosynthesis promotes light limitation of photosynthesis in the model at most radiation values. Again, the model has a bias towards over predicting observations at the highest values of D_C and air temperature. Simulated $\delta^{13}\text{C}$ is on the higher end the observed range at this site (Table 7). Additionally, under this parameterisation the model under predicts observed evaporative fraction in the early morning and late afternoon conditions.

4 Scaling up to basin level

An attempt to produce a canopy scale function to scale up simulations to basin level is also presented. Such scaling up is made based on the model calibration and evaluation for the five tower sites. The scaling function can be used to parameterise the model at other sites, using the linear regression obtained from relating the best V_{\max} obtained from parameterisations at each of the 5 sites to its correspondent foliar N. Further, given that (1) tropical forest might be phosphorus instead of nitrogen limited and (2) that phosphorus (P) plays a regulatory role in the partition of the products of photosynthesis directly affecting the activity of the Calvin cycle and therefore affecting the activity of Rubisco carboxylation, relationships between V_{\max} and foliar P, and the ratio of foliar N to foliar P are also explored.

Phosphate limitation to photosynthesis may be linked to a failure in the capacity of starch and sucrose synthesis to match the capacity of the production of triose phosphates in the Calvin cycle, usually when both C_i and light are high. The result is an inadequate rate of release of inorganic phosphate in the chloroplast to recycle the P sequestered in the production of triose phosphates. In this case inorganic phosphorus can limit photosynthesis (Harley and Sharkey, 1991). This has been included in the biochemistry of C3 photosynthesis (Collatz et al., 1991). Implementation of the TPU limitation to photosynthesis in the Collatz et al. (1991) model suggests that the capacity of export or utilization of the products of photosynthesis is approximately equal to the maximum photosynthetic uptake at saturating irradiance, defining the TPU limitation as

$$J_s = V_{\max}/2 \quad (9)$$

where J_s is the export capacity or utilization of the photosynthetic products (most likely sucrose synthesis). Additionally, several studies have shown P deficiency to reduce photosynthesis (Terry and Ulrich, 1973; Sharkey, 1985; Brooks, 1986; Jacob and Lawlor, 1992; Campbell and Sage, 2006). However, the mechanisms by which P deficiency affects photosynthesis are not well understood (Campbell and Sage, 2006). Other possible mechanisms by which P deficiency affects photosynthesis are via reductions in Rubisco activity (Brooks, 1986; Brooks et al., 1988), reductions in the rate of RuBP regeneration, reduction of quantum yield due to photoinhibition caused by reduced efficiency of RuBP regeneration at low irradiances, and reductions in the Calvin cycle activity due to reductions in key regulatory enzymes of the cycle (Brooks, 1986; Campbell and Sage, 2006). From the above limited understanding on the mechanisms by which leaf phosphorus affects photosynthesis, we decided to relate V_{\max} obtained in this study to leaf P for the 5 study sites.

Two methods were used to scale up to the basin level, one based on leaf N and the other on leaf P. Linear relationships between mean leaf N (and leaf P) content per dry weight and per leaf area, using data outlined in Fyllas et al. (2009) were used, and best fitted V_{\max} at each site were used. However, because Φ (Eq. A6) affects to some extent the efficiency of V_{\max} , to relate V_{\max} to leaf N and leaf P under the same conditions, V_{\max} and J_{\max}/V_{\max} were fitted again with a fixed quantum yield of 0.4, with results shown in Table 8.

Using this parameterisation for the sun/shade model, similar results are obtained at all sites compared to what was obtained except at Tapajos. At Tapajos, the model overestimated gross photosynthesis by 15% and 20% during the dry and wet season, respectively, with a better comparison to observed evaporative fraction than obtained earlier (Fig. B3). The reason for this overestimation is linked to a much higher quantum yield (0.4 vs. 0.16) and a much lower J_{\max}/V_{\max} ratio (1.88 vs. 2.96) than obtained in the site-specific simulations.

Table 8. Fitted V_{\max} , J_{\max}/V_{\max} with ($\Phi=0.4$) at each site.

Site	V_{\max} top fitted again ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	original V_{\max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	J_{\max}/V_{\max} fitted again	J_{\max}/V_{\max} original	Φ fixed	Φ original
Manaus C14		52.3		1.92		0.4
Manaus K34	37.8	40	1.9	1.9	0.4	0.35
Jaru	51.89	51.89	1.82	1.82	0.4	0.4
Caxiuana	39.44	32.11	1.83	1.9	0.4	0.5
Tapajos	40	47.49	1.88	1.96	0.4	0.16

Table 9. Statistics of regressions of V_{\max} (with $\Phi=0.4$) in area and DW basis against foliar N and foliar P, with a and b as intercept and slope of the linear relationship obtained.

		a	b	R^2	p -value
V_{\max} ($\mu\text{mol g}^{-1} \text{s}^{-1}$)	vs. P (mg g^{-1})	0.18	0.38	0.3	0.45
V_{\max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	vs. P (g m^{-2})	17.9	386.9	0.41	0.4
V_{\max} ($\mu\text{mol g}^{-1} \text{s}^{-1}$)	vs. N (mg g^{-1})	0.36	0.004	0.03	0.35
V_{\max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	vs. N (g m^{-2})	30.8	6.2	0.08	0.32
V_{\max} ($\mu\text{mol g}^{-1} \text{s}^{-1}$)	vs. N (mg g^{-1})/P (mg g^{-1})	1.2	-0.02	0.65	0.033
V_{\max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	vs. N (g m^{-2})/P (g m^{-2})	95.1	-1.59	0.49	0.052

For all sites V_{\max} in dry weight (DW) and on an area basis from Table 8 were then related to upper canopy average foliar N and P (DW and area basis) as in Fyllas et al. (2009) as shown in Fig. 5 (see data used in Table 4), with statistics from the correspondent linear relationships shown in Table 9.

V_{\max} on an area basis was converted to (DW) using associated measurements of leaf mass per unit area (Fyllas et al., 2009). The highest correlations was then for leaf P with V_{\max} on area basis. In contrast to the relationships of V_{\max} with leaf N which have very low regression coefficients and almost a flat slope indicating no change of V_{\max} with leaf N. Additionally, relationships of V_{\max} in area and DW basis against leaf P and against the ratio of leaf N to leaf P are shown in Fig. 5 (bottom row) with associated statistics given in Table 9. The V_{\max} vs. Leaf P (DW and area basis) and V_{\max} vs. leaf N/leaf P predict increasing V_{\max} and both are statistically significant.

4.1 Model parameterization

Figure 6 shows a comparison of top of the canopy V_{\max} obtained in this study (with quantum yield fixed to 0.4) against available measurements (3 sites) and against V_{\max} calculated from linear regressions of V_{\max} against leaf N obtained from measurements at individual sites (available at 3 sites). The V_{\max} values obtained (circles on Fig. 6) are on the upper end of the range, closest to the measured values at the 3 avail-

able sites (crosses in Fig. 6). The large difference between V_{\max} obtained in this study and those predicted from linear relationships of V_{\max} and leaf N from other sites, are likely due to the relationships reflecting specific conditions of the individual sites. In any case, V_{\max} predicted from the regression based on the Carswell et al. (2000) data is the lowest in Fig. 6. This is because leaf N from this data set are the highest from all data sets considered here. Moreover, leaf N reported from Carswell et al. (2000) for Man C14 are also higher than those measured at the same site by Ometto et al. (2006) and reported by Fyllas et al. (2009).


The quantum efficiency for CO_2 fixation “quantum yield”, used in Eq. (A6) to estimate the irradiance absorbed by photosystem II, was one of the parameters initially fitted at each site. Farquhar et al. (1980) defined a theoretical upper limit to quantum yield (when $f=0$ in Eq. A6) in the absence of oxygenation set by NADPH or by ATP requirement of 0.5 and 0.4, respectively. Equation A6 can be rewritten using α , which intrinsically includes f and Φ , this providing an alternative definition of the intrinsic quantum efficiency for CO_2 uptake. The form of Eq. (10) is the most commonly used in modelling studies and α is the parameter value usually reported.

$$I_2 = I_0 \times (1 - r - t)\alpha \quad (10)$$

Even though the theoretical upper limit for Φ is 0.4–0.5 (i.e. α of 0.125–0.11), measured values of α range considerably;

Table 10. Comparison of α as reported in various studies and obtained in this study. Unit conversion is done assuming that to produce 1 mol of CO₂, 4 electrons are needed.

Study	Use	α [mol CO ₂ mol ⁻¹ PAR]	α [mol electrons mol ⁻¹ photons]	θ
Ehleringer and Bjorkman (1977)	measured	0.073	0.29	
Ehleringer and Pearcy (1983)	measured	0.052	0.22	
Long et al. (1993)	measured	0.093	0.372	
Farquhar et al. (1980)	theoretical	0.125	0.5	
Thornley (2002)	upper limit of α	0.11	0.44	
	used for model parameterisation	0.05	0.2	
Wand and Leuning (1998)	used for model parameterisation	0.096	0.385 ^a	
Harley et al. (1992)	used for model parameterisation	0.06	0.24	
Collatz et al. (1991)	used for model parameterisation	0.08	0.32	0.95
Leuning (1990)	fitted to leaf gas exchange measurements	[0.032–0.066] ^a	0.128–0.264	0.95
Leuning (1995)	used for model parameterisation	0.05 ^a	0.2 ^a	0.95
Leuning et al. (1995)	parameterisation		0.9	
Medlyn et al. (2002)	used for model parameterisation	0.093	0.372	0.9
de Pury and Farquhar (1997)	used for model parameterisation	0.074 ^b	0.29	
	used for model parameterisation	0.106	0.425	0.7
This study	used for model parameterisation			
Man C14 ($\Phi=0.40$)	parameterisation	0.085	0.34	0.7
Man K34 ($\Phi=0.35$)		0.074	0.3	0.7
Jaru ($\Phi=0.35$)		0.074	0.3	0.7
Tapajos ($\Phi=0.30$)		0.063	0.25	0.7
Caxiuana ($\Phi=0.50$)		0.106	0.425	0.7

^a Value including absorbed photon irradiance^b Including leaf absorbance of 0.8**Fig. 5.** Linear regressions obtained from best fitted V_{\max} in DW and in area basis (with $\Phi=0.4$) against foliar N and foliar P in DW at five rainforest sites.

for example 0.073 ($\Phi=0.29$) (Ehleringer and Björkman, 1977), 0.052 ($\Phi=0.22$) (Ehleringer and Pearcy, 1983) and 0.093 ($\Phi=0.372$) (Long et al., 1993). A list of reported α values used or determined in modelling studies is listed in Table 10. This shows that values obtained in this study are indeed within the published range of variation, with α for Caxiuana close to the theoretical upper limit for quantum yield. According to Leuning (1990), when fitting α and curvature parameter θ (Eq. A5) to gas exchange measurements, the obtained variation in α has little physiological meaning due to the interaction between α and θ that occurs during the non-linear curve fitting procedure. The curvature parameter θ has no mechanistic basis (Collatz et al., 1990), although it can be viewed as an indicator of the extent to which co-limitation (of light and Rubisco limitations) of photosynthesis is present with θ approaching 1 meaning that co-limitation is minimized. But a certain level of co-limitation is always present (Woodrow and Berry, 1988). Values of θ in some of the modelling studies (listed in Table 10), include variations between 0.7 and 0.95. Moreover, a recent




Fig. 6. Top leaf V_{max} . Obtained from numerical fitting in this study are circles, measured at Manaus C14, Tapajos and Caxiuana, crosses, calculated from linear regression between measured V_{max} and leaf N at 3 at different canopy levels at 3 of the studied sites: Manaus C14 (Carswell et al., 2000), Caxiuana (Vale et al., 2000) and Tapajos (Domingues et al., 2005) (asterisks, triangles and rhombuses sign, respectively). See linear regressions used for calculations of V_{max} in Table B1.

modelling study using data for Jaru and Manaus C14 optimized parameters of the C_3 photosynthesis model and obtained an Φ of 0.15 and 0.2 and θ of 0.8 and 0.9, respectively (Simon et al., 2005). Domingues et al. (2005) measured leaf photosynthesis at the same Tapajos forest site as used in this study. They adopted a fixed Φ of 0.19 following Ehleringer and Björkman (1977) fitting light response measurements of photosynthesis to a non-rectangular hyperbolic model. The curvature factors derived from such a fitting exercise ranged between 0.45 and 0.9 for most of the cases.

Variation in the parameterised value of Φ for the different sites of this study could thus simply be a result of a mathematical interaction between Φ and θ (here fixed to 0.7). On the other hand, as it was shown in Fig. B1, there do seem to be real differences in the slope of the light response of measured N_E among sites which could have some physiological meaning. For example, differences in efficiency of photosynthetic uptake due to nutrient or other stresses which, from the analysis here not appear to be related to soil water deficits. Detailed measurements would be required to examine variation of Φ across sites. In summary, our fitting procedure produced parameter values within the ranges derived from theory, observation and values adopted by other modelling studies.

4.2 Data

Moreover, model parameters obtained from this fitting exercise and the model evaluation presented in this study are subject to the fidelity of the data used for model parameterisation and evaluation. The main limitations to this data-model parameterisation and evaluation are associated with difficulties in estimating ecosystem respiration and the lack of energy balance closure. Due to spatial and seasonal vari-

ability in soil respiration (Chambers et al., 2004; Silva de Souza, 2004), and with varying meteorological conditions (mainly precipitation regimes) during the measurement and model comparison periods, it is more than likely that there were differences between the measured respiration and those which we used at 3 of the studied sites. Another source of uncertainty in the ecosystem respiration is associated with the leaf respiration term. There is increasing evidence to show that leaf respiration rates are lower when plants are photosynthetically active (Brooks and Farquhar, 1985; Hoefnagel et al., 1998; Atkin et al., 1998, 2000), especially at high leaf temperatures. Unfortunately, biochemical models of gas exchange do not yet include these effects and neither did the leaf level parameters used in this study for comparisons against our obtained parameters. A recent study (Wohlfahrt et al., 2005) assessing daytime ecosystem respiration in a mountain meadow found that a failure to include light inhibition of canopy respiration resulted in an overestimation of daily estimates of ecosystem respiration and hence gross primary productivity from eddy correlation measurements. Their results suggest a reduction in estimated GPP from eddy correlation measurements on the order of 11–13% and 13–17% for a low and high estimate of the simulated maximum leaf-level reduction of dark respiration, respectively. Another study using eddy correlation data from various sites within Europe (Janssens et al., 2001) reports up to a 15% reduction in estimated total ecosystem respiration when considering daytime inhibition of leaf respiration. An alternative method to refine estimates of ecosystem respiration using eddy correlation measurements (Van Gorsel et al., 2007, 2008) is yet to be tested for the Amazon rainforest sites. The method uses the maximum of the sum of the turbulent flux and storage flux of CO_2 in the early evening, and has been shown to be in close agreement with measurements from soil and plant respiration chambers at a moderately complex topography. Such maximum, which is unaffected by advection (Van Gorsel et al., 2008) due to stable stratification conditions after sunset, is used to construct relationships between night time ecosystem exchange measurements, soil temperature and humidity which then can be used to predict ecosystem respiration at all times of day.

Furthermore, there are considerable uncertainties in eddy covariance measurements, especially above rain forest vegetation. For instance, there is a large sensitivity (10–25% annually) to the treatment of low frequencies and non-horizontal flow and delay corrections and data spikes (Kruijt et al., 2004). The total uncertainty in daytime measurements estimated for the Manaus K34 and Jaru site are ± 12 and 32%, respectively (Kruijt et al., 2004). In addition to these uncertainties in N_E , the estimates of total ecosystem respiration and use of modelled storage flux imposes an additional uncertainty to the comparison presented. Further, non-closure of the energy balance in the eddy covariance data (Aubinet et al., 2002; Massman and Lee, 2002) forced us to consider only energy partition in terms of evaporative

fraction. Due to the high hour-to-hour variability in the measurements we also decided to test the model against mean diurnal cycles only.

In order to close the energy balance, fluxes of carbon and energy have been subject to different modes of data processing and post processing correction at each site (Malhi et al., 2002; Finnigan et al., 2003; Gash and Dolman, 2003; van der Molen et al., 2004). With respect to linear detrending, and accounting for low frequency contributions and “angle of attack dependent calibrations”, in 2 out of 5 cases the energy balance improved with the magnitude of the CO₂ fluxes increasing as well (Malhi et al., 2002; von Randow et al., 2004). In general, non-closure of the energy balance at four of the five sites implies a missing CO₂ which may have contributed to the model/data mismatch. However, the extent to which this has occurred is unknown. After recalibration of fluxes for angle of attack and low frequency contributions, the energy and carbon fluxes increased (compared to data without this calibration) (Gash and Dolman, 2003; van der Molen et al., 2004). However, the level of improvement in percentage closure of energy fluxes differs from those improvements for carbon fluxes. Furthermore, A. Araújo (personal communication, 2004) obtained an increase of 8% on average for sensible, latent and CO₂ fluxes after recalculating the fluxes including the angle of attack correction for the Man K34 site.

Higher rates of litterfall towards the end of the wet season/onset of dry season are well documented for some sites in the Amazon region (Klinge, 1968; Luizao and Schubart, 1987; Nepstad et al., 2002; Goulden et al., 2004). However the expected concurrent decrease in LAI has still not been measured. The phenology behind this phenomenon still needs better understanding. Besides satellite derived LAI, in situ measurements of seasonality in LAI are very scarce for the Amazon region. Nevertheless, according to Doughty and Goulden (2008a), in situ LAI seasonality is actually opposite to that derived from satellite measurements. Lacking in situ measurements for most sites, LAI seasonality was not included in any of the sites unless there were available LAI data (used only at the Caxiuana site). Seasonality in LAI may play a major role in driving seasonality of gross uptake at Caxiuana and Tapajos (Carswell et al. 2002; Goulden et al., 2004). A recent study using field work observations at one of the rainforest Tapajos sites (Doughty and Goulden, 2008a) reports that measured seasonality of GPP at this site could only be explained when taking into account, seasonal changes in (i) LAI, (ii) leaf age and (iii) leaf photosynthesis. Another study, measuring leaf level photosynthesis at a well drained and droughted forest at Caxiuana, during various dry and wet periods, concluded that the observed seasonality in maximum rates of photosynthesis at both well drained and droughted forest was not accompanied by any significant change in the photosynthetic parameters V_{\max} , R_D , J_{\max} , in most species, suggesting that the inhibition observed in CO₂ carbon uptake in the dry season was mostly dictated by stom-

atal closure (Vale et al., 2003). More experiments of this type are needed at different sites across the Amazon basin to understand the main factors controlling seasonality in carbon and water exchange at the different sites.

4.3 Model evaluation

Model bias towards overestimation of measured carbon uptake at high D_C and temperature was usually linked to mid-day and afternoon values, the exceptions being Man C14 and Caxiuana. This indicates a model insensitivity to stomatal closure under these conditions. However, the observed decrease in CO₂ uptake at high D_C and air temperatures may not be caused by plant water stress per se because at least for the period studied, the original investigators have indicated that there were no indications of soil water stress at Tapajos (da Rocha et al., 2004; Goulden et al., 2004; Saleska et al., 2003) and at Jaru (von Randow et al., 2004). Manaus K34 was studied here only during the wet and end of dry periods. Furthermore, Mercado (2007) compared the performance of the model used here (also coupled to the sun and shade model) against two versions of the Ball and Berry (Ball et al., 1987) stomatal conductance model. The first version is the modified scheme by Leuning (1990) with dependence on relative humidity, and second version included a dependence on D_C as modified by Leuning (Leuning, 1995). Both stomatal conductance models were coupled to the sun and shade model. Using the “Ball and Berry” or the “Lambda” model made no major difference to the simulated carbon uptake or to the quality of the residuals with either photosynthesis model. Lloyd et al. (1995) obtained similar results when comparing the same two stomatal conductance models coupled to a big leaf model at the Jaru site. This was not unexpected because theoretically both models embody the same principles. They incorporate the well known correlations between photosynthesis and stomatal conductance and describe G_s using similar variables (i.e. D_C or relative humidity at the leaf surface, CO₂ uptake, and CO₂ concentration at the leaf surface). Finally, Tuzet et al. (2003) suggest that model insensitivity to afternoon conditions (high D_C and air temperatures) may result from inadequate/or in this case no coupling of stomatal conductance to the dynamics of water transport from soil to the roots and leaves. This may suggest the need for a stomatal conductance formulation that includes stomatal regulation taking into account both the external environment (demand) but also the dynamics of water movement from the soil/root to the leaf (supply) (Tyree, 2003; Buckley, 2005; Fisher et al., 2007). From soil moisture measurements at a neighboring site to that studied here at Tapajos, da Rocha et al. (2004) found night time water recharge and day time water withdrawal from roots down to 60 cm up to the top 5 cm during dry and wet periods, with soil at 2 m depth remaining moist all year round implying full access to soil water throughout the dry season.

Another explanation for the model mismatch at high temperature and D_C conditions could be linked to decrease in forest photosynthesis due to high temperatures. Doughty and Goulden (2008b) measured leaf level canopy temperatures in parallel with canopy and leaf level photosynthesis and evaporation. From their observations, these authors suggest that the forest (neighbor to the Tapajos km 67 studied here) appears to be close to a high temperature threshold above which photosynthesis declines. However, Lloyd and Farquhar (2008) using observations, model parameterizations of leaf level photosynthesis with known temperature sensitivities, and climate model output conclude that there is no evidence for tropical forest being close to a temperature threshold or to the upper temperature limit for optimum photosynthesis. Their model suggests decreases in photosynthesis above 30°C, which the authors explain as a stomatal conductance response to high D_C . The simulations presented in this study used a value of 39°C as the maximum optimum temperature dependency of electron transport rate, obtained fitting eddy correlation measurements to the sun and shade model at the Manaus C14 site (Mercado et al., 2006). Lloyd et al. (1995) obtained a value of 43°C when fitting a big leaf model to eddy correlation measurements at the Jaru site. More measurements of leaf level temperature response at different rainforest sites is needed to check if temperature threshold measured at Tapajos (Doughty and Goulden, 2008b) prevails elsewhere. Finally, another possible explanation for the decline in photosynthesis and stomatal closure at high D_C and temperature conditions might be related to a reduction in the activation state of Rubisco due to the high temperature being above its thermal optimum (Sage and Kubien, 2007). If this was the case, the capacity of deactivated Rubisco would be needed to be incorporated into the photosynthesis models.

The underestimation of simulated latent heat fluxes during the rainy season at 3 of 5 sites could mean the model needs larger simulated stomatal conductances to account for the observations. For instance, the best simulated energy partitioning was obtained when the fit for carbon uptake was on the upper range (Table 5). This was the case for end of dry and dry seasons at Manaus C14, end of dry season at Manaus K34 and the dry season in Jaru. On the other hand, when the fit for carbon uptake was on the lower range there was usually a model tendency to underestimate the mean observed diurnal cycle of evaporative fraction. This was the case during all seasons at Caxiuana, Tapajos and rainy seasons at Manaus C14 and K34 and rainy season at Jaru. Furthermore, at Tapajos, after refitting with a fixed quantum yield of 0.4, the fit for carbon uptake was overestimated observations by 20 and 15% during the rainy and wet seasons, respectively, whilst obtaining a better fit to evaporative fraction (Fig. B3).

Furthermore, a larger simulated stomatal conductance could be obtained either by increasing the λ parameter from the stomatal conductance model or by increasing the simulated assimilation rate. Sensitivity analysis showed that by

increasing λ , the C_i/C_a ratio increased and therefore the simulated $\delta^{13}\text{C}$ decreased (becoming more negative). In general, this led to model underestimation of $\delta^{13}\text{C}$. The second option would be to increase simulated G_P which automatically would lead to model overestimation with respect to the observations. However, associated with a failure to close the energy balance is a probably also a non-measured CO_2 flux signal (Aranibar et al., 2006), implying a higher G_P than currently measured. A consequence of a higher simulated G_P is a concurrent increase in simulated latent heat fluxes which would improve the model comparison to evaporative fraction measurements.

Generally, besides the D_C bias towards overestimation (found at 3 of 5 sites), the sun and shade model performed well in simulating the carbon uptake at all tower sites, a finding supported by the $\delta^{13}\text{C}$ evaluation, with a model tendency to underestimate energy partition at some sites during the rainy season. Morales et al. (2005) evaluated 4 process-based models against eddy correlation flux data at 15 European sites and found no single model performed well for both carbon and water fluxes. According to Morales et al. (2005), two models performed better at simulating carbon fluxes, the remaining two models in simulating water fluxes. Another modelling study using the land surface scheme of the Hadley Centre GCM carried out an evaluation with the same data set as used here for the Man K34 site, and obtained better results for carbon uptake than for energy partition (Mercado et al., 2007).

4.4 Scaling up to basin level

Unfortunately there were only five data points to derive relationships between V_{max} and leaf N and leaf P, and the slope and high correlation coefficient for the correlations between V_{max} and leaf P (in area and DW basis) is due to a one data point which has the highest leaf P in the data set, corresponding to the Jaru site. However, the linear relationships of V_{max} against leaf P (in area basis) from this study are comparable to those from leaf level measurements from tropical forest in West Africa with more fertile soil types (Domingues, unpublished data) (Fig. 7).

These results suggest the possibility that foliar P or the ratio of Foliar N to P may emerge as good predictors for the photosynthetic capacity for Amazonian forests. This needs to be verified by field observations and unfortunately, the eddy correlation systems have yet to be installed at sites where leaf phosphorus concentrations are higher than at the five Brazilian sites used here, all of which were on relatively infertile soils. This is important, because Fyllas et al. (2009) substantially higher average foliar P for sites in western Amazonia such as in South Peru and Ecuador, than the current Brazilian sites used in this study.




Fig. 7. Linear relationships obtained in this study (in circles) and obtained from tropical forest in Africa (in squares, Domingues, unpublished data).

5 Summary

Using a sun and shade model, the carbon uptake at 5 rainforest sites in the Brazilian Amazon was relatively well simulated. This result is supported by the ability of the model to reproduce values of $\delta^{13}\text{C}$ discrimination, although this was obtained with a poor partition of energy at some sites, mainly for wet season measurements. Additionally, the sun and shade model coupled to a model for stomatal conductance based on that of Cowan and Farquhar (1977) was unable to accurately capture carbon and energy fluxes at high D_C , over predicting measurements. There is still a need to understand the physiology of the stomatal closure under the afternoon conditions in these forests before it can be modelled. Nevertheless, a stomatal conductance description including hydraulic supply and demand might improve the model response under such conditions (Buckley, 2005).

Major constraints to model-data evaluation exercises such as attempted here are, however, imposed by the uncertainty in estimates of ecosystem respiration, mainly due to the seasonality of soil respiration, but also CO_2 storage fluxes and uncertainties associated with measurements from the eddy correlation N_E and non-closure of the energy balance.

Finally we note that this modelling exercise can be used to extrapolate simulations to the basin level. We have developed empirical relations between best fits of V_{max} and foliar N and P concentrations that can be tested to scale up to the basin level. Furthermore, the relationship between of V_{max} and P suggests the possibility of foliar P, or maybe N/P ratio a better predictor for canopy photosynthetic capacity than only foliar N in these forests. And given that most sites in western amazon have higher average foliar P (Fyllas et al., 2009), this might imply a gradient of GPP across Amazonia with higher GPP where leaf P is highest. Implications of these relationships will be explored further in an accompanying paper (Mercado et al., 2009) investigating the sensitivity of GPP to V_{max} parameterised using leaf P and leaf N at 35 sites across the Amazon Basin.

Results from this study highlight the need for field studies to further investigate the relationship between photosynthesis and leaf phosphorus in Amazonian forests.

Appendix A

Equations for canopy photosynthesis using the sun and shade model from de Pury and Farquhar (1997)

Following de Pury and Farquhar (1997), canopy level photosynthetic capacity is calculated as the integral of leaf photosynthetic capacity over the entire canopy, as parameterised with leaf level data for the Manaus C14 site in Mercado et al. (2006):

$$V_{\text{max_canopy}} = \int_0^{\text{LAI}} V_{\text{max_leaf}} dl \quad (\text{A1})$$

$$V_{\text{max_canopy}} = V_{\text{max_top}} [1 - \exp(-k_n)] / k_n \quad (\text{A2})$$

where $V_{\text{max_leaf}}$ is the maximum carboxylation activity of Rubisco at the leaf level in area basis, $V_{\text{max_canopy}}$ is the canopy level value for V_{max} , $V_{\text{max_top}}$, is the V_{max} value for top leaves. The cumulative leaf area index from the top of the canopy down to any level in the canopy is l , LAI is the total canopy leaf area index and k_n is a nitrogen or V_{max} allocation coefficient, defining the vertical decrease of $V_{\text{max_leaf}}$ with cumulative leaf area within the canopy.

Photosynthetic capacity of the sunlit and shaded fractions of the canopy is calculated by integrating the leaf photosynthetic capacity and the sunlit and shaded leaf area fractions, respectively. Photosynthesis of the sunlit and shaded fractions is then calculated separately using the leaf level biochemistry model of Farquhar and Caemmerer (1982), as two big leaves, with the absorbed irradiance and photosynthetic capacity, light saturated rate of electron transport and leaf dark respiration of each fraction used instead of the equivalent leaf level variables. Finally canopy photosynthesis is calculated by adding the individual contributions from the sunlit and shaded photosynthesis.

A1 Leaf biochemistry

The CO_2 assimilation rate (A) in units ($\mu\text{mol m}^2 \text{s}^{-1}$) is controlled by the rate of carboxylation when Rubisco activity is limiting (A_v) at low intercellular partial pressure of CO_2 and/or at high irradiances, and by the rate of electron transfer when Ribulose bisphosphate (RuBP) regeneration is limiting (A_J) at high intercellular partial pressure of CO_2 and/or at low irradiances. The rate of CO_2 assimilation is modelled as the minimum of A_v and A_J . The Rubisco-limited rate, A_v , and electron transport-limited rate, A_J , both in units ($\mu\text{mol m}^2 \text{s}^{-1}$) are defined as:

$$A_v = V_{\text{max}} \left(\frac{C_c - \Gamma^*}{K_C (1 + p\text{O}_2/K_0) + C_c} \right) - R_C \quad (\text{A3})$$

$$A_J = \frac{J}{4} \left(\frac{C_c - \Gamma^*}{C_c + 2\Gamma^*} \right) - R_C \quad (\text{A4})$$

where V_{max} in ($\mu\text{mol m}^2 \text{s}^{-1}$) is the maximum rate of Rubisco activity, K_0 and K_c in (Pa) are the Michaelis-Menton

constants for carboxylation and oxygenation by Rubisco. C_c in (Pa) is the partial pressure of CO_2 in the chloroplast, Γ^* in (Pa) is the CO_2 compensation point in the absence of mitochondrial respiration and R_C in ($\mu\text{mol m}^{-2} \text{s}^{-1}$) on a ground area basis, is the leaf dark respiration in the light. The Rubisco Michaelis constants for CO_2 and O_2 are described to follow an Arrhenius type temperature dependency as in Lloyd et al. (1995).

The potential rate of electron transport, J in ($\mu\text{mol m}^{-2} \text{s}^{-1}$), is modelled as a non-rectangular hyperbolic function of the absorbed quantum flux with I_2 in ($\mu\text{mol quanta m}^{-2} \text{s}^{-1}$) the absorbed irradiance that reaches photosystem II, J_{max} in ($\mu\text{mol m}^{-2} \text{s}^{-1}$), as saturating value and θ as the curvature factor:

$$\theta J^2 - (I_2 + J_{\text{max}})J + I_2 J_{\text{max}} = 0 \quad (\text{A5})$$

$$I_2 = I_0(1 - r - t)(1 - f)\Phi \quad (\text{A6})$$

r and t are the canopy reflectance and transmittance for PAR, respectively and f is the fraction of light lost as absorption by other than chloroplast lamellae, which increases with leaf thickness (Farquhar et al., 1980). I_0 is the PAR reaching the leaf or canopy surface in ($\mu\text{mol quanta m}^{-2} \text{s}^{-1}$). Φ is the efficiency of CO_2 fixation, so called quantum yield, defined as the initial slope of the relationship between assimilation rate A and irradiance. It describes the efficiency with which light is converted into fixed carbon.

The temperature sensitivities for Rubisco activity and electron transport are given by Farquhar and von Caemmerer (1982) as presented by Lloyd et al. (1995):

$$V_{\text{max}} = V_{\text{max},25} \exp\left(\frac{E_v}{298.2R} \left(1 - \frac{292.2}{T_C}\right)\right) \quad (\text{A7})$$

$$J_{\text{max}} = J_{\text{max},25} \frac{\exp\left[\frac{(T_C/298.2-1)E_J}{RT_C}\right] \left[1 + \exp\left(\frac{298.2S_J - H_J}{298.2R}\right)\right]}{1 + \exp\left(\frac{S_J T_C - H_J}{RT_C}\right)} \quad (\text{A8})$$

where T_C is absolute temperature (K) of the leaf or canopy, R is the universal gas constant ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$), $V_{\text{max},25}$ and $J_{\text{max},25}$ are Rubisco activity and electron transport capacity at 25°C in ($\mu\text{mol quanta m}^{-2} \text{s}^{-1}$). E_v and E_J in (J mol^{-1}) are activation energies. H_J in (J mol^{-1}) and S_J in ($\text{J mol}^{-1} \text{ K}^{-1}$) control maximum and minimum optimum temperature dependencies of the electron transport rate. The temperature optimum (T_{opt}) of J_{max} is known to acclimate in different environments, and can be estimated from Farquhar et al. (1980):

$$T_{\text{opt}} = \frac{H_J}{(S + R \ln(H_J/E_J - 1))} \quad (\text{A9})$$

The temperature dependency of leaf respiration is taken as presented by Lloyd et al. (1995):

$$R_D = R_{D,25} \exp\left(308.45 \left(\frac{1}{71.02} - \frac{1}{T_C - 227.13}\right)\right) \quad (\text{A10})$$




Fig. B1. Light response of N_E measured by eddy correlation averaged over $200 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ bins at the different tower sites. Data used correspond to the period July–December. Δ Tapajos, $*$ Man K34, \diamond Jaru, \square Caxiuana, $+$ Man C14.

where R_D is the rate of canopy dark respiration at T_C and $R_{D,25}$ is the rate of canopy dark respiration at 25°C , both are given in ($\mu\text{mol m}^{-2} \text{s}^{-1}$).

A2 Stomatal conductance

Stomatal conductance is calculated with the equation from Cowan and Farquhar (1977), called the “Lambda” model:

$$G_S = A \sqrt{\frac{1.6\lambda P}{(C_a - \Gamma^*)D_C}} \quad (\text{A11})$$



where A is the CO_2 assimilation rate in ($\text{mol m}^{-2} \text{s}^{-1}$), D_C vapour pressure deficit in mol fraction, P atmospheric pressure and C_a ambient partial pressure of CO_2 , and Γ^* is the CO_2 compensation partial pressure in the absence of dark respiration, all expressed in (mol mol^{-1}). The lambda parameter (λ) in ($\text{mol H}_2\text{O mol}^{-1} \text{CO}_2$) is a Lagrangian multiplier representing the marginal water cost of plant carbon gain.

Appendix B

See the table and figures.

Table B1. Calculated top of the canopy V_{\max} in ($\mu\text{mol m}^{-2} \text{s}^{-1}$) using linear regressions of V_{\max} against foliar N from different sources and reported values of V_{\max} from top canopy leaves.

	Jaru	Caxiuana	Tapajos	Man K34	Man C14
Leaf N (g m^{-2})	2.6 ^a	1.75 ^a	2.16 ^a	2.36 ^a	1.98 ^a
V_{\max} calculated with Carswell et al. (2000) regression $V_{\max}=12.99\text{N}-3.84$	29.97	18.8	24.1	26.8	21.9
V_{\max} calculated with Vale et al. (2003) regression $V_{\max}=23\text{N}-17.2$	42.7	22.9	32.4	37.0	28.5
V_{\max} calculated with Domingues et al. (2005) regression $V_{\max}=23\text{N}-7.02$	52.8	33.1	42.6	47.2	38.6
$V_{\max}=f(\text{N,P})$ Developed from other tropical forest (Domingues, unpublished)	45.9	30.17	39.72	35.92	38.03
V_{\max} measured and reported for single sites at top of the canopy		42.61 ^b	[28–75] ^d		42.8 ^c
V_{\max} fitted in this study. $\Phi=0.4$	51.9	39.4	40.0	37.8	52.3

^a Fyllas et al. (2009)^b Mean value at 30 m reported by Vale et al. (2003)^c Carswell et al. (2000) at 24 m^d Reported by Domingues et al. (2005) for top canopy leaves**Fig. B2.** (a) Observed and (b) simulated storage flux at Man C14 during the period October–November 1995. (c) Standardised residuals of observed and simulated storage flux at Man C14.**Fig. B3.** Mean diurnal cycle of observed (black) and simulated (grey) evaporative fraction at Tapajos using $V_{\max}=40 \mu\text{mol m}^{-2} \text{s}^{-1}$ and $\Phi=0.4$.

Acknowledgements. The authors wish to thank the following investigators for providing meteorology and eddy correlation flux data for the different sites: S. Saleska and N. Restrepo Coupe (University of Arizona), Tapajos site, A. Araujo (INPA), Manaus K34 site, C. von Randow (CPTEC/INPE) and B. Kruijt (Alterra), Jaru site, and A. Lola da Costa (Universidade Federal do Pará), Caxiuana and Y. Malhi (University of Oxford), Caxiuana site and Manaus C14 sites. We thank T. Domingues for supplying correlations of photosynthetic capacity and leaf nutrients in West Africa. Financial support for L. M. M. was provided through the EU Project “Carbonsink-LBA Proposal No. EVK 1999-0191” with most of this work being done at the Max-Planck-Institute for Biogeochemistry, Jena-Germany. The final analysis was also supported through salary support to L. M. M. through the UK National Environment Research Council funded Tropical Biomes in Transition (TROBIT) project. S. S. was supported by the

DEFRA and MoD integrated Climate Programme – GA01101, BC/2B/0417_Annex C5.r.

The service charges for this open access publication have been covered by the Max Planck Society.

Edited by: J. Grace

References

- Aranibar, J. N., Berry, J. A., Riley, W. J., Pataki, D. E., Law, B. E., and Ehleringer, J. R.: Combining meteorology, eddy fluxes, isotope measurements, and modeling to understand environmental controls of carbon isotope discrimination at the canopy scale, *Glob. Change Biol.*, 12, 710–730, 2006.
- Araújo, A. C., Nobre, A. D., Kruijt, B., Elbers, J. A., Dal-larosa, R., Stefani, P., von Randow, C., Manzi, A. O., Culf, A. D., Gash, J. H. C., Valentini, R., and Kabat, P.: Comparative measurements of carbon dioxide fluxes from two nearby towers in a central Amazonian rainforest: The Manaus LBA site, *J. Geophys. Res. Atmos.*, 107, 8090, doi:10.1029/2001JD000676, 2002.
- Asner, G. P., Nepstad, D., Cardinot, G., and Ray, D.: Drought stress and carbon uptake in an Amazon forest measured with spaceborne imaging spectroscopy, *Proc. Natl. Acad. Sci. USA*, 101, 6039–6044, 2004.
- Atkin, O. K., Evans, J. R., and Siebke, K.: Relationship between the inhibition of leaf respiration by light and enhancement of leaf dark respiration following light treatment, *Aust. J. Plant Physiol.*, 25, 437–443, 1998.
- Atkin, O. K., Evans, J. R., Ball, M. C., Lambers, H., and Pons, T. L.: Leaf respiration of snow gum in the light and dark. interactions between temperature and irradiance, *Plant Physiol.*, 122, 915–923, 2000.
- Aubinet, M., Heinesch, B., and Longdoz, B.: Estimation of the carbon sequestration by a heterogeneous forest: night flux corrections, heterogeneity of the site and inter-annual variability, *Glob. Change Biol.*, 8, 1053–1071, 2002.
- Baker, T. R., Phillips, O. L., Malhi, Y., Almeida, S., Arroyo, L., Di Fiore, A., Erwin, T., Higuchi, N., Killeen, T. J., Laurance, S. G., Laurance, W. F., Lewis, S. L., Monteagudo, A., Neill, D. A., Vargas, P. N., Pitman, N. C. A., Silva, J. N. M., and Martinez, R. V.: Increasing biomass in Amazonian forest plots, *Philos. Trans. R. Soc. Lond. Ser. B-Biol. Sci.*, 359, 353–365, 2004.
- Baldocchi, D. D.: Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future, *Glob. Change Biol.*, 9, 479–492, 2003.
- Ball, J. T., Woodrow, I. E., and Berry, J. A.: A model predicting stomatal conductance and its contribution to the control of photosynthesis under different environmental conditions, in: *Progress in Photosynthesis Research*, edited by: Martinus Nijhoff, M. J. B., Dordrecht, 221–224, 1987.
- Brooks, A. and Farquhar, G. D.: Effect of temperature on the CO₂/O₂ specificity of ribulose 1,5-bisphosphate carboxylase oxygenase and the rate of respiration in the light: estimates from gas exchange measurements on spinach, *Planta*, 165, 397–406, 1985.
- Brooks, A.: Effects of Phosphorus-Nutrition on Ribulose-1,5-Bisphosphate Carboxylase Activation, *Photosynthetic Quantum Yield and Amounts of Some Calvin-Cycle Metabolites in Spinach Leaves*, *Aust. J. Plant Physiol.*, 13, 221–237, 1986.
- Brooks, A., Woo, K. C., and Wong, S. C.: Effects of Phosphorus-Nutrition on the Response of Photosynthesis to CO₂ and O₂, Activation of Ribulose Bisphosphate Carboxylase and Amounts of Ribulose Bisphosphate and 3-Phosphoglycerate in Spinach Leaves, *Photosynth. Res.*, 15, 133–141, 1988.
- Buckley, T. N.: The control of stomata by water balance, *New Phytol.*, 168, 275–291, doi:10.1111/j.1469-8137.2005.01543.x, 2005.
- Campbell, C. D. and Sage, R. F.: Interactions between the effects of atmospheric CO₂ content and P nutrition on photosynthesis in white lupin (*Lupinus albus* L.), *Plant Cell Environ.*, 29, 844–853, doi:10.1111/j.1365-3040.2005.01464.x, 2006.
- Carswell, F. E., Meir, P., Wandelli, E. V., Bonates, L. C. M., Kruijt, B., Barbosa, E. M., Nobre, A. D., Grace, J., and Jarvis, P. G.: Photosynthetic capacity in a central Amazonian rain forest, *Tree Physiol.*, 20, 179–186, 2000.
- Carswell, F. E., Costa, A. L., Palheta, M., Malhi, Y., Meir, P., Costa, J. D. R., Ruivo, M. D., Leal, L. D. M., Costa, J. M. N., Clement, R. J., and Grace, J.: Seasonality in CO₂ and H₂O flux at an eastern Amazonian rain forest, *J. Geophys. Res. Atmos.*, 107(D20), 8076, doi:10.1029/2000JD000284, 2002.
- Chambers, J. Q., Higuchi, N., Schimel, J. P., Ferreira, L. V., and Melack, J. M.: Decomposition and carbon cycling of dead trees in tropical forests of the central Amazon, *Oecologia*, 122, 380–388, 2000.
- Chambers, J., Tribuzy, E. S., Toledo, L. C., Crispim, B. F., Higuchi, N., dos Santos, J., Araujo, A. C., Kruijt, B., Nobre, A., and Trumbore, S.: Respiration from a tropical forest ecosystem: partitioning of sources and low carbon use efficiency, *Ecol. Appl.*, 14, S72–S88, 2004.
- Chambers, J. Q. and Silver, W. L.: Some aspects of ecophysiological and biogeochemical responses of tropical forests to atmospheric change, *Philos. T. Roy. Soc. B*, 359, 463–476, 2004.
- Collatz, G. J., Berry, J. A., Farquhar, G., and Pierce, J.: The relationship between the Rubisco reaction mechanism and models of photosynthesis, *Plant Cell Environ.*, 13, 219–225, 1990.
- Collatz, G. J., Ball, J. T., Grivet, C., and Berry, J. A.: Physiological and environmental regulation of stomatal conductance, photosynthesis and transpiration: a model that includes a laminar boundary layer, *Agr. Forest Meteorol.*, 54, 107–136, 1991.
- Cowan, I. R. and Farquhar, G. D.: Stomatal function in relation to leaf metabolism and environment, *Symposium of the Society for Experimental Biology*, 471–505, 1977.
- Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A., and Totterdell, I. J.: Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model, *Nature*, 408, 750–750, 2000.
- da Rocha, H. R., Goulden, M. L., Miller, S. D., Menton, M. C., Pinto, L., de Freitas, H. C., and Figueira, A.: Seasonality of water and heat fluxes over a tropical forest in eastern Amazonia, *Ecol. Appl.*, 14, S22–S32, 2004.
- Davidson, E. A., Verchot, L. V., Cattanio, J. H., Ackerman, I. L., and Carvalho, J. E. M.: Effects of soil water content on soil respiration in forests and cattle pastures of eastern Amazonia, *Biogeochemistry*, 48, 53–69, 2000.
- de Pury, D. G. G. and Farquhar, G. D.: Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf

- models, *Plant Cell Environ.*, 20, 537–557, 1997.
- Domingues, T. F., Berry, J. A., Martinelli, L. A., Ometto, J. P. H. B., and Ehleringer, J. R.: Parameterization of canopy structure and leaf-level gas exchange for an Eastern Amazonian Tropical Rain Forest (Tapajos National Forest, Para, Brazil), *Earth Interact.*, 9, 1–23, 2005.
- Doughty, C. E. and Goulden, M. L.: Seasonal patterns of tropical forest leaf area index and CO₂ exchange, *J. Geophys. Res.-Biogeosci.*, 113, G00b06, doi:10.1029/2007jg000590, 2008a.
- Doughty, C. E. and Goulden, M. L.: Are tropical forests near a high temperature threshold?, *J. Geophys. Res.-Biogeosci.*, 113, G00b07, doi:10.1029/2007jg000632, 2008b.
- Ehleringer, J. and Björkman, O.: Quantum Yields for CO₂ Uptake in C₃ and C₄ Plants: Dependence on Temperature, CO₂, and O₂ Concentration, *Plant Physiol.*, 59, 86–90, 1977.
- Ehleringer, J. and Pearcy, R. W.: Variation in Quantum Yield for CO₂ Uptake among C₃ and C₄ plants, *Plant Physiol.*, 73, 555–559, 1983.
- Ehleringer, J. R., Lin, Z. F., Field, C. B., Sun, G. C., and Kuo, C. Y.: Leaf carbon isotope ratios of plants from a subtropical monsoon forest, *Oecologia*, 72, 109–114, 1987.
- Evans, J. R.: Photosynthetic Acclimation and Nitrogen Partitioning within a Lucerne Canopy 1. Canopy Characteristics, *Aust. J. Plant Physiol.*, 20, 55–67, 1993.
- Farquhar, G. D., Caemmerer, S. V., and Berry, J. A.: A Biochemical-Model of Photosynthetic CO₂ Assimilation in Leaves of C-3 Species, *Planta*, 149, 78–90, 1980.
- Farquhar, G. D., O'Leary, M. H., and Berry, J. A.: On the Relationship between Carbon Isotope Discrimination and the Inter-Cellular Carbon-Dioxide Concentration in Leaves, *Aust. J. Plant Physiol.*, 9, 121–137, 1982.
- Farquhar, G. D. and von Caemmerer, S.: Modelling of photosynthetic response to environmental conditions, in: *Encyclopedia of Plant physiology*, edited by: Lange, O. L., Nobel, P. S., Osmond, C. B., and Ziegler, H., Springer-Verlag, Berlin, 549–587, 1982.
- Finnigan, J. J., Clement, R., Malhi, Y., Leuning, R., and Cleugh, H. A.: A re-evaluation of long-term flux measurement techniques – Part I: Averaging and coordinate rotation, *Bound.-Lay. Meteorol.*, 107, 1–48, 2003.
- Fisher, R. A., Williams, M., Da Costa, A. L., Malhi, Y., Da Costa, R. F., Almeida, S., and Meir, P.: The response of an Eastern Amazonian rain forest to drought stress: results and modelling analyses from a throughfall exclusion experiment, *Glob. Change Biol.*, 13, 2361–2378, 2007.
- Fyllas, N. M., Patiño, S., Baker, T. R., Bielefeld Nardoto, G., Martinelli, L. A., Quesada, C. A., Paiva, R., Schwarz, M., Horna, V., Mercado, L. M., Santos, A., Arroyo, L., Jiménez, E. M., Luizão, F. J., Neill, D. A., Silva, N., Prieto, A., Rudas, A., Silveira, M., Vieira, I. C. G., Lopez-Gonzalez, G., Malhi, Y., Phillips, O. L., and Lloyd, J.: Basin-wide variations in foliar properties of Amazonian forest: phylogeny, soils and climate, *Biogeosciences Discuss.*, 6, 3707–3769, 2009, <http://www.biogeosciences-discuss.net/6/3707/2009/>.
- Gash, J. H. C. and Dolman, A. J.: Sonic anemometer (co)sine response and flux measurement I. The potential for (co)sine error to affect sonic anemometer-based flux measurements, *Agr. Forest Meteorol.*, 119, 195–207, 2003.
- Goudriaan, J.: *Crop micrometeorology: A simulation study*, Centre for Agricultural Publishing and Documentation, Wageningen, The Netherlands, 249 pp., 1977.
- Goulden, M. L., Miller, S. D., da Rocha, H. R., Menton, M. C., de Freitas, H. C., Figueira, A., and de Sousa, C. A. D.: Diel and seasonal patterns of tropical forest CO₂ exchange, *Ecol. Appl.*, 14, S42–S54, 2004.
- Grace, J., Lloyd, J., McIntyre, J., Miranda, A. C., Meir, P., Miranda, H. S., Nobre, C., Moncrieff, J., Massheder, J., Malhi, Y., Wright, I., and Gash, J.: Carbon-dioxide uptake by an undisturbed tropical rain-forest in Southwest Amazonia, 1992 to 1993, *Science*, 270, 778–780, 1995a.
- Grace, J., Lloyd, J., McIntyre, J., Miranda, A., Meir, P., Miranda, H., Moncrieff, J., Massheder, J., Wright, I., and Gash, J.: Fluxes of carbon-dioxide and water-vapor over an undisturbed tropical forest in South-West Amazonia, *Glob. Change Biol.*, 1, 1–12, 1995b.
- Grace, J., Malhi, Y., Higuchi, N., and Meir, P.: Productivity and carbon fluxes of tropical rain forest, in: *Global Terrestrial Productivity: Past, Present and Future*, edited by: Mooney, H. and Saugier, B., Academic Press, San Diego, 401–426, 2001.
- Harley, P. C. and Sharkey, T. D.: An Improved Model of C₃ Photosynthesis at High CO₂ – Reversed O₂ Sensitivity Explained by Lack of Glycerate Reentry into the Chloroplast, *Photosynth. Res.*, 27, 169–178, 1991.
- Harley, P. C., Thomas, R. B., Reynolds, J. F., and Strain, B. R.: Modeling Photosynthesis of Cotton Grown in Elevated CO₂, *Plant Cell Environ.*, 15, 271–282, 1992.
- Hirose, T. and Werger, M. J. A.: Maximizing daily canopy photosynthesis with respect to the leaf nitrogen allocation pattern in the canopy, *Oecologia*, 72, 520–526, 1987.
- Hoefnagel, M. H. N., Atkin, O. K., and Wiskich, J. T.: Interdependence between chloroplasts and mitochondria in the light and the dark, *BBA-Bioenergetics*, 1366, 235–255, 1998.
- Hutyra, L. R., Munger, J. W., Hammond-Pyle, E., Saleska, S. R., Restrepo-Coupe, N., de Camargo, P. B., and Wofsy, S. C.: Resolving systematic errors in estimates of net ecosystem exchange of CO₂ and ecosystem respiration in a tall-stature forest: application to a tropical forest biome, *Agr. Forest Meteorol.*, 148, 1266–1279, doi:10.1016/j.agrformet.2008.08.03, 2007.
- Iwata, H., Malhi, Y., and von Randow, C.: Gap-filling measurements of carbon dioxide storage in tropical rainforest canopy airspace, *Agr. Forest Meteorol.*, 132, 305–314, doi:10.1016/j.agrformet.2005.08.005, 2005.
- Jacob, J. and Lawlor, D. W.: Dependence of Photosynthesis of Sunflower and Maize leaves on phosphate supply, ribulose-1,5-bisphosphate carboxylase/oxygenase activity, and ribulose-1,5-bisphosphate pool size, *Plant Physiol.*, 98, 801–807, 1992.
- Janssens, I. A., Lankreijer, H., Matteucci, G., Kowalski, A. S., Buchmann, N., Epron, D., Pilegaard, K., Kutsch, W., Longdoz, B., Grunwald, T., Montagnani, L., Dore, S., Rebmann, C., Moors, E. J., Grelle, A., Rannik, U., Morgenstern, K., Oltchev, S., Clement, R., Gudmundsson, J., Minerbi, S., Berbigier, P., Ibrom, A., Moncrieff, J., Aubinet, M., Bernhofer, C., Jensen, N. O., Vesala, T., Granier, A., Schulze, E.-D., Lindroth, A., Dolman, A. J., Jarvis, P. G., Ceulemans, R., and Valentini, R.: Productivity overshadows temperature in determining soil and ecosystem respiration across European forests, *Glob. Change Biol.*, 7, 269–278, 2001.
- Jones, H. G.: *Plants and Microclimate*, 2nd edn., Cambridge Uni-

- versity Press, Cambridge, 1992.
- Katul, G. G., Ellsworth, D. S., and Lai, C.-T.: Modelling assimilation and intercellular CO₂ from measured conductance: a synthesis of approaches, *Plant Cell Environ.*, 23, 1313–1328, 2000.
- Keeling, H. C. and Phillips, O. L.: A calibration method for the crown illumination index for assessing forest light environments, *Forest Ecol. Manag.*, 242, 431–437, doi:10.1016/j.foreco.2007.01.060, 2007.
- Klinge, H.: Litter production in an area of Amazonian Terra firme forest. Part I. Litter-fall, organic carbon and total nitrogen contents of litter, *Amazoniana*, 1, 287–302, 1968.
- Kruijt, B., Elbers, J. A., von Randow, C., Araujo, A. C., Oliveira, P. J., Culf, A., Manzi, A. O., Nobre, A. D., Kabat, P., and Moors, E. J.: The robustness of eddy correlation fluxes for Amazon rain forest conditions, *Ecol. Appl.*, 14, S101–S113, 2004.
- Leuning, R.: Modeling stomatal behavior and photosynthesis of *Eucalyptus grandis*, *Aust. J. Plant Physiol.*, 17, 159–175, 1990.
- Leuning, R.: A Critical-Appraisal of a Combined Stomatal-Photosynthesis Model for C-3 Plants, *Plant Cell Environ.*, 18, 339–355, 1995.
- Leuning, R., Kelliher, F. M., Depury, D. G. G., and Schulze, E. D.: Leaf nitrogen, photosynthesis, conductance and transpiration: scaling from leaves to canopies, *Plant Cell Environ.*, 18, 1183–1200, 1995.
- Lewis, S. L., Malhi, Y., and Phillips, O. L.: Fingerprinting the impacts of global change on tropical forests, *Philos. T. Roy. Soc. B*, 359, 437–462, 2004.
- Lloyd, J. and Farquhar, G. D.: C-13 Discrimination during CO₂ assimilation by the terrestrial biosphere, *Oecologia*, 99, 201–215, 1994.
- Lloyd, J. and Taylor, J. A.: On the temperature dependence of soil respiration, *Funct. Ecol.*, 8, 315–323, 1994.
- Lloyd, J., Grace, J., Miranda, A. C., Meir, P., Wong, S. C., Miranda, B. S., Wright, I. R., Gash, J. H. C., and McIntyre, J.: A Simple Calibrated Model of Amazon Rain-Forest Productivity Based on Leaf Biochemical-Properties, *Plant Cell Environ.*, 18, 1129–1145, 1995.
- Lloyd, J. and Farquhar, G. D.: The CO₂ dependence of photosynthesis, plant growth responses to elevated atmospheric CO₂ concentrations and their interaction with soil nutrient status 1. General principles and forest ecosystems, *Funct. Ecol.*, 10, 4–32, 1996.
- Lloyd, J., Kruijt, B., Hollinger, D., Grace, J., Francey, R. J., Wong, S. C., Kelliher, F. M., Miranda, A. C., Farquhar, G. D., Gash, J. H. C., Vygodskaya, N. N., Wright, I. R., Miranda, H. S., and Schulze, E. D.: Vegetation effects on the isotopic composition of the atmospheric CO₂ at local and regional scales: Theoretical aspects and a comparison between rain forest in Amazonia and a boreal forest in Siberia, *Aust. J. Plant Physiol.*, 23, 371–399, 1996.
- Lloyd, J., Bird, M. I., Veenendaal, E. M., and Kruijt, B.: Should Phosphorous Availability be Constraining Moist Tropical Forest responses to increasing CO₂ concentrations?, in: *Global biogeochemical cycles in the climate system*, edited by: Schulze, E. D., Schimel, D. S., Heimann, M., Harrison, S. P., Holland, E. A., Lloyd, J., and Prentice, I. C., Academic Press, San Diego, 95–114, 2001.
- Lloyd, J. and Farquhar, G. D.: Effects of rising temperatures and [CO₂] on the physiology of tropical forest trees, *Philos. T. Roy. Soc. B*, 363, 1811–1817, doi:10.1098/rstb.2007.0032, 2008.
- Lloyd, J., Patiño, S., Paiva, R. Q., Nardoto, G. B., Quesada, C. A., Santos, A. J. B., Baker, T. R., Brand, W. A., Hilke, I., Giemann, H., Raessler, M., Luizao, F. J., Martinelli, L. A., and Mercado, L. M.: Gradients in leaf physiological properties within Amazon forest canopies, *Biogeosciences Discuss.*, in preparation, 2009.
- Long, S. P., Postl, W. F., and Bolharnordenkamp, H. R.: Quantum Yields for Uptake of Carbon-Dioxide in C-3 Vascular Plants of Contrasting Habitats and Taxonomic Groupings, *Planta*, 189, 226–234, 1993.
- Luizao, F. and Schubart, H. O. R.: Litter production and decomposition in a terra-firme forest of Central Amazonia, *Experientia*, 43, 259–265, 1987.
- Malhi, Y., Nobre, A. D., Grace, J., Kruijt, B., Pereira, M. G. P., Culf, A., and Scott, S.: Carbon dioxide transfer over a Central Amazonian rain forest, *J. Geophys. Res. Atmos.*, 103, 31593–31612, 1998.
- Malhi, Y., Pegoraro, E., Nobre, A. D., Pereira, M. G. P., Grace, J., Culf, A. D., and Clement, R.: Energy and water dynamics of a central Amazonian rain forest, *J. Geophys. Res. Atmos.*, 107, 8061, doi:10.1029/2001JD000623, 2002.
- Massman, W. J. and Lee, X.: Eddy covariance flux corrections and uncertainties in long-term studies of carbon and energy exchanges, *Agr. Forest Meteorol.*, 113, 121–144, 2002.
- Medlyn, B. E., Dreyer, E., Ellsworth, D., Forstreuter, M., Harley, P. C., Kirschbaum, M. U. F., Le Roux, X., Montpied, P., Strassmeyer, J., Walcroft, A., Wang, K., and Loustau, D.: Temperature response of parameters of a biochemically based model of photosynthesis. II. A review of experimental data, *Plant Cell Environ.*, 25, 1167–1179, doi:10.1046/j.1365-3040.2002.00891.x, 2002.
- Meir, P., Grace, J., Miranda, A. C., and Lloyd, J.: Soil Respiration in a rainforest in Amazonia and in cerrado in central Brazil, in: *Amazon Deforestation and Climate*, edited by: Gash, J. H. C., Nobre, C. A., Roberts, J., and Victoria, R. L., John Wiley & Sons, Chichester, UK, 319–329, 1996.
- Meir, P., Grace, J., and Miranda, A. C.: Photographic method to measure the vertical distribution of leaf area density in forests, *Agr. Forest Meteorol.*, 102, 105–111, 2000.
- Meir, P., Grace, J., and Miranda, A. C.: Leaf respiration in two tropical rainforests: constraints on physiology by phosphorus, nitrogen and temperature, *Funct. Ecol.*, 15, 378–387, 2001.
- Meir, P. and Grace, J.: Scaling relationships for woody tissue respiration in two tropical rain forests, *Plant Cell Environ.*, 25, 963–973, 2002.
- Meir, P., Metcalfe, D. B., Costa, A. C. L., and Fisher, R. A.: The fate of assimilated carbon during drought: impacts on respiration in Amazon rainforests, *Philos. T. Roy. Soc. B*, 363, 1849–1855, doi:10.1098/rstb.2007.0021, 2008.
- Mercado, L. M., Lloyd, J., Carswell, F., Malhi, Y., Meir, P., and Nobre, A.: Modelling Amazonian forest eddy covariance data: a comparison of big leaf versus sun/shade model for the C-14 tower at Manaus I. Canopy photosynthesis, *Acta Amazonica*, 36, 69–82, 2006.
- Mercado, L. M.: Modelling Rainforest Canopy Photosynthesis across the Amazon Basin, Ph.D. thesis, Free University of Amsterdam, Amsterdam, 2007.

- Mercado, L. M., Lloyd, J., Dolman, A. J., Sitch, S., and Patiño, S.: Modelling basin-wide variations in Amazon forest productivity – Part 2: Application of the model on a Basin-wide scale, in preparation, 2009.
- Mercado, L. M., Huntingford, C., Gash, J. H. C., Cox, P. M., and Jogireddy, V.: Improving the representation of radiation interception and photosynthesis for climate model applications, *Tellus B*, 59, 553–565, 2007.
- Miller, S. D., Goulden, M. L., Menton, M. C., da Rocha, H. R., de Freitas, H. C., Figueira, A., and de Sousa, C. A. D.: Biometric and micrometeorological measurements of tropical forest carbon balance, *Ecol. Appl.*, 14, S114–S126, 2004.
- Morales, P., Sykes, M. T., Prentice, I. C., Smith, P., Smith, B., Bugmann, H., Zierl, B., Friedlingstein, P., Viovy, N., Sabate, S., Sanchez, A., Pla, E., Gracia, C. A., Sitch, S., Arneth, A., and Ogee, J.: Comparing and evaluating process-based ecosystem model predictions of carbon and water fluxes in major European forest biomes, *Glob. Change Biol.*, 11, 2211–2233, 2005.
- Nelder, J. A. and Mead, R.: A Simplex-Method for Function Minimization, *Comput. J.*, 7, 308–313, 1965.
- Nepstad, D. C., Moutinho, P., Dias, M. B., Davidson, E., Cardinot, G., Markewitz, D., Figueiredo, R., Vianna, N., Chambers, J., Ray, D., Guerreiros, J. B., Lefebvre, P., Sternberg, L., Moreira, M., Barros, L., Ishida, F. Y., Tohlver, I., Belk, E., Kalif, K., and Schwalbe, K.: The effects of partial throughfall exclusion on canopy processes, aboveground production, and biogeochemistry of an Amazon forest, *J. Geophys. Res. Atmos.*, 107, 8085, doi:10.1029/2001JD000360, 2002.
- Ometto, J. P. H. B., Ehleringer, J. H., Domingues, T. F., Berry, J. A., Ishida, F. Y., Mazzi, E., Higuchi, N., Flanagan, L., Nardoto, G. B., and Martinelli, L. A.: The stable carbon and nitrogen isotopic composition of vegetation in tropical forests of the Amazon region, Brazil, *Biogeochemistry*, 79, 251–274, 2006.
- Pattey, E., Strachan, I. B., Desjardins, R. L., and Massheder, J.: Measuring nighttime CO₂ flux over terrestrial ecosystems using eddy covariance and nocturnal boundary layer methods, *Agr. Forest Meteorol.*, 113, 145–158, 2002.
- Phillips, O. L., Malhi, Y., Higuchi, N., Laurance, W. F., Nunez, P. V., Vasquez, R. M., Laurance, S. G., Ferreira, L. V., Stern, M., Brown, S., and Grace, J.: Changes in the carbon balance of tropical forests: Evidence from long-term plots, *Science*, 282, 439–442, 1998.
- Poorter, L., Oberbauer, S. F., and Clark, D. B.: Leaf optical-properties along a vertical gradient in a tropical rain-forest canopy in Costa-Rica, *Am. J. Bot.*, 82, 1257–1263, 1995.
- Prentice, I. C. and Lloyd, J.: C-quest in the Amazon Basin, *Nature*, 396, 619–620, 1998.
- Quesada, C. A., Lloyd, J., Anderson, L. O., Fyllas, N. M., Schwarz, M., and Czimczik, C. I.: Soils of amazonia with particular reference to the rainfor sites, *Biogeosciences Discuss.*, 6, 3851–3921, 2009, <http://www.biogeosciences-discuss.net/6/3851/2009/>.
- Restrepo-Coupe, N., da Rocha, H., Saleska, S. R., Aguiar, R. G., Araujo, A. C., Borma, L. S., Cabral, O. M. R., de Camargo, P. B., Cardoso, F. L., da Costa, L. A. C., Fitzjarrald, D. R., Goulden, M. L., Hutyrá, L. R., Kruijt, B., Maia, J. M. F., Malhi, Y., Manzi, A. O., Miller, S. D., Nobre, A. D., von Randow, C., da Abreu Sá, L. D., Sakai, R. K., Tota, J., Wofsy, S. C., and Zanchi, F. B.: What controls the seasonality of photosynthesis across the Amazon basin? A cross-site analysis of eddy flux tower measurements from the Brasil flux network, in preparation, 2009.
- Sage, R. F. and Kubien, D. S.: The temperature response of C-3 and C-4 photosynthesis, *Plant Cell Environ.*, 30, 1086–1106, 10.1111/j.1365–3040.2007.01682.x, 2007.
- Saleska, S. R., Miller, S. D., Matross, D. M., Goulden, M. L., Wofsy, S. C., da Rocha, H. R., de Camargo, P. B., Crill, P., Daube, B. C., de Freitas, H. C., Hutyrá, L., Keller, M., Kirchhoff, V., Menton, M., Munger, J. W., Pyle, E. H., Rice, A. H., and Silva, H.: Carbon in amazon forests: Unexpected seasonal fluxes and disturbance-induced losses, *Science*, 302, 1554–1557, 2003.
- Sharkey, T. D.: O₂-insensitive photosynthesis in C3 plants: Its occurrence and a possible explanation, *Plant Physiol.*, 78, 71–75, 1985.
- Shuttleworth, W. J.: Micrometeorology of temperate and tropical forest, *Philos. T. Roy. Soc. B*, 324, 299–334, 1989.
- Silva de Souza, J.: Dinamica espacial e temporal do fluxo de CO₂ do solo em floresta de terra firme na amazonia central, Master thesis, Universidad Federal do Amazonas, Manaus, 2004.
- Simon, E., Meixner, F. X., Ganzeveld, L., and Kesselmeier, J.: Coupled carbon-water exchange of the Amazon rain forest, I. Model description, parameterization and sensitivity analysis, *Biogeosciences*, 2, 231–253, 2005, <http://www.biogeosciences.net/2/231/2005/>.
- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Glob. Change Biol.*, 9, 161–185, 2003.
- Sotta, E. D., Meir, P., Malhi, Y., Nobre, A. D., Hodnett, M., and Grace, J.: Soil CO₂ efflux in a tropical forest in the central Amazon, *Glob. Change Biol.*, 10, 601–617, 2004.
- Terry, N. and Ulrich, A.: Effects of phosphorus deficiency on the photosynthesis and respiration of leaves of sugar beet, *Plant Physiol.*, 51, 43–47, 1973.
- Thornley, J. H. M.: Instantaneous canopy photosynthesis: Analytical expressions for sun and shade leaves based on exponential light decay down the canopy and an acclimated non-rectangular hyperbola for leaf photosynthesis, *Ann. Bot.*, 89, 451–458, 2002.
- Tian, H. Q., Melillo, J. M., Kicklighter, D. W., McGuire, A. D., Helfrich III, J. V. K., Moore III, B., and Vörösmarty, C. J.: Effect of interannual climate variability on carbon storage in Amazonian ecosystems, *Nature*, 396, 664–667, 1998.
- Tuzet, A., Perrier, A., and Leuning, R.: A coupled model of stomatal conductance, photosynthesis and transpiration, *Plant Cell Environ.*, 26, 1097–1116, doi:10.1046/j.1365–3040.2003.01035.x, 2003.
- Tyree, M. T.: Hydraulic limits on tree performance: transpiration, carbon gain and growth of trees, *Trees-Struct. Funct.*, 17, 95–100, 2003.
- Vale, R. L., Maroco, J. P., Carvalho, C. R. J., Almeida, S., Meir, P., Grace, J., Pereira, J. S., and Chaves, M. M.: Carbon assimilation in an amazonian rainforest: a rain exclusion experiment, in: Comparative Biochemistry and Physiology, Abstracts of the Annual Main Meeting of the Society for Experimental Biology, Southampton, UK, S1–S238, 2003.
- van der Molen, M. K., Gash, J. H. C., and Elbers, J. A.: Sonic

- anemometer (co)sine response and flux measurement – II. The effect of introducing an angle of attack dependent calibration, *Agr. Forest Meteorol.*, 122, 95–109, 2004.
- van Gorsel, E., Leuning, R., Cleugh, H. A., Keith, H., and Suni, T.: Nocturnal carbon efflux: reconciliation of eddy covariance and chamber measurements using an alternative to the u^* -threshold filtering technique, *Tellus B*, 59, 397–403, 2007.
- van Gorsel, E., Leuning, R., Cleugh, H. A., Keith, H., Kirschbaum, M. U. F., and Suni, T.: Application of an alternative method to derive reliable estimates of nighttime respiration from eddy covariance measurements in moderately complex topography, *Agr. Forest Meteorol.*, 148, 1174–1180, 2008.
- Vitousek, P. M. and Sanford, R. L.: Nutrient cycling in moist tropical forest, *Annu. Rev. Ecol. Syst.*, 17, 137–167, 1986.
- von Randow, C., Manzi, A. O., Kruijt, B., de Oliveira, P. J., Zanchi, F. B., Silva, R. L., Hodnett, M. G., Gash, J. H. C., Elbers, J. A., Waterloo, M. J., Cardoso, F. L., and Kabat, P.: Comparative measurements and seasonal variations in energy and carbon exchange over forest and pasture in South West Amazonia, *Theor. Appl. Climatol.*, 78, 5–26, 2004.
- Wang, Y. P. and Leuning, R.: A two-leaf model for canopy conductance, photosynthesis and partitioning of available energy I: Model description and comparison with a multi-layered model, *Agr. Forest Meteorol.*, 91, 89–111, 1998.
- Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., Bernhofer, C., Ceulemans, R., Dolman, H., Field, C., Grelle, A., Ibrom, A., Law, B. E., Kowalski, A., Meyers, T., Moncrieff, J., Monson, R., Oechel, W., Tenhunen, J., Valentini, R., and Verma, S.: Energy balance closure at FLUXNET sites, *Agr. Forest Meteorol.*, 113, 223–243, 2002.
- Wohlfahrt, G., Bahn, M., Haslwanter, A., Newesely, C., and Cernusca, A.: Estimation of daytime ecosystem respiration to determine gross primary production of a mountain meadow, *Agr. Forest Meteorol.*, 130, 13–25, 2005.
- Woodrow, I. E. and Berry, J. A.: Enzymatic regulation of photosynthetic CO_2 fixation in C-3 plants, *Annu. Rev. Plant Phys.*, 39, 533–594, 1988.