

Washington University School of Medicine

Digital Commons@Becker

2020-Current year OA Pubs

Open Access Publications

5-12-2023

A phase 1b randomized clinical trial of CT1812 to measure A β oligomer displacement in Alzheimer's disease using an indwelling CSF catheter

Kelsie M LaBarbera
Cognition Therapeutics Inc.

Carla M Yuede
Washington University School of Medicine in St. Louis

Hannah M Edwards
Washington University School of Medicine in St. Louis

Woodrow D Gardiner
Washington University School of Medicine in St. Louis

John R Cirrito
Washington University School of Medicine in St. Louis

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/oa_4



Part of the [Medicine and Health Sciences Commons](#)

Please let us know how this document benefits you.

Recommended Citation

LaBarbera, Kelsie M; Yuede, Carla M; Edwards, Hannah M; Gardiner, Woodrow D; Cirrito, John R; and et al., "A phase 1b randomized clinical trial of CT1812 to measure A β oligomer displacement in Alzheimer's disease using an indwelling CSF catheter." *Translational Neurodegeneration*. 12, 1. 24 (2023).
https://digitalcommons.wustl.edu/oa_4/2817

This Open Access Publication is brought to you for free and open access by the Open Access Publications at Digital Commons@Becker. It has been accepted for inclusion in 2020-Current year OA Pubs by an authorized administrator of Digital Commons@Becker. For more information, please contact vanam@wustl.edu.

Authors

Kelsie M LaBarbera, Carla M Yuede, Hannah M Edwards, Woodrow D Gardiner, John R Cirrito, and et al.

LETTER

Open Access



A phase 1b randomized clinical trial of CT1812 to measure A β oligomer displacement in Alzheimer's disease using an indwelling CSF catheter

Kelsie M. LaBarbera¹, Yvette I. Sheline², Nicholas J. Izzo¹, Carla M. Yuede³, Lora Waybright¹, Raymond Yurko¹, Hannah M. Edwards³, Woodrow D. Gardiner³, Kaj Blennow⁴, Henrik Zetterberg^{4,5,6,7,8,9,10}, Anne Börjesson-Hanson¹¹, Roger Morgan¹², Charles S. Davis¹³, Robert J. Guttendorf¹⁴, Lon S. Schneider¹⁵, Steven DeKosky¹⁶, Harry LeVine III¹⁷, Michael Grundman^{18,19}, Anthony O. Caggiano¹, John R. Cirrito³, Susan M. Catalano^{1†} and Mary E. Hamby^{1*†} 

Trial Registration: May 11th, 2018 ClinicalTrials.gov Identifier: NCT03522129 <https://clinicaltrials.gov/ct2/show/NCT03522129>.

Investigational therapies for Alzheimer's disease (AD) target a wide range of mechanisms, yet promising disease-modifying therapies remain a huge unmet need. Much evidence indicates that the oligomeric form of amyloid-beta (A β) is a toxic species contributing to AD through synaptic damage and neuronal toxicity [1]. In support of this, A β oligomer reduction in an AD mouse

model leads to memory preservation [2, 3], and clinical benefit was observed in trials of lecanemab, which targets A β oligomers and protofibrils [4], in AD patients, encouraging the continued development of A β oligomer-targeting therapies.

CT1812 is a novel, small-molecule, brain-penetrant sigma-2 receptor (S2R) modulator that selectively prevents and displaces A β oligomers from binding to neuronal synapses, thereby mitigating downstream toxicity. This is thought to occur through allosteric modulation

[†]Susan M Catalano and Mary E Hamby are dual senior authors. Susan M Catalano is no longer affiliated with Cognition Therapeutics Inc

*Correspondence:

Mary E. Hamby
mhamby@cogrx.com

¹ Cognition Therapeutics Inc., Pittsburgh, PA, USA

² University of Pennsylvania, Philadelphia, USA

³ Washington University, St. Louis, USA

⁴ University of Gothenburg, Mölndal, Sweden

⁵ Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, The Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

⁶ Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

⁷ Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, UK

⁸ UK Dementia Research Institute at UCL, London, UK

⁹ Hong Kong Center for Neurodegenerative Diseases, Clear Water Bay, Hong Kong, China

¹⁰ Wisconsin Alzheimer's Disease Research Center, University of Wisconsin School of Medicine and Public Health, University of Wisconsin-Madison, Madison, WI, USA

¹¹ Karolinska University Hospital, Stockholm, Sweden

¹² MedSurgPI, LLC, Raleigh, NC, USA

¹³ CSD Biostatistics, Inc., Oro Valley, AZ, USA

¹⁴ Aclairo Pharmaceutical Development Group, Inc, Vienna, VA, USA

¹⁵ Keck School of Medicine of USC, Los Angeles, CA, USA

¹⁶ McKnight Brain Institute, University of Florida, Gainesville, FL, USA

¹⁷ Sanders-Brown Center on Aging, University of Kentucky, Lexington, KY, USA

¹⁸ Global R&D Partners, LLC, San Diego, CA, USA

¹⁹ Department of Neurosciences, University of California, San Diego, USA



of the oligomer receptor, cellular prion protein (PrP^c), by direct interaction between PrP^c and S2R at synapses [3, 5]. CT1812 rescues synaptic loss and neuronal function in primary neuronal cultures, and restores cognition in a mouse model of AD to levels of healthy controls [3].

Demonstration of target engagement, successful binding of a compound to its intended receptor, is important for increasing probability of success in drug development. Given the mechanism of action of CT1812, we postulated that A β oligomers could be a proximal indicator of S2R target engagement. Preclinical studies supported this, with CT1812 impacting A β oligomers in AD transgenic mice [3]. A β oligomer levels in hippocampal interstitial fluid and lateral ventricle cerebrospinal fluid (CSF) were measured in vivo using a microimmuno-electrode (MIE) coated with an oligomer-selective antibody following a single administration of CT1812. A significant, dose-dependent rise in A β oligomers was detected after administration, consistent with A β oligomer displacement from neurons and clearance into CSF. In contrast, A β monomer levels were not affected [3].

To determine whether this approach could demonstrate target engagement in patients with AD, a phase 1b, randomized, double-blind, placebo-controlled trial (NCT03522129) was designed. Eligibility criteria for the study included a diagnosis of mild-moderate AD (Mini-Mental State Examination score 18–26) and either an AD-positive amyloid positron emission tomography scan or a positive CSF result for AD biomarkers (Additional file 1) within 12 months before or at screening. Participants were randomized 2:1 to receive either a single oral dose of CT1812 (560 mg) or identically appearing placebo capsules. An indwelling catheter was placed in the lumbar subarachnoid space and 4–6 ml CSF samples were collected hourly for 28 h (five samples pre-dose and 24 samples following drug or placebo administration). Plasma was taken at the same intervals for pharmacokinetic analysis.

Although intended enrollment was 18 patients, recruitment was challenged by the 28-h spinal catheter procedure and the single-dose nature of the study without optional open-label extension. Fifteen participants were screened and three were randomized and completed the trial before it was ceased. Baseline demographics are shown in Fig. 1a. There were no deaths and no subjects were withdrawn from the study due to treatment-emergent adverse events.

To assess target engagement, A β oligomer levels in CSF samples were measured at all time points (Fig. 1b–g) using MIE methods similar to those described previously [3], as well as using non-denaturing western blotting quantifying A β oligomers ranging from 25 to 99 kDa [3]. CSF A β -40 and A β -42 monomer levels were

also measured at all time points (Lumipulse immunoassay [6]). Plasma and CSF CT1812 concentrations were measured (LC–MS/MS [3]) to understand the pharmacokinetic/pharmacodynamic relationship across the total 24 h of exposure. For each patient, the C_{max} was determined and total drug exposure was calculated as area under the curve (AUC_{0–last}).

Given the semi-quantitative nature of the A β oligomer assays, the percent change of A β oligomer level relative to baseline (average of the pre-dose values) was calculated for each patient (Fig. 1b–g). A β oligomer levels in the two CT1812-treated patients, assessed by MIE, increased by >250%–500% above baseline over time, but did not increase in the placebo-treated patient (Fig. 1b–d). A similar pattern was observed using the independent native gel Western blot method (Fig. 1e–g). A high degree of congruence between these distinct assays was observed (Pearson correlation coefficient $r=0.74$; Fig. 1h).

The CT1812 exposure level in patient 1 (AUC_{0–last} = 120 h*ng/ml, C_{max} = 24.9 ng/ml) was more than twofold higher than that of patient 3 (AUC_{0–last} = 46.4 h*ng/ml, C_{max} = 7.27 ng/ml, Fig. 1i), allowing for a preliminary gauge of exposure-dependence. Similarly, CSF A β oligomer levels were also higher in patient 1 than in patient 3 (Fig. 1i). Together with the preclinical studies [3], this observation is consistent with a drug exposure-dependent impact of CT1812 on A β oligomers after a single dose.

Importantly, there was little to no change (<50% increase from baseline) in A β 40 and A β 42 levels over the 24-h period irrespective of treatment. The slight increase observed in some patients may be due to the decreased overall CSF volume caused by multiple CSF collections within a short period [7, 8]. The selective increase in CSF A β oligomers but not monomers in the two patients with CT1812 treatment is consistent with preclinical evidence that CT1812 selectively displaces oligomeric over monomeric A β [3].

This is the first study evaluating hourly changes of A β oligomer concentrations in AD patient CSF and examining how this pharmacodynamic biomarker is impacted by the drug candidate CT1812, which selectively displaces A β oligomers allosterically through the S2R. The observed selective increase in CSF A β oligomers also suggests that there is a displaceable pool of A β oligomers in the AD central nervous system and provides early evidence that the mechanism of action of CT1812 elucidated preclinically translates to the clinic. The observation that the degree of change in A β oligomers aligned with the exposure level of CT1812 supports the use of A β oligomers as a measure of target engagement in future studies. As the relative increase in A β oligomers




Fig. 1 CT1812 treatment results in increases in CSF Aβ oligomers, but not monomers, in an exposure-dependent manner. **a** Subject demographics. **b–g** CSF Aβ oligomer or monomer concentrations were measured in individual AD patient samples collected hourly before (–4 to 0 h as baseline) and 24 h after a single dose of 560 mg CT1812 or placebo. Aβ monomers (ELISA) increased by < 50% of baseline during the post-dose period, but Aβ oligomers (MIE) increased by > 250% (Patient 3) and > 500% (Patient 1) of baseline (**b–d**). Non-denaturing western blotting revealed similar increases of CSF Aβ oligomers in CT1812 patients (**e–g**). **h** The oligomer level by western blot correlated with that measured by MIE (Pearson $r = 0.74$; $n = 70$ samples) **i** CSF total exposure levels of CT1812 (data shown in Table S1) were related to the total AUC_{0-last} of CSF Aβ oligomer concentration (MIE or Western blot) across patients

over time did not reach a plateau by the end of the study, assessment with longer intervals may be warranted to establish the peak time of this pharmacodynamic response. Despite the small cohort size analyzed in this trial, the unique design provides early proof of principle that target engagement can be measured in patients after a single dose of CT1812. Given that toxic A β oligomers play a critical role in AD pathogenesis, these findings support continued clinical development of CT1812, which is under further investigation in ongoing phase 2 clinical trials for AD (NCT03507790, NCT04735536) and dementia with Lewy bodies (NCT05225415).

Abbreviations

MIE	Microimmunoelectrode
PrP ^C	Cellular prion protein
S2R	Sigma-2 receptor

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40035-023-00358-w>.

Additional file 1. Methods. Table S1: CT1812 pharmacokinetic parameters in plasma and CSF of two AD patients after a single 560-mg oral dose.

Acknowledgements

Hilary North prepared the manuscript.

Author contributions

Trial, analysis conceived by SMC, SD, LSS, and MG (YIS assisted). ABH and YIS conducted trial operations. AOC, RM, and MG monitored trial safety. KB and HZ conducted A β monomer ELISA in CSF; MEH analyzed data. CMY, HME, WDG, and JRC developed, validated A β O MIE assay and conducted CSF sample measurements and analysis. KML, NJI, LW, and RY developed, validated A β O gel electrophoresis/native WB assay, and KML, NJI, LW, RY, and MEH conducted CSF sample measurements and data analysis. RJG performed the pharmacokinetic analysis. All authors interpreted data. CSD contributed to statistical analysis. SMC, KML, MEH, MG, and NJI wrote the paper. All authors read and approved the final manuscript.

Funding

This work was supported by grants from the National Institute on Aging (AG057780 to SMC) and by Cognition Therapeutics, Inc. Content is solely the authors' and does not represent the National Institutes of Health.

Availability of data and materials

All data needed to evaluate the conclusions are presented here or available upon request.

Declarations

Ethics approval and consent to participate

Informed consent form was required to be signed by the subject and caregiver or study partner before exposure to any study-related procedure, including screening tests for eligibility. The Institutional Review Board (IRB; Advarra IRB, Columbia, MD) reviewed the protocol and informed consent form. The Investigators were required to conduct all the study aspects in accordance with applicable Regulatory national, state and local laws.

Consent to publication

Not applicable.

Competing interests

Current/former employees of Cognition Therapeutics (Cognition): NJI, KML, RY, LW, MEH, SMC, AOC. Cognition paid consultants: CMY, JRC, RJG, CSD, MG, RM. LSS grants/fees: Eli Lilly, Merck, Roche/Genentech, Avraham, Boehringer Ingelheim, Neurim, Neuronix, Cognition, Eisai, Takeda, vTv, Abbott, Biogen, Novartis, Biohaven, outside this work. SD Chair of: Acumen Medical Scientific Advisory Board, Biogen Drug Monitoring Committee, Cognition Medical Advisory Board, DSMB for Prevail Pharmaceuticals and for pepinemab (Vaccinex, Inc.), Associate Editor of Neurotherapeutics, Editor for Dementia for UpToDate. HZ: SABs/as consultant for Abbvie, Acumen, Alektor, Alzinova, ALZPath, Annexon, Apellis, Artery Therapeutics, AZTherapeutics, Cognition, Denali, Eisai, Nervgen, Novo Nordisk, Optoceutics, Passage Bio, Pinteon Therapeutics, Prothena, Red Abbey Labs, reMYND, Roche, Samumed, Siemens Healthineers, Triplet Therapeutics, Wave, has given lectures in symposia sponsored by Cellectricon, Fujirebio, Alzecure, Biogen, Roche, is co-founder of Brain Biomarker Solutions (BBS, of GU Ventures Incubator; outside this work). KB: consultant, advisory boards, data monitoring committees Abcam, Axon, Biogen, Cognition, Julius Clinical, Lilly, MagQu, Novartis, Siemens Healthineers, Roche Diagnostics, is co-founder of BBS. International Patent WO 15/116923 pertains to this paper. YIS received a grant to the University of Pennsylvania supporting this research. ABH has lectured or served at SABs for Abbvie, Alektor, BioArctic, Biogen, Eisai, Janssen, Novo Nordisk, Roche.

Received: 23 January 2023 Accepted: 22 April 2023

Published: 12 May 2023

References

- Viola KL, Klein WL. Amyloid beta oligomers in Alzheimer's disease pathogenesis, treatment, and diagnosis. *Acta Neuropathol.* 2015;129:183–206.
- Cleary JP, Walsh DM, Hofmeister JJ, Shankar GM, Kuskowski MA, Selkoe DJ, et al. Natural oligomers of the amyloid- β protein specifically disrupt cognitive function. *Nat Neurosci.* 2005;8(1):79–84.
- Izzo NJ, Yuede CM, LaBarbera KM, Limegrover CS, Rehak C, Yurko R, et al. Preclinical and clinical biomarker studies of CT1812: a novel approach to Alzheimer's disease modification. *Alzheimers Dement.* 2021;17(8):1365–82.
- Gandy S, Ehrlich ME. Moving the needle on Alzheimer's disease with an anti-oligomer antibody. *New Engl J Med.* 2023;388(1):80–1.
- Lizama BN, Kahle J, Catalano SM, Caggiano AO, Grundman M, Hamby ME. Sigma-2 receptors—from basic biology to therapeutic target: a focus on age-related degenerative diseases. *Int J Mol Sci.* 2023;24(7):6251.
- Gobom J, Parnetti L, Rosa-Neto P, Vyhnaek M, Gauthier S, Cataldi S, et al. Validation of the LUMIPULSE automated immunoassay for the measurement of core AD biomarkers in cerebrospinal fluid. *Clin Chem Lab Med.* 2021;60(2):207–19.
- Havugimana PC, Hart GT, Nepusz T, Yang H, Turinsky AL, Li Z, et al. A census of human soluble protein complexes. *Cell.* 2012;150(5):1068–81.
- Bateman RJ, Wen G, Morris JC, Holtzman DM. Fluctuations of CSF amyloid- β levels: Implications for a diagnostic and therapeutic biomarker. *Neurology.* 2007;68(9):666–9.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

