



UvA-DARE (Digital Academic Repository)

Mechanisms Underlying Alcohol Approach Bias and Its Modification

Wiers, R.W.; van Dessel, P.

DOI

[10.1016/j.bpsgos.2023.02.010](https://doi.org/10.1016/j.bpsgos.2023.02.010)

Publication date

2023

Document Version

Final published version

Published in

Biological Psychiatry: Global Open Science

License

CC BY-NC-ND

[Link to publication](#)

Citation for published version (APA):

Wiers, R. W., & van Dessel, P. (2023). Mechanisms Underlying Alcohol Approach Bias and Its Modification. *Biological Psychiatry: Global Open Science*, 3(3), 317-318. <https://doi.org/10.1016/j.bpsgos.2023.02.010>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: <https://uba.uva.nl/en/contact>, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Mechanisms Underlying Alcohol Approach Bias and Its Modification

Reinout W. Wiers and Pieter van Dessel

In the current issue of *Biological Psychiatry: Global Open Science*, an interesting article by Chen *et al.* (1) argues that the mechanisms underlying alcohol approach bias may overlap with those underlying Pavlovian-to-instrumental transfer (PIT). Patients who showed a stronger PIT effect in a task without alcohol cues showed a stronger tendency to approach alcohol cues in an indirect alcohol approach avoidance task, and this correlation was stronger in individuals with more severe alcohol dependence and in individuals who scored high on trait impulsivity. The strength of the behavioral approach bias was also related to neural activation of the nucleus accumbens in the PIT task, an area involved in reinforcement learning, processing of alcohol cues, and craving. What do these findings tell us about mechanisms underlying an approach bias for alcohol in alcohol dependence and what is the clinical relevance of these findings?

Alcohol-dependent patients, as well as heavy drinkers, have often been demonstrated to show an automatically activated tendency to approach alcohol cues, the so-called alcohol approach bias. This is clinically relevant, as a number of well-powered clinical randomized controlled trials have demonstrated that this approach bias can be modified by approach bias modification (ApBM), which has yielded a consistent reduction in relapse rates of approximately 10% 1 year after treatment discharge, when added to abstinence-oriented clinical treatment [for review see (2) and references 30–33 in (1)]. While clinically relevant, we should acknowledge that this is a small effect, similar to the effect size of the most effective medication in this domain. The small effect size also makes clear that ApBM is not a silver bullet solution and that its effects are not yet well understood. For instance, it is unclear why this training is not effective in changing drinking behavior in healthy volunteers or in problem drinkers who wish to reduce their drinking in an online training (3); that is to say, ApBM helps them reduce their drinking, but not better than sham training (in contrast to the clinical findings). A better understanding of the cognitive (and neurological) mechanisms underlying the alcohol approach bias might aid the development of more effective varieties of training.

Chen *et al.* (1) interpret their findings from dual process models that inspired the original development of ApBM and other varieties of cognitive bias modification that differentiate between impulsive/associative processes and reflective processes [e.g., (4)]. Specifically, it is argued that both approach bias and PIT effects may depend on the activation of impulsive processes that also underpin alcohol dependence. From this perspective, the results can be interpreted as indicating that the (neuro)cognitive processes of (a subgroup of) addicted

people are qualitatively different from nonaddicted people. This is related to discussions surrounding the chronic brain disease model of addiction (5) as it may seem that (a subgroup of) addicted people have lost their ability to voluntarily make choices when faced with conditioned stimuli related to their addiction.

Notably, basic cognitive research on ApBM in healthy volunteers has yielded results that better fit with a single-process inferential perspective than with a dual-process perspective (6). For example, training effects require conscious awareness and can sometimes be generated by instruction only, rather than requiring repeated training. From an inferential account, behavior is the result of inferences (e.g., causal inferences or predictions) that are goal directed, and this is also true if the behavior is self-defeating in the long run, as is the case with addictive behaviors. The predictive brain evokes behavior (e.g., drinking alcohol) because it predicts that this behavior can be a means to achieve a goal (e.g., relaxation after stress). Importantly, these inferential processes can often be highly automatic and context dependent, such that a given situational cue (e.g., stress) may readily evoke goal-directed inferences to engage in (addictive) behavior. This alternative account suggests that it is important to include contextual antecedents as well as goals in training, as proposed in a new variety of cognitive bias modification: ABC training (7). In this new variety of training, patients train in personally relevant antecedent contexts (A) (e.g., coming home stressed) to achieve their goals (consequences [C]) (e.g., relaxation) in another way than by drinking alcohol (behavioral alternatives [B]) (e.g., going for a walk). This new variety has yielded promising results in healthy volunteers (8) but still awaits testing in patients.

It is important to note that from the inferential perspective, addiction is not the result of qualitative differences in processing for (subgroups of) addictive people. Instead, differences in (addictive) behavior result from people having learned to apply different automatic inferences throughout their (ontogenetic and epigenetic) learning history (their learned inferential network) (Figure 1). It is important to study which inferences underlie addictive behavior and to what extent practicing alternative inferences allows to better bring behavior under voluntary control by helping patients automatize behaviors that are more in line with their long-term objectives.

From this perspective, the results of Chen *et al.* (1) might be informative because they indicate that for alcohol-dependent patients, alcohol approach bias and PIT rely on similar inferences. For instance, both PIT and alcohol approach bias may depend on cue-based inferences that a certain outcome is wanted [goal activation inferences that also play a role in

SEE CORRESPONDING ARTICLE ON PAGE 443




Figure 1. Illustration of the mental processes underlying Pavlovian-to-instrumental transfer (PIT) and alcohol approach bias according to dual-process and inferential accounts. In the dual-process account, a cue (e.g., an image of a beer or a fractal image previously paired with a wanted outcome) may lead to the automatic activation of mental associations (e.g., the association between beer and approach or between the fractal image and a response that previously led to the wanted outcome). Depending on the dominance of their impulsive system (trait impulsivity), this activation may determine approach bias (e.g., faster responses to beer cues) and PIT responding (e.g., more responses that previously led to the wanted outcome). In contrast, in

the inferential account, the same cues are thought to evoke automatic inferences such as the inference that a certain outcome is wanted and that one is likely to readily engage in actions that previously served this outcome. These inferences depend on a person's network of learned inferences and may determine approach bias and PIT responding.

nudging effects; see (9)] or inferences that one is likely to readily engage in behavior that fosters these outcomes [goal-directed active inferences; see (10)]. These inferences may relate to both alcohol dependence and trait impulsivity and partly rely on neurological activation in networks that play a role in reward prediction, including the nucleus accumbens. Furthermore, changes in these inferences might relate to treatment success, which might explain the observed stronger association between approach bias and PIT in patients who relapsed (although the evidence for this effect was weak and the effect warrants replication). It can therefore be useful to assess whether targeting such inferences (as is done in ABC training) provides opportunities for intervention.

In sum, we agree that testing the (neuro-)cognitive mechanisms underlying the approach bias for alcohol, as well as other biases in information processing that play a role in addiction, is important for a better scientific understanding of the puzzles surrounding addiction. As such, the results of Chen *et al.* (1) provide information that could also be used for improving treatment, especially when this information is integrated with other empirical findings in reference to recent theorizing. Ultimately, it remains to be tested whether inference-based cognitive training indeed has stronger effects in patients as would be predicted from theory and promising findings, and how this relates to PIT and its neural underpinnings.

Acknowledgments and Disclosures

RWW acknowledges funding from the Centre for Urban Mental Health at the University of Amsterdam and a national gravitation grant from the Dutch Research Council (NWO Grant No. 024.004.016, "New Science of Mental Disorders"). Pvd acknowledges funding from Ghent University (Grant No. BOF/STA/202202/004).

The authors report no biomedical financial interests or potential conflicts of interest.

Article Information

From the Addiction Development and Psychopathology Lab, Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands (RWW);

Centre for Urban Mental Health, University of Amsterdam, Amsterdam, The Netherlands (RWW); and Department of Experimental-Clinical and Health Psychology, Ghent University, Ghent, Belgium (PvD).

Address correspondence to Reinout W. Wiers, Ph.D., at r.wiers@uva.nl.

Received Feb 27, 2023; accepted Feb 27, 2023.

References

- Chen K, Garbusow M, Sebold M, Kuitunen-Paul S, Smolka MN, Huys QJM, *et al.* (2023): Alcohol approach bias is associated with both behavioral and neural Pavlovian-to-instrumental transfer effects in alcohol-dependent patients. *Biol Psychiatry Glob Open Sci* 3:443–450.
- Wiers RW, Pan T, Van Dessel P, Rinck M, Lindenmeyer J (2023): Approach bias retraining and other training interventions as add-on in the treatment of AUD patients. In: Sommer WH, Spanagel R, editors. *Current Topics in Behavioral Neuroscience*. Berlin/Heidelberg, Germany: Springer. Available at: https://link.springer.com/chapter/10.1007/7854_2023_421.
- Wiers RW, Boffo M, Field M (2018): What's in a trial? On the importance of distinguishing between experimental lab studies and randomized controlled trials: The case of cognitive bias modification and alcohol use disorders. *J Stud Alcohol Drugs* 79:333–343.
- Stacy AW, Wiers RW (2010): Implicit cognition and addiction: A tool for explaining paradoxical behavior. *Annu Rev Clin Psychol* 6:551–575.
- Heather N, Field M, Moss AC, Satel S (2022): *Evaluating the Brain Disease Model of Addiction*. Abingdon, UK: Routledge.
- Van Dessel P, Hughes S, De Houwer J (2019): How do actions influence attitudes? An inferential account of the impact of action performance on stimulus evaluation. *Personal Soc Psychol Rev* 23:267–284.
- Wiers RW, Van Dessel P, Köpzet C (2020): ABC training: A new theory-based form of cognitive-bias modification to foster automatization of alternative choices in the treatment of addiction and related disorders. *Curr Dir Psychol Sci* 29:499–505.
- Van Dessel P, Cummins J, Wiers RW (in press): ABC-training as a new intervention for hazardous alcohol drinking: Two proof-of-principle randomized pilot studies. *Addiction*.
- Van Dessel P, Boddez Y, Hughes S (2022): Nudging societally relevant behavior by promoting cognitive inferences. *Sci Rep* 12:1–11.
- Parr T, Pezzulo G, Friston K (2022): *Active Inference: The Free Energy Principle in Mind, Brain, and Behavior*. Cambridge, MA: MIT Press.