



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

FeSiO₄H₂ stabilized at subducting slab conditions: A geologically viable water carrier into the Earth's lower mantle

Citation for published version:

Shao, S, Lv, J, Liu, H, Lu, W, Liu, P, Hermann, A, Wang, Y, Chen, C & Ma, Y 2023, 'FeSiO₄H₂ stabilized at subducting slab conditions: A geologically viable water carrier into the Earth's lower mantle', *Physical Review B*, vol. 108, no. 21, 214107, pp. 1-7. <https://doi.org/10.1103/PhysRevB.108.214107>

Digital Object Identifier (DOI):

[10.1103/PhysRevB.108.214107](https://doi.org/10.1103/PhysRevB.108.214107)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Physical Review B

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



FeSiO₄H₂ Stabilized at Subducting Slab Conditions: A Geologically Viable Water Carrier into the Earth's Lower Mantle

Sen Shao,^{1,2,*} Jian Lv,^{1,2,*} Hanyu Liu,^{1,2,3,*} Wencheng Lu,^{1,2} Peng Liu,^{1,2}
Andreas Hermann,⁴ Yanchao Wang,^{1,2,†} Changfeng Chen,^{5,‡} and Yanming Ma^{1,2,3,§}

¹Key Laboratory of Material Simulation Methods & Software of Ministry of Education,
College of Physics, Jilin University, Changchun 130012, China

²State Key Laboratory of Superhard Materials, College of Physics, Jilin University, Changchun 130012, China

³International Center of Future Science, Jilin University, Changchun 130012, China

⁴Centre for Science at Extreme Conditions and Scottish Universities Physics Alliance,
School of Physics and Astronomy, The University of Edinburgh, Edinburgh EH9 3FD, UK

⁵Department of Physics and Astronomy, University of Nevada, Las Vegas, NV 89154, USA

Hydrous minerals hold the key to unlocking the enduring mystery of water cycle deep inside Earth. Tremendous efforts have been devoted to identifying geologically viable minerals meeting stringent pressure-temperature-density stability requirements for descent into deep Earth, and such pursuits remain active. Here, we identify two hydrous iron silicates, α - and β -FeSiO₄H₂, formed by reaction of Earth-abundant FeSiO₃ and H₂O and stabilized at the pressure-temperature conditions in cold subducting slabs. These phases have sufficiently high density for stable descent into Earth's lower mantle, and then decompose to release water after reaching equilibrium with the mantle geotherm. Moreover, Mg(Fe)SiO₄H₂ solutions are found to be more stable than the pure substances and can serve as effective carriers to transport substantial amounts of water to lower-mantle regions via the cold subduction zones. These findings establish a viable and robust material basis for deep-Earth water cycle, with major implications for elucidation of many prominent geological processes.

Water (H₂O) has tremendous influence on physical and chemical properties of the Earth's constituent compositions [1–3]. The knowledge about the transport of water into the Earth's interior is crucial to understanding its evolution and dynamics. It was estimated that a large amount of water ($\sim 10^{11}$ kg/year) is continuously transported into the Earth's interior through subduction zones, while only a fraction of the water degases to the Earth's surface [4, 5]. This imbalance indicates that a large amount of water is delivered and stored in the Earth's interior by deep, global water circulation [6–11]. The transition zone is known to be a major water reservoir [12, 13]. There is also evidence of significant amounts of water stored in the lower mantle; for example, the phenomenon of partial melting that offers a reasonable interpretation of the observed low velocity regions near the top of lower mantle and at the core-mantle boundary can occur only in the presence of abundant water [14].

It is widely accepted that water is carried into the Earth's interior by hydrous minerals in the the uppermost of the descending cold plate [15, 16]. Extensive past studies focused on various dense hydrous magnesium silicates, such as phase A (Mg₇Si₂O₁₄H₆), phase E (Mg₂SiO₆H₄), superhydrous phase B (Mg₁₀Si₃O₁₈H₄), phase D (MgSi₂O₆H₂), and phase H (MgSiO₄H₂) [17–27], as potential deep-Earth water carriers [28]. However, these dense hydrous magnesium silicates are not fully compatible with the stability requirements in the geologically relevant environments, as some dissociate into an assemblage of nominally anhydrous phases plus water at the lower-mantle pressure and temperature conditions (<1,500 km) [16], while all of them possess lower den-

sity compared to the Preliminary Reference Earth Model (PREM) data. When these low-density hydrous minerals located at the uppermost of the subducting slab, one possible scenario might arise, wherein low-density hydrous minerals may undergo upward transportation through geological activities, encompassing diapirs. [29–32]. As a result, certain portions of the low-density hydrous minerals have the potential to be recycled to the overriding plate instead of being further transported into the deep interior of the Earth. Recently, several hydroxide phases (e.g., δ -AlOOH and Ca(OH)₂) and their solid solutions with hydrous silicates were found stable in a wider pressure-temperature stability range [27, 33–39], and these hydrous minerals were proposed as potential carriers to transport water into the lower mantle depths. However, challenges remain in finding more minerals with high gravitational stability and Earth abundance that have major impact on water transport into the deep lower mantle [40]. Therefore, finding geologically viable carriers for deep water circulation remains an outstanding problem that requires new insights and further exploration from materials physics perspectives.

Bridgmanite (Mg,Fe)SiO₃ is considered the most abundant mineral in Earth's lower mantle, and it was thought to have very low water storage capacities [41, 42]. Previous studies revealed that the presence of Fe, which is an increasingly more abundant element at greater depths, may significantly change the stability of silicates and their water solubility [43–47]. An important issue, therefore, is whether FeSiO₃ can react with water to form hydrous phases at relevant pressure and temperature conditions. In this work, we address this is-

sue by evaluating the stability of H_2O -saturated FeSiO_3 under pressure, using an advanced structure prediction method in conjunction with first-principle calculations. Our search found two hitherto unknown Fe-bearing hydrous silicate phases that are stable at a wide range of pressures (18-61 GPa) and temperatures ($<1,450$ K). Moreover, these hydrous Fe silicates have consistently notable higher density compared to the PREM data, giving them superior gravitational stability during descent into the lower-mantle depths. The stability at the pressure-temperature-density conditions of the subducting slabs makes FeSiO_4H_2 geologically viable carrier for transporting significant amounts of water into the Earth's deep interior.

To identify stable FeSiO_4H_2 phases, we employed an advanced structure search method and its same name code CALYPSO [48–51], which has been successfully employed in predicting the crystal structures of a variety of multiple-component minerals [34, 36, 52]. We carried out variable-cell calculations at 20, 50 and 100 GPa with one to four formula units per simulated cell, where 60% lowest enthalpy structures were retained to produce the next-generation structures by a particle swarm optimization algorithm and the remaining 40% structures were randomly generated under the symmetry constraint. Our *ab initio* calculations were performed in the framework of density-functional theory within the Perdew-Burke-Ernzerh (PBE) generalized gradient approximation [53], as implemented in the VASP code [54]. The all-electron projector-augmented wave (PAW) method [55] was adopted with $3p^6 3d^4 4s^1$, $3s^2 3p^2$, $2s^2 2p^4$ and $1s^1$ treated as valence electrons for Fe, Si, O and H atoms, respectively. The GGA+U approach [56] was used to describe the correlation effects among the localized Fe $3d$ electrons, adopting the recently proposed values for the on-site Coulomb interaction $U=5.0$ eV and Hund's coupling $J=0.8$ eV [39, 57]. The plane-wave basis set cutoff energy of 1,000 eV and Monkhorst-Pack Brillouin zone sampling grid of $2\pi \times 0.03 \text{ \AA}^{-1}$ were used to ensure the enthalpy convergence of better than 1 meV/atom. To establish the pressure-temperature phase diagram, we calculated the Gibbs free energy in the frame work of quasi-harmonic approximation as implemented in PHONOPY code [58]. We simulated the Gibbs free energy of liquid water using the thermodynamic properties and equations of states [see Supplemental Materials (SM) [59]].

Our structure search yielded two hitherto unknown Fe-based hydrous silicate phases, α - and β - FeSiO_4H_2 (See Table S1 for detailed structural information [59]). The α - FeSiO_4H_2 is in an antiferromagnetic ordering (Fig. S1) with the space group of $P2_1/c$. The crystal lattice parameters are $a=5.12 \text{ \AA}$, $b=4.69 \text{ \AA}$, $c=9.36 \text{ \AA}$, $\alpha=\gamma=90^\circ$, and $\beta=96.9^\circ$ at 20 GPa. This structure is similar to that of δ - AlOOH , where Al is replaced by Fe and Si atoms, which are octahedrally coordinated with six O atoms [Fig. 1(a)]. The SiO_6 and FeO_6 octahedrons




FIG. 1. The crystal structure of (a) α - FeSiO_4H_2 and (b) β - FeSiO_4H_2 . The Fe, Si, O and H atoms are represented by brown, blue, red and pink spheres, respectively.

are connected by edge or vertex sharing, and H atoms occupy SiO_6 and FeO_6 octahedral interstices and bond with O atoms to form OH dipoles in the crystal lattice. The β - FeSiO_4H_2 phase also adopts an antiferromagnetic ordering with the same space group, and its crystal lattice parameters are $a=4.77 \text{ \AA}$, $b=9.13 \text{ \AA}$, $c=4.67 \text{ \AA}$, $\alpha=\gamma=90^\circ$, and $\beta=85.1^\circ$ at 40 GPa. While β - FeSiO_4H_2 retains the main structural features of α - FeSiO_4H_2 [Fig. 1(b)], the octahedrons formed by SiO_6 and FeO_6 in β - FeSiO_4H_2 are only connected by vertex sharing, leading to denser packing. These structures of FeSiO_4H_2 do not contain symmetric hydrogen bonds, in contrast to many hydrous minerals stable at lower mantle pressure conditions (e.g., MgSiO_4H_2 , FeOOH , and AlOOH), which all contain symmetric hydrogen bonds.

To evaluate the stability of FeSiO_4H_2 , we constructed a four-component convex hull, which takes into account all known stable compounds formed by Fe, Si, O, and H elements. The convex hull results (Figs. S2 and S3) show that FeSiO_4H_2 remains stable when many decomposing reaction paths are taken into account. We also considered various related minerals in assessing the stability of FeSiO_4H_2 with respect to decomposition at high pressures [Fig. 2(a) and Fig. S4]. In the current stability assessment, we considered only the minerals known to be major components in the Earth, including Fe_2O_3 , FeO , SiO_2 , FeSiO_3 , Fe_2SiO_4 , and H_2O . Our calculations show [Fig. 2(a)] that α - FeSiO_4H_2 becomes energetically more favorable above 18 GPa, then transforms at 35 GPa into β - FeSiO_4H_2 , which remains stable up to ~ 61 GPa, before decomposing to FeSiO_3 and H_2O . Furthermore, calculated phonon spectra show no imaginary modes in the pressure range of 18~61 GPa [Fig. S6], confirming the dynamic stability of FeSiO_4H_2 in the entire pressure range where it is energetically stable.

Dense hydrous silicates are known to be thermody-





FIG. 2. (a) Calculated enthalpy as a function of pressure of β -FeSiO₄H₂, Fe+Fe₂O₃+3SiO₂+3H₂O, FeO+SiO₂+H₂O, FeSiO₃+H₂O and Fe₂SiO₄+H₂O-FeO measured relative to α -FeSiO₄H₂. Negative relative enthalpy indicates that FeSiO₄H₂ is stable in the pressure ranges of 18-61 GPa, which spans the pressures in the upper mantle (UM), mantle transition zone (MTZ) and lower mantle (LM), as indicated by the distinct color-shaded zones. (b) The pressure-temperature phase diagram of FeSiO₄H₂. The boundary between α - and β -FeSiO₄H₂ phases is shown by the gray solid line. Purple and orange dashed lines represent the dissociation boundary where FeSiO₄H₂ decomposes into FeO+SiO₂+H₂O and FeSiO₃+H₂O, respectively. The gray zone represents the geothermal conditions of the subducting slab, while the blue line indicates the geothermal conditions of the mantle. The pressure boundary of the mantle transition zone (MTZ) and the lower mantle (LM) is about 25 GPa as indicated at the bottom of the panel.

namically stable at pressure-temperature conditions relevant to the subducting slabs, which are lower compared to those along the normal mantle geotherm. To examine the stability fields of FeSiO₄H₂, we calculated its Gibbs free energy, especially along the following two possible routes for the dissociation of FeSiO₄H₂ involving several minerals known to be among the most

abundant in the Earth's mantle and H₂O as reactants:



Water turns into liquid phase at elevated temperatures, so we simulated the Gibbs free energy of liquid water using the thermodynamic properties and equations of states from previous work [59]. Results in Fig. 2(b) show that α -FeSiO₄H₂ is stabilized up to \sim 35 GPa at 0 K, and thermal stability of α -FeSiO₄H₂ is achieved at 20, 25, 30, and 32 GPa around 1,190, 1,230, 1,310, and 1,350 K, respectively, indicating a positive Clapeyron slope. The threshold pressure for the phase transition from α - to β -FeSiO₄H₂ decreases as the temperature rises. Meanwhile, the β -FeSiO₄H₂ phase spans a wider pressure-temperature stability field, from 32-61 GPa and up to 1,450 K. The threshold temperature for the thermal stability of the FeSiO₄H₂ phases first increases with rising pressure, then decreases steeply when pressure exceeds 40 GPa, which is similar to the behavior of phase H MgSiO₄H₂ under pressure [24–26].

To further assess the structural stability of the hydrous Fe silicates, we performed *ab initio* molecular dynamics simulations to evaluate the mean square displacements (MSD) (Fig. S7), which indicate that all the atoms in α -FeSiO₄H₂ at 20 GPa & 1,000 K and β -FeSiO₄H₂ at 40 GPa & 1,500 K fluctuate within a small range around the equilibrium positions, indicating that these atoms remain near their lattice sites. These MSD results offer compelling evidence for the structural stability of FeSiO₄H₂ at pertinent high-pressure high-temperature conditions.

The existence of lower-mantle water reservoirs hinges on availability of hydrous minerals that can transport water into the lower mantle without premature dehydration. Our calculations show that the FeSiO₄H₂ phases are thermodynamically stable at pressures from 18 to 61 GPa and temperatures up to 1,450 K. The temperature for the dissociation boundary of FeSiO₄H₂ is lower than those of typical mantle geotherm and, as a result, these hydrous Fe silicates cannot form the basis for long-term water storage in the lower mantle. It is, however, important to note that FeSiO₄H₂, which accommodate much larger amount of water (\sim 12 wt%) than the water carriers in the transition zone (\sim 1-3 wt% water in wadsleyite and ringwoodite) [13, 60], can serve as an effective, albeit transient, water carrier deep into the lower mantle via the transport of cold subducting slabs.

Thermodynamic stability is a widely considered key requirement for minerals to serve as potential water carriers into the deep Earth. Meanwhile, another equally important but less considered requirement is gravitational stability, having an influence on the depth to which hydrous minerals descend into the deep Earth's interior alongside the subducting slab. In the uppermost of the subduct-




FIG. 3. (a) Gibbs free energies of $\text{Fe}_x\text{Mg}_{1-x}\text{SiO}_4\text{H}_2$ relative to those of pure substances MgSiO_4H_2 and FeSiO_4H_2 at 35 GPa and 0, 500 and 1000 K. (b) Density of hydrous minerals AlOOH , MgSiO_4H_2 , FeSiO_4H_2 , $(\text{Fe}_{0.45}\text{Mg}_{0.55})\text{SiO}_4\text{H}_2$ and $(\text{Fe}_{0.7}\text{Mg}_{0.3})\text{SiO}_4\text{H}_2$ compared with that of Earth's mantle according to the Preliminary Reference Earth Model [61].

ing slab, a portion of hydrous minerals with densities smaller than that of PREM[61] (Fig. 3) have the potential to form diapirs, which may facilitate the return of some subducting materials to the overriding plate [29–32]. In this crucial regard, FeSiO_4H_2 has the notable advantage of possessing clearly higher density compared to that of the PREM, making it superior in gravitational stability for descent into deep lower mantle regions.

The subducting slab is composed of ocean sediments, basaltic oceanic crust and peridotitic mantle. The composition of peridotitic mantle is similar to that of normal mantle, where magnesium silicates are prevalent, while the ocean sediments and basaltic oceanic crust (uppermost of the subducting slab) possess Fe-rich environment[62, 63]. Additional, previous investigations have provided compelling evidence supporting the presence of Fe-rich minerals in the Mg-rich environment interior of the Earth [64–68]. It is therefore likely that Fe incorporated hydrous mineral $\text{Mg}(\text{Fe})\text{SiO}_4\text{H}_2$ will form in the

uppermost of the subducting slab. To assess the viability of this scenario, we evaluated the Gibbs free energy for the formation of solid solutions of MgSiO_4H_2 and FeSiO_4H_2 at pertinent pressure-temperature conditions. It is noted that configurational entropy exerts a dominant influence on the thermodynamic stability in solution systems, while the electronic and vibrational entropy have negligible contributions [69–71]. We calculated relative Gibbs free energy (ΔG) defined by

$$\Delta G = \Delta H - TS_{conf} \quad (3)$$

$$\begin{aligned} \Delta H = & H_{\text{Fe}_x\text{Mg}_{1-x}\text{SiO}_4\text{H}_2} - xH_{\text{FeSiO}_4\text{H}_2} \\ & - (1-x)H_{\text{MgSiO}_4\text{H}_2}, \end{aligned} \quad (4)$$

where T and S_{conf} in Eq. (3) are temperature and configurational entropy of $\text{Fe}_x\text{Mg}_{1-x}\text{SiO}_4\text{H}_2$ solution, and the three terms on the right-hand side in Eq. (4) are enthalpies of the solution, MgSiO_4H_2 and FeSiO_4H_2 , respectively. The calculated ΔG is negative for the solution in a full range of Fe/Mg ratios [Fig. 3(a)], indicating the stability of $\text{Mg}(\text{Fe})\text{SiO}_4\text{H}_2$ at rising temperatures.

We further evaluated the gravitational stability of $\text{Mg}(\text{Fe})\text{SiO}_4\text{H}_2$ solutions. The results in Fig. 3(b) show that the density of $\text{Fe}_{0.45}\text{Mg}_{0.55}\text{SiO}_4\text{H}_2$ becomes comparable to that of PREM at pressures corresponding to the depths of the mantle transition zone and the uppermost of the lower mantle; increasing Fe content in $\text{Fe}_{0.7}\text{Mg}_{0.3}\text{SiO}_4\text{H}_2$ extends its gravitational stability at the depths further inside the lower mantle. These results demonstrate that Fe-rich $\text{Mg}(\text{Fe})\text{SiO}_4\text{H}_2$ solutions, serving as potential carriers in the uppermost of the subducting slab, are likely to restrain their involvement in the formation of diapirs and have great advantage to transport water into Earth's deep lower mantle. The Fe-poor $\text{Mg}(\text{Fe})\text{SiO}_4\text{H}_2$ solutions might exist in the peridotitic mantle of subducting slab and carry water in deep interior of Earth for its good thermodynamic stability under subducting slab conditions.

Geological studies suggest a nearly dry Bridgmanite dominated environment in vast lower mantle regions [41]. This scenario stems from a lack of known hydrous minerals that can be stabilized in the deep lower-mantle geotherm conditions [24–27]. There is, however, experimental evidence showing that Al could enhance the thermal stability of hydrous magnesium silicates [72, 73]. For example, it was observed that Al-bearing phase $\text{H MgSiO}_4\text{H}_2$ remains stable at pressure and temperature conditions corresponding to the depths greater than 2,600 km in the subducting slab [72, 73]. One may expect that the same phenomenon could occur in Al incorporated FeSiO_4H_2 given its structural similarity with hydrous magnesium silicate. An expanded range of thermodynamic stability combined with the gravitational stability makes hydrous Fe silicates geologically viable water carriers into the deep lower mantle.

In summary, we have pursued a rational design idea leading to the identification of hydrous iron silicates as geologically viable water carrier into the Earth's lower mantle, based on the consideration of both thermodynamic and gravitational stability of the targeted minerals together with their Earth abundance as key criteria. We employed crystal structure search in conjunction with first-principles energetic calculations to predict two distinct structural phases of FeSiO_4H_2 that are stabilized in the pressure range of 18-61 GPa and temperatures up to $\sim 1,450$ K. The superior gravitational stability and favorable thermodynamic stability make the newly identified FeSiO_4H_2 promising carrier to transport this hydrous mineral in subducting slabs then release water deep into the lower mantle. Our study also suggests that a wide range of hydrous Fe/Mg silicates may serve as a viable and robust material basis for water cycle in Earth's deep interior, with major implications for understanding the evolution of constituent compositions in deep Earth.

This work was supported by National Natural Science Foundation of China (Grant Nos. 52090024, 12074138, T2225013 and 12174142), Jilin Province Science and Technology Development Program (Grant No. YDZJ202102CXJD016), Program for Jilin University Science and Technology Innovative Research Team (2021TD-05), Program for Jilin University Computational Interdisciplinary Innovative Platform, the Strategic Priority Research Program of Chinese Academy of Sciences (Grant No. XDB33000000), and the High-Performance Computing Centre of Jilin University.

* These authors contribute equally

† wyc@calypso.cn

‡ changfeng.chen@unlv.edu

§ mym@jlu.edu.cn

- [1] S. I. Karato, The role of hydrogen in the electrical conductivity of the upper mantle, *Nature* **347**, 272 (1990).
- [2] T. Kubo, E. Ohtani, T. Kato, T. Shinmei, and K. Fujino, Effects of water on the α - β transformation kinetics in San Carlos olivine, *Science* **281**, 85 (1998).
- [3] E. Ohtani and K. Litasov, The effect of water on mantle phase transitions, *Rev. Mineral. Geochem.* **62**, 397 (2006).
- [4] P. E. van Keken, B. R. Hacker, E. M. Syracuse, and G. A. Abers, Subduction factory: 4. Depth-dependent flux of H_2O from subducting slabs worldwide, *J. Geophys. Res.* **116**, B1 (2011).
- [5] S. A. Peacock, Fluid processes in subduction zones, *Science* **248**, 329 (1990).
- [6] T. J. Ahrens, Water storage in the mantle, *Nature* **342**, 122 (1989).
- [7] Q. Bai and D. Kohlstedt, Substantial hydrogen solubility in olivine and implications for water storage in the mantle, *Nature* **357**, 672 (1992).
- [8] D. R. Bell and G. R. Rossman, Water in Earth's mantle: the role of nominally anhydrous minerals, *Science* **255**, 1391 (1992).
- [9] M. Murakami, K. Hirose, H. Yurimoto, S. Nakashima, and N. Takafuji, Water in Earth's lower mantle, *Science* **295**, 1885 (2002).
- [10] A. V. Sobolev, E. V. Asaflov, A. A. Gurenko, N. T. Arndt, V. G. Batanova, M. V. Portnyagin, D. Garbe-Schönberg, A. H. Wilson, and G. R. Byerly, Deep hydrous mantle reservoir provides evidence for crustal recycling before 3.3 billion years ago, *Nature* **571**, 555 (2019).
- [11] L. J. Hallis, G. R. Huss, K. Nagashima, G. J. Taylor, S. A. Halldórsson, D. R. Hilton, M. J. Mottl, and K. J. Meech, Evidence for primordial water in Earth's deep mantle, *Science* **350**, 795 (2015).
- [12] X. Huang, Y. Xu, and S. I. Karato, Water content in the transition zone from electrical conductivity of wadsleyite and ringwoodite, *Nature* **434**, 746 (2005).
- [13] D. Pearson, F. Brenker, F. Nestola, J. McNeill, L. Nasdala, M. Hutchison, S. Matveev, K. Mather, G. Silversmit, S. Schmitz, *et al.*, Hydrous mantle transition zone indicated by ringwoodite included within diamond, *Nature* **507**, 221 (2014).
- [14] A. H. Peslier, M. Schönbacher, H. Busemann, and S. I. Karato, Water in the Earth's interior: distribution and origin, *Space Sci. Rev.* **212**, 743 (2017).
- [15] H. W. Green II, W.-P. Chen, and M. R. Brudzinski, Seismic evidence of negligible water carried below 400-km depth in subducting lithosphere, *Nature* **467**, 828 (2010).
- [16] E. Ohtani, The role of water in Earth's mantle, *Natl. Sci. Rev.* **7**, 224 (2020).
- [17] A. Ringwood and A. Major, High-pressure reconnaissance investigations in the system Mg_2SiO_4 - MgO - H_2O , *Earth Planet. Sci. Lett.* **2**, 130 (1967).
- [18] M. Kanzaki, Stability of hydrous magnesium silicates in the mantle transition zone, *Phys. Earth Planet. Inter.* **66**, 307 (1991).
- [19] E. Ohtani, H. Mizobata, Y. Kudoh, T. Nagase, H. Arashi, H. Yurimoto, and I. Miyagi, A new hydrous silicate, a water reservoir, in the upper part of the lower mantle, *Geophys. Res. Lett.* **24**, 1047 (1997).
- [20] E. Ohtani, H. Mizobata, and H. Yurimoto, Stability of dense hydrous magnesium silicate phases in the systems Mg_2SiO_4 - H_2O and MgSiO_3 - H_2O at pressures up to 27 GPa, *Phys. Chem. Minerals* **27**, 533 (2000).
- [21] D. J. Frost and Y. Fei, Stability of phase D at high pressure and high temperature, *J. Geophys. Res. Solid Earth* **103**, 7463 (1998).
- [22] S. R. Shieh, H. K. Mao, R. J. Hemley, and L. C. Ming, Decomposition of phase D in the lower mantle and the fate of dense hydrous silicates in subducting slabs, *Earth Planet. Sci. Lett.* **159**, 13 (1998).
- [23] R. E. Pacalo and J. B. Parise, Crystal structure of superhydrous B, a hydrous magnesium silicate synthesized at 1400 °C and 20 GPa, *Am. Mineral.* **77**, 681 (1992).
- [24] J. Tsuchiya, First principles prediction of a new high-pressure phase of dense hydrous magnesium silicates in the lower mantle, *Geophys. Res. Lett.* **40**, 4570 (2013).
- [25] E. Ohtani, Y. Amaike, S. Kamada, T. Sakamaki, and N. Hirao, Stability of hydrous phase H MgSiO_4H_2 under lower mantle conditions, *Geophys. Res. Lett.* **41**, 8283 (2014).
- [26] M. Nishi, T. Irifune, J. Tsuchiya, Y. Tange, Y. Nishihara, K. Fujino, and Y. Higo, Stability of hydrous silicate at high pressures and water transport to the deep lower mantle, *Nat. Geosci.* **7**, 224 (2014).

- [27] L. Bindi, M. Nishi, J. Tsuchiya, and T. Irifune, Crystal chemistry of dense hydrous magnesium silicates: The structure of phase H, MgSiH_2O_4 , synthesized at 45 GPa and 1000 °C, *Am. Mineral.* **99**, 1802 (2014).
- [28] E. Ohtani, Hydrous minerals and the storage of water in the deep mantle, *Chem. Geol.* **418**, 6 (2015).
- [29] M. D. Behn, P. B. Kelemen, G. Hirth, B. R. Hacker, and H.-J. Massonne, Diapirs as the source of the sediment signature in arc lavas, *Nat. Geosci.* **4**, 641 (2011).
- [30] H. R. Marschall and J. C. Schumacher, Arc magmas sourced from mélange diapirs in subduction zones, *Nat. Geosci.* **5**, 862 (2012).
- [31] A. Schaarschmidt, K. Haase, P. Voudouris, V. Melfos, and R. Klemm, Migration of arc magmatism above mantle wedge diapirs with variable sediment contribution in the aegaeon, *Geochemistry Geophys. Geosystems* **22**, e2020GC009565 (2021).
- [32] I. Safonova, S. Maruyama, and K. Litasov, Generation of hydrous-carbonated plumes in the mantle transition zone linked to tectonic erosion and subduction, *Tectonophysics* **662**, 454 (2015).
- [33] L. Zhang, H. Yuan, Y. Meng, and H.-K. Mao, Discovery of a hexagonal ultradense hydrous phase in $(\text{Fe}, \text{Al})\text{OOH}$, *Proc. Natl. Acad. Sci.* **115**, 2908 (2018).
- [34] X. Zhong, A. Hermann, Y. Wang, and Y. Ma, Monoclinic high-pressure polymorph of AlOOH predicted from first principles, *Phys. Rev. B* **94**, 224110 (2016).
- [35] X. Su, C. Zhao, C. Lv, Y. Zhuang, N. Salke, L. Xu, H. Tang, H. Gou, X. Yu, Q. Sun, *et al.*, The effect of iron on the sound velocities of $\delta\text{-AlOOH}$ up to 135 GPa, *Geosci. Front.* **12**, 937 (2021).
- [36] S. Shao, J. Bi, P. Gao, G. Liu, M. Zhou, J. Lv, Y. Xie, and Y. Wang, Stability of $\text{Ca}(\text{OH})_2$ at Earth's deep lower mantle conditions, *Phys. Rev. B* **104**, 014107 (2021).
- [37] Q. Hu, D. Y. Kim, W. Yang, L. Yang, Y. Meng, L. Zhang, and H. K. Mao, FeO_2 and FeOOH under deep lower-mantle conditions and earth's oxygen-hydrogen cycles, *Nature* **534**, 241 (2016).
- [38] J. Liu, Q. Hu, D. Young Kim, Z. Wu, W. Wang, Y. Xiao, P. Chow, Y. Meng, V. B. Prakapenka, H. K. Mao, *et al.*, Hydrogen-bearing iron peroxide and the origin of ultralow-velocity zones, *Nature* **551**, 494 (2017).
- [39] C. Lu and C. Chen, High-pressure evolution of crystal bonding structures and properties of FeOOH , *J. Phys. Chem. Lett.* **9**, 2181 (2018).
- [40] J. Tsuchiya and E. C. Thompson, The role of hydrogen bonds in hydrous minerals stable at lower mantle pressure conditions, *Prog. Earth Planet. Sci.* **9**, 1 (2022).
- [41] Z. Liu, H. Fei, L. Chen, C. McCammon, L. Wang, R. Liu, F. Wang, B. Liu, and T. Katsura, Bridgmanite is nearly dry at the top of the lower mantle, *Earth Planet. Sci. Lett.* **570**, 117088 (2021).
- [42] S. Fu, J. Yang, S. I. Karato, A. Vasiliev, M. Y. Presniakov, A. G. Gavriluk, A. G. Ivanova, E. H. Hauri, T. Okuchi, N. Purevjav, *et al.*, Water concentration in single-crystal (Al, Fe) -bearing Bridgmanite grown from the hydrous melt: Implications for dehydration melting at the topmost lower mantle, *Geophys. Res. Lett.* **46**, 10346 (2019).
- [43] C. Xu, T. Inoue, J. Gao, M. Noda, and S. Kakizawa, Melting phase relation of Fe-bearing Phase D up to the uppermost lower mantle, *Am. Mineral.* **107**, 343 (2022).
- [44] G. Ganskow and F. Langenhorst, Stability and crystal chemistry of iron-bearing dense hydrous magnesium silicates, *Geochemistry* **74**, 489 (2014).
- [45] B. B. Karki, D. B. Ghosh, and S. I. Karato, Behavior and properties of water in silicate melts under deep mantle conditions, *Sci. Rep.* **11**, 10588 (2021).
- [46] K. D. Litasov and E. Ohtani, Effect of water on the phase relations in Earth's mantle and deep water cycle, *Advances in High-Pressure Mineralogy* **421**, 115 (2007).
- [47] Y. H. Zhao, S. Ginsberg, and D. L. Kohlstedt, Solubility of hydrogen in olivine: dependence on temperature and iron content, *Contrib. Mineral. Petrol.* **147**, 155 (2004).
- [48] Y. Wang, J. Lv, L. Zhu, and Y. Ma, Crystal structure prediction via particle-swarm optimization, *Phys. Rev. B* **82**, 094116 (2010).
- [49] Y. Wang, J. Lv, L. Zhu, and Y. Ma, CALYPSO: A method for crystal structure prediction, *Comput. Phys. Commun.* **183**, 2063 (2012).
- [50] B. Gao, P. Gao, S. Lu, J. Lv, Y. Wang, and Y. Ma, Interface structure prediction via CALYPSO method, *Sci. Bull.* **64**, 301 (2019).
- [51] X. Shao, J. Lv, P. Liu, S. Shao, P. Gao, H. Liu, Y. Wang, and Y. Ma, A symmetry-orientated divide-and-conquer method for crystal structure prediction, *J. Chem. Phys.* **156**, 014105 (2022).
- [52] J. Zhang, J. Lv, H. Li, X. Feng, C. Lu, S. A. Redfern, H. Liu, C. Chen, and Y. Ma, Rare helium-bearing compound FeO_2He stabilized at deep-earth conditions, *Phys. Rev. Lett.* **121**, 255703 (2018).
- [53] J. P. Perdew, J. Chevary, S. Vosko, K. A. Jackson, M. R. Pederson, D. Singh, and C. Fiolhais, Erratum: Atoms, molecules, solids, and surfaces: Applications of the generalized gradient approximation for exchange and correlation, *Phys. Rev. B* **48**, 4978 (1993).
- [54] G. Kresse and J. Furthmüller, Efficient iterative schemes for ab initio total-energy calculations using a plane-wave basis set, *Phys. Rev. B* **54**, 11169 (1996).
- [55] P. E. Blöchl, Projector augmented-wave method, *Phys. Rev. B* **50**, 17953 (1994).
- [56] L. Wang, T. Maxisch, and G. Ceder, Oxidation energies of transition metal oxides within the GGA+U framework, *Phys. Rev. B* **73**, 195107 (2006).
- [57] Q. Hu, D. Y. Kim, J. Liu, Y. Meng, L. Yang, D. Zhang, W. L. Mao, and H.-K. Mao, Dehydrogenation of goethite in Earth's deep lower mantle, *Proc. Natl. Acad. Sci.* **114**, 1498 (2017).
- [58] A. Togo, F. Oba, and I. Tanaka, First-principles calculations of the ferroelastic transition between rutile-type and CaCl_2 -type SiO_2 at high pressures, *Phys. Rev. B* **78**, 134106 (2008).
- [59] See supplemental material at url for detailed first principles calculations and experimental method, detailed structural information and phonon spectra.
- [60] J. R. Smyth, $\beta\text{-Mg}_2\text{SiO}_4$; a potential host for water in the mantle?, *Am. Mineral.* **72**, 1051 (1987).
- [61] A. M. Dziewonski and D. L. Anderson, Preliminary reference Earth model, *Phys. Earth and Planet. Inter.* **25**, 297 (1981).
- [62] Z. Chemia, D. Dolejš, and G. Steinle-Neumann, Thermal effects of variable material properties and metamorphic reactions in a three-component subducting slab, *J. Geophys. Res. Solid Earth* **120**, 6823 (2015).
- [63] R. J. Stern, Subduction zones, *Rev. Geophys.* **40**, 3 (2002).
- [64] S. M. Dorfman, Y. Meng, V. B. Prakapenka, and T. S. Duffy, Effects of Fe-enrichment on the equation of state

- and stability of (Mg, Fe)SiO₃ perovskite, *Earth Planet. Sci. Lett.* **361**, 249 (2013).
- [65] L. Bindi, S.-H. Shim, T. G. Sharp, and X. Xie, Evidence for the charge disproportionation of iron in extraterrestrial bridgmanite, *Sci. Adv.* **6**, eaay7893 (2020).
- [66] W. L. Mao, Y. Meng, G. Shen, V. B. Prakapenka, A. J. Campbell, D. L. Heinz, J. Shu, R. Caracas, R. E. Cohen, Y. Fei, *et al.*, Iron-rich silicates in the Earth's D'' layer, *Proc. Natl. Acad. Sci.* **102**, 9751 (2005).
- [67] P. Nimis, F. Nestola, M. Schiazza, R. Reali, G. Agrosi, D. Mele, G. Tempesta, D. Howell, M. T. Hutchison, and R. Spiess, Fe-rich ferropiclasite and magnesiowüstite inclusions reflecting diamond formation rather than ambient mantle, *Geology* **47**, 27 (2019).
- [68] R. Wirth, L. Dobrzhinetskaya, B. Harte, A. Schreiber, and H. Green, High-Fe (Mg, Fe) O inclusion in diamond apparently from the lowermost mantle, *Earth and Planetary Science Letters* **404**, 365 (2014).
- [69] A. Benisek and E. Dachs, The vibrational and configurational entropy of disordering in Cu₃Au, *J. Alloys Compd.* **632**, 585 (2015).
- [70] W. R. Panero and R. Caracas, Stability of phase H in the MgSiO₄H₂-AlOOH-SiO₂ system, *Earth Planet. Sci. Lett.* **463**, 171 (2017).
- [71] A. Manzoor, S. Pandey, D. Chakraborty, S. R. Phillpot, and D. S. Aidhy, Entropy contributions to phase stability in binary random solid solutions, *NPJ Comput. Mater.* **4**, 47 (2018).
- [72] I. Ohira, E. Ohtani, T. Sakai, M. Miyahara, N. Hirao, Y. Ohishi, and M. Nishijima, Stability of a hydrous δ -phase, AlOOH-MgSiO₂(OH)₂, and a mechanism for water transport into the base of lower mantle, *Earth Planet. Sci. Lett.* **401**, 12 (2014).
- [73] S. Ghosh and M. W. Schmidt, Melting of phase D in the lower mantle and implications for recycling and storage of H₂O in the deep mantle, *Geochim. Cosmochim. Acta* **145**, 72 (2014).