

Original citation:

Elliott, Charles M. and Fritz, Hans. (2015) Time-periodic solutions of advection-diffusion equations on moving hypersurfaces. *SIAM Journal on Mathematical Analysis*, 47 (3). pp. 1693-1718.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/83695>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

First Published in *SIAM Journal on Mathematical Analysis* in 47 (3), 2015 published by the Society for Industrial and Applied Mathematics (SIAM). Copyright © by SIAM. Unauthorized reproduction of this article is prohibited.

A note on versions:

The version presented in WRAP is the published version or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

TIME-PERIODIC SOLUTIONS OF ADVECTION-DIFFUSION EQUATIONS ON MOVING HYPERSURFACES*

CHARLES M. ELLIOTT[†] AND HANS FRITZ[†]

Abstract. In this paper we study time-periodic solutions to advection-diffusion equations of a scalar quantity u on a periodically moving n -dimensional hypersurface $\Gamma(t) \subset \mathbb{R}^{n+1}$. We prove existence and uniqueness of solutions in suitable Hölder spaces.

Key words. periodic solutions, advection diffusion, moving surface, existence and uniqueness

AMS subject classifications. 35A01, 35A02, 35B10, 35R01, 58J35

DOI. 10.1137/140977060

1. Introduction. In this paper we consider the advection and diffusion equation

$$(1.1) \quad \Delta_{\Gamma(t)} u - u \nabla_{\Gamma(t)} \cdot v - \partial^\bullet u = f$$

for a scalar quantity $u : \overline{\mathcal{G}}_t \rightarrow \mathbb{R}$ on the space-time hypersurface $\mathcal{G}_t := \bigcup_{t \in (0, T)} \Gamma(t) \times \{t\} \subset \mathbb{R}^{n+2}$, where $\Gamma(t)$ denotes a closed n -dimensional moving hypersurface $\Gamma(t) \subset \mathbb{R}^{n+1}$. Such equations arise in many applications such as processes on biological cell surfaces [10] and the transport of surfactants on fluid interfaces [14]. The variational Hilbert space theory for the initial value problem has been considered in [1].

Here the motion of the hypersurface is assumed to be periodic in time with period T in the sense that $\Gamma(0) = \Gamma(T)$. The velocity, $v : \mathcal{G}_t \rightarrow \mathbb{R}^{n+1}$, of the moving hypersurface $\Gamma(t)$ is given by $v = v_N + v_T$, where v_N denotes the normal velocity of $\Gamma(t)$ and v_T is an advective velocity field tangential to $\Gamma(t)$. The tangential gradient and the Laplace–Beltrami operator on $\Gamma(t)$ are denoted by $\nabla_{\Gamma(t)}$ and $\Delta_{\Gamma(t)}$, respectively. The material derivative $\partial^\bullet u$ of a function u defined in a neighborhood of $\Gamma(t)$ is given by

$$(1.2) \quad \partial^\bullet u = u_t + v_N \cdot \nabla u + v_T \cdot \nabla_{\Gamma(t)} u,$$

where ∇ denotes the usual gradient in \mathbb{R}^{n+1} . A solution $u : \overline{\mathcal{G}}_t \rightarrow \mathbb{R}$ of the advection-diffusion equation (1.1) is called periodic if

$$(1.3) \quad u(\cdot, 0) = u(\cdot, T) \quad \text{on} \quad \Gamma(0).$$

The aim of this work is to establish the existence and uniqueness of periodic solutions. Since the mass $m(t) := \int_{\Gamma(t)} u(\cdot, t) d\sigma(t)$ of a solution u of the advection-diffusion equation (1.1) evolves according to

$$m'(t) = - \int_{\Gamma(t)} f(\cdot, t) d\sigma(t),$$

*Received by the editors July 10, 2014; accepted for publication (in revised form) February 6, 2015; published electronically May 5, 2015. The research of the second author was supported by the Alexander von Humboldt Foundation, Germany, and by a Feodor Lynen Research Fellowship in collaboration with the University of Warwick, UK.

<http://www.siam.org/journals/sima/47-3/97706.html>

[†]Mathematics Institute, University of Warwick, Coventry, CV4 7AL, UK (C.M.Elliott@warwick.ac.uk, H.Fritz@warwick.ac.uk).

a necessary condition for u to be periodic is that

$$(1.4) \quad \int_0^T \int_{\Gamma(t)} f d\sigma(t) dt = 0.$$

However, please note that neither f nor the velocity v is supposed to be periodic. That is, in general $\lim_{t \searrow 0} f(\cdot, t) \neq \lim_{t \nearrow T} f(\cdot, t)$ as well as $\lim_{t \searrow 0} v(\cdot, t) \neq \lim_{t \nearrow T} v(\cdot, t)$. Although we prove a slightly more general result, the main result of this paper is that for each given mass $m_0 \in \mathbb{R}$ there exists a unique periodic solution $u \in \mathcal{H}_{2+\alpha}(\mathcal{G}_t)$ of the advection-diffusion equation (1.1) with initial mass $m(0) = m_0$. Here $\mathcal{H}_{2+\alpha}(\mathcal{G}_t)$ denotes a suitable Hölder space, which we introduce below.

Since for $f = 0$ the mass $m(t)$ of a solution u is constant in time, u describes a conservative scalar quantity in this case. In order to prove existence and uniqueness of periodic solutions to the advection-diffusion equation (1.1) we have to look at slightly more general linear parabolic PDEs on \mathcal{G}_t . In fact, we have to consider advection-diffusion equations of the form

$$(1.5) \quad \Delta_{\Gamma(t)} u - \mathbf{c}u - \partial^\bullet u = f,$$

where $\mathbf{c} : \mathcal{G}_t \rightarrow \mathbb{R}$ is a scalar function on the space-time hypersurface \mathcal{G}_t that is not required to be periodic. Unfortunately, it turns out that the periodicity condition in (1.3) is too restrictive in this case even if we assume (1.4), since in general the mass of a solution u to (1.5) cannot be periodic in time, that is, $m(0) \neq m(T)$. The reason for this is the zero-order term in (1.5). For example, for $f = 0$ and $\mathbf{c} = \nabla_{\Gamma(t)} \cdot v + \alpha$ with $\alpha \in (0, +\infty)$ the mass decays exponentially, that is, $m(t) = m(0) \exp(-\alpha t)$. Hence, a periodic solution would have to satisfy $m(0) = \int_{\Gamma(0)} u(\cdot, 0) d\sigma(0) = 0$ in this case, which is, however, a much too restrictive assumption for applications. Therefore, we slightly relax the notion of periodicity to

$$(1.6) \quad u(\cdot, 0) - \int_{\Gamma(0)} u(\cdot, 0) d\sigma(0) = u(\cdot, T) - \int_{\Gamma(0)} u(\cdot, T) d\sigma(0) \quad \text{on } \Gamma(0),$$

where $\int_{\Gamma} u d\sigma := \frac{1}{|\Gamma|} \int_{\Gamma} u d\sigma$ denotes the mean value of u on Γ . Fortunately, for the advection-diffusion equation (1.1) this condition is equivalent to the condition (1.3) as long as the constraint (1.4) is satisfied.

The study of time-periodic solutions to both linear and nonlinear parabolic equations has a long history; see, for example, [16] and references therein, and in particular [12], [15], and [21], as well as [2] and [17]. However, we are not aware of any previous analytic work that studies the periodic problem on moving hypersurfaces. Our methodology reformulates the problem on a moving closed hypersurface to a problem on a fixed hypersurface with time varying coefficients. In order to obtain some analytic results it is then useful to relate the hypersurface equation to a Neumann-type boundary problem on a flat domain; see the appendix for further explanations. Observe that the problem in our case is slightly more involved than the results on the oblique derivative problems for flat domains in [16], since we cannot assume that the zero-order term \mathbf{c} is nonnegative. For example for the choice $\mathbf{c} = \nabla_{\Gamma(t)} \cdot v$ this is certainly not true in general. The fact that the zero-order term \mathbf{c} can be negative is indeed the reason why this paper is beyond the scope of previous works. Furthermore, the slightly weaker periodicity condition (1.6), which ensures the existence of periodic solutions for any initial mass $m(0) = m_0 \in \mathbb{R}$, does not seem to have been used in the literature so far. Anyway, it is still possible to adopt the techniques used

by Lieberman in [16] to our problem. Therefore, our existence proof mainly relies on a simple fixed point iteration and standard Fredholm theory.

Our work is partially motivated by numerical simulations of periodic solutions of advection-diffusion equations on moving two-dimensional surfaces for $f = 0$ performed in [11] using an evolving surface finite element method. The periodic solutions were obtained by computing the initial value problem for arbitrary chosen initial conditions. Indeed, the numerical solutions for different initial conditions with same initial mass appear to converge very quickly to the same time-periodic solution. We would like to emphasize that in this numerical work the formulation (1.1), which entirely avoids the use of local coordinates and surface parametrisations, is very suitable for the evolving surface finite element method.

This paper is organized as follows. First, we introduce the notation and the parabolic Hölder spaces on the space-time hypersurface \mathcal{G}_t as well as their associated norms. Then we rewrite the advection-diffusion equation on the moving hypersurface $\Gamma(t)$ as an advection-diffusion equation with time-dependent coefficients on a fixed reference hypersurface \mathcal{M} . This formulation is more amenable for our purposes, since it can be easily related to a (nondegenerate) Neumann-type boundary problem on a flat domain. We derive this Neumann boundary problem in the appendix. In section 2, we summarize the results of this paper. This also serves as a reader’s guide to the proof of the main result in Theorem 3.5. The detailed proofs of all results given in section 2 can be found in section 3. In the appendix, we discuss the technique to extend a surface PDE to a nondegenerate PDE on an extended neighborhood. This result is used in sections 2 and 3 to prove existence and uniqueness to the initial boundary value problem on hypersurfaces without using any local parametrizations of the hypersurface.

2. Preliminaries. We make use of the convention to sum over repeated indices.

2.1. Hypersurfaces. Henceforward, we assume that $\Gamma(t) \subset \mathbb{R}^{n+1}$ is a family of closed (that is, compact and without boundary), orientable, connected, n -dimensional, embedded hypersurfaces of class C_1^l , with $l \in \mathbb{N}$, and that there is a closed, orientable, connected, n -dimensional, embedded hypersurface $\mathcal{M} \subset \mathbb{R}^{n+1}$ of the same class and a C_1^l -embedding $X : \bar{\mathcal{G}} \rightarrow \mathbb{R}^{n+2}$ from the closure of the cylinder $\mathcal{G} := \mathcal{M} \times (0, T) \subset \mathbb{R}^{n+2}$ onto \mathbb{R}^{n+2} such that for any $t \in [0, T]$ the map $X(\cdot, t)$ is a bijection from \mathcal{M} onto $\Gamma(t)$. This implies that $\mathcal{G}_t := \bigcup_{t \in (0, T)} \Gamma(t) \times \{t\} \subset \mathbb{R}^{n+2}$ is an $(n + 1)$ -dimensional space-time hypersurface of class C_1^l . Here, C_1^l refers to l -times continuously differentiable functions, whose derivatives are Lipschitz. Note that we use the notation $C^{2,1}$ for functions continuously differentiable in time and twice continuously differentiable in space. We also assume that the motion of $\Gamma(t)$ is periodic in time in the sense that $\Gamma(0) = \Gamma(T)$ and $X(\cdot, 0) = X(\cdot, T)$, respectively.

2.2. Tangential gradient and material derivative. Let $\mathcal{N} \subset \mathbb{R}^{n+1}$ be an arbitrary hypersurface. The tangent space to \mathcal{N} at the point $x \in \mathcal{N}$ is the linear space

$$T_x \mathcal{N} := \{ \tau \in \mathbb{R}^{n+1} \mid \exists \gamma \in C^1((-\epsilon, \epsilon), \mathbb{R}^{n+1}), \gamma((-\epsilon, \epsilon)) \subset \mathcal{N}, \gamma(0) = x, \gamma'(0) = \tau \};$$

see [9]. For a function f on an arbitrary hypersurface \mathcal{N} differentiable at $x \in \mathcal{N}$, we define the tangential gradient of f at $x \in \mathcal{N}$ by

$$\nabla_{\mathcal{N}} f(x) := \nabla \hat{f}(x) - \nu(x) \nu(x) \cdot \nabla \hat{f}(x) = P(x) \nabla \hat{f}(x),$$

where \hat{f} denotes a differentiable extension of f to an open neighborhood $U \subset \mathbb{R}^{n+1}$ of x , such that $\hat{f}|_{\mathcal{N} \cap U} = f|_{\mathcal{N} \cap U}$; see [9] for more details. Here, \cdot denotes the Euclidean scalar product in the ambient space, $\nu(x)$ is a unit normal of \mathcal{N} at x , and $P(x) = \mathbb{1} - \nu(x) \otimes \nu(x)$ is the projection onto the tangent space of \mathcal{N} at x . It can be easily shown that the tangential gradient only depends on the values of f on \mathcal{N} and hence is independent of the chosen extension \hat{f} . The components of the tangential gradient are denoted by

$$\begin{pmatrix} \underline{D}_1 f \\ \vdots \\ \underline{D}_{n+1} f \end{pmatrix} := \nabla_{\mathcal{N}} f.$$

For a twice continuously differentiable function f on \mathcal{N} we have the commutator rule

$$(2.1) \quad \underline{D}_\alpha \underline{D}_\beta f - \underline{D}_\beta \underline{D}_\alpha f = (\mathcal{H}_{\beta\eta} \nu_\alpha - \mathcal{H}_{\alpha\eta} \nu_\beta) \underline{D}_\eta f,$$

where $\mathcal{H} := \nabla_{\mathcal{N}} \nu$ denotes the (extended) Weingarten map on \mathcal{N} . The Laplace-Beltrami operator on \mathcal{N} is defined by

$$\Delta_{\mathcal{N}} f := \nabla_{\mathcal{N}} \cdot \nabla_{\mathcal{N}} f.$$

These definitions can be easily generalized to moving hypersurfaces $\Gamma(t)$, as well as to the space-derivatives on the space-time hypersurfaces \mathcal{G} and \mathcal{G}_t . Since \mathcal{G} is a cylinder the definition of the time-derivative of a function f on \mathcal{G} is obvious. On the space-time hypersurface \mathcal{G}_t we define the material derivative $\partial^\bullet f$ of a function $f : \mathcal{G}_t \rightarrow \mathbb{R}$ by

$$\partial^\bullet f := (f \circ X)_t \circ X^{-1}.$$

Since the velocity v of $\Gamma(t)$ is given by

$$v := X_t \circ X^{-1},$$

this definition is consistent with formula (1.2).

2.3. Hölder spaces. For a function $f : \mathcal{G} \rightarrow \mathbb{R}$ we define the norm

$$|f|_{0,\mathcal{G}} := \sup_{(x,t) \in \mathcal{G}} |f(x,t)|,$$

and we say that f is Hölder continuous in \mathcal{G} with exponent $\alpha \in (0, 1]$ if the seminorm

$$H_{\alpha,\mathcal{G}}(f) := \sup_{(x,t) \in \mathcal{G}} \sup_{(y,s) \in \mathcal{G} \setminus \{(x,t)\}} \frac{|f(x,t) - f(y,s)|}{|(x,t) - (y,s)|^\alpha}$$

is finite. Here $|(x,t)| := \max\{|x|, |t|^{\frac{1}{2}}\}$ with $|x| := \left(\sum_{\alpha=1}^{n+1} x_\alpha^2\right)^{\frac{1}{2}}$ is the parabolic distance in \mathbb{R}^{n+2} . Furthermore, we define the norms

$$\begin{aligned} |f|_{\alpha,\mathcal{G}} &:= |f|_{0,\mathcal{G}} + H_{\alpha,\mathcal{G}}(f), \\ |f|_{1+\alpha,\mathcal{G}} &:= |f|_{0,\mathcal{G}} + \langle f \rangle_{1+\alpha,\mathcal{G}} + |\nabla_{\mathcal{M}} f|_{\alpha,\mathcal{G}}, \\ |f|_{2+\alpha,\mathcal{G}} &:= |f|_{0,\mathcal{G}} + |\nabla_{\mathcal{M}} f|_{0,\mathcal{G}} + \langle \nabla_{\mathcal{M}} f \rangle_{1+\alpha,\mathcal{G}} + |\nabla_{\mathcal{M}}^2 f|_{\alpha,\mathcal{G}} + |f_t|_{\alpha,\mathcal{G}}, \end{aligned}$$

where

$$\langle f \rangle_{1+\alpha, \mathcal{G}} := \sup_{(x,t) \in \mathcal{G}} \sup_{(x,s) \in \mathcal{G} \setminus \{(x,t)\}} \frac{|f(x,t) - f(x,s)|}{|t - s|^{\frac{1+\alpha}{2}}}.$$

For $k = 0, 1, 2$, we introduce the following Hölder spaces on \mathcal{G} :

$$\mathcal{H}_{k+\alpha}(\mathcal{G}) := \{f : \mathcal{G} \rightarrow \mathbb{R} \mid |f|_{k+\alpha, \mathcal{G}} < \infty\}.$$

Obvious modifications of the above definitions lead to the definition of $|\cdot|_{k+\alpha, \mathcal{M}}$ for $k = 0, 1, 2$. For a function f on \mathcal{G} the following inequality holds:

$$|f(\cdot, t)|_{k+\alpha, \mathcal{M}} \leq |f|_{k+\alpha, \mathcal{G}} \quad \forall t \in [0, T], \forall \alpha \in (0, 1], k = 0, 1, 2.$$

Let the hypersurfaces \mathcal{M} and $\Gamma(t)$ as well as the embedding X be of class C^2_1 ; then the norm $|\cdot|_{k+\alpha, \mathcal{G}_t}$ on the linear space $\mathcal{H}_{k+\alpha}(\mathcal{G}_t) := \{f : \mathcal{G}_t \rightarrow \mathbb{R} \mid f \circ X \in \mathcal{H}_{k+\alpha}(\mathcal{G})\}$ is defined by

$$|f|_{k+\alpha, \mathcal{G}_t} := |f \circ X|_{k+\alpha, \mathcal{G}} \quad \text{for } k = 0, 1, 2.$$

Henceforward, d denotes the oriented distance function to $\mathcal{M} \subset \mathbb{R}^{n+1}$; see, for example, [7]. There exists $\delta > 0$ such that the decomposition

$$x = a(x) + d(x)\nu(a(x))$$

with $a(x) \in \mathcal{M}$ is unique for all $x \in \mathcal{N}_\delta$, where

$$(2.2) \quad \mathcal{N}_\delta := \{x \in \mathbb{R}^{n+1} \mid |d(x)| < \delta\}.$$

Please note that for a C^l -hypersurface, $l \geq 2$, the oriented distance function d is also of class C^l , whereas the projection $a : \mathcal{N}_\delta \rightarrow \mathcal{M}$ is of class C^{l-1} . The oriented distance function d is also Lipschitz continuous on \mathcal{N}_δ and so is the projection a , provided that the width δ is chosen sufficiently small and $l \geq 2$. This can be seen as follows:

$$\begin{aligned} |a(x) - a(y)| &\leq |x - y| + |d(x) - d(y)| + |d(y)| |\nu(a(x)) - \nu(a(y))| \\ &\leq C|x - y| + c\delta|a(x) - a(y)|. \end{aligned}$$

The extensions of the unit normal ν , of the projection P , and of the Weingarten map \mathcal{H} on \mathcal{M} to the neighborhood \mathcal{N}_δ are defined by $\nu(x) := \nabla d(x) = \nu(a(x))$, $P(x) := \mathbb{1} - \nu(x) \otimes \nu(x)$ and by $\mathcal{H}(x) := \nabla^2 d(x)$. For a function f on \mathcal{M} we define the lift to \mathcal{N}_δ by $f^l(x) := f(a(x))$. This is actually the extension of f to \mathcal{N}_δ that is constant along the normal direction to the hypersurface \mathcal{M} . A direct calculation yields

$$(2.3) \quad \nabla f^l = (\mathbb{1} - d\mathcal{H})(\nabla_{\mathcal{M}} f)^l,$$

$$(2.4) \quad (\nabla_{\mathcal{M}} f)^l = (\mathbb{1} - d\mathcal{H})^{-1} \nabla f^l;$$

compare to (2.2.13) and (2.2.16) in [8]. The definitions of the norms $|\cdot|_{k+\alpha, \mathcal{G}_\delta}$ and of the spaces $\mathcal{H}_{k+\alpha}(\mathcal{G}_\delta)$ for $k = 0, 1, 2$ on the cylinder $\mathcal{G}_\delta := \mathcal{N}_\delta \times (0, T)$ are obvious. In order to prove the norm equivalence

$$\frac{1}{C_{k,\alpha}} |f|_{k+\alpha, \mathcal{G}} \leq |f^l|_{k+\alpha, \mathcal{G}_\delta} \leq C_{k,\alpha} |f|_{k+\alpha, \mathcal{G}}$$

for functions $f : \mathcal{G} \rightarrow \mathbb{R}$, we need the following statement.

LEMMA 2.1. For functions $f, g : \mathcal{G} \rightarrow \mathbb{R}$ the following inequalities hold:

$$\begin{aligned} H_{\alpha, \mathcal{G}}(fg) &\leq H_{\alpha, \mathcal{G}}(f)|g|_{0, \mathcal{G}} + |f|_{0, \mathcal{G}}H_{\alpha, \mathcal{G}}(g), \\ |fg|_{\alpha, \mathcal{G}} &\leq |f|_{\alpha, \mathcal{G}}|g|_{0, \mathcal{G}} + |f|_{0, \mathcal{G}}|g|_{\alpha, \mathcal{G}}. \end{aligned}$$

Analogous estimates hold for the norms $|\cdot|_{k+\alpha, \mathcal{G}_\delta}$ on \mathcal{G}_δ .

Proof. The first inequality easily follows from the definition of the Hölder coefficient. The second inequality is then a direct result. \square

LEMMA 2.2. For \mathcal{M} of class C_1^3 there exist constants $C_{k, \alpha} > 0$ such that

$$(2.5) \quad \frac{1}{C_{k, \alpha}}|f|_{k+\alpha, \mathcal{G}} \leq |f^l|_{k+\alpha, \mathcal{G}_\delta} \leq C_{k, \alpha}|f|_{k+\alpha, \mathcal{G}}$$

for $k = 0, 1, 2$ and $\alpha \in (0, 1]$.

Proof. Since the lifted function f^l is constant along the normal direction, we obtain

$$|f|_{0, \mathcal{G}} = |f^l|_{0, \mathcal{G}_\delta}.$$

For the Hölder coefficient it is obvious that $H_{\alpha, \mathcal{G}}(f) \leq H_{\alpha, \mathcal{G}_\delta}(f^l)$. Furthermore, since the projection a is Lipschitz-continuous on \mathcal{N}_δ for $\delta > 0$ sufficiently small, it follows that

$$\begin{aligned} H_{\alpha, \mathcal{G}_\delta}(f^l) &= \sup_{\substack{(x,t), (y,s) \in \mathcal{G}_\delta \\ (x,t) \neq (y,s)}} \frac{|f^l(x,t) - f^l(y,s)|}{|(x,t) - (y,s)|^\alpha} \\ &\leq \sup_{\substack{(x,t), (y,s) \in \mathcal{G}_\delta \\ (x,t) \neq (y,s)}} \frac{|(a(x), t) - (a(y), s)|^\alpha}{|(x,t) - (y,s)|^\alpha} \sup_{\substack{(x,t), (y,s) \in \mathcal{G}_\delta \\ (x,t) \neq (y,s)}} \frac{|f(a(x), t) - f(a(y), s)|}{|(a(x), t) - (a(y), s)|^\alpha} \\ &\leq CH_{\alpha, \mathcal{G}}(f). \end{aligned}$$

This already establishes $|f|_{\alpha, \mathcal{G}} \leq |f^l|_{\alpha, \mathcal{G}_\delta} \leq C|f|_{\alpha, \mathcal{G}}$. From Lemma 2.1 and formulas (2.3) and (2.4) we then obtain

$$\begin{aligned} |f|_{1+\alpha, \mathcal{G}} &= |f|_{0, \mathcal{G}} + \langle f \rangle_{1+\alpha, \mathcal{G}} + |\nabla_{\mathcal{M}} f|_{\alpha, \mathcal{G}} \leq |f^l|_{0, \mathcal{G}_\delta} + \langle f^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |(\nabla_{\mathcal{M}} f)^l|_{\alpha, \mathcal{G}_\delta} \\ &\leq |f^l|_{0, \mathcal{G}_\delta} + \langle f^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |A^{-1} \nabla f^l|_{\alpha, \mathcal{G}_\delta} \\ &\leq C(|f^l|_{0, \mathcal{G}_\delta} + \langle f^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |\nabla f^l|_{\alpha, \mathcal{G}_\delta}), \end{aligned}$$

and conversely,

$$\begin{aligned} |f^l|_{1+\alpha, \mathcal{G}_\delta} &= |f^l|_{0, \mathcal{G}_\delta} + \langle f^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |\nabla f^l|_{\alpha, \mathcal{G}_\delta} = |f^l|_{0, \mathcal{G}_\delta} + \langle f^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |A(\nabla_{\mathcal{M}} f)^l|_{\alpha, \mathcal{G}_\delta} \\ &\leq C(|f^l|_{0, \mathcal{G}_\delta} + \langle f^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |(\nabla_{\mathcal{M}} f)^l|_{\alpha, \mathcal{G}_\delta}) \\ &\leq C(|f|_{0, \mathcal{G}} + \langle f^l \rangle_{1+\alpha, \mathcal{G}} + |\nabla_{\mathcal{M}} f|_{\alpha, \mathcal{G}}), \end{aligned}$$

where $A := \mathbb{1} - d\mathcal{H}$. Similarly, the result for $k = 2$ can be deduced from

$$\begin{aligned} (\nabla^2 f^l)_{\alpha\beta} &= A_{\alpha\delta} A_{\beta\gamma} (D_\delta D_\gamma f)^l + \nabla_\alpha A_{\beta\gamma} (D_\gamma f)^l, \\ (\nabla_{\mathcal{M}}^2 f)_{\alpha\beta}^l &= (A^{-1})^{\alpha\delta} (A^{-1})^{\beta\gamma} \nabla_\delta \nabla_\gamma f^l + (A^{-1})^{\alpha\delta} \nabla_\delta (A^{-1})^{\beta\gamma} \nabla_\gamma f^l, \end{aligned}$$

which follows from formulas (2.3) and (2.4). For $|\cdot|_{2+\alpha, \mathcal{G}}$ we then obtain

$$\begin{aligned} |f|_{2+\alpha, \mathcal{G}} &= |f|_{0, \mathcal{G}} + |\nabla_{\mathcal{M}} f|_{0, \mathcal{G}} + \langle \nabla_{\mathcal{M}} f \rangle_{1+\alpha, \mathcal{G}} + |\nabla_{\mathcal{M}}^2 f|_{\alpha, \mathcal{G}} + |f_t|_{\alpha, \mathcal{G}} \\ &\leq C(|f^l|_{0, \mathcal{G}_\delta} + |(\nabla_{\mathcal{M}} f)^l|_{0, \mathcal{G}_\delta} + \langle (\nabla_{\mathcal{M}} f)^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |(\nabla_{\mathcal{M}}^2 f)^l|_{\alpha, \mathcal{G}_\delta} + |f_t^l|_{\alpha, \mathcal{G}_\delta}) \\ &\leq C(|f^l|_{0, \mathcal{G}_\delta} + |A^{-1} \nabla f^l|_{0, \mathcal{G}_\delta} + \langle A^{-1} \nabla f^l \rangle_{1+\alpha, \mathcal{G}_\delta} \\ &\quad + |((A^{-1})^{\alpha\delta} (A^{-1})^{\beta\gamma} \nabla_\delta \nabla_\gamma f^l)|_{\alpha, \mathcal{G}_\delta} \\ &\quad + |((A^{-1})^{\alpha\delta} \nabla_\delta (A^{-1})^{\beta\gamma} \nabla_\gamma f^l)|_{\alpha, \mathcal{G}_\delta} + |f_t^l|_{\alpha, \mathcal{G}_\delta}) \\ &\leq C(|f^l|_{0, \mathcal{G}_\delta} + |\nabla f^l|_{0, \mathcal{G}_\delta} + \langle \nabla f^l \rangle_{1+\alpha, \mathcal{G}_\delta} + |\nabla^2 f^l|_{\alpha, \mathcal{G}_\delta} + |\nabla f^l|_{\alpha, \mathcal{G}_\delta} + |f_t^l|_{\alpha, \mathcal{G}_\delta}). \end{aligned}$$

Since $H_{\alpha, \mathcal{G}_\delta}(\nabla f^l) \leq C(|\nabla^2 f^l|_{\alpha, \mathcal{G}_\delta} + \langle \nabla f^l \rangle_{1+\alpha, \mathcal{G}_\delta})$, we can conclude that $|f|_{2+\alpha, \mathcal{G}} \leq C|f^l|_{2+\alpha, \mathcal{G}_\delta}$. The opposite direction follows in the same way. \square

LEMMA 2.3. *The spaces $\mathcal{H}_{k+\alpha}(\mathcal{G})$, with $k \in \{0, 1, 2\}$ and $\alpha \in (0, 1]$, are Banach spaces.*

Proof. The statement follows directly from the equivalence (2.5) and the fact that the spaces $\mathcal{H}_{k+\alpha, \mathcal{G}_\delta}$, $k = 0, 1, 2$, are Banach spaces. \square

LEMMA 2.4. *Let $\alpha \in (0, 1)$. For each $\epsilon > 0$ there is a constant $C(\epsilon)$ such that*

$$(2.6) \quad |f|_{\alpha, \mathcal{G}} \leq c\epsilon^{1-\alpha}|f|_{2+\alpha, \mathcal{G}} + C(\epsilon)|f|_{0, \mathcal{G}} \quad \forall f \in \mathcal{H}_{2+\alpha}(\mathcal{G}),$$

where the constant c does not depend on ϵ .

Proof. Without loss of generality we can assume that $\epsilon < \min\{\frac{\delta}{2}, 1\}$. In the case that $|(x, t) - (y, s)| \geq \epsilon$, we directly obtain

$$\frac{|f(x, t) - f(y, s)|}{|(x, t) - (y, s)|^\alpha} \leq \frac{2|f|_{0, \mathcal{G}}}{\epsilon^\alpha}.$$

For $|(x, t) - (y, s)| < \epsilon$, we have

$$\lambda(x, t) + (1 - \lambda)(y, s) \in \mathcal{N}_{\frac{\delta}{2}} \times [0, T] \quad \forall \lambda \in [0, 1],$$

and there is some $(\xi, \chi) = \lambda^*(x, t) + (1 - \lambda^*)(y, s)$ with $\lambda^* \in [0, 1]$ such that

$$\begin{aligned} |f(x, t) - f(y, s)| &= |f^l(x, t) - f^l(y, s)| = |\nabla f^l(\xi, \chi) \cdot (x - y) + f_t^l(\xi, \chi)(t - s)| \\ &\leq |\nabla f^l|_{0, \mathcal{G}_\delta}|x - y| + |f_t^l|_{0, \mathcal{G}_\delta}|t - s| \\ &\leq |\nabla f^l|_{0, \mathcal{G}_\delta}|x - y| + |f_t^l|_{0, \mathcal{G}_\delta}|t - s|^{\frac{1}{2}} \\ &\leq (|\nabla f^l|_{0, \mathcal{G}_\delta} + |f_t^l|_{0, \mathcal{G}_\delta})|(x, t) - (y, s)| \\ &\leq |f^l|_{2+\alpha, \mathcal{G}_\delta}|(x, t) - (y, s)| \\ &\leq c|f|_{2+\alpha, \mathcal{G}}|(x, t) - (y, s)|, \end{aligned}$$

where we have used (2.5) in the last step. Hence, we obtain

$$\frac{|f(x, t) - f(y, s)|}{|(x, t) - (y, s)|^\alpha} \leq c|f|_{2+\alpha, \mathcal{G}}|(x, t) - (y, s)|^{1-\alpha} \leq c\epsilon^{1-\alpha}|f|_{2+\alpha, \mathcal{G}}. \quad \square$$

2.4. Reformulation on a stationary hypersurface. In Lemma 2.5 of this subsection we reformulate the equation on the moving space-time hypersurface \mathcal{G}_t to the fixed space-time cylinder \mathcal{G} . In order to do this, we introduce a time-dependent, symmetric, and positive definite map $G : \overline{\mathcal{G}} \rightarrow \mathbb{R}^{(n+1) \times (n+1)}$ defined by

$$(2.7) \quad G := (G_{\alpha\beta})_{\alpha, \beta=1, \dots, n+1} \quad \text{with} \quad G_{\alpha\beta} := \underline{D}_\alpha X \cdot \underline{D}_\beta X + \nu_\alpha \nu_\beta.$$

Since X is time-periodic, we have $G(0) = G(T)$. The term $\underline{D}_\alpha X \cdot \underline{D}_\beta X$ is a kind of Cartesian representation of the Riemannian metric (first fundamental form) induced by the embedding X on \mathcal{M} at time t . We explain in subsection 2.4.1 below how this definition can be generalized to arbitrary Riemannian metrics on \mathcal{M} .

Definition (2.7) is motivated by the following observations. For $a \in \mathcal{M}$ the map $\hat{G} := ((\nabla_{\mathcal{M}} X)^T \nabla_{\mathcal{M}} X)(a) : T_a \mathcal{M} \rightarrow T_a \mathcal{M}$ is a bijective linear map on $T_a \mathcal{M}$, because X is an embedding. By adding a term $(\nu \otimes \nu)(a)$ to \hat{G} in the definition of $G(a)$, the map \hat{G} is extended to a bijective map on \mathbb{R}^{n+1} . We write $G^{\alpha\beta}$ for the components of the inverse G^{-1} . Please also note that the eigenvalues of G are the eigenvalues of the map \hat{G} and the eigenvalue 1 with corresponding eigenvector ν .

Henceforward, the volume form $d\sigma(t)$ on $\Gamma(t)$ is given by the n -dimensional Hausdorff measure, whereas the volume form $do(t)$ on \mathcal{M} is the corresponding volume form weighted by the density $\sqrt{\det G(t)}$. That is, $do(t) := \sqrt{\det G(t)} d\sigma$, where $d\sigma$ is the volume form on \mathcal{M} induced by the n -dimensional Hausdorff measure. We also use the notation do_0 instead of $do(0)$. Because of the periodicity we have $do_0 = do(T)$. From Jacobi's formula we immediately obtain that $\frac{d}{dt} \sqrt{\det G(t)} = \frac{1}{2} \sqrt{\det G(t)} G^{\alpha\beta} G_{t\alpha\beta}(t)$ and hence

$$(2.8) \quad \frac{d}{dt} \int_{\mathcal{M}} \tilde{f}(\cdot, t) do(t) = \int_{\mathcal{M}} \tilde{f}_t(\cdot, t) + \frac{1}{2} G^{\alpha\beta} G_{t\alpha\beta} \tilde{f}(\cdot, t) do(t).$$

Below, it will become clear how this formula is related to the transport formula on the moving hypersurface $\Gamma(t)$, (see Theorem 5.1 of [9]),

$$(2.9) \quad \frac{d}{dt} \int_{\Gamma(t)} f(\cdot, t) d\sigma(t) = \int_{\Gamma(t)} \partial^\bullet f(\cdot, t) + (\nabla_{\Gamma(t)} \cdot v) f(\cdot, t) d\sigma(t).$$

We define the linear elliptic operator $\Delta_{g(t)}$ by

$$(2.10) \quad \Delta_{g(t)} \tilde{u} := \underline{D}_\alpha (G^{\alpha\beta} \underline{D}_\beta \tilde{u}) + \frac{1}{2} P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta G_{\alpha\eta} \underline{D}_\rho \tilde{u},$$

where $P_{\alpha\gamma}$ denotes the components of the projection onto the tangent space of \mathcal{M} . In section 2.4.1, we show that this operator is exactly the Laplace operator on \mathcal{M} with respect to a Riemannian metric $g(t)$, which in this section is given by the metric induced by the embedding X at time t . It then follows from the invariance of the Laplace operator under reparametrizations (see p. 117 in [6]) that for a function u on $\Gamma(t)$ we have $\Delta_{g(t)}(u \circ X) = (\Delta_{\Gamma(t)} u) \circ X = (\nabla_{\Gamma(t)} \cdot \nabla_{\Gamma(t)} u) \circ X$. The advantage of the above definition, however, is that the operator is defined in the global coordinate system of the ambient space, whereas the Laplace operator with respect to a certain metric is usually defined in local coordinates. This global perspective can simplify proofs, since it is then no longer necessary to patch together local results, such as estimates, to global ones. Below, we demonstrate the practicability of our global approach by proving the identity $\Delta_{g(t)}(u \circ X) = (\Delta_{\Gamma(t)} u) \circ X$ directly in the global coordinate frame.

LEMMA 2.5. *Let $c \in \mathbb{R}$. Suppose $\mathfrak{c}, f : \mathcal{G}_t \rightarrow \mathbb{R}$ and $\tilde{\mathfrak{c}}, \tilde{f} : \mathcal{G} \rightarrow \mathbb{R}$ are such that $\tilde{\mathfrak{c}} = \mathfrak{c} \circ X$ and $\tilde{f} = f \circ X$. Then a function $u \in \mathcal{H}_{2+\alpha}(\mathcal{G}_t)$ is a solution of*

$$\begin{aligned} \Delta_{\Gamma(t)} u - cu - \partial^\bullet u &= f \quad \text{in } \mathcal{G}_t, \\ \int_{\Gamma(0)} u(\cdot, 0) d\sigma(0) &= c, \\ u(\cdot, 0) &= u(\cdot, T) - \left(\int_{\Gamma(0)} u(\cdot, T) d\sigma(0) - c \right) \quad \text{on } \Gamma(0) \end{aligned}$$

if and only if $\tilde{u} = u \circ X \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ is a solution of

$$\begin{aligned} \Delta_{g(t)}\tilde{u} - \tilde{c}\tilde{u} - \tilde{u}_t &= \tilde{f} \quad \text{in } \mathcal{G}, \\ \int_{\mathcal{M}} \tilde{u}(\cdot, 0) d\sigma_0 &= c, \\ \tilde{u}(\cdot, 0) &= \tilde{u}(\cdot, T) - \left(\int_{\mathcal{M}} \tilde{u}(\cdot, T) d\sigma_0 - c \right) \quad \text{on } \mathcal{M}. \end{aligned}$$

Proof. From the definition of the material derivative we have $\tilde{u}_t = (\partial^\bullet u) \circ X$. Now, let $\underline{D}'_\alpha u$, $\alpha = 1, \dots, n + 1$, denote the components of the tangential gradient $\nabla_{\Gamma(t)} u$. Then we have

$$(2.11) \quad \underline{D}_\beta \tilde{u} = \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X.$$

The projection $P' : \bar{\mathcal{G}}_t \rightarrow \mathbb{R}^{(n+1) \times (n+1)}$ onto the tangent space of $\Gamma(t)$ satisfies

$$(2.12) \quad P'_{\rho\eta} \circ X = G^{\alpha\beta} \underline{D}_\alpha X_\rho \underline{D}_\beta X_\eta \quad \text{on } \bar{\mathcal{G}}.$$

We thus obtain

$$\begin{aligned} \Delta_{g(t)}\tilde{u} &= G^{\alpha\beta} \underline{D}_\alpha \underline{D}_\beta \tilde{u} - G^{\alpha\gamma} G^{\beta\delta} \underline{D}_\alpha G_{\gamma\delta} \underline{D}_\beta \tilde{u} + \frac{1}{2} P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta G_{\alpha\eta} \underline{D}_\rho \tilde{u} \\ &= G^{\alpha\beta} (\underline{D}_\alpha X_\rho \underline{D}_\beta X_\eta) (\underline{D}'_\rho \underline{D}'_\eta u) \circ X + G^{\alpha\beta} \underline{D}_\alpha \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X \\ &\quad - G^{\alpha\gamma} G^{\beta\delta} \underline{D}_\alpha G_{\gamma\delta} \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X + \frac{1}{2} P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta G_{\alpha\eta} \underline{D}_\rho X_\kappa (\underline{D}'_\kappa u) \circ X \\ &= P'_{\rho\eta} \circ X (\underline{D}'_\rho \underline{D}'_\eta u) \circ X + G^{\alpha\beta} \underline{D}_\alpha \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X \\ &\quad - G^{\alpha\gamma} G^{\beta\delta} \underline{D}_\alpha \underline{D}_\gamma X_\rho \underline{D}_\delta X_\rho \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X \\ &\quad - G^{\alpha\gamma} G^{\beta\delta} \underline{D}_\gamma X_\rho \underline{D}_\alpha \underline{D}_\delta X_\rho \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X \\ &\quad + \frac{1}{2} P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} (\underline{D}_\beta \underline{D}_\alpha X_\iota \underline{D}_\eta X_\iota + \underline{D}_\alpha X_\iota \underline{D}_\beta \underline{D}_\eta X_\iota) \underline{D}_\rho X_\kappa (\underline{D}'_\kappa u) \circ X \\ &= \Delta_{\Gamma(t)} u \circ X + G^{\alpha\beta} \underline{D}_\alpha \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X - G^{\alpha\gamma} \underline{D}_\alpha \underline{D}_\gamma X_\rho (\underline{D}'_\rho u) \circ X \\ &\quad - G^{\alpha\gamma} G^{\beta\delta} \underline{D}_\alpha \underline{D}_\delta X_\rho \underline{D}_\gamma X_\rho \underline{D}_\beta X_\eta (\underline{D}'_\eta u) \circ X \\ &\quad + P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta \underline{D}_\alpha X_\iota \underline{D}_\eta X_\iota \underline{D}_\rho X_\kappa (\underline{D}'_\kappa u) \circ X \\ &= \Delta_{\Gamma(t)} u \circ X + P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} (\underline{D}_\beta \underline{D}_\alpha X_\iota - \underline{D}_\alpha \underline{D}_\beta X_\iota) \underline{D}_\eta X_\iota \underline{D}_\rho X_\kappa (\underline{D}'_\kappa u) \circ X, \end{aligned}$$

where we have used the fact that $G^{-1}\nu = \nu$ and that PG^{-1} is symmetric. From the commutator rule (2.1) it then directly follows that

$$\Delta_{g(t)}\tilde{u} = \Delta_{\Gamma(t)} u \circ X.$$

The rest of the proof is obvious. \square

Due to the result in Lemma 2.5 it is sufficient to consider the periodic problem on the reference cylinder \mathcal{G} . In particular, if $\mathbf{c} = \nabla_{\Gamma(t)} \cdot v$, then $\tilde{\mathbf{c}}$ on \mathcal{M} is given by $\tilde{\mathbf{c}} = \frac{1}{2}G^{\alpha\beta}G_{t\alpha\beta} =: \frac{1}{2}\text{tr}_{g(t)}(g_t)$. This can be seen as follows:

$$\begin{aligned} \frac{1}{2}G^{\alpha\beta}G_{t\alpha\beta} &= \frac{1}{2}G^{\alpha\beta}(\underline{D}_\alpha X_{t\eta}\underline{D}_\beta X_\eta + \underline{D}_\alpha X_\eta\underline{D}_\beta X_{t\eta}) \\ &= \frac{1}{2}G^{\alpha\beta}(\underline{D}_\alpha X_\kappa(\underline{D}'_\kappa v_\eta) \circ X \underline{D}_\beta X_\eta + \underline{D}_\alpha X_\eta\underline{D}_\beta X_\kappa(\underline{D}'_\kappa v_\eta) \circ X) \\ &= P'_{\kappa\eta} \circ X(\underline{D}'_\kappa v_\eta) \circ X = (\nabla_{\Gamma(t)} \cdot v) \circ X, \end{aligned}$$

where we have used (2.11) with $\tilde{u} = X_{t\eta}$ and $u = X_{t\eta} \circ X^{-1} = v_\eta$ as well as (2.12). This also shows the connection between formula (2.8) and the transport formula (2.9).

2.4.1. An arbitrary Riemannian metric. The main results of this paper in section 3 are also valid for maps $G : \overline{\mathcal{G}} \rightarrow \mathbb{R}^{(n+1) \times (n+1)}$ defined by

$$(2.13) \quad \begin{aligned} G(a, t)X \cdot Y &:= g(a, t)(P(a)X, P(a)Y) + (\nu(a) \cdot X)(\nu(a) \cdot Y) \\ \forall X, Y \in \mathbb{R}^{n+1}, \forall (a, t) \in \mathcal{M} \times [0, T], \end{aligned}$$

where $g(t)$ is an arbitrary (sufficiently smooth) time-dependent Riemannian metric on \mathcal{M} ; see also [13] for further details. Indeed, $G(t)$ is a kind of Cartesian representation of the metric $g(t)$. In particular, if we choose $g(t) := X^*h$ to be the (periodic) pull-back metric of the Riemannian metric h on $\Gamma(t) \subset \mathbb{R}^{n+1}$ that is induced by the Euclidean metric in \mathbb{R}^{n+1} , then definition (2.13) coincides with definition (2.7). Using integration by parts on closed hypersurfaces (see Theorem 2.10 of [9]) leads to Green’s formula for the operator $\Delta_{g(t)}$ defined in (2.10),

$$\begin{aligned} \int_{\mathcal{M}} G^{-1}(t)\nabla_{\mathcal{M}}u \cdot \nabla_{\mathcal{M}}w do(t) &= \int_{\mathcal{M}} G^{\alpha\beta}\underline{D}_\alpha u \underline{D}_\beta w \sqrt{\det G} d\sigma \\ &= - \int_{\mathcal{M}} u \underline{D}_\alpha (G^{\alpha\beta}\underline{D}_\beta w) \sqrt{\det G} + u G^{\alpha\beta}\underline{D}_\alpha \sqrt{\det G} \underline{D}_\beta w d\sigma \\ &= - \int_{\mathcal{M}} u \underline{D}_\alpha (G^{\alpha\beta}\underline{D}_\beta w) \sqrt{\det G} + \frac{1}{2}G^{\alpha\beta}G^{\rho\eta}\underline{D}_\alpha G_{\rho\eta}\underline{D}_\beta w \sqrt{\det G} d\sigma \\ &= - \int_{\mathcal{M}} u \left(\underline{D}_\alpha (G^{\alpha\beta}\underline{D}_\beta w) + \frac{1}{2}G^{\alpha\beta}G^{\rho\eta}\underline{D}_\alpha G_{\rho\eta}\underline{D}_\beta w \right) do(t) \\ &= - \int_{\mathcal{M}} u \Delta_{g(t)} w do(t), \end{aligned}$$

where we have used the fact that

$$\begin{aligned} G^{\rho\eta}\underline{D}_\alpha G_{\rho\eta} &= P_{\rho\gamma}G^{\gamma\eta}\underline{D}_\alpha G_{\rho\eta} + \nu_\rho\nu_\gamma G^{\gamma\eta}\underline{D}_\alpha G_{\rho\eta} \\ &= P_{\rho\gamma}G^{\gamma\eta}\underline{D}_\alpha G_{\rho\eta} + \nu_\rho\nu_\eta\underline{D}_\alpha G_{\rho\eta} \\ &= P_{\rho\gamma}G^{\gamma\eta}\underline{D}_\alpha G_{\rho\eta} - \underline{D}_\alpha(\nu_\rho\nu_\eta)G_{\rho\eta} \\ &= P_{\rho\gamma}G^{\gamma\eta}\underline{D}_\alpha G_{\rho\eta} - \underline{D}_\alpha\nu_\rho\nu_\rho - \nu_\eta\underline{D}_\alpha\nu_\eta \\ &= P_{\rho\gamma}G^{\gamma\eta}\underline{D}_\alpha G_{\rho\eta}. \end{aligned}$$

The last identity also shows that $\Delta_{g(t)}\tilde{u} = \underline{D}_\alpha (G^{\alpha\beta}\underline{D}_\beta \tilde{u}) + \frac{1}{2}G^{\alpha\eta}G^{\beta\rho}\underline{D}_\beta G_{\alpha\eta}\underline{D}_\rho \tilde{u}$. However, we keep formula (2.10), since it is also valid, if we would replace (2.7) by

$G_{\alpha\beta} := \underline{D}_\alpha X \cdot \underline{D}_\beta X + \lambda \nu_\alpha \nu_\beta$ with $\lambda : \mathcal{M} \rightarrow (0, +\infty)$ continuously differentiable. It can be shown that the elliptic operator $\Delta_{g(t)}$ defined in (2.10) is the usual Laplace operator on \mathcal{M} with respect to the Riemannian metric $g(t)$. In order to see this, assume that $\varphi : \Omega \subset \mathbb{R}^n \rightarrow \mathcal{M} \subset \mathbb{R}^{n+1}$ is a local parametrization of \mathcal{M} and let $\tilde{g}_{ij}(\theta) := \frac{\partial \varphi}{\partial \theta^i}(\theta) \cdot \frac{\partial \varphi}{\partial \theta^j}(\theta)$. Furthermore, let \tilde{g}^{ij} denote the components of the inverse of $\tilde{g} = (\tilde{g}_{ij})_{i,j=1,\dots,n}$. Since $P_{\alpha\beta}(\varphi(\theta)) = \tilde{g}^{ij}(\theta) \frac{\partial \varphi_\alpha}{\partial \theta^i}(\theta) \frac{\partial \varphi_\beta}{\partial \theta^j}(\theta)$, we immediately obtain from definition (2.13) that

$$(2.14) \quad G_{\alpha\beta}(\varphi(\theta), t) := \frac{\partial \varphi_\alpha}{\partial \theta^i}(\theta) \tilde{g}^{ij}(\theta) g_{jk}(\theta, t) \tilde{g}^{kl}(\theta) \frac{\partial \varphi_\beta}{\partial \theta^l}(\theta) + \nu_\alpha(\varphi(\theta)) \nu_\beta(\varphi(\theta)),$$

$$(2.15) \quad G^{\alpha\beta}(\varphi(\theta), t) := \frac{\partial \varphi_\alpha}{\partial \theta^i}(\theta) g^{ij}(\theta, t) \frac{\partial \varphi_\beta}{\partial \theta^j}(\theta) + \nu_\alpha(\varphi(\theta)) \nu_\beta(\varphi(\theta)),$$

where $g_{ij}(\theta, t) := g(\varphi(\theta), t)(\frac{\partial \varphi}{\partial \theta^i}(\theta), \frac{\partial \varphi}{\partial \theta^j}(\theta))$ are the components of the metric with respect to local coordinates and $(g^{ij}(\theta, t))_{i,j=1,\dots,n} = (g_{ij}(\theta, t))_{i,j=1,\dots,n}^{-1}$ are the components of the inverse matrix. For a function $u : \mathcal{M} \rightarrow \mathbb{R}^{n+1}$ the tangential gradient on \mathcal{M} satisfies

$$(\nabla_{\mathcal{M}} u) \circ \varphi = \tilde{g}^{ij} \frac{\partial U}{\partial \theta^i} \frac{\partial \varphi}{\partial \theta^j} \quad \text{on } \Omega,$$

where $U := u \circ \varphi$; see, for example, in [7]. Using relation (2.15), we obtain

$$\begin{aligned} \underline{D}_\alpha(G^{\alpha\beta} \underline{D}_\beta u)(\varphi(\theta)) &= \tilde{g}^{ij}(\theta) \frac{\partial \varphi_\alpha}{\partial \theta^i}(\theta) \frac{\partial}{\partial \theta^j} ((G^{\alpha\beta} \underline{D}_\beta u)(\varphi(\theta))) \\ &= \tilde{g}^{ij}(\theta) \frac{\partial \varphi_\alpha}{\partial \theta^i}(\theta) \frac{\partial}{\partial \theta^j} \left(\frac{\partial \varphi_\alpha}{\partial \theta^m} g^{mn} \frac{\partial U}{\partial \theta^n} \right) (\theta) \\ &= \frac{\partial}{\partial \theta^m} \left(g^{mn} \frac{\partial U}{\partial \theta^n} \right) (\theta) + \tilde{g}^{ij}(\theta) \frac{\partial \varphi_\alpha}{\partial \theta^i}(\theta) \cdot \frac{\partial^2 \varphi_\alpha}{\partial \theta^j \partial \theta^m}(\theta) g^{mn}(\theta) \frac{\partial U}{\partial \theta^n}(\theta). \end{aligned}$$

The same procedure gives

$$\begin{aligned} &(P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta G_{\alpha\eta} \underline{D}_\rho u) \circ \varphi \\ &= \frac{\partial \varphi_\alpha}{\partial \theta^i} g^{ij} \frac{\partial \varphi_\eta}{\partial \theta^j} \frac{\partial \varphi_\beta}{\partial \theta^m} g^{mn} \frac{\partial \varphi_\rho}{\partial \theta^n} \frac{\partial \varphi_\beta}{\partial \theta^k} \tilde{g}^{kl} \frac{\partial (G_{\alpha\eta} \circ \varphi)}{\partial \theta^l} \frac{\partial \varphi_\rho}{\partial \theta^s} \tilde{g}^{st} \frac{\partial U}{\partial \theta^t} \\ &= \frac{\partial \varphi_\alpha}{\partial \theta^i} g^{ij} \frac{\partial \varphi_\eta}{\partial \theta^j} g^{lt} \frac{\partial (G_{\alpha\eta} \circ \varphi)}{\partial \theta^l} \frac{\partial U}{\partial \theta^t} \\ &= \frac{\partial \varphi_\alpha}{\partial \theta^i} g^{ij} \frac{\partial \varphi_\eta}{\partial \theta^j} g^{lk} \frac{\partial}{\partial \theta^l} \left(\frac{\partial \varphi_\alpha}{\partial \theta^n} \tilde{g}^{nm} g_{mu} \tilde{g}^{uv} \frac{\partial \varphi_\eta}{\partial \theta^v} + (\nu_\alpha \nu_\eta) \circ \varphi \right) \frac{\partial U}{\partial \theta^k} \\ &= g^{mu} g^{lk} \frac{\partial g_{mu}}{\partial \theta^l} \frac{\partial U}{\partial \theta^k} + \frac{\partial \varphi_\alpha}{\partial \theta^i} g^{ij} \frac{\partial \varphi_\eta}{\partial \theta^j} g^{lk} g_{mu} \frac{\partial}{\partial \theta^l} \left(\frac{\partial \varphi_\alpha}{\partial \theta^n} \tilde{g}^{nm} \tilde{g}^{uv} \frac{\partial \varphi_\eta}{\partial \theta^v} \right) \frac{\partial U}{\partial \theta^k} \\ &= g^{mu} g^{lk} \frac{\partial g_{mu}}{\partial \theta^l} \frac{\partial U}{\partial \theta^k} + 2 \frac{\partial \varphi_\eta}{\partial \theta^u} g^{lk} \frac{\partial}{\partial \theta^l} \left(\tilde{g}^{uv} \frac{\partial \varphi_\eta}{\partial \theta^v} \right) \frac{\partial U}{\partial \theta^k} \\ &= g^{mu} g^{lk} \frac{\partial g_{mu}}{\partial \theta^l} \frac{\partial U}{\partial \theta^k} - 2 g^{lk} \tilde{g}^{uv} \frac{\partial \varphi_\eta}{\partial \theta^v} \frac{\partial^2 \varphi_\eta}{\partial \theta^l \partial \theta^u} \frac{\partial U}{\partial \theta^k}. \end{aligned}$$

Altogether, we obtain

$$\begin{aligned} & \left(\underline{D}_\alpha (G^{\alpha\beta} \underline{D}_\beta u) + \frac{1}{2} P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta G_{\alpha\eta} \underline{D}_\rho u \right) \circ \varphi \\ &= \frac{\partial}{\partial \theta^m} \left(g^{mn} \frac{\partial U}{\partial \theta^n} \right) + \frac{1}{2} g^{mu} g^{lk} \frac{\partial g_{mu}}{\partial \theta^l} \frac{\partial U}{\partial \theta^k} \\ &= \frac{1}{\sqrt{\det(g_{ij})}} \frac{\partial}{\partial \theta^m} \left(\sqrt{\det(g_{ij})} g^{mn} \frac{\partial U}{\partial \theta^n} \right), \end{aligned}$$

which shows that $\Delta_{g(t)}$ is indeed the Laplace operator on \mathcal{M} with respect to $g(t)$.

Our approach to define the Laplace operator with respect to $g(t)$ via the Cartesian representation $G(t)$ of the metric in (2.10) instead of using the usual definition of the Laplacian based on local coordinates is advantageous for two reasons. First, by the global perspective we can avoid the need to patch together local results, such as Schauder estimates, to global ones. Second, we believe that the global approach is more advantageous for numerical simulations; see, for example, in [13] and [19] for two different problems that are both reformulated in a global coordinate frame. See also Remark 5.1 at the end of this paper.

2.5. Strong maximum principle. The (strong) maximum principle for parabolic PDEs in flat domains (see, for example, in [20]), is also valid for parabolic PDEs on closed hypersurfaces.

LEMMA 2.6. *Suppose that the hypotheses of Theorem 3.1 hold and that \mathcal{M} is connected. Furthermore, suppose that $\Delta_{g(t)}u - \mathbf{c}u - u_t \geq 0$ in \mathcal{G} and that $u(x^*, t^*) = \max_{\overline{\mathcal{G}}} u =: M$ for some $(x^*, t^*) \in \overline{\mathcal{G}}$ with $t^* > 0$. Then $u = M$ on $\mathcal{M} \times [0, t^*]$ if $\mathbf{c} = 0$, or if $\mathbf{c} \geq 0$ and $M \geq 0$.*

Proof. Although the result is known, here we sketch a proof that is based on our global perspective of the problem. We use the maximum principle in flat domains by observing that a linear parabolic operator L on $\mathcal{M} \subset \mathbb{R}^{n+1}$ can be extended to a linear parabolic operator \hat{L} on an open strip $\mathcal{N}_\delta \subset \mathbb{R}^{n+1}$ about \mathcal{M} such that

$$\hat{L}u^l(x, t) = Lu(a(x), t) \quad \forall (x, t) \in \mathcal{N}_\delta \times [0, T];$$

see the appendix for more details. Hence, if $Lu \geq 0$, we also have $\hat{L}u^l \geq 0$. Moreover, $u(x^*, t^*) = \max_{\overline{\mathcal{G}}} u =: M$ for some $(x^*, t^*) \in \overline{\mathcal{G}}$ if and only if $u^l(x^*, t^*) = \max_{\overline{\mathcal{G}_\delta}} u^l$. From the strong maximum principle in flat domains (see, for example, [20]), it therefore follows that the set $S := \{x \in \mathcal{M} \mid u(x, t) = M\}$ for any fixed $t \in [0, T]$ must be open, provided that the zero-order term \mathbf{c} satisfies $\mathbf{c} = 0$ or that $\mathbf{c} \geq 0$ and $M \geq 0$. Since S is also closed and \mathcal{M} is connected, we have either $S = \emptyset$ or $S = \mathcal{M}$. See also [4, Chapter 12] and [5] for a detailed discussion of the topic. \square

3. Periodic solutions: Results. The starting point for the study of time-periodic solutions on hypersurfaces is the following result on the existence and uniqueness of solutions to the corresponding initial value problem.

THEOREM 3.1. *Let $\mathcal{M} \subset \mathbb{R}^{n+1}$ be a closed, orientable, n -dimensional hypersurface of class C_1^3 , and let $g(t)$, $t \in [0, T]$, be a family of Riemannian metrics on \mathcal{M} such that the map G defined in (2.13) is of class $\mathcal{H}_{1+\alpha}(\mathcal{G})$ for some $\alpha \in (0, 1)$. Furthermore, let $\mathbf{c} \in \mathcal{H}_\alpha(\mathcal{G})$. Then for any $f \in \mathcal{H}_\alpha(\mathcal{G})$ and $u_0 \in C^0(\mathcal{M})$ there is a unique solution of*

$$(S_I) = \begin{cases} \Delta_{g(t)}u - \mathbf{c}u - u_t = f & \text{in } \mathcal{M} \times (0, T), \\ u(\cdot, 0) = u_0 & \text{on } \mathcal{M}. \end{cases}$$

If $u_0 \in \mathcal{H}_{2+\alpha}(\mathcal{M})$, then $u \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ and there is a constant C such that

$$(3.1) \quad |u|_{2+\alpha, \mathcal{G}} \leq C(|f|_{\alpha, \mathcal{G}} + |u_0|_{2+\alpha, \mathcal{M}}).$$

Using this result and a fixed point argument, it is possible to prove the existence of periodic solutions for advection-diffusion equations on \mathcal{M} with an explicit lower bound on the zero-order term \mathfrak{c} .

PROPOSITION 3.2. *Let $\mathcal{M} \subset \mathbb{R}^{n+1}$ be a closed, orientable, and connected, n -dimensional hypersurface of class C_1^3 , and let $g(t)$, $t \in [0, T]$, be a family of Riemannian metrics on \mathcal{M} such that the map G defined in (2.13) is of class $\mathcal{H}_{1+\alpha}(\mathcal{G})$ for some $\alpha \in (0, 1)$. Furthermore, let $\mathfrak{c}, f \in \mathcal{H}_\alpha(\mathcal{G})$. If $\mathfrak{c} \geq c_0 > \frac{\ln 2}{T}$, then there is a unique solution $u \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ of*

$$(S_P) = \begin{cases} \Delta_{g(t)}u - \mathfrak{c}u - u_t = f & \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} u(\cdot, 0) d\sigma_0 = 0, \\ u(\cdot, 0) = u(\cdot, T) - \int_{\mathcal{M}} u(\cdot, T) d\sigma_0 & \text{on } \mathcal{M} \end{cases}$$

with

$$(3.2) \quad |u|_{2+\alpha, \mathcal{G}} \leq C|f|_{\alpha, \mathcal{G}}$$

for some constant C depending on \mathcal{M} , g , and \mathfrak{c} .

This result is sufficient to establish conditional existence for the periodic problem without a lower bound on \mathfrak{c} .

PROPOSITION 3.3 (Fredholm alternative). *Suppose that the hypotheses of Proposition 3.2 hold. Then either the homogeneous problem*

$$\begin{aligned} \Delta_{g(t)}u - \mathfrak{c}u - u_t &= 0 & \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} u(\cdot, 0) d\sigma_0 &= 0, \\ u(\cdot, 0) &= u(\cdot, T) - \int_{\mathcal{M}} u(\cdot, T) d\sigma_0 & \text{on } \mathcal{M} \end{aligned}$$

has zero as its only solution, in which case the problem

$$\begin{aligned} \Delta_{g(t)}u - \mathfrak{c}u - u_t &= f & \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} u(\cdot, 0) d\sigma_0 &= c, \\ u(\cdot, 0) &= u(\cdot, T) - \left(\int_{\mathcal{M}} u(\cdot, T) d\sigma_0 - c \right) & \text{on } \mathcal{M} \end{aligned}$$

is solvable in the class $\mathcal{H}_{2+\alpha}(\mathcal{G})$ for all $f \in \mathcal{H}_\alpha(\mathcal{G})$ and $c \in \mathbb{R}$, or the homogeneous problem has nonzero solutions, in which case the nonhomogeneous problem cannot be solved for some choices of $f \in \mathcal{H}_\alpha(\mathcal{G})$ and $c \in \mathbb{R}$.

Using this proposition, one can prove existence for the special choice $\mathfrak{c} = 0$.

COROLLARY 3.4. *Suppose that the hypotheses of Proposition 3.2 hold. Then for all $f \in \mathcal{H}_\alpha(\mathcal{G})$ and $c \in \mathbb{R}$ there exists a unique solution $u \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ of*

$$\begin{aligned} \Delta_{g(t)}u - u_t &= f \quad \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} u(\cdot, 0) d\sigma_0 &= c, \\ u(\cdot, 0) &= u(\cdot, T) - \left(\int_{\mathcal{M}} u(\cdot, T) d\sigma_0 - c \right) \quad \text{on } \mathcal{M}. \end{aligned}$$

Existence of solutions for the adjoint operator $L^*u := \Delta_{g(t)}u(x, t) + u_t(x, t)$ of the operator $Lu := \Delta_{g(t)}u - \frac{1}{2}\text{tr}_g(g_t)u - u_t$ can now be established in the following way. First, we define $\underline{g}(t) := g(T-t)$ and $\underline{f}(\cdot, t) := f(\cdot, T-t)$ for all $t \in [0, T]$. Here $f \in \mathcal{H}_\alpha(\mathcal{G})$ and $g(t)$ is assumed to be a given family of Riemannian metrics with $g(0) = g(T)$ and with $G(t)$, defined as in (2.13), of class $\mathcal{H}_{1+\alpha}(\mathcal{G})$. From Corollary 3.4 it follows that there exists a unique solution $\underline{u} \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ of

$$\begin{aligned} \Delta_{\underline{g}(t)}\underline{u} - \underline{u}_t &= \underline{f} \quad \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} \underline{u}(\cdot, 0) d\sigma_0 &= 0, \\ \underline{u}(\cdot, 0) &= \underline{u}(\cdot, T) - \int_{\mathcal{M}} \underline{u}(\cdot, T) d\sigma_0 \quad \text{on } \mathcal{M}. \end{aligned}$$

Next, we define $u(\cdot, t) := \underline{u}(\cdot, T-t)$. Obviously, we have $u_t(\cdot, t) = -\underline{u}_t(\cdot, T-t)$. Furthermore, it follows that

(3.3)

$$\begin{aligned} \Delta_{g(t)}u(x, t) + u_t(x, t) &= \Delta_{\underline{g}(T-t)}\underline{u}(x, T-t) - \underline{u}_t(x, T-t) = \underline{f}(x, T-t) = f(x, t), \\ \int_{\mathcal{M}} u(\cdot, T) d\sigma_0 &= 0, \\ u(\cdot, T) &= u(\cdot, 0) - \int_{\mathcal{M}} u(\cdot, 0) d\sigma_0 \quad \text{on } \mathcal{M}. \end{aligned}$$

We use this result below to prove uniqueness of periodic solutions for the choice $\mathbf{c} = \frac{1}{2}\text{tr}_g(g_t)$ of the zero-order term. That is, for the operator $Lu := \Delta_{g(t)}u - \frac{1}{2}\text{tr}_g(g_t)u - u_t$. The Fredholm alternative in Proposition 3.3 then gives the main theorem of the paper.

THEOREM 3.5. *Suppose that the hypotheses of Proposition 3.2 hold. Then for all $f \in \mathcal{H}_\alpha(\mathcal{G})$ and $c \in \mathbb{R}$ there exists a unique solution $u \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ of*

$$\begin{aligned} \Delta_{g(t)}u - \frac{1}{2}\text{tr}_g(g_t)u - u_t &= f \quad \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} u(\cdot, 0) d\sigma_0 &= c, \\ u(\cdot, 0) &= u(\cdot, T) - \left(\int_{\mathcal{M}} u(\cdot, T) d\sigma_0 - c \right) \quad \text{on } \mathcal{M}. \end{aligned}$$

In particular, for all $c \in \mathbb{R}$ and for all $f \in \mathcal{H}_\alpha(\mathcal{G})$ with $\int_0^T \int_{\mathcal{M}} f(\cdot, t) do(t) dt = 0$ there exists a unique solution $u \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ of

$$\begin{aligned} \Delta_{g(t)} u - \frac{1}{2} \text{tr}_g(g_t) u - u_t &= f \quad \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} u(\cdot, 0) do_0 &= c, \\ u(\cdot, 0) &= u(\cdot, T) \quad \text{on } \mathcal{M}. \end{aligned}$$

Finally, we obtain the following existence and uniqueness result for time-periodic solutions to advection-diffusion equations on moving hypersurfaces.

THEOREM 3.6. *Let $\Gamma(t) \subset \mathbb{R}^{n+1}$, $t \in [0, T]$, be a family of closed, orientable, connected, n -dimensional hypersurfaces of class C_1^3 with $\Gamma(0) = \Gamma(T)$, such that there exist a hypersurface \mathcal{M} satisfying the hypotheses of Proposition 3.2 and a C_1^3 -embedding $X : \overline{\mathcal{G}} \rightarrow \overline{\mathcal{G}_t}$ in the sense of section 2. Then for all $c \in \mathbb{R}$ and for all $f \in \mathcal{H}_\alpha(\mathcal{G}_t)$ with $\int_0^T \int_{\Gamma(t)} f(\cdot, t) d\sigma(t) dt = 0$ there exists a unique solution $u \in \mathcal{H}_{2+\alpha}(\mathcal{G}_t)$ of*

$$\begin{aligned} \Delta_{\Gamma(t)} u - u \nabla_{\Gamma(t)} \cdot v - \partial^\bullet u &= f \quad \text{in } \mathcal{G}_t, \\ \int_{\Gamma(0)} u(\cdot, 0) d\sigma(0) &= c, \\ u(\cdot, 0) &= u(\cdot, T) \quad \text{on } \Gamma(0). \end{aligned}$$

4. Periodic solutions: Proofs.

Proof of Theorem 3.1. Again we use the fact that the advection-diffusion equation on \mathcal{M} can be reformulated as a (nondegenerate) PDE on a neighborhood \mathcal{N}_δ of \mathcal{M} ; see the appendix for more details. The theorem is a consequence of the norm equivalence (2.5) and Theorem 5.18 in [18]. More precisely, the norm equivalence ensures that the lifted data on \mathcal{N}_δ is sufficiently smooth and bounded. Hence, there exists a unique solution \hat{u} of the Neumann boundary problem on \mathcal{N}_δ satisfying a parabolic Schauder estimate. A solution u to the problem (S_I) can be easily constructed from this solution \hat{u} ; see the appendix for details. Moreover, the solution u to the problem (S_I) is unique, since it has to satisfy $u^l = \hat{u}$. Finally, the Schauder estimate for u follows from the norm equivalence (2.5) and the corresponding estimate for \hat{u} . See also section 3.1.3 in [3] for an alternative proof of the global Schauder estimates that is based on local Schauder estimates for an atlas which are patched together. \square

Proof of Proposition 3.2. We divide the proof in to two steps. First, we show that there is a unique solution $u \in C^0(\mathcal{M} \times [0, T]) \cap C^{2,1}(\mathcal{M} \times (0, T))$ of (S_P) by applying a contraction argument. Here, $C^{2,1}$ refers to functions that are continuously differentiable with respect to time and twice continuously differentiable with respect to the space coordinates. Then in a second step, we choose a special series that converges against the periodic solution with respect to the $|\cdot|_{2+\alpha, \mathcal{G}}$ -norm in order to establish the Schauder estimate (3.2).

Step (i) We define $J_0 : C^0(\mathcal{M}) \rightarrow C^0(\mathcal{M} \times [0, T]) \cap C^{2,1}(\mathcal{M} \times (0, T))$ by $J_0(u_0) := u$, where u is the unique solution to the initial value problem (S_I) ; see Theorem 3.1. Then we define $J : C^0(\mathcal{M}) \rightarrow C^0(\mathcal{M})$ by $J(u_0)(x) := J_0(u_0)(x, T) = u(x, T)$ for all $x \in \mathcal{M}$. In the following we show that J is a contraction with contraction constant $\theta_J < \frac{1}{2}$. Let $w_1, w_2 \in C^0(\mathcal{M})$ and $w := w_1 - w_2$, $v := J(w_1) - J(w_2)$. We thus have to show that

$$|v|_{0, \mathcal{M}} \leq \theta_J |w|_{0, \mathcal{M}}$$

for some $0 \leq \theta_J < \frac{1}{2}$. Henceforward, the linear second-order operator in (S_P) is denoted by L , that is,

$$Lu := \Delta_{g(t)}u - \mathbf{c}u - u_t.$$

Let $\epsilon \in (\frac{\ln 2}{T}, c_0)$; then we obtain

$$\begin{aligned} Le^{-\epsilon t} &= \epsilon e^{-\epsilon t} - \mathbf{c}e^{-\epsilon t} = (\epsilon - \mathbf{c})e^{-\epsilon t} \\ &\leq (\epsilon - c_0)e^{-\epsilon t} < 0. \end{aligned}$$

Hence $U^\pm := \pm(J_0(w_1) - J_0(w_2)) - |w|_{0,\mathcal{M}}e^{-\epsilon t}$ satisfies the conditions

$$\begin{aligned} LU^\pm &= \pm(LJ_0(w_1) - LJ_0(w_2)) - |w|_{0,\mathcal{M}}Le^{-\epsilon t} \\ &= \pm(f - f) - |w|_{0,\mathcal{M}}Le^{-\epsilon t} \\ &= -|w|_{0,\mathcal{M}}Le^{-\epsilon t} \geq 0 \quad \text{in } \mathcal{M} \times (0, T). \end{aligned}$$

Furthermore, we have

$$\begin{aligned} U^\pm(\cdot, 0) &= \pm(J_0(w_1) - J_0(w_2))(\cdot, 0) - |w|_{0,\mathcal{M}} \\ &= \pm(w_1 - w_2) - |w|_{0,\mathcal{M}} \\ (4.1) \quad &= \pm w - |w|_{0,\mathcal{M}} \leq 0 \quad \text{in } \mathcal{M}. \end{aligned}$$

Now we suppose that $M^\pm := \max_{(x,t) \in \mathcal{M} \times [0,T]} U^\pm(x,t) > 0$. Then M^\pm must be attained at a point $(x^*, t^*) \in \mathcal{M} \times (0, T]$. It follows from the maximum principle (see Lemma 2.6) that $U^\pm(x^*, 0) = M^\pm > 0$, which contradicts (4.1). Hence $M^\pm \leq 0$, which means that $U^\pm \leq 0$ in $\mathcal{M} \times [0, T]$. It follows that $\pm v \leq |w|_{0,\mathcal{M}}e^{-\epsilon T}$ on \mathcal{M} , and

$$|v|_{0,\mathcal{M}} \leq e^{-\epsilon T}|w|_{0,\mathcal{M}}.$$

This shows that J is a contraction with constant $\theta_J := e^{-\epsilon T} < e^{-\frac{\ln 2}{T}T} = \frac{1}{2}$. Now we define

$$\mathcal{B} := \left\{ u \in C^0(\mathcal{M}) : \int_{\mathcal{M}} u d\mathbf{o}_0 = 0 \right\}$$

and the operator $K : \mathcal{B} \rightarrow C^0(\mathcal{M})$ by

$$K(u_0) = J(u_0) - \int_{\mathcal{M}} J(u_0) d\mathbf{o}_0.$$

In fact we have

$$\int_{\mathcal{M}} K(u_0) d\mathbf{o}_0 = \left(1 - \int_{\mathcal{M}} 1 d\mathbf{o}_0 \right) \int_{\mathcal{M}} J(u_0) d\mathbf{o}_0 = 0.$$

Hence $K : \mathcal{B} \rightarrow \mathcal{B}$. Obviously, $(\mathcal{B}, |\cdot|_{0,\mathcal{M}})$ is a (nonempty) Banach space. In the following we show that K is a contraction. Let $w_1, w_2 \in \mathcal{B}$. We thus have to show that

$$|K(w_1) - K(w_2)|_{0,\mathcal{M}} \leq \theta_K |w_1 - w_2|_{0,\mathcal{M}}$$

for some $0 \leq \theta_K < 1$. Using the fact that J is a contraction we obtain

$$\begin{aligned} |K(w_1) - K(w_2)|_{0,\mathcal{M}} &= \left| J(w_1) - \int_{\mathcal{M}} J(w_1) d\omega_0 - \left(J(w_2) - \int_{\mathcal{M}} J(w_2) d\omega_0 \right) \right|_{0,\mathcal{M}} \\ &\leq |J(w_1) - J(w_2)|_{0,\mathcal{M}} + \left| \int_{\mathcal{M}} J(w_1) - J(w_2) d\omega_0 \right|_{0,\mathcal{M}} \\ &\leq \left(1 + \int_{\mathcal{M}} 1 d\omega_0 \right) |J(w_1) - J(w_2)|_{0,\mathcal{M}} \\ &\leq 2\theta_J |w_1 - w_2|_{0,\mathcal{M}}. \end{aligned}$$

We set $\theta_K := 2\theta_J < 1$. Since K is a contraction with constant θ_K , it follows that there is unique function $u_0^* \in \mathcal{B}$ with $K(u_0^*) = u_0^*$, that is,

$$J(u_0^*) - \int_{\mathcal{M}} J(u_0^*) d\omega_0 = u_0^*.$$

Now let $u^* := J_0(u_0^*)$. We then have

$$\begin{aligned} Lu^* &= f \quad \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} u^*(\cdot, 0) d\omega_0 &= 0, \\ u^*(\cdot, 0) &= u_0^*(\cdot) = J(u_0^*) - \int_{\mathcal{M}} J(u_0^*) d\omega_0 = u^*(\cdot, T) - \int_{\mathcal{M}} u^*(\cdot, T) d\omega_0. \end{aligned}$$

Now suppose $\hat{u} \in C^0(\mathcal{M} \times [0, T]) \cap C^{2,1}(\mathcal{M} \times (0, T))$ is a solution of (S_P) . Then we have $\hat{u}_0 := \hat{u}(\cdot, 0) \in \mathcal{B}$ and $\hat{u} = J_0(\hat{u}_0)$ as well as $\hat{u}(\cdot, T) = J(\hat{u}_0)$. Moreover, it follows that

$$K(\hat{u}_0) = J(\hat{u}_0) - \int_{\mathcal{M}} J(\hat{u}_0) d\omega_0 = \hat{u}(\cdot, T) - \int_{\mathcal{M}} \hat{u}(\cdot, T) d\omega_0 = \hat{u}_0.$$

Therefore, we have $\hat{u} = u^*$, which completes the first step of the proof.

Step (ii) We now define $w_0 := 0$ and $w_{k+1} := K(w_k)$ for $k \in \mathbb{N}_0$. By induction it follows from Theorem 3.1 that $J_0(w_k) \in \mathcal{H}_{2+\alpha}(\mathcal{G})$. Moreover, we have

$$\begin{aligned} |K(w_{k+m}) - K(w_k)|_{2+\alpha,\mathcal{M}} &\leq |J(w_{k+m}) - J(w_k)|_{2+\alpha,\mathcal{M}} + \left| \int_{\mathcal{M}} J(w_{k+m}) - J(w_k) d\omega_0 \right|_{0,\mathcal{M}} \\ &\leq |J(w_{k+m}) - J(w_k)|_{2+\alpha,\mathcal{M}} + |J(w_{k+m}) - J(w_k)|_{0,\mathcal{M}} \\ &\leq 2|J(w_{k+m}) - J(w_k)|_{2+\alpha,\mathcal{M}} \\ &\leq 2|J_0(w_{k+m}) - J_0(w_k)|_{2+\alpha,\mathcal{G}}. \end{aligned}$$

In the following we show that $|J_0(w_{k+m}) - J_0(w_k)|_{2+\alpha,\mathcal{G}} \rightarrow 0$ for $k \rightarrow \infty$ and hence $u_0^* = \lim_{k \rightarrow \infty} w_k \in \mathcal{H}_{2+\alpha}(\mathcal{M})$ as well as $u^* = J_0(u_0^*) \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ according to Theorem 3.1. Furthermore, we then have $u^* = \lim_{k \rightarrow \infty} J_0(w_k)$. We now choose $\zeta \in C^\infty([0, T])$ with $\zeta = 0$ on $[0, \frac{1}{3}T]$, $\zeta = 1$ on $[\frac{2}{3}T, T]$, and $\zeta' \geq 0$. The function $\zeta(J_0(w_{k+1}) - J_0(w_k))$ then satisfies

$$\begin{aligned} L(\zeta(J_0(w_{k+1}) - J_0(w_k))) &= -\zeta'(J_0(w_{k+1}) - J_0(w_k)) \quad \text{in } \mathcal{M} \times (0, T), \\ (\zeta(J_0(w_{k+1}) - J_0(w_k)))(&\cdot, 0) = 0 \quad \text{on } \mathcal{M}. \end{aligned}$$

From the Schauder estimate (3.1) it follows that

$$\begin{aligned} |\zeta(J_0(w_{k+1}) - J_0(w_k))|_{2+\alpha, \mathcal{G}} &\leq C|\zeta'(J_0(w_{k+1}) - J_0(w_k))|_{\alpha, \mathcal{G}} \\ &\leq C(\zeta)|J_0(w_{k+1}) - J_0(w_k)|_{\alpha, \mathcal{G}}, \end{aligned}$$

and hence

$$(4.2) \quad \begin{aligned} |J(w_{k+1}) - J(w_k)|_{2+\alpha, \mathcal{M}} &\leq |\zeta(J_0(w_{k+1}) - J_0(w_k))|_{2+\alpha, \mathcal{G}} \\ &\leq C(\zeta)|J_0(w_{k+1}) - J_0(w_k)|_{\alpha, \mathcal{G}}. \end{aligned}$$

Since $L(J_0(w_{k+2}) - J_0(w_{k+1})) = 0$, the estimate (3.1) also gives

$$\begin{aligned} |J_0(w_{k+2}) - J_0(w_{k+1})|_{2+\alpha, \mathcal{G}} &\leq C|w_{k+2} - w_{k+1}|_{2+\alpha, \mathcal{M}} \leq C|K(w_{k+1}) - K(w_k)|_{2+\alpha, \mathcal{M}} \\ &\leq C|J(w_{k+1}) - J(w_k)|_{2+\alpha, \mathcal{M}} \\ &\leq C(\zeta)|J_0(w_{k+1}) - J_0(w_k)|_{\alpha, \mathcal{G}}, \end{aligned}$$

where we have used (4.2) in the last step. The interpolation estimate (2.6) then yields

$$\begin{aligned} |J_0(w_{k+2}) - J_0(w_{k+1})|_{2+\alpha, \mathcal{G}} \\ \leq C(\zeta)\epsilon^{1-\alpha}|J_0(w_{k+1}) - J_0(w_k)|_{2+\alpha, \mathcal{G}} + C(\zeta, \epsilon)|J_0(w_{k+1}) - J_0(w_k)|_{0, \mathcal{G}}. \end{aligned}$$

Moreover, the maximum principle gives

$$|J_0(w_{k+1}) - J_0(w_k)|_{0, \mathcal{G}} \leq |w_{k+1} - w_k|_{0, \mathcal{M}},$$

and hence

$$\begin{aligned} |J_0(w_{k+2}) - J_0(w_{k+1})|_{2+\alpha, \mathcal{G}} \\ \leq C(\zeta)\epsilon^{1-\alpha}|J_0(w_{k+1}) - J_0(w_k)|_{2+\alpha, \mathcal{G}} + C(\zeta, \epsilon)|w_{k+1} - w_k|_{0, \mathcal{M}}. \end{aligned}$$

Choosing $\epsilon > 0$ such that $C(\zeta)\epsilon^{1-\alpha} = \theta_K$ and setting $C^* := C(\zeta, \epsilon)$ for this choice of ϵ leads to

$$|J_0(w_{k+2}) - J_0(w_{k+1})|_{2+\alpha, \mathcal{G}} \leq \theta_K|J_0(w_{k+1}) - J_0(w_k)|_{2+\alpha, \mathcal{G}} + C^*|w_{k+1} - w_k|_{0, \mathcal{M}}.$$

Since K is a contraction, we obtain $|w_{k+j+1} - w_{k+j}|_{0, \mathcal{M}} \leq \theta_K^{k+j}|w_1 - w_0|_{0, \mathcal{M}}$ and

$$\begin{aligned} |J_0(w_{k+m+1}) - J_0(w_{k+1})|_{2+\alpha, \mathcal{G}} &\leq \sum_{j=0}^{m-1} |J_0(w_{k+j+2}) - J_0(w_{k+j+1})|_{2+\alpha, \mathcal{G}} \\ &\leq \sum_{j=0}^{m-1} \left\{ \theta_K |J_0(w_{k+j+1}) - J_0(w_{k+j})|_{2+\alpha, \mathcal{G}} + C^* \theta_K^{k+j} |w_1 - w_0|_{0, \mathcal{M}} \right\} \\ &\leq \sum_{j=0}^{m-1} \left\{ \theta_K^{k+j+1} |J_0(w_1) - J_0(w_0)|_{2+\alpha, \mathcal{G}} + C^* (k+j+1) \theta_K^{k+j} |w_1 - w_0|_{0, \mathcal{M}} \right\} \\ &\leq |J_0(w_1) - J_0(w_0)|_{2+\alpha, \mathcal{G}} \frac{\theta_K^{k+1} - \theta_K^{k+m+1}}{1 - \theta_K} + C^* |w_1 - w_0|_{0, \mathcal{M}} \frac{\partial}{\partial \theta_K} \frac{\theta_K^{k+1} - \theta_K^{k+m+1}}{1 - \theta_K} \\ &\leq |J_0(w_1) - J_0(w_0)|_{2+\alpha, \mathcal{G}} \frac{\theta_K^{k+1}}{1 - \theta_K} + C^* |w_1 - w_0|_{0, \mathcal{M}} \frac{(k+1)\theta_K^k}{(1 - \theta_K)^2}, \end{aligned}$$

which converges to 0 for $k \rightarrow \infty$. Therefore, the periodic solution $u^* = J_0(u_0^*)$ is in $\mathcal{H}_{2+\alpha}(\mathcal{G})$ and

$$|J_0(u_0^*) - J_0(w_1)|_{2+\alpha, \mathcal{G}} \leq |J_0(w_1) - J_0(w_0)|_{2+\alpha, \mathcal{G}} \frac{\theta_K}{1 - \theta_K} + |w_1 - w_0|_{0, \mathcal{M}} \frac{C^*}{(1 - \theta_K)^2}.$$

Since $w_0 = 0$ and $|w_1|_{2+\alpha, \mathcal{M}} \leq C|J_0(w_0)|_{2+\alpha, \mathcal{G}} \leq C|f|_{\alpha, \mathcal{G}}$, as well as

$$\begin{aligned} |J_0(w_1)|_{2+\alpha, \mathcal{G}} &\leq C(|f|_{\alpha, \mathcal{G}} + |w_1|_{2+\alpha, \mathcal{M}}) \leq C|f|_{\alpha, \mathcal{G}}, \\ |J_0(w_1) - J_0(w_0)|_{2+\alpha, \mathcal{G}} &\leq C|w_1|_{2+\alpha, \mathcal{M}} \leq C|f|_{\alpha, \mathcal{G}}, \end{aligned}$$

which hold because of (3.1), we finally obtain the estimate

$$|u^*|_{2+\alpha, \mathcal{G}} \leq C|f|_{\alpha, \mathcal{G}}. \quad \square$$

Proof of Proposition 3.3. Since u solves the nonhomogeneous problem if and only if $\tilde{u} := u - c$ solves the problem

$$\begin{aligned} \Delta_{g(t)} \tilde{u} - \mathbf{c} \tilde{u} - \tilde{u}_t &= f + \mathbf{c} \mathbf{c}, \\ \int_{\mathcal{M}} \tilde{u}(\cdot, 0) do_0 &= 0, \\ \tilde{u}(\cdot, 0) &= \tilde{u}(\cdot, T) - \int_{\mathcal{M}} \tilde{u}(\cdot, T) do_0, \end{aligned}$$

we can assume without loss of generality that $c = 0$. Now, let $\hat{\mathcal{B}} := \{u \in \mathcal{H}_{2+\alpha}(\mathcal{G}) : \int_{\mathcal{M}} u(\cdot, 0) do_0 = 0 \text{ and } u(\cdot, 0) = u(\cdot, T) - \int_{\mathcal{M}} u(\cdot, T) do_0 \text{ on } \mathcal{M}\}$ and let $L_T : \hat{\mathcal{B}} \rightarrow \mathcal{H}_\alpha(\mathcal{G})$ be the linear second-order operator defined by

$$L_T u = \Delta_{g(t)} u - \frac{1}{T} u - u_t.$$

According to Proposition 3.2 the operator L_T is invertible and the inverse operator L_T^{-1} is continuous. Hence, we can define the operator $\mathcal{K} : \mathcal{H}_\alpha(\mathcal{G}) \rightarrow \hat{\mathcal{B}} \subset \mathcal{H}_\alpha(\mathcal{G})$ by $\mathcal{K}u := L_T^{-1}(\frac{1}{T}u - \mathbf{c}u)$. Because of the Schauder estimate (3.2) and the fact that $\hat{\mathcal{B}}$ is compactly embedded in $\mathcal{H}_\alpha(\mathcal{G})$, this is a compact operator. The equation $\Delta_{g(t)} u - \mathbf{c}u - u_t = f$ is equivalent to

$$\begin{aligned} L_T u + \left(\frac{1}{T} - \mathbf{c}\right) u &= f \\ \Leftrightarrow u + \mathcal{K}u &= L_T^{-1} f. \end{aligned}$$

Since $\mathbf{1} + \mathcal{K}$ is a Fredholm operator, the second equation has a solution if and only if $(\mathbf{1} + \mathcal{K})u = 0$ implies $u = 0$. The standard Fredholm theory therefore gives the result. \square

Proof of Corollary 3.4. Because of the Fredholm alternative in Proposition 3.3 we only have to establish uniqueness of the homogeneous problem. Suppose we have a solution $\varphi \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ of the homogeneous problem. Because of the maximum principle (see Lemma 2.6), we then have

$$\begin{aligned} \max_{\mathcal{M}} \varphi(\cdot, 0) &\geq \max_{\mathcal{M}} \varphi(\cdot, T) = \max_{\mathcal{M}} \left(\varphi(\cdot, 0) + \int_{\mathcal{M}} \varphi(\cdot, T) do_0 \right) \\ &= \max_{\mathcal{M}} \varphi(\cdot, 0) + \int_{\mathcal{M}} \varphi(\cdot, T) do_0, \end{aligned}$$

and hence $\int_{\mathcal{M}} \varphi(\cdot, T) do_0 \leq 0$. In the same way we obtain

$$\max_{\mathcal{M}} (-\varphi(\cdot, 0)) \geq \max_{\mathcal{M}} (-\varphi(\cdot, 0)) - \int_{\mathcal{M}} \varphi(\cdot, T) do_0$$

and $\int_{\mathcal{M}} \varphi(\cdot, T) do_0 \geq 0$. It follows that $\int_{\mathcal{M}} \varphi(\cdot, T) do_0 = 0$ and hence $\varphi(\cdot, 0) = \varphi(\cdot, T)$ on \mathcal{M} . Moreover, we know from the strong maximum principle that either φ is constant or $\max_{\mathcal{M}} \varphi(\cdot, T) < \max_{\mathcal{M}} \varphi(\cdot, 0)$. Therefore, φ has to be constant. From $\int_{\mathcal{M}} \varphi(\cdot, 0) do_0 = 0$ it then follows that $\varphi = 0$. \square

Proof of Theorem 3.5. The second statement easily follows from the first statement and the fact that

$$\begin{aligned} \int_{\mathcal{M}} u(\cdot, T) do_0 &= \int_{\mathcal{M}} u(\cdot, T) do(T) = \int_{\mathcal{M}} u(\cdot, 0) do(0) + \int_0^T \frac{d}{dt} \int_{\mathcal{M}} u(\cdot, t) do(t) dt \\ &= c|\mathcal{M}| + \int_0^T \frac{d}{dt} \int_{\mathcal{M}} u(\cdot, t) do(t) dt \\ &= c|\mathcal{M}| + \int_0^T \int_{\mathcal{M}} u_t + \frac{1}{2} \text{tr}_g(g_t) u do(t) dt \\ &= c|\mathcal{M}| + \int_0^T \int_{\mathcal{M}} \Delta_{g(t)} u do(t) - \int_0^T \int_{\mathcal{M}} f do(t) = c|\mathcal{M}|, \end{aligned}$$

that is, $\int_{\mathcal{M}} u(\cdot, T) do_0 = c$ and hence $u(\cdot, 0) = u(\cdot, T)$. In order to prove the first statement, we mention that according to Proposition 3.3, it suffices to prove the uniqueness result for the homogeneous equation, that is, $f = 0$ and $c = 0$. Let $u \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ be a solution of the homogeneous problem. As above, we obtain $u(\cdot, 0) = u(\cdot, T)$ on \mathcal{M} . Next, we choose $\varphi \in \mathcal{H}_{2+\alpha}(\mathcal{G})$ such that

$$\begin{aligned} \Delta_{g(t)} \varphi + \varphi_t &= u \quad \text{in } \mathcal{M} \times (0, T), \\ \int_{\mathcal{M}} \varphi(\cdot, T) do_0 &= 0, \\ \varphi(\cdot, T) &= \varphi(\cdot, 0) - \int_{\mathcal{M}} \varphi(\cdot, 0) do_0 \quad \text{on } \mathcal{M}. \end{aligned}$$

According to Corollary 3.4 such a solution exists if we choose $f = u$ in (3.3). We then obtain

$$\begin{aligned} 0 &= \int_0^T \int_{\mathcal{M}} (\Delta_{g(t)} u - \frac{1}{2} \text{tr}_g(g_t) u - u_t) \varphi do(t) dt \\ &= \int_0^T \int_{\mathcal{M}} u (\Delta_{g(t)} \varphi + \varphi_t) do(t) dt - \int_0^T \int_{\mathcal{M}} \frac{1}{2} \text{tr}_g(g_t) u \varphi + (u \varphi)_t do(t) dt \\ &= \int_0^T \int_{\mathcal{M}} u (\Delta_{g(t)} \varphi + \varphi_t) do(t) dt - \int_0^T \frac{d}{dt} \int_{\mathcal{M}} u \varphi do(t) dt \\ &= \int_0^T \int_{\mathcal{M}} |u|^2 do(t) dt - \int_{\mathcal{M}} u(\cdot, T) \varphi(\cdot, T) do(T) + \int_{\mathcal{M}} u(\cdot, 0) \varphi(\cdot, 0) do(0) \\ &= \int_0^T \int_{\mathcal{M}} |u|^2 do(t) dt - \int_{\mathcal{M}} u(\cdot, 0) \varphi(\cdot, T) do(0) + \int_{\mathcal{M}} u(\cdot, 0) \varphi(\cdot, 0) do(0) \end{aligned}$$

$$\begin{aligned} &= \int_0^T \int_{\mathcal{M}} |u|^2 do(t) dt - \int_{\mathcal{M}} u(\cdot, 0)(\varphi(\cdot, T) - \varphi(\cdot, 0)) do(0) \\ &= \int_0^T \int_{\mathcal{M}} |u|^2 do(t) dt + \int_{\mathcal{M}} u(\cdot, 0) do_0 \int_{\mathcal{M}} \varphi(\cdot, 0) do_0 = \int_0^T \int_{\mathcal{M}} |u|^2 do(t) dt. \end{aligned}$$

Hence, we have $u = 0$. This completes the proof of the claim. \square

Proof of Theorem 3.6. The theorem directly follows from Lemma 2.5 and Theorem 3.5. \square

5. Appendix. In this section, we show that solving a parabolic PDE on the hypersurface $\mathcal{M} \subset \mathbb{R}^{n+1}$ is, in a certain sense, equivalent to solving a related parabolic PDE on an open neighborhood of $\mathcal{M} \subset \mathbb{R}^{n+1}$. The main advantage of this approach is that the well-established machinery of parabolic PDEs on $(n+1)$ -dimensional domains of \mathbb{R}^{n+1} can be immediately applied after a suitable PDE on the open neighborhood has been found. The main task therefore remains to formulate a PDE on an open neighborhood of \mathcal{M} from which the solution of the PDE on \mathcal{M} can be extracted.

To motivate the idea, we first have a look at the following second-order parabolic PDE on the n -dimensional hyperplane $\mathcal{M}_0 := \{x \in \mathbb{R}^{n+1} : x_{n+1} = 0\}$:

$$(P) = \begin{cases} \Delta_{\mathcal{M}_0} u - u_t = f & \text{in } \mathcal{M}_0 \times (0, T), \\ u(\cdot, 0) = u_0(\cdot) & \text{on } \mathcal{M}_0, \end{cases}$$

where $\Delta_{\mathcal{M}_0} u = \sum_{i=1}^n u_{x_i x_i}$ denotes the standard Laplacian on \mathcal{M}_0 . Now, let $u \in C^0(\mathcal{M}_0 \times [0, T]) \cap C^{2,1}(\mathcal{M}_0 \times (0, T))$ be a solution to this problem. Furthermore, let $\mathcal{N}_\delta := \{x \in \mathbb{R}^{n+1} : |x_{n+1}| < \delta\}$ be an open neighborhood of width δ around \mathcal{M}_0 . The function $u^l : \overline{\mathcal{N}_\delta} \times [0, T] \rightarrow \mathbb{R}$ defined by $u^l(x, t) := u(\underline{x}, t)$ with $x = (\underline{x}, x_{n+1})$ is then a solution to the following (strongly) parabolic initial value boundary problem:

$$(P^l) = \begin{cases} \Delta \hat{u} - \hat{u}_t = f^l & \text{in } \mathcal{N}_\delta \times (0, T), \\ \hat{u}_{x_{n+1}} = 0 & \text{on } \partial \mathcal{N}_\delta \times (0, T), \\ \hat{u}(\cdot, 0) = u_0^l(\cdot) & \text{on } \mathcal{N}_\delta. \end{cases}$$

Here, $f^l : \mathcal{N}_\delta \times (0, T) \rightarrow \mathbb{R}$ and $u_0^l : \mathcal{N}_\delta \rightarrow \mathbb{R}$ are defined by $f^l(x, t) := f(\underline{x}, t)$ and $u_0^l(x) := u_0(\underline{x})$, respectively. $\Delta \hat{u} = \sum_{i=0}^{n+1} \hat{u}_{x_i x_i}$ denotes the standard Laplacian on \mathbb{R}^{n+1} . Obviously, we have $\Delta u^l(x, t) = \Delta_{\mathcal{M}_0} u(\underline{x}, t) + u_{x_{n+1} x_{n+1}}^l(x, t) = \Delta_{\mathcal{M}_0} u(\underline{x}, t)$ since $u_{x_{n+1}}^l = 0$ on $\mathcal{N}_\delta \times (0, T)$. Conversely, let $\hat{u} \in C^0(\overline{\mathcal{N}_\delta} \times [0, T]) \cap C^{2,1}(\overline{\mathcal{N}_\delta} \times (0, T))$ be a solution of the lifted PDE (P^l) . Then the function $u : \mathcal{M}_0 \times [0, T] \rightarrow \mathbb{R}$ defined by $u(\underline{x}, t) := \frac{1}{2\delta} \int_{-\delta}^\delta \hat{u}(\underline{x}, s, t) ds$ for all $(\underline{x}, t) \in \mathcal{M}_0 \times [0, T]$ is a solution to (P) . This easily follows from $f(\underline{x}, t) = \frac{1}{2\delta} \int_{-\delta}^\delta f^l(\underline{x}, s, t) ds$, $u_0(\underline{x}) = \frac{1}{2\delta} \int_{-\delta}^\delta u_0^l(\underline{x}, s) ds$ and the fact that

$$\begin{aligned} &\frac{1}{2\delta} \int_{-\delta}^\delta \Delta \hat{u}(\underline{x}, s, t) - \hat{u}_t(\underline{x}, s, t) ds \\ &= \frac{1}{2\delta} \sum_{i=1}^{n+1} \int_{-\delta}^\delta \hat{u}_{x_i x_i}(\underline{x}, s, t) ds - \frac{1}{2\delta} \int_{-\delta}^\delta \hat{u}_t(\underline{x}, s, t) ds \\ &= \frac{1}{2\delta} \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} \int_{-\delta}^\delta \hat{u}(\underline{x}, s, t) ds + \frac{1}{2\delta} (\hat{u}_{x_{n+1}}(\underline{x}, \delta, t) - \hat{u}_{x_{n+1}}(\underline{x}, -\delta, t)) \end{aligned}$$

$$\begin{aligned}
 & -\frac{1}{2\delta} \frac{\partial}{\partial t} \int_{-\delta}^{\delta} \hat{u}(\underline{x}, s), t) ds \\
 & = \sum_{i=1}^n u_{x_i x_i}(\underline{x}, t) - u_t(\underline{x}, t)
 \end{aligned}$$

for all $(\underline{x}, t) \in \mathcal{M}_0 \times (0, T)$.

Now we want to use this idea for the following second-order parabolic PDE on the closed hypersurface $\mathcal{M} \subset \mathbb{R}^{n+1}$:

$$(S) = \begin{cases} Lu = f & \text{in } \mathcal{M} \times (0, T), \\ u(\cdot, 0) = u_0(\cdot) & \text{on } \mathcal{M}, \end{cases}$$

where

$$Lu := \Delta_{g(t)}u + w \cdot \nabla_{\mathcal{M}}u - cu - u_t.$$

Here $\Delta_{g(t)}u = \underline{D}_\alpha (G^{\alpha\beta} \underline{D}_\beta u) + \frac{1}{2} P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta G_{\alpha\eta} \underline{D}_\rho u$ denotes the Laplacian with respect to a (sufficiently) smooth time-dependent Riemannian metric $g(t)$ on \mathcal{M} , where the map $G(t)$ is the Cartesian representation of $g(t)$ as defined in (2.13). w is a given vector field on \mathcal{M} . The corresponding lifted second-order parabolic PDE is then given by

$$(S^l) = \begin{cases} \hat{L}\hat{u} = f^l & \text{in } \mathcal{N}_\delta \times (0, T), \\ \frac{\partial \hat{u}}{\partial \nu} = 0 & \text{on } \partial\mathcal{N}_\delta \times (0, T), \\ \hat{u}(\cdot, 0) = u_0^l(\cdot) & \text{on } \mathcal{N}_\delta, \end{cases}$$

where \mathcal{N}_δ is the open strip of width δ around \mathcal{M} defined in (2.2), and $f^l(x, t) := f(a(x), t)$ as well as $u_0^l(x, t) := u(a(x), t)$ are the lifted data. In order to define an appropriate parabolic operator \hat{L} on \mathcal{N}_δ , we first introduce the parallel hypersurfaces $\mathcal{M}_s := \{a + s\nu(a) : a \in \mathcal{M}\}$ for $|s| < \delta$ and the bijective projections $a_s : \mathcal{M}_s \rightarrow \mathcal{M}$ defined by $a_s := a|_{\mathcal{M}_s}$. Obviously, we have $\mathcal{N}_\delta = \bigcup_{|s| < \delta} \mathcal{M}_s$. On \mathcal{M}_s we introduce the rescaled tangential gradient for differentiable functions \hat{u} by

$$\tilde{\nabla}_{\mathcal{M}_s} \hat{u}(x_s) := (\mathbf{1} - d(x_s)\mathcal{H}(x_s))^{-1} \nabla_{\mathcal{M}_s} \hat{u}(x_s) \quad \forall x_s \in \mathcal{M}_s, \forall s \in (-\delta, \delta).$$

For $\delta > 0$ sufficiently small the map $(\mathbf{1} - d\mathcal{H})(x_s)$ is indeed invertible; see [8] for further details. The Cartesian components of the rescaled tangential gradient $\tilde{\nabla}_{\mathcal{M}_s} \hat{u}$ are denoted by

$$\begin{pmatrix} \tilde{D}_1 \hat{u} \\ \vdots \\ \tilde{D}_{n+1} \hat{u} \end{pmatrix} := \tilde{\nabla}_{\mathcal{M}_s} \hat{u}.$$

We then define the parabolic operator \hat{L} by

$$(5.1) \quad \hat{L}\hat{u} = \tilde{D}_\alpha (G^{l\alpha\beta} \tilde{D}_\beta \hat{u}) + \frac{1}{2} P_{\alpha\gamma} G^{l\gamma\eta} G^{l\beta\rho} \tilde{D}_\beta G_{\alpha\eta}^l \tilde{D}_\rho \hat{u} + \frac{\partial^2 \hat{u}}{\partial \nu^2} + w^l \cdot \tilde{\nabla}_{\mathcal{M}_s} \hat{u} - c^l \hat{u} - \hat{u}_t,$$

where $G^{l\alpha\beta}$ denotes the components of $(G^l)^{-1}$ and $G_{\alpha\beta}^l$ denotes the components of $G^l(x, t) := G(a(x), t)$. Furthermore, $w^l(x, t) := w(a(x), t)$ and $c^l(x, t) := c(a(x), t)$. Below we show that for $\hat{u} = u^l(x, s) := u(a(x), t)$ we have

$$(5.2) \quad \hat{L}u^l(x, t) = Lu(a(x), t) \quad \forall x \in \mathcal{N}_\delta.$$

Moreover, we have $\frac{\partial u^l}{\partial \nu} = 0$, since $u^l(a + s\nu(a)) = u(a)$ for all $a \in \mathcal{M}$, $|s| < \delta$. Using these facts, it is easy to show that u^l is a solution to (S^l) if u is a solution to (S) .

Now suppose that $\hat{u} : \overline{\mathcal{N}_\delta} \times [0, T] \rightarrow \mathbb{R}^{n+1}$ is a solution to (S^l) . In this case we define $u : \mathcal{M} \times [0, T] \rightarrow \mathbb{R}$ by

$$u(a, t) := \frac{1}{2\delta} \int_{-\delta}^{\delta} \hat{u}(a + s\nu(a), t) ds \quad \forall (a, t) \in \mathcal{M} \times [0, T].$$

In order to see that u satisfies (S) , we introduce the family of functions $\tilde{u}_s : \mathcal{M} \times [0, T] \rightarrow \mathbb{R}$ for $s \in [-\delta, \delta]$ defined by

$$(5.3) \quad \tilde{u}_s(a, t) := \hat{u}(a + s\nu(a), t) \quad \forall (a, t) \in \mathcal{M} \times [0, T], \forall s \in [-\delta, \delta].$$

We need this definition, because it is a priori not clear whether \hat{u} is constant in the normal direction. Obviously, we have

$$(5.4) \quad \hat{u}|_{\mathcal{M}_s} = \tilde{u}_s|_{\mathcal{M}_s}.$$

Since the tangential gradient $\tilde{\nabla}_{\mathcal{M}_s}$ only depends on the values of \hat{u} on \mathcal{M}_s , we obtain

$$\begin{aligned} \hat{L}\hat{u}(x_s, t) &= \hat{L}\tilde{u}_s^l(x_s, t) - \frac{\partial^2 \tilde{u}_s^l}{\partial \nu^2}(x_s, t) + \frac{\partial^2 \hat{u}}{\partial \nu^2}(x_s, t) \\ &= L\tilde{u}_s(a(x_s), t) + \frac{\partial^2 \tilde{u}_s}{\partial s^2}(a(x_s), t) \quad \forall x_s \in \mathcal{M}_s, \forall s \in (-\delta, \delta), \end{aligned}$$

where we have used formula (5.2), $\frac{\partial \tilde{u}_s^l}{\partial \nu} = 0$ and $\frac{\partial^2 \hat{u}}{\partial \nu^2}(x_s, t) = \frac{\partial^2 \tilde{u}_s}{\partial s^2}(a(x_s), t)$ in the last step. It follows that for all $a \in \mathcal{M}$ and $t \in (0, T)$ we have

$$\begin{aligned} f(a, t) &= \frac{1}{2\delta} \int_{-\delta}^{\delta} f^l(a + s\nu(a), t) ds = \frac{1}{2\delta} \int_{-\delta}^{\delta} \hat{L}\hat{u}(a + s\nu(a), t) ds \\ &= \frac{1}{2\delta} \int_{-\delta}^{\delta} L\tilde{u}_s(a, t) ds + \frac{1}{2\delta} \int_{-\delta}^{\delta} \frac{\partial^2 \tilde{u}_s}{\partial s^2}(a, t) ds \\ &= \frac{1}{2\delta} L \int_{-\delta}^{\delta} \tilde{u}_s(a, t) ds \\ &= (Lu)(a, t), \end{aligned}$$

where the term $\int_{-\delta}^{\delta} \frac{\partial^2 \tilde{u}_s}{\partial s^2} ds$ vanishes, because of the Neumann boundary condition in (S^l) . Hence, $u = \frac{1}{2\delta} \int_{-\delta}^{\delta} \tilde{u}_s ds$ is a solution to (S) if \hat{u} is a solution to the lifted problem (S^l) . From the uniqueness of solutions to (S^l) it follows that $\hat{u} = u^l$, which shows that \hat{u} has to be constant in the normal direction.

In the following, we prove formula (5.2) and show that \hat{L} is in fact a strongly parabolic second-order operator on \mathcal{N}_δ . First, it is easy to show that the tangential gradient of the projection a is given by

$$\nabla_{\mathcal{M}_s} a(x_s) = P(x_s) - d(x_s)\mathcal{H}(x_s) \quad \forall x_s \in \mathcal{M}_s.$$

Using the fact that the $\nu(x) = \nu(a(x))$ and $P(x) = P(a(x))$, it follows that for $u^l(\cdot, t) := u(a(\cdot), t)$ the following identity holds

$$\nabla_{\mathcal{M}_s} u^l(x_s, t) = (\mathbb{1} - d(x_s)\mathcal{H}(x_s))\nabla_{\mathcal{M}} u(a(x_s), t).$$

Hence, for all $(x_s, t) \in \mathcal{M}_s \times [0, T]$ we have

$$(5.5) \quad \nabla_{\mathcal{M}} u(a(x_s), t) = (\mathbb{1} - d(x_s)\mathcal{H}(x_s))^{-1}\nabla_{\mathcal{M}_s} u^l(x_s, t) = \tilde{\nabla}_{\mathcal{M}_s} u^l(x_s, t),$$

or $(\nabla_{\mathcal{M}} u)^l = \tilde{\nabla}_{\mathcal{M}_s} u^l$, respectively. From this result and the fact that $\frac{\partial u^l}{\partial \nu} = 0$ we directly obtain that

$$\begin{aligned} \hat{L}u^l(x, t) &= \tilde{D}_\alpha \left(G^{l\alpha\beta} \tilde{D}_\beta u^l \right) (x, t) + \frac{1}{2} (P_{\alpha\gamma} G^{l\gamma\eta} G^{l\beta\rho} \tilde{D}_\beta G_{\alpha\eta}^l \tilde{D}_\rho u^l) (x, t) \\ &\quad + w^l(x, t) \cdot \tilde{\nabla}_{\mathcal{M}_s} u^l(x, t) - \mathbf{c}^l(x, t) u^l(x, t) - u_t^l(x, t) \\ &= \underline{D}_\alpha (G^{\alpha\beta} \underline{D}_\beta u) (a(x), t) + \frac{1}{2} (P_{\alpha\gamma} G^{\gamma\eta} G^{\beta\rho} \underline{D}_\beta G_{\alpha\eta} \underline{D}_\rho u) (a(x), t) \\ &\quad + w(a(x), t) \cdot \nabla_{\mathcal{M}} u(a(x), t) - \mathbf{c}(a(x), t) u(a(x), t) - u_t(a(x), t) \\ &= Lu(a(x), t). \end{aligned}$$

In order to see that the operator \hat{L} is strongly parabolic, we consider the second-order terms in (5.1). Using the notation $A(x) := (\mathbb{1} - d\mathcal{H})(x)$ for all $x \in \mathcal{N}_\delta$ as well as

$$(A_{\alpha\beta})_{\alpha, \beta=1, \dots, n+1} := A, \quad (A^{\alpha\beta})_{\alpha, \beta=1, \dots, n+1} := A^{-1},$$

for the components of A and A^{-1} , we obtain

$$\begin{aligned} \tilde{D}_\alpha (G^{l\alpha\beta} \tilde{D}_\beta \hat{u}) + \frac{\partial^2 \hat{u}}{\partial \nu^2} &= A^{\alpha\rho} P_{\rho\eta} D_\eta (G^{l\alpha\beta} A^{\beta\kappa} P_{\kappa i} D_i \hat{u}) + \frac{\partial^2 \hat{u}}{\partial \nu^2} \\ &= A^{\alpha\rho} D_\rho (G^{l\alpha\beta} A^{\beta i} D_i \hat{u}) - A^{\alpha\rho} \nu_\rho \nu_\eta D_\eta (G^{l\alpha\beta} A^{\beta i} D_i \hat{u}) - A^{\alpha\rho} D_\rho (G^{l\alpha\beta} A^{\beta\kappa} \nu_\kappa \nu_i D_i \hat{u}) \\ &\quad + A^{\alpha\rho} \nu_\rho \nu_\eta D_\eta (G^{l\alpha\beta} A^{\beta\kappa} \nu_\kappa \nu_i D_i \hat{u}) + \frac{\partial^2 \hat{u}}{\partial \nu^2} \\ &= A^{\alpha\rho} D_\rho (G^{l\alpha\beta} A^{\beta i} D_i \hat{u}) - \nu_\alpha \nu_\eta D_\eta (G^{l\alpha\beta} A^{\beta i} D_i \hat{u}) - A^{\alpha\rho} D_\rho (\nu_\alpha \nu_i D_i \hat{u}) \\ &\quad + \nu_\alpha \nu_\eta D_\eta (\nu_\alpha \nu_i D_i \hat{u}) + \frac{\partial^2 \hat{u}}{\partial \nu^2} \\ &= A^{\alpha\rho} D_\rho (G^{l\alpha\beta} A^{\beta i} D_i \hat{u}) - \nu_\eta D_\eta (\nu_i D_i \hat{u}) + G^{l\alpha\beta} A^{\beta i} \nu_\eta D_\eta \nu_\alpha D_i \hat{u} - \nu_\rho D_\rho (\nu_i D_i \hat{u}) \\ &\quad - A^{\alpha\rho} D_\rho \nu_\alpha \nu_i D_i \hat{u} + \nu_\eta D_\eta (\nu_i D_i \hat{u}) + \nu_\alpha \nu_\eta D_\eta \nu_\alpha \nu_i D_i \hat{u} + \frac{\partial^2 \hat{u}}{\partial \nu^2} \\ &= A^{\alpha\rho} D_\rho (G^{l\alpha\beta} A^{\beta i} D_i \hat{u}) - A^{\alpha\rho} D_\rho \nu_\alpha \nu_i D_i \hat{u} \\ &= A^{\alpha\rho} D_\rho (G^{l\alpha\beta} A^{\beta i} D_i \hat{u}) - (\tilde{\nabla}_{\mathcal{M}_s} \cdot \nu) \frac{\partial}{\partial \nu} \hat{u} \\ &= A^{\alpha\rho} G^{l\alpha\beta} A^{\beta i} D_\rho D_i \hat{u} + A^{\alpha\rho} D_\rho (G^{l\alpha\beta} A^{\beta i}) D_i \hat{u} - (\tilde{\nabla}_{\mathcal{M}_s} \cdot \nu) \frac{\partial}{\partial \nu} \hat{u}. \end{aligned}$$

Here we have used the fact that $(G^l)^{-1}\nu = \nu$ and $A^{-1}\nu = \nu$ as well as $\frac{\partial \nu}{\partial \nu} = 0$. For $G = \mathbb{1}$, this identity simplifies to

$$(5.6) \quad \tilde{D}_\alpha \tilde{D}_\alpha \hat{u} + \frac{\partial^2 \hat{u}}{\partial \nu^2} = A^{\alpha\rho} A^{\alpha i} D_\rho D_i \hat{u} + A^{\alpha\rho} D_\rho A^{\alpha i} D_i \hat{u} - (\tilde{\nabla}_{\mathcal{M}_s} \cdot \nu) \frac{\partial}{\partial \nu} \hat{u}.$$

Remark 5.1. A similar extension idea has been used only recently to develop numerical schemes for the simulation of geometric PDEs on surfaces; see [19]. The model problem considered in [19] is the following elliptic problem:

$$\Delta_\Gamma u - cu = f \quad \text{on } \Gamma.$$

Using our notation, the authors propose the following nondegenerate extended equations:

$$\begin{aligned} \nabla \cdot (\mu A^{-2} \nabla \hat{u}) - \mu c^l \hat{u} &= \mu f^l \quad \text{in } \mathcal{N}_\delta, \\ \frac{\partial \hat{u}}{\partial \nu} &= 0 \quad \text{on } \partial \mathcal{N}_\delta, \end{aligned}$$

where $\mu = \det A$. This problem is of course equivalent to the problem

$$\frac{1}{\mu} \nabla \cdot (\mu A^{-2} \nabla \hat{u}) - c^l \hat{u} = f^l \quad \text{in } \mathcal{N}_\delta$$

with zero Neumann boundary conditions. A long calculation now reveals that in the case of $G = \mathbb{1}$ and $w = 0$ the resulting elliptic part of our parabolic operator \hat{L} in (5.1) indeed reduces to the operator $\frac{1}{\mu} \nabla \cdot (\mu A^{-2} \nabla \hat{u}) - c^l \hat{u}$,

$$\begin{aligned} \frac{1}{\mu} \nabla \cdot (\mu A^{-2} \nabla \hat{u}) &= \frac{1}{\mu} D_\alpha (\mu A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u}) \\ &= A^{\alpha\beta} A^{\beta\gamma} D_\alpha D_\gamma \hat{u} + A^{\alpha\beta} D_\alpha A^{\beta\gamma} D_\gamma \hat{u} + D_\alpha A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} + \frac{1}{\mu} D_\alpha \mu A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} \\ &= A^{\alpha\beta} A^{\beta\gamma} D_\alpha D_\gamma \hat{u} + A^{\alpha\beta} D_\alpha A^{\beta\gamma} D_\gamma \hat{u} - A^{\kappa\iota} D_\kappa \nu_\iota \frac{\partial}{\partial \nu} \hat{u}, \end{aligned}$$

since

$$\begin{aligned} D_\alpha A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} + \frac{1}{\mu} D_\alpha \mu A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} &= -A^{\alpha\kappa} A^{\beta\iota} D_\alpha A_{\kappa\iota} A^{\beta\gamma} D_\gamma \hat{u} + A^{\kappa\iota} D_\alpha A_{\kappa\iota} A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} \\ &= A^{\alpha\kappa} D_\alpha (d\mathcal{H}_{\kappa\iota}) A^{\iota\beta} A^{\beta\gamma} D_\gamma \hat{u} - A^{\kappa\iota} D_\alpha (d\mathcal{H}_{\kappa\iota}) A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} \\ &= A^{\alpha\kappa} \nu_\alpha \mathcal{H}_{\kappa\iota} A^{\iota\beta} A^{\beta\gamma} D_\gamma \hat{u} + dA^{\alpha\kappa} D_\alpha D_\kappa D_\iota dA^{\iota\beta} A^{\beta\gamma} D_\gamma \hat{u} \\ &\quad - A^{\kappa\iota} \nu_\alpha \mathcal{H}_{\kappa\iota} A^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} - dA^{\kappa\iota} D_\alpha D_\kappa D_\iota dA^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} \\ &= dA^{\alpha\kappa} D_\iota D_\alpha D_\kappa dA^{\iota\beta} A^{\beta\gamma} D_\gamma \hat{u} - A^{\kappa\iota} \mathcal{H}_{\kappa\iota} \nu_\gamma D_\gamma \hat{u} - dA^{\kappa\iota} D_\alpha D_\kappa D_\iota dA^{\alpha\beta} A^{\beta\gamma} D_\gamma \hat{u} \\ &= -A^{\kappa\iota} D_\kappa \nu_\iota \frac{\partial}{\partial \nu} \hat{u}. \end{aligned}$$

From (5.6) it hence follows that

$$\frac{1}{\mu} \nabla \cdot (\mu A^{-2} \nabla \hat{u}) = \tilde{D}_\alpha \tilde{D}_\alpha \hat{u} + \frac{\partial^2 \hat{u}}{\partial \nu^2}.$$

Acknowledgments. This paper resulted from a stay of both authors at the Isaac Newton Institute of Mathematical Sciences in Cambridge, UK, as participants of the research program Free Boundary Problems and Related Topics. The authors would like to thank the institute and the organizers of the program for their invitation and their kind hospitality.

REFERENCES

- [1] A. ALPHONSE, C. M. ELLIOTT, AND B. STINNER, *An Abstract Framework for Parabolic PDEs on Evolving Spaces*, preprint, arXiv:1403.4500, 2014.
- [2] M. BOSTON, *Periodic Solutions for Evolution Equations*, Electron. J. Differ. Equ. Monogr. 3, Department of Math, Texas, State University, San Marcos, 2002.
- [3] C. BAKER, *The Mean Curvature Flow of Submanifolds of High Codimension*, Ph.D. thesis, arXiv:1104.4409, 2011.
- [4] B. CHOW, S.-C. CHU, D. GLICKENSTEIN, C. GUENTHER, J. ISENBERG, T. IVEY, D. KNOPF, P. LU, F. LUO, AND L. NI, *The Ricci Flow: Techniques and Applications. Part II*, Math. Surveys Monogr., AMS, Providence, RI, 2008.
- [5] B. CHOW AND D. KNOPF, *The Ricci Flow: An Introduction*, Math. Surveys Monogr., AMS, Providence, RI, 2004.
- [6] B. CHOW, P. LU, AND L. NI, *Hamilton's Ricci Flow*. Grad. Stud. Math. 77, AMS, Providence, RI, 2006.
- [7] K. DECKELNICK, G. DZIUK, AND C. M. ELLIOTT, *Computation of geometric partial differential equations and mean curvature flow*, Acta Numer., 14 (2005), pp. 139–232
- [8] A. DEMLOW AND G. DZIUK, *An adaptive finite element method for the Laplace–Beltrami operator on implicitly defined surfaces*, SIAM J. Numer. Anal., 45 (2007), pp. 421–442.
- [9] G. DZIUK AND C. M. ELLIOTT, *Finite element methods for surface PDEs*, Acta Numer., 22 (2013), pp. 289–396.
- [10] C. M. ELLIOTT, B. STINNER, AND C. VENKATARAMAN, *Modelling cell motility and chemotaxis with evolving surface finite elements*, J. Roy. Soc. Interface, 9 (2012), pp. 3027–3044.
- [11] C. M. ELLIOTT AND C. VENKATARAMAN, *Error Analysis for an ALE Evolving Surface Finite Element Method*, Preprint NII4032-FRB, Isaac Newton Institute for Mathematical Sciences, 2014.
- [12] P. C. FIFE, *Solutions of parabolic boundary problems existing for all time*, Arch. Ration. Mech. Anal., 16 (1964), pp. 155–186.
- [13] H. FRITZ, *Numerical Ricci–DeTurck flow*, Numer. Math. (2014), DOI:10.1007/s00211-014-0690-5.
- [14] H. GARCKE, K. F. LAM, AND B. STINNER, *Diffuse interface modelling of soluble surfactants in two-phase flow*, Commun. Math. Sci., 12 (2014), pp. 1475–1522.
- [15] P. HESS, *Periodic-Parabolic Boundary Value Problems and Positivity*, Pitman, Boston, 1991.
- [16] G. M. LIEBERMAN, *Time-periodic solutions of linear parabolic differential equations*, Comm. Partial Differential Equations, 24 (1999), pp. 631–663.
- [17] G. M. LIEBERMAN, *Time-periodic solutions of quasilinear parabolic differential equations*, J. Math. Anal. Appl., 264 (2001), pp. 617–638.
- [18] G. M. LIEBERMAN, *Second Order Parabolic Differential Equations*, World Scientific, River Edge, NJ, 1996.
- [19] M. A. OLSHANSKII AND D. SAFIN, *A Narrow-Band Unfitted Finite Element Method for Elliptic PDEs Posed on Surfaces*, preprint, arXiv:1401.7697, 2014.
- [20] M. H. PROTTER AND H. F. WEINBERGER, *Maximum Principles in Differential Equations*, Springer, New York, 1984.
- [21] O. VEJVODA, *Partial Differential Equations: Time-Periodic Solutions*, Martinus Nijhoff, The Hague, 1982.