

Transcriptomics-driven drug repositioning for the treatment of diabetic foot ulcer

by Universitas Ahmad Dahlan Yogyakarta 39

Submission date: 12-Jan-2024 09:05AM (UTC+0700)

Submission ID: 2269696615

File name: 25.pdf (1.44M)

Word count: 5160

Character count: 29048



OPEN Transcriptomics-driven drug repositioning for the treatment of diabetic foot ulcer

Wirawan Adikusuma^{1,2,3✉}, Zainul Amiruddin Zakaria^{1✉}, Lalu Muhammad Irham^{4,5}, Baiq Leny Nopitasari², Anna Pradingsih², Firdayani Firdayani³, Abdi Wira Septama⁵ & Rockie Chong⁶

Diabetic foot ulcers (DFUs) are a common complication of diabetes and can lead to severe disability and even amputation. Despite advances in treatment, there is currently no cure for DFUs and available drugs for treatment are limited. This study aimed to identify new candidate drugs and repurpose existing drugs to treat DFUs based on transcriptomics analysis. A total of 31 differentially expressed genes (DEGs) were identified and used to prioritize the biological risk genes for DFUs. Further investigation using the database DGIdb revealed 12 druggable target genes among 50 biological DFU risk genes, corresponding to 31 drugs. Interestingly, we highlighted that two drugs (urokinase and lidocaine) are under clinical investigation for DFU and 29 drugs are potential candidates to be repurposed for DFU therapy. The top 5 potential biomarkers for DFU from our findings are *IL6ST*, *CXCL9*, *IL1R1*, *CXCR2*, and *IL10*. This study highlights *IL1R1* as a highly promising biomarker for DFU due to its high systemic score in functional annotations, that can be targeted with an existing drug, Anakinra. Our study proposed that the integration of transcriptomic and bioinformatic-based approaches has the potential to drive drug repurposing for DFUs. Further research will further examine the mechanisms by which targeting *IL1R1* can be used to treat DFU.

26 Abbreviations

BP	Biological processes
CC	Cellular components
DEG	Differentially expressed genes
DFU	Diabetic foot ulcer
DGIdb	Drug-gene interaction database
27	Diabetes mellitus
GO	Gene Ontology
KEGG	Kyoto Encyclopedia of Genes and Genomes
KMP	Knockout mouse phenotype
Log FC	Log fold change
MF	Molecular functions
MP	Mammalian Phenotype Ontology
ORA	Over-representation analysis
PID	Primary immunodeficiency

Diabetes mellitus (DM) is a common chronic condition affecting 451 million people globally. By 2045, it is predicted that the number of DM patients would increase to 693 million¹. Treatment for DM thus far has only been able to regulate blood sugar levels; unfortunately, it does not totally cure the condition. Improper management

¹Borneo Research on Algesia, Inflammation, and Neurodegeneration (BRAIN) Group, Department of Biomedical Sciences, Faculty of Medicines and Health Sciences, University Malaysia Sabah, Jalan UMS, 88400 Kota Kinabalu, Sabah, Malaysia. ²Departement of Pharmacy, University of Muhammadiyah Mataram, Mataram, Indonesia. ³Research Center for Accine and Drugs, National Research and Innovation Agency (BRIN), 10 Jth Tangerang, Indonesia. ⁴Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta, Indonesia. ⁵Research Center for Pharmaceutical Ingredients and Traditional Medicine, National Research and Innovation Agency (BRIN), South Tangerang, Indonesia. ⁶Department of Chemistry and Biochemistry, University of California, Los Angeles, USA. ✉email: adikusuma28@gmail.com; zaz@ums.edu.my

of DM can result in chronic problems; 15–25% of DM patients will experience diabetic foot ulcer (DFU), and 50–70% of relapses will occur within five years^{2,3}. Amputation, death, and hospitalization rates are frequently high because of the refractory wound of DFUs⁴. DFU occurs in 85% of diabetic amputees before developing significant gangrene or infection⁵. Early detection is necessary to avoid long-term consequences. Hospitalization, disability, and mortality in DM patients will decrease if the risk of DFU can be identified early. Biomarkers are crucial for early clinical diagnosis, illness prevention, and disease progression prediction. In addition, finding and developing novel therapeutic medications or assessing therapeutic approaches is also essential^{6,7}.

Genomic and biomedical information in the form of databases has rapidly become available, thanks to technological advances in experimental and computational biology. These comprise transcriptome data (such as human patient gene expression profiles, animal models of human diseases, small molecule treatments, etc.), other molecular profiling technologies, and publicly available databases, providing a previously unheard-of chance to enhance rational drug design in combination with the implementation of the network idea of drug targets and the power of phenotypic screening^{8,9}. Transcriptome data can enable the identification and prioritization of biomarkers that can be used as potential therapeutic candidates⁷. In the past, several studies successfully identified promising results for various indications, such as inflammatory bowel disease¹⁰, dermatomyositis¹¹, cancer^{12–14}, and preterm birth¹⁵, by applying a transcriptomics-based computational drug repositioning pipeline. Consequently, we postulate that transcriptomic data can also help identify biomarkers for DFU and potential drug candidates for this disease.

Integrating transcriptomics and bioinformatics could provide new insights into DFU pathogenesis by developing common transcription features. Here, we applied a transcriptomics-based computational drug repositioning pipeline to identify potential candidate therapy for DFU. This study was designed to discover the common genes among DFU patients. Detailed information on the study workflow is depicted in Fig. 1. The common DEGs and their roles in DFU were examined. Additionally, biological DFU risk genes were identified based on six functional annotations. As intended, higher scores that are given to candidate genes by the annotation process represent a more prominent biological influence that we termed “biological risk genes” for the pathogenesis of DFU. Subsequently, we used an *in-silico* workflow for the analyses from multiple databases to find prospective drugs or biomarkers for DFU. This information will be an important tool to not only drive biomarker discovery but can also drive drug repurposing for DFUs.

Results

Identification of differentially expressed genes for DFU. This study leveraged the previously published study datasets GSE37265 and GSE80178.

The GSE37265 and GSE80178 expression profile were identified using the R limma package to determine the differentially expressed genes (DEGs). We chose these two datasets with the following rationale. Firstly, we ensured that all the subjects included in our studies were human. Secondly, the sample type was healthy or diseased tissue from patients rather than a specific cell type, which was consistent with our goal for drug repositioning. Thirdly, we ensured that the datasets included in our study had complete data for analysis and had obtained




Figure 1. A workflow of the bioinformatics-driven drug repositioning pipeline for diabetic foot ulcer (DFU). This figure was created with BioRender.com under agreement number “QL24UF44NA”.

ethical approval. Finally, we included diseased or normal skin samples and a control group without systemic or autoimmune diseases or relevant family history as inclusion criteria for our study.

A total of 358 DEGs in the GSE37265 dataset were successfully identified, including 28 upregulated and 330 downregulated, as demonstrated in the volcano plots in Fig. 2A. Next, we identified the DEGs from the GSE80178 datasets between six DFUs and three normal skin tissues. This step, 734 DEGs were identified (175 upregulated genes and 559 downregulated genes), as illustrated in the volcano plots in Fig. 2B. To increase the stringency of the identified risk genes, all DEGs were collected based on the section of the analysis results of two datasets. Thirty-one overlapping genes were obtained from both groups, as shown in Fig. 2C and Supplementary Table S1.

Biological DFU risk genes. A strict functional annotation procedure was used to find biological DFU risk genes. To obtain more prospective drug target genes, we enlarge the networking of DEGs through the STRING database prior to identifying biological DFU risk genes. We set the threshold of 50 interactions, inputted 31 DEG from the phases before, and obtained 81 DEGs for further analysis (Fig. 2D; Supplementary Table S2). Furthermore, in order to prioritize genes for drug discovery for the present study, we utilized a scoring system previously employed in other research studies, as mentioned in the previous publications^{16–19}. Following are the scoring results of the six functional annotations: (1) gene prioritized by Knockout mouse phenotype (MP) (n = 31); (2) gene prioritized by Primary immunodeficiency (PID) 2019 (n = 7); (3) gene prioritized by Kyoto Encyclopedia of Genes and Genomes (KEGG) (n = 28); (4) gene prioritized by Biological processes (BP) (n = 47); (5) gene prioritized by Cellular components (CC) (n = 22); and (6) gene prioritized by Molecular functions (MF) (n = 38). Figure 3A and 3B shows a distribution score for each criterion. Ultimately, 50 biological DFU risk genes met the requirements with a score of 2 or higher. From further analysis of the gene scores, our findings revealed that the top five genes, *IL6ST*, *CXCL9*, *IL1R1*, *CXCR2*, and *IL10*, all having scores higher than 4 (Fig. 3C).




Figure 2. Detection of differentially expressed genes (DEGs) for diabetic foot ulcer. (A) A volcano plot corresponding to the GSE37265. (B) Volcano plot representing the identification of DEGs in the GSE80178 dataset. (C) Venn diagrams showing DEGs based on the intersection between GSE37265 and GSE80178 datasets. (D) Gene network expansion for the identified DEGs using STRING database. The color of the nodes reflects the degree of connectivity, with darker colors indicating a higher interaction score.




Figure 3. Prioritizing biological risk genes for diabetic foot ulcer (DFU) using a scoring system based on functional annotations. **(A)** A diagram pie chart shows the distribution score in each functional annotation for all 6 annotations. **(B)** An overview of gene score distribution for the 6 functional annotations. **(C)** Satisfied criteria for 50 biological DFU risk genes with a score of two or higher were indicated by the colors across each of the 6 functional annotations. A white box indicates the lack of functional annotations.

Discovery candidate drugs that potentially for DFU therapeutic. Next, we mapped 50 biological DFU risk genes into DGIdb (https://www.dgldb.org/search_interactions, accessed on November 12, 2022). We investigated whether gene profiles from biological DFU risk genes could be targeted by drugs with known interactions type. However, not all of the obtained biological DFU risk genes from the pipeline are druggable. Only 12 DFU-risk genes correspond to 31 drugs based on DGIdb (Supplementary Table S3). Hence, these drugs were considered as potential candidate drugs for DFU therapy. It is important to note among these 31 drugs, two drugs (urokinase; NCT01108120 and lidocaine; NCT04154046) were under clinical investigation for DFU, and 29 drugs were potential candidates drugs for DFU that have not been reported to DFU therapy (Fig. 4). In particular, our study identified *IL1R1* overlapping anakinra was screened as a highly promising DFU target, since it also achieved a high systemic score in functional annotations. Our proof-of-principle study demonstrated that transcriptomic data not only can be utilized for biomarker discovery for DFUs, but it can also drive drug repurposing for this devastating disease.

Discussion

Bioinformatics-driven drug repositioning is a method that uses computational techniques to identify new potential uses for existing drugs²⁰. Drug repositioning is a promising strategy for treating DFU, a common complication of diabetes that can lead to amputation. By identifying existing drugs that could target the underlying molecular disease mechanisms, drug repositioning could provide an alternative, promising approach that quickly identifies and prioritize new potential treatment options for DFU patients. This study uses transcriptomic data to conduct a bioinformatics-based drug repositioning analysis for DFU. The study could analyze the gene expression profiles of tissue samples from patients with DFU and compare them to normal samples. This could help identify upregulated or downregulated genes in the ulcer tissue, which could suggest potential targets for drug repositioning. Our next step was prioritizing the DEGs for candidate drug prediction using a scoring system based on six functional annotations. Through this pipeline, we successfully identified 31 drugs targeting 12 genes, two of which (urokinase and lidocaine) are in clinical trials for DFU.

Our research identified the top 5 potential markers in biological DFU risk genes: *IL6ST*, *CXCL9*, *IL1R1*, *CXCR2*, and *IL10*. *IL6ST*, also known as gp130, plays a key role in the inflammatory response and has been found to have higher expression in diabetic wounds^{20–22}. *CXCL9* is a chemokine involved in the immune response that coordinates cells in the wound-healing process²³. *CXCR2* is a chemokine receptor that is involved in inflammation. Keratinocytes and endothelial cells express *CXCR2* during wound healing in regions where epithelialization and neovascularization occur. Leukocyte recruitment, keratinocyte migration and proliferation, and angiogenesis are necessary for wound healing²⁴. *IL10* is an anti-inflammatory cytokine that helps to regulate the immune response and reduce inflammation, with decreased levels in DFUs, suggesting that a lack of *IL10* may contribute to the progression of these wounds²⁵. Lastly, *IL1R1* is a receptor for the pro-inflammatory cytokine *IL-1*. It is increased in DFUs, indicating an overactive *IL-1* response may contribute to the development of these wounds^{26,27}. Of note, while our study focused specifically on gene expression differences between DFUs and healthy skin tissue, it is important to note that DFUs are a specific type of chronic wound with distinct

28 Methods

Identification of differentially expressed genes. The gene expression profiling datasets GSE37265 and GSE80178 were retrieved from the GEO database (<https://www.ncbi.nlm.nih.gov/geo/>, accessed on November 12, 2022). The GSE37265 expression profile comprised 14 DFU and 14 normal tissues, which were paired samples from the same patients, and was analyzed by the Affymetrix Human Genome U133 Plus 2.0 Array. The GSE80178 expression profile comprised six DFUs and three normal skin tissues, which were unpaired samples, and was analyzed by the Affymetrix Human Gene 2.0 ST Array. The downloaded data had already been normalized. Here, we use the *limma* package³² in R to identify DEGs. We applied a paired model matrix for the paired samples in GSE37265 to account for the correlation between the DFU and normal tissue samples from the same patients, and an unpaired model matrix for the unpaired samples in GSE80178 that did not account for pairing information. DEGs were screened using the following criteria: $|\log \text{fold change (log FC)}| > 2$ and p -value 0.05. Only samples from human skin tissue were included in our study, and we ensured that the datasets had complete data for analysis and had obtained ethical approval. We also included DFU or normal samples and a control group without systemic or autoimmune diseases or a relevant family history, which met our necessary inclusion criteria. Finally, we visualized the DEGs that met the intersection between GSE37265 and GSE80178 using a Venn diagram.

Expansion of the network of differentially expressed genes by using the STRING Database. The DEG was enlarged using the STRING database to get more potential drug target genes. The STRING database (<https://string-db.org>, accessed on November 12, 2022) was created to integrate protein-protein interactions with functional connections related to protein expression³³. We entered the list of DEGs chosen in the preceding phases and established the threshold of 50 interactions in order to increase the number of the initial set of DEGs. The rationale is a higher potential to discover new disease treatment targets with more disease-protein networks³⁴.

Prioritizing biological DFU risk genes. Next, we used the network-expanded list of DEGs, leveraging the STRING database for further functional annotation to explore deep insights into the pathogenesis of DFU and identify DFU therapeutic targets. The six functional annotations were applied to the filter of DEGs by a scoring system, with the criteria as follows: (1) KMP: To determine whether the gene contributes to certain mouse phenotypic diseases. Genes were prioritized using Mammalian Phenotype Ontology (MP) from WebGestalt (2019) with an FDR $q < 0.05$ and were considered significant³⁵; (2) PID: DFU is associated with innate immune diseases. Noteworthy, genetic variants overlapping with PID genes may contribute to DFU pathogenesis. PID genes were collected from the 2019 Update of the IUIS Phenotypical Classification³⁶. A hypergeometric test was used to conduct enrichment analysis on the data; the significance threshold for this step was p -value < 0.05 ; (3) KEGG: to identify the molecular pathway³⁷. KEGG prioritized genes in the Webgestalt³⁵. The significance threshold was set at a q -value (FDR) < 0.05 ; (4) Gene Ontology (GO): Generally, GO categories can be broken down as follows: BP, CC, and MF to identify specific biological functions involved in DFU. A GO enrichment analysis was performed using from WebGestalt³⁵, and FDR $q < 0.05$ was used as a significance threshold. Genes scored more than equal to 2 were considered biological DFU-risk genes. Genes with higher annotation scores represent genes with a more prominent biological impact on DFU pathogenesis which we termed “biological risk genes”.

Drug gene interactions for DFU. In order to identify DFU candidate drugs, we analyzed the set of biological DFU risk genes using the drug-gene interaction database (DGIdb) (https://www.dgldb.org/search_interactions). The DGIdb is an online database that compiles data on drug-gene interactions and druggable genes from articles, databases, and other online resources for drug discovery. The information about drugs, genes, and interactions is normalized and combined into conceptual groups³⁸. From DGIdb, candidate drugs are selected based on FDA-approved drugs with clear interaction types.

Statistical analysis. This study was conducted using R (version 4.2.1) for all statistical analyses. The *limma* package was used to identify DEGs³². The WebGestalt 2019 R package was used to perform over-representation analysis (ORA), including KMP, GO, and KEGG pathway enrichment analysis³⁵. PID was analyzed using the hypergeometric test with a significance threshold of p -value < 0.05 . The *ggplot2* package (R v4.2.1)³⁹ was used to visualize KMP, GO, and KEGG. The Venn diagram package (R v4.2.1)⁴⁰ was applied to visualize all DEGs based on the intersection between GSE37265 and GSE80178. In addition, DFU candidate drugs were visualized using a chord diagram generated by the *circlize* package in R⁴¹.

4 Data availability

All data generated or analyzed during this study are included in this article and its supplementary information files.

Received: 5 March 2023; Accepted: 15 June 2023

Published online: 20 June 2023

References

1. Cho, N. H. *et al.* IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res. Clin. Pract.* **138**, 271–281. <https://doi.org/10.1016/j.diabres.2018.02.023> (2018).

2. Alavi, A. *et al.* Diabetic foot ulcers: Part I. Pathophysiology and prevention. *J. Am. Acad. Dermatol.* **70**, 1 e1–18. <https://doi.org/10.1016/j.jaad.2013.06.055> (2014).
3. Armstrong, D. G., Boulton, A. J. M. & Bus, S. A. Diabetic foot ulcers and their recurrence. *N Engl. J. Med.* **376**, 2367–2375. <https://doi.org/10.1056/NEJMra1615439> (2017).
4. Prompers, L. *et al.* Resource utilisation and costs associated with the treatment of diabetic foot ulcers. Prospective data from the Eurodiale Study. *Diabetologia* **51**, 1826–1834. <https://doi.org/10.1007/s00125-008-1089-6> (2008).
5. Zhang, P. *et al.* Global epidemiology of diabetic foot ulceration: a systematic review and meta-analysis (dagger). *Ann. Med.* **49**, 106–116. <https://doi.org/10.1080/07853890.2016.1231932> (2017).
6. Qian, L. *et al.* Integrated bioinformatics-based identification of potential diagnostic biomarkers associated with diabetic foot ulcer development. *J. Diabetes Res.* **2021**, 5445349. <https://doi.org/10.1155/2021/5445349> (2021).
7. Wang, Y. *et al.* An update on potential biomarkers for diagnosing diabetic foot ulcer at early stage. *Biomed. Pharmacother.* **133**, 110991. <https://doi.org/10.1016/j.biopha.2020.110991> (2021).
8. Atta, L. & Fan, J. Computational challenges and opportunities in spatially resolved transcriptomic data analysis. *Nat. Commun.* **12**, 5283. <https://doi.org/10.1038/s41467-021-25557-9> (2021).
9. Le, B. L. *et al.* Transcriptomics-based drug repositioning pipeline identifies therapeutic candidates for COVID-19. *Sci. Rep.* **11**, 12310. <https://doi.org/10.1038/s41598-021-91625-1> (2021).
10. Dudley, J. T. *et al.* Computational repositioning of the anticonvulsant topiramate for inflammatory bowel disease. *Sci. Transl. Med.* **3**, 96ra76. <https://doi.org/10.1126/scitranslmed.3002648> (2011).
11. Cho, H. G., Fiorentino, D., Lewis, M., Sirota, M. & Sarin, K. Y. Identification of alpha-adrenergic agonists as potential therapeutic agents for dermatomyositis through drug-repurposing using public expression datasets. *J. Invest. Dermatol.* **136**, 1517–1520. <https://doi.org/10.1016/j.jid.2016.03.001> (2016).
12. Chen, B. *et al.* Computational discovery of niclosamide ethanolamine, a repurposed drug candidate that reduces growth of hepatocellular carcinoma cells in vitro and in mice by inhibiting cell division cycle 37 signaling. *Gastroenterology* **152**, 2022–2036. <https://doi.org/10.1053/j.gastro.2017.02.039> (2017).
13. Chen, B. *et al.* Reversal of cancer gene expression correlates with drug efficacy and reveals therapeutic targets. *Nat. Commun.* **8**, 16022. <https://doi.org/10.1038/ncomms16022> (2017).
14. Jahchan, N. S. *et al.* A drug repositioning approach identifies tricyclic antidepressants as inhibitors of small cell lung cancer and other neuroendocrine tumors. *Cancer Discov.* **3**, 1364–1377. <https://doi.org/10.1158/2159-8290.CD-13-0183> (2013).
15. Le, B. L., Iwatani, S., Wong, R. J., Stevenson, D. K. & Sirota, M. Computational discovery of therapeutic candidates for preventing preterm birth. *JCI Insight* <https://doi.org/10.1172/jci.insight.133761> (2020).
16. Adikusuma, W. *et al.* Identification of druggable genes for asthma by integrated genomic network analysis. *Biomedicines* <https://doi.org/10.3390/biomedicines10010113> (2022).
17. Adikusuma, W. *et al.* Drug repurposing for atopic dermatitis by integration of gene networking and genomic information. *Front. Immunol.* **12**, 724277. <https://doi.org/10.3389/fimmu.2021.724277> (2021).
18. Okada, Y. *et al.* Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature* **506**, 376–381. <https://doi.org/10.1038/nature12873> (2014).
19. Imamura, M. *et al.* Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes. *Nat. Commun.* **7**, 10531. <https://doi.org/10.1038/ncomms10531> (2016).
20. Reay, W. R. & Cairns, M. J. Advancing the use of genome-wide association studies for drug repurposing. *Nat. Rev. Genet.* **22**, 658–671. <https://doi.org/10.1038/s41576-021-00387-z> (2021).
21. Pradhan, L. *et al.* Gene expression of pro-inflammatory cytokines and neuropeptides in diabetic wound healing. *J. Surg. Res.* **167**, 336–342. <https://doi.org/10.1016/j.jss.2009.09.012> (2011).
22. Niemi, J. P. *et al.* Injury-induced gp130 cytokine signaling in peripheral ganglia is reduced in diabetes mellitus. *Exp. Neurol.* **296**, 1–15. <https://doi.org/10.1016/j.expneurol.2017.06.020> (2017).
23. Ridandries, A., Tan, J. T. M. & Bursill, C. A. The role of chemokines in wound healing. *Int. J. Mol. Sci.* <https://doi.org/10.3390/ijms19103217> (2018).
24. Devalaraja, R. M. *et al.* Delayed wound healing in CXCR2 knockout mice. *J. Invest. Dermatol.* **115**, 234–244. <https://doi.org/10.1046/j.1523-1747.2000.00034.x> (2000).
25. Nanda, R., Patel, S., Ghosh, A., Asha, K. S. & Mohapatra, E. A study of apolipoprotein A1 (ApoA1) and interleukin-10 (IL-10) in diabetes with foot ulcers. *Biomedicine (Taipei)* **12**, 30–38. <https://doi.org/10.37796/2211-8039.1279> (2022).
26. Thomay, A. A. *et al.* Disruption of interleukin-1 signaling improves the quality of wound healing. *Am. J. Pathol.* **174**, 2129–2136. <https://doi.org/10.2353/ajpath.2009.080765> (2009).
27. Tan, J. L. *et al.* Restoration of the healing microenvironment in diabetic wounds with matrix-binding IL-1 receptor antagonist. *Commun. Biol.* **4**, 422. <https://doi.org/10.1038/s42003-021-01913-9> (2021).
28. Ramirez, J. & Canete, J. D. Anakinra for the treatment of rheumatoid arthritis: A safety evaluation. *Expert Opin. Drug Saf.* **17**, 727–732. <https://doi.org/10.1080/14740338.2018.1486819> (2018).
29. Fleischmann, R. M. *et al.* Anakinra, a recombinant human interleukin-1 receptor antagonist (r-metHuIL-1ra), in patients with rheumatoid arthritis: A large, international, multicenter, placebo-controlled trial. *Arthritis Rheum.* **48**, 927–934. <https://doi.org/10.1002/art.10870> (2003).
30. Theocharidis, G. *et al.* Single cell transcriptomic landscape of diabetic foot ulcers. *Nat. Commun.* **13**, 181. <https://doi.org/10.1038/s41467-021-27801-8> (2022).
31. Li, Y. *et al.* Characterization of the microenvironment of diabetic foot ulcers and potential drug identification based on scRNA-seq. *Front. Endocrinol. (Lausanne)* **13**, 997880. <https://doi.org/10.3389/fendo.2022.997880> (2022).
32. Ritchie, M. E. *et al.* Limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47. <https://doi.org/10.1093/nar/gkv007> (2015).
33. Szklarczyk, D. *et al.* STRING v11: Protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* **47**, D607–D613. <https://doi.org/10.1093/nar/gky1131> (2019).
34. Ruffner, H., Bauer, A. & Bouwmeester, T. Human protein-protein interaction networks and the value for drug discovery. *Drug Discov. Today* **12**, 709–716. <https://doi.org/10.1016/j.drudis.2007.07.011> (2007).
35. Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z. & Zhang, B. WebGestalt 2019: Gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Res.* **47**, W199–W205. <https://doi.org/10.1093/nar/gkz401> (2019).
36. Bousfiha, A. *et al.* Human inborn errors of immunity: 2019 update of the IUIS phenotypical classification. *J. Clin. Immunol.* **40**, 66–81. <https://doi.org/10.1007/s10875-020-00758-x> (2020).
37. Kanehisa, M. & Goto, S. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**, 27–30. <https://doi.org/10.1093/nar/28.1.27> (2000).
38. Freshour, S. L. *et al.* Integration of the Drug-Gene Interaction Database (DGIdb 4.0) with open crowdsourcing efforts. *Nucleic Acids Res.* **49**, D1144–D1151. <https://doi.org/10.1093/nar/gkaa1084> (2021).
39. Wickham H. Ggplot2: Elegant Graphics for Data Analysis. 2nd edn (Springer, 2009). <https://doi.org/10.1007/978-0-387-98141-3>.
40. Chen, H. & Boutros, P. C. VennDiagram: a package for the generation of highly-customizable Venn and Euler diagrams in R. *BMC Bioinform.* **12**, 35. <https://doi.org/10.1186/1471-2105-12-35> (2011).

41. Gu, Z., Gu, L., Eils, R., Schlesner, M. & Brors, B. circlize Implements and enhances circular visualization in R. *Bioinformatics* **30**, 2811–2812. <https://doi.org/10.1093/bioinformatics/btu393> (2014).

Acknowledgements

This project was supported by grants from Riset Muhammadiyah (RisetMu) Batch VI 2022 Majelis DIKTI-LITBANG PP Muhammadiyah & Universitas Muhammadiyah Mataram (No: 1687.032/PP/I.3/D/2022).

Author contributions

Conceptualized, conceived, and designed the study: W.A., Z.A.Z and L.M.I. Performed the analysis: W.A. Curated the data: W.A., and L.M.I. Wrote the original draft: WA. Interpreted the data: W.A., M.I., B.L.N., A.P., F.F., A.W.S., R.C and Z.A.Z. Revised the manuscript: W.A., L.M.I., B.L.N., A.P., F.F., A.W.S., R.C and Z.A.Z. All authors have read and agreed to the published version of the manuscript.

6

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-023-37120-1>.

23

Correspondence and requests for materials should be addressed to W.A. or Z.A.Z.

5

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License¹⁶ which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

Transcriptomics-driven drug repositioning for the treatment of diabetic foot ulcer

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	researchmgt.monash.edu Internet Source	1%
2	mdpi-res.com Internet Source	1%
3	Petteri Stenroos, Isabelle Guillemain, Federico Tesler, Olivier Montigon et al. "How Absence Seizures Impair Sensory Perception: Insights from Awake fMRI and Simulation Studies in Rats", Cold Spring Harbor Laboratory, 2024 Publication	1%
4	assets.researchsquare.com Internet Source	1%
5	Lauren M. Sanders, Ryan T. Scott, Jason H. Yang, Amina Ann Qutub et al. "Biological research and self-driving labs in deep space supported by artificial intelligence", Nature Machine Intelligence, 2023 Publication	1%

6

Nicole Boivin, Susanne Täuber, Ulrike Beisiegel, Ursula Keller, Janet G. Hering. "Sexism in academia is bad for science and a waste of public funding", Nature Reviews Materials, 2023

Publication

1 %

7

Jessica J. Y. Lee, Michael M. Gottlieb, Jake Lever, Steven J. M. Jones et al. "Text-based phenotypic profiles incorporating biochemical phenotypes of inborn errors of metabolism improve phenomics-based diagnosis", Journal of Inherited Metabolic Disease, 2018

Publication

1 %

8

Lalu Muhammad Irham, Henry Sung-Ching Wong, Wan-Hsuan Chou, Wirawan Adikusuma, Eko Mugiyanto, Wan-Chen Huang, Wei-Chiao Chang. "Integration of Genetic Variants and Gene Network for Drug Repurposing in Colorectal Cancer", Pharmacological Research, 2020

Publication

1 %

9

www.journal.uad.ac.id

Internet Source

1 %

10

Marissa Angelina, Muhammad Hanafi, Tri Yuliani, Franciscus Suyatna, Tjahjani Mirawati Soediro, Beti Ernawati Dewi. "The inhibitory activity of Cassia alata leaves extract on denv-2 replication infected mice", Pharmacia, 2022

<1 %

11

Yuqing Wang, Tengteng Shao, Jialin Wang, Xiaoting Huang, Xiaofei Deng, Yemin Cao, Mingmei Zhou, Cheng Zhao. "An update on potential biomarkers for diagnosing diabetic foot ulcer at early stage", Biomedicine & Pharmacotherapy, 2021

Publication

<1 %

12

Arwa A. Al-Qahtani, Fatimah S. Alhamlan, Ahmed Ali Al-Qahtani. "Pro-Inflammatory and Anti-Inflammatory Interleukins in Infectious Diseases: A Comprehensive Review", Tropical Medicine and Infectious Disease, 2024

Publication

<1 %

13

www.omicsdi.org

Internet Source

<1 %

14

Submitted to Taylor's Education Group

Student Paper

<1 %

15

hrcak.srce.hr

Internet Source

<1 %

16

www.intechopen.com

Internet Source

<1 %

17

www.saffronbooks.com

Internet Source

<1 %

18

www.dovepress.com

Internet Source

<1 %

19 Submitted to Fayetteville State University <1 %
Student Paper

20 Long Qian, Zhipeng Xia, Ming Zhang, Qiong Han, Die Hu, Sha Qi, Danmou Xing, Yan Chen, Xin Zhao. "Integrated Bioinformatics-Based Identification of Potential Diagnostic Biomarkers Associated with Diabetic Foot Ulcer Development", Journal of Diabetes Research, 2021 <1 %
Publication

21 Submitted to University of Queensland <1 %
Student Paper

22 hub.hku.hk <1 %
Internet Source

23 research.vu.nl <1 %
Internet Source

24 www.biorxiv.org <1 %
Internet Source

25 Lalu Muhammad Irham, Wirawan Adikusuma, Lolita Lolita, Anisa Nova Puspitaningrum et al. "Investigation of susceptibility genes for chickenpox disease across multiple continents", Biochemistry and Biophysics Reports, 2023 <1 %
Publication

26 journals.lww.com

Internet Source

<1 %

27

www.aging-us.com

Internet Source

<1 %

28

www.frontiersin.org

Internet Source

<1 %

29

www.researchsquare.com

Internet Source

<1 %

30

Felipe Andre Oliveira Freitas, Luiz F. Brito, Simara Larissa Fanalli, Janaína Lustosa Gonçalves et al. "Identification of eQTL using different sets of single nucleotide polymorphisms associated with carcass and body composition traits in pigs", Research Square Platform LLC, 2023

Publication

<1 %

31

Yuzhang Hu, Mei Dang, Murtala Bindawa Isah, Saeed Yakhkeshi, Chen Chen, Xiaoying Zhang. "Comparative analysis of muscle profiles in silky fowl and white Leghorn Chicken: Insights from multi-omics and experimental approaches", LWT, 2023

Publication

<1 %

32

journals.biologists.com

Internet Source

<1 %

33

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

34

www.azolifesciences.com

Internet Source

<1 %

35

Eileen M. Weinheimer-Haus, Rita E. Mirza, Timothy J. Koh. "Nod-Like Receptor Protein-3 Inflammasome Plays an Important Role during Early Stages of Wound Healing", PLOS ONE, 2015

Publication

<1 %

36

Guanghua Chen, Jia Yan. "Integrated bioinformatics-based identification of potential diagnostic biomarkers associated with atopic dermatitis", Advances in Dermatology and Allergology, 2022

Publication

<1 %

37

Huaxiong Zhang, Yiya Zhang, Yangfan Li, Yaling Wang, Sha Yan, San Xu, Zhili Deng, Xinling Yang, Hongfu Xie, Ji Li. "Bioinformatics and Network Pharmacology Identify the Therapeutic Role and Potential Mechanism of Melatonin in AD and Rosacea", Frontiers in Immunology, 2021

Publication

<1 %

38

Nicola Capuano, Pasquale Foggia, Luca Greco, Pierluigi Ritrovato. "A Linked Data Application for Harmonizing Heterogeneous

<1 %

39

Ok Hyung Nam, Tae Jun Oh, Jae-Hyung Lee, Yu-Shik Hwang, Sung Chul Choi. "Differential gene expression profiles of human periodontal ligament cells preserved in Hank's balanced salt solution and milk", Dental Traumatology, 2019

Publication

<1 %

40

journal.umpr.ac.id

Internet Source

<1 %

41

www.cell.com

Internet Source

<1 %

42

Arief Rahman Afief, Lalu Muhammad Irham, Wirawan Adikusuma, Dyah Aryani Perwitasari, Ageng Brahmadhi, Rocky Cheung. "Integration of genomic variants and bioinformatic-based approach to drive drug repurposing for multiple sclerosis", Biochemistry and Biophysics Reports, 2022

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On