

University of Groningen

Correction

Ersoydan, Selin; Yakar, Derya; Kasalak, Ömer; Kwee, Thomas C

Published in:
European Radiology

DOI:
[10.1007/s00330-023-09821-8](https://doi.org/10.1007/s00330-023-09821-8)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Ersoydan, S., Yakar, D., Kasalak, Ö., & Kwee, T. C. (2023). Correction: Did medical doctors who order abdominal CT scans during on-call hours truly become worse at clinical reasoning? Yes, they did. *European Radiology*, 33, 8396–8397. <https://doi.org/10.1007/s00330-023-09821-8>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Correction: Did medical doctors who order abdominal CT scans during on-call hours truly become worse at clinical reasoning? Yes, they did

Selin Ersoydan¹ · Derya Yakar¹ · Ömer Kasalak¹ · Thomas C. Kwee¹

Published online: 20 June 2023
© The Author(s) 2023

Correction: European Radiology (2022) 33:1015-1021
<https://doi.org/10.1007/s00330-022-09121-7>

The original version of this article, published on 7 September 2022, unfortunately contained some mistakes.

The authors noted an error in the calculation of the annual clinical reasoning scores. The original legend with the explanation of this calculation and the display of the upper panel of Fig. 2 were incorrect, and have been replaced with a corrected version. Some sentences in the results of the abstract and main text have also been corrected based on this recalculation (see below). However, these corrections do not change the interpretation of the results and the conclusions we have drawn.

Abstract, results section, corrected first two sentences:
The median annual clinical reasoning score was 14.7% (interquartile range: 12.2 to 16.0%; range: 7.7 to 34.6%). The quality of clinical reasoning significantly decreased between 2005 and 2019 (Mann–Kendall Tau of -0.390 , $p=0.048$), while the number of abdominal CT scans significantly increased (Mann–Kendall tau of 0.790 , $p<0.001$).

Main text, results section, corrected paragraph “Clinical reasoning quality and number of CT scans: temporal changes”:

The median annual clinical reasoning score was 14.7% (IQR: 12.2 to 16.0%; range: 7.7 to 34.6%). Clinical reasoning scores significantly decreased over time (Mann–Kendall tau of -0.390 , $p<0.048$) (Fig. 2). The median annual number of abdominal CT scans during on-call hours on the 36 randomly sampled unique calendar days was 35 (IQR: 23.25 to 46.75; range: 17–55). The number of abdominal CT scans significantly increased over time (Mann–Kendall tau of 0.790 , $p<0.001$) (Fig. 2).

Figure 2, Upper panel

Clinical reasoning scores per year (gray line), calculated as $(a/b) \times 100\%$, where a denotes the number of differential diagnoses on the CT request forms that matched the CT diagnoses and b denotes the total number of differential diagnoses on the CT request forms, for each year. The blue line represents the non-parametric locally estimated scatterplot smoothing (LOESS) fit (Mann–Kendall Tau of -0.390 , $P=0.048$).


The upper panel of Fig. 2 has also been replaced with a newer version.

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1007/s00330-022-09121-7>.

✉ Thomas C. Kwee
thomaskwee@gmail.com

¹ Department of Radiology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not

permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.