

University of Groningen

Corrigendum to “Novel spirooxindole based benzimidazole scaffold: In vitro, nanoformulation and in vivo studies on anticancer and antimetastatic activity of breast adenocarcinoma”

Barakat, Assem; Alshahrani, Saeed; Al-Majid, Abdullah Mohammed; Alamar, Abdullah Saleh; Haukka, Matti; Abu-Serie, Marwa M.; Dömling, Alexander; Mazyed, Eman A.; Badria, Farid A.; El-Senduny, Fardous F.

Published in:
 Bioorganic Chemistry

DOI:
[10.1016/j.bioorg.2023.107012](https://doi.org/10.1016/j.bioorg.2023.107012)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2024

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Barakat, A., Alshahrani, S., Al-Majid, A. M., Alamar, A. S., Haukka, M., Abu-Serie, M. M., Dömling, A., Mazyed, E. A., Badria, F. A., & El-Senduny, F. F. (2024). Corrigendum to “Novel spirooxindole based benzimidazole scaffold: In vitro, nanoformulation and in vivo studies on anticancer and antimetastatic activity of breast adenocarcinoma”. *Bioorganic Chemistry*, 143, Article 107012. <https://doi.org/10.1016/j.bioorg.2023.107012>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the “Taverne” license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Corrigendum

Corrigendum to “Novel spirooxindole based benzimidazole scaffold: *In vitro*, nanoformulation and *in vivo* studies on anticancer and antimetastatic activity of breast adenocarcinoma” [Bioorg. Chem. 129 (2022) 106124]

Assem Barakat^{a,*}, Saeed Alshahrani^a, Abdullah Mohammed Al-Majid^a,
Abdullah Saleh Alamaray^a, Matti Haukka^b, Marwa M. Abu-Serie^c, Alexander Dömling^d,
Eman A. Mazyed^e, Farid A. Badria^{f,*}, Fardous F. El-Senduny^g

^a Department of Chemistry, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

^b Department of Chemistry, University of Jyväskylä, P.O. Box 35, FI-40014 Jyväskylä, Finland

^c Medical Biotechnology Department, Genetic Engineering and Biotechnology Research Institute, City of Scientific Research and Technological Applications (SRTA-City), Egypt

^d Department of Drug Design, Groningen Research Institute of Pharmacy, University of Groningen, 9713 AV Groningen, The Netherlands

^e Department of Pharmaceutical Technology, Faculty of Pharmacy, Kaferelsheikh University, Kaferelsheikh, P.O. Box 33516, Egypt

^f Department of Pharmacognosy, Faculty of Pharmacy, Mansoura University, Mansoura 35516, Egypt

^g Department of Chemistry, Faculty of Science, Mansura University, Mansura 35516, Egypt

This is a critical needed correction. In Fig. 15, the image corresponding to the lung from mouse treated with 25 mg/kg was taken from the same section of the mouse that was treated with 10 mg/kg.

The corrected Fig. 15

(red arrows), peribronchial and interstitial aggregation (black arrows) of tumour cells admixed with MNCs. Microscopic pictures of H&E stained lungs sections from treated groups (B) 10 or (c) 25 mg/kg) showing disappeared congestion with decreased numbers of perivascular and interstitial infiltration of tumour cells. Increasing dose of treatment 25




Fig. 15. Microscopic pictures of H&E stained lung sections from groups received (A) 4×10^6 cells/200 μ L showing congested blood vessels

mg/kg was more efficient than 10 mg/kg. Low magnification X: 100 with 100 μ m scale bar.

DOI of original article: <https://doi.org/10.1016/j.bioorg.2022.106124>.

* Corresponding authors.

E-mail addresses: ambarakat@ksu.edu.sa (A. Barakat), badri002@mans.edu.eg (F.A. Badria).

<https://doi.org/10.1016/j.bioorg.2023.107012>

Available online 8 December 2023

0045-2068/© 2023 Elsevier Inc. All rights reserved.