

University of Groningen

The role of depressive symptoms and symptom dimensions in actigraphy-assessed sleep, circadian rhythm, and physical activity

Difrancesco, Sonia; Penninx, Brenda W J H; Riese, Harriëtte; Giltay, Erik J; Lamers, Femke

Published in:
 Psychological Medicine

DOI:
[10.1017/S0033291720004870](https://doi.org/10.1017/S0033291720004870)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Difrancesco, S., Penninx, B. W. J. H., Riese, H., Giltay, E. J., & Lamers, F. (2021). The role of depressive symptoms and symptom dimensions in actigraphy-assessed sleep, circadian rhythm, and physical activity. *Psychological Medicine*, 52(13), 2760–2766. <https://doi.org/10.1017/S0033291720004870>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Original Article

Cite this article: Difrancesco S, Penninx BWJH, Riese H, Giltay EJ, Lamers F (2021). The role of depressive symptoms and symptom dimensions in actigraphy-assessed sleep, circadian rhythm, and physical activity. *Psychological Medicine* 1–7. <https://doi.org/10.1017/S0033291720004870>

Received: 7 April 2020

Revised: 9 November 2020

Accepted: 23 November 2020

Key words:


Actigraphy; major depressive disorder; symptom dimensions

Author for correspondence:

Sonia Difrancesco,

E-mail: s.difrancesco@ggzingeest.nl

The role of depressive symptoms and symptom dimensions in actigraphy-assessed sleep, circadian rhythm, and physical activity

Sonia Difrancesco¹ , Brenda W. J.H. Penninx¹, Harriëtte Riese², Erik J. Giltay³ and Femke Lamers¹

¹Department of Psychiatry, Amsterdam UMC, Vrije Universiteit, Amsterdam Public Health research institute, Amsterdam, The Netherlands; ²Department of Psychiatry, University of Groningen, University Medical Center Groningen, Interdisciplinary Center for Psychopathology and Emotion regulation, Groningen, The Netherlands and ³Department of Psychiatry, Leiden University Medical Center, Leiden, The Netherlands

Abstract

Background. Considering the heterogeneity of depression, distinct depressive symptom dimensions may be differentially associated with more objective actigraphy-based estimates of physical activity (PA), sleep and circadian rhythm (CR). We examined the association between PA, sleep, and CR assessed with actigraphy and symptom dimensions (i.e. mood/cognition, somatic/vegetative, sleep).

Methods. Fourteen-day actigraphy data of 359 participants were obtained from the Netherlands Study of Depression and Anxiety. PA, sleep, and CR estimates included gross motor activity (GMA), sleep duration (SD), sleep efficiency (SE), relative amplitude between daytime and night-time activity (RA) and sleep midpoint. The 30-item Inventory of Depressive Symptomatology was used to assess depressive symptoms, which were categorised in three depression dimensions: mood/cognition, somatic/vegetative, and sleep.

Results. GMA and RA were negatively associated with higher score on all three symptom dimensions: mood/cognition (GMA: $\beta = -0.155$, $p < 0.001$; RA: $\beta = -0.116$, $p = 0.002$), somatic/vegetative (GMA: $\beta = -0.165$, $p < 0.001$; RA: $\beta = -0.133$, $p < 0.001$), sleep (GMA: $\beta = -0.169$, $p < 0.001$; RA: $\beta = -0.190$, $p < 0.001$). The association with sleep was more pronounced for two depression dimensions: longer SD was linked to somatic/vegetative ($\beta = 0.115$, $p = 0.015$) dimension and lower SE was linked to sleep ($\beta = -0.101$, $p = 0.011$) dimension.

Conclusion. As three symptom dimensions were associated with actigraphy-based low PA and dampened CR, these seem to be general indicators of depression. Sleep disturbances appeared more linked to the somatic/vegetative and sleep dimensions; the effectiveness of sleep interventions in patients reporting somatic/vegetative symptoms may be explored, as well as the potential of actigraphy to monitor treatment response to such interventions.

Introduction

Major depressive disorder (MDD) is a highly prevalent disorder, associated with high disability (Murray et al., 2012). It often has a chronic course (Verduijn et al., 2017) and a third of the patients experience poor treatment outcomes (Gaynes et al., 2009). Despite the challenges in finding consensus regarding classification, diagnosis and treatment, there has been a recent and significant increase in research to identify novel methods to measure, unravel aetiology, and treat depressive disorders. Actigraphy, an ecologically valid method to objectively measure disturbances in sleep, circadian rhythm (CR) and physical activity (PA), has become widely used in depression research.

Several studies, including the systematic review of Burton et al. (2013) and our previous work (Difrancesco et al. 2019; Minaeva et al. 2020), have shown that depressive disorders and symptoms are associated with the lower daily activity (Helgadóttir, Forsell, & Ekblom, 2015; Hori et al., 2016; Stubbs et al., 2016) and CR amplitude (Lyll et al., 2018). The association between actigraphy measures of sleep and depression is less clear. Although night-time activity level appears to be higher in patients with depression (Burton et al., 2013), this is not always reflected in actigraphy measures of sleep duration (SD) and sleep efficiency (SE). Despite studies assessing sleep with self-reported questionnaires have shown that both insomnia and hypersomnia are associated with depression (Nutt, Wilson, & Paterson, 2008; van Mill, Hoogendijk, Vogelzangs, Van Dyck, & Penninx, 2010), our previous results with actigraphy have found that higher severity of depressive symptom is associated with longer (but not shorter) SD (Difrancesco et al., 2019). Depression is however heterogeneous in its presentation and these results may not be generalizable to all patients.

© The Author(s) 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article.

CAMBRIDGE
UNIVERSITY PRESS

Symptoms vary substantially in patients with depression and opposite clinical presentations occur. For instance, both insomnia and hypersomnia (Nutt et al., 2008) and psychomotor retardation and agitation (Avery & Silverman, 1984) are symptoms of depression. Considering that 1030 unique Diagnostic and Statistical Manual of Mental Disorders-Fifth Edition (DSM-5) symptom profiles have been found among 3703 patients with MDD (Fried & Nesse, 2015), substantial symptom variation occurs among individuals who all qualify for the same DSM-5 diagnosis. Thus, as the 'depression' label provides limited information about the particular problems experienced by a patient, some researchers have moved away from traditional methods of diagnosis such as the DSM-5 and have focused on symptom dimensions (Fried, 2017). Although approaches to identify symptom dimensions vary, empirical methods such as factor analyses have been employed to identify symptoms dimensions or factor structures underlying self-reported questionnaires on depressive symptoms in a population-based sample, such as the often used Inventory of Depressive Symptomatology (IDS) (Hegeman et al., 2012; Rush, Gullion, Basco, Jarrett, & Trivedi, 1996; Wardenaar et al., 2010). Most studies illustrate the existence of three major symptom dimensions: mood/cognition, somatic, and sleep symptom dimensions (Hegeman et al., 2012; Rush et al., 1996; Wardenaar et al., 2010).

Although it is clear that depression is heterogeneous, little research has focused on the role of symptoms dimensions assessed with severity measures in sleep and CR disturbances and physical inactivity as assessed objectively with actigraphy. As cognitive impairments have been linked to sleep problems (Mantua & Simonelli, 2019) and physical inactivity has been related to somatic/energy-related symptoms (Puetz, 2006), a differential association with symptom dimensions may be expected. However, this has not properly been investigated so far. A better understanding would be of benefit for the diagnosis and treatment of depression. For instance, pharmacological treatment, psychotherapy and chronotherapies to treat sleep and CR problems can be especially administered to patients with specific symptom dimensions.

The aim of this study was to examine the association between actigraphy-based PA, sleep, and CR with symptom dimensions (i.e. cognition/mood, somatic, sleep). The association with individual depressive symptoms was also explored to further check on the consistency of findings for symptoms within dimensions. To address these questions, we used data from the Netherlands Study of Depression and Anxiety (NESDA) which includes persons without and with current or remitted depressive and/or anxiety disorders. As depression and anxiety are highly comorbid disorders (Lamers et al., 2011; Rodney et al., 1997) with the previously mentioned limitations in traditional diagnosis method and using the full sample gives the opportunity to investigate the associations on a larger spectrum of severity.

Method

Sample

Participants from the NESDA were selected to participate in the Ecological Momentary Assessment (EMA) and Actigraphy sub-study (NESDA-EMAA). Details about NESDA have been provided extensively before (Penninx et al., 2008). NESDA was designed to investigate the course of depressive and anxiety disorders over a period of several years and the factors that influence

the development and prognosis of such disorders. NESDA participants were initially included at the baseline assessment in 2004–2007 ($n = 2981$) and seen for the fifth time at the 9-year follow-up assessment wave (2014–2017, $n = 1776$) for a follow-up interview, during which 367 siblings of NESDA participants were added, bringing the 9-year follow-up sample to 2143 subjects. At this wave, 384 participants enrolled for the EMAA sub-study. The NESDA study, including NESDA-EMAA, was approved by the VUmc ethical committee (reference number 2003/183) and all respondents gave informed consent for both the regular interview and the EMAA sub-study. See for a flow-chart of the NESDA-EMAA in our previous work (Difrancesco et al., 2019). Participants were eligible if they had a smartphone or were willing to use a smartphone provided by the study, and were willing to wear a wrist-worn actigraphy device. Participants to the EMAA-sub-study were provided with a GENEActiv device (Activinsights Ltd, Kimbolton, UK). Participants wore the wrist-worn GENEActiv actigraphy device for 14 days on their non-dominant wrist. Of the 384 participants included in the NESDA-EMAA study, 14 had no available actigraphy data for several reasons, such as technical failure, resulting in 370 (96.4%) participants with available data. According to previously published criteria (da Silva et al., 2014), participant's actigraphy data were included in analyses if at least 1-week day and one weekend day of usable data was available, with at least 16 h recorded per day and per night. The final sample was composed of 359 (93.5%) participants with an average $13.68 \pm \text{s.e. } 1.26$ valid days, of whom 90% completed the protocol for 14 days.

Depressive symptoms and symptom dimensions

The Dutch version of the 30-item IDS, self-report (Rush et al., 1996) was used to assess depressive symptoms (0 = no problems to 3 = severe problems) in the previous 7 days. A total sum score can be calculated (range, 0–84), with higher scores indicating higher levels of depressive symptomatology. Validity and reliability have been shown before, with Cronbach's alpha ranging from 0.92 to 0.94 (Rush et al., 1996). We used three-dimensional depression symptom sum scores based in prior factor analyses in NESDA (Wardenaar et al., 2010): mood/cognition, somatic/vegetative, and sleep. The mood/cognition dimension consisted of the 16 items: feeling irritable, interpersonal sensitivity, feeling sad, diminished quality of mood, feeling anxious or tense, diminished capacity of pleasure/enjoyment, diminished reactivity of mood, diminished interest in people/activities, suicidal thoughts, future pessimism, concentration/decision-making problems, self-criticism and blame, psychomotor retardation, reduced interest in sex, low energy level/fatigability, leaden paralysis. The somatic/vegetative domain included nine items: panic/phobic symptoms, psychomotor agitation, decreased weight, increase in appetite, other bodily symptoms, decrease in appetite, increased weight, constipation/diarrhoea, aches, and pains. Finally, the sleep domain included four items: early morning awakening, problems sleeping during the night, problems falling asleep, sleeping too much. The item 'diurnal variation (worse in the morning)' did not belong to any domain in the factor analyses and was therefore used only in our exploratory analyses with individual symptoms.

Actigraphy estimates of sleep, CR, and PA

In this study, the accelerometer was set to sample at 30 Hz and raw actigraphy data were analysed using an open source R

package, GGIR (version 1.5–18). Actigraphy data were gathered after the depressive symptoms assessment with a median number of 23 days between the two assessments. As described before (Difrancesco et al., 2019), we used objective indicators of sleep, CR, and PA. Sleep was assessed as total SD per night [in hh:mm] and SE per night [%]. CR was assessed by the relative amplitude (RA) between daytime and night-time activity per day and sleep midpoint [clock time]. RA describes the amplitude between the activity during the day and the night; lower RA means a dampened CR amplitude suggesting lower activity during the day and higher activity during the night. Sleep midpoint is a proxy of chronotype or preference for morningness/eveningness; later sleep midpoint means a preference for evening chronotype. Exploratory analyses showed that RA and sleep midpoint have low correlation and therefore they describe different aspects of CR (online Supplementary material Table S1). PA is indexed/expressed given as gross motor activity (GMA) per day [milligravity (mg), $1\text{ g} = 9.81\text{ m/s}^2$]. Daily estimates of each variable were used in the current study.

Covariates and descriptive variables

Covariates were age, sex, and education level expressed in years at the time of the NESDA EMEA sub-study. These covariates were selected as they have an established theoretical association with depression and with sleep, CR and PA levels, and have been regularly used in similar studies (Droomers, Schrijvers, & Mackenbach, 2001; Stamatakis, Kaplan, & Roberts, 2007).

DSM-IV based diagnoses of depressive disorders (dysthymia and MDD) and anxiety (social anxiety disorder, panic disorder with and without agoraphobia, agoraphobia, and generalized anxiety disorder) were established with the Composite International Diagnostic Interview (CIDI, version 2.1) (Wittchen, 1994). The interviews were conducted at the regular interview by specially trained clinical research staff. Participants were divided into three groups for descriptive purposes only: (1) a group with no lifetime depressive and/or anxiety disorders, (2) a group with remitted depressive and/or anxiety disorders (having a lifetime, but not current (6-month) diagnosis), and (3) a group with current depressive or anxiety disorder in the past 6 months.

Antidepressant use was based on drug container inspection, and medications were coded according to the World Health Organization Anatomical Therapeutic Chemical (ATC) classification. Antidepressant use was considered present if participants reported using it more than 50% of the time. Antidepressants included were selective serotonin reuptake inhibitors (SSRIs, ATC code N06AB), tricyclic antidepressant (TCA, ATC code N06AA) and other antidepressants (ATC codes N06AF, N06AG, N06AX).

Statistical analyses

Distributions of all variables were checked on normality with QQ plots. Non-normally distributed outcomes were transformed with Log-transformation (GMA) or with Box-Cox transformation (SD, SE, RA between daytime and night-time activity level and sleep midpoint). Correlation of symptom dimensions was explored using Pearson correlation coefficient to assess the degree of inter-correlation of symptom dimensions. The association of actigraphy estimates with symptom dimensions and individual symptoms was tested in separate models using Generalized Estimating Equations (GEE) corrected for covariates and with each

actigraphy estimate as an outcome. GEE was used to account for correlations between repeated days per person. Separate models were run for each symptoms dimension and each individual symptoms; each model was adjusted for age, sex, and education level. As having a current/remitted diagnosis of depression and/or anxiety is highly correlated with the severity of symptoms, adjustment for diagnostic groups was not performed.

All analyses were performed with the statistical software R (version 1.0.143). A p value < 0.05 was considered statistically significant. Correction for multiple testing by using false discovery rate was applied when testing the association with symptom dimensions adjusting for the number of outcomes ($n = 5$) and symptom dimensions ($n = 3$). This was not done for individual symptoms, as our aim was to visually explore overall patterns to help understand whether symptoms follow the same patterns as the dimension they belong to or not. Specifically, we visually assessed the univariate associations of individual symptoms with actigraphy measures by using a forest plot presenting the standardised coefficients and the 95% confidence intervals. In this way, it was possible to visualise whether the associations with individual symptoms followed the same direction as the dimension they belong to and whether some symptoms were outliers.

Results

Demographics and descriptive

Table 1 shows demographics and descriptive in our sample. The average age of the sample was 49.5 ± 12.6 years, of whom 63.7% was female and with a duration of education of 12.9 ± 3.1 years. Most of the persons included had a lifetime diagnosis of depressive and/or anxiety disorders: 93 (26.0%) persons had current depressive and/or anxiety disorders, 176 (49.0%) persons had remitted depressive and/or anxiety disorders and only 90 (25.0%) persons had no depressive and/or anxiety disorders. The average severity of depressive symptoms in the sample was 14.2 ± 11.7 on the IDS total score.

Symptoms dimensions were moderately to strongly correlated (correlation ranged from 0.41 for the somatic/vegetative and sleep dimensions, to 0.75 for mood/cognition and somatic/vegetative dimensions, not tabulated and available upon request).

Association of sleep, CR, and PA with dimensions

A 14-day lower GMA and RA between daytime and night-time activity level were significantly associated with all three symptom dimensions (Table 2, all $p < 0.05$). The associations with actigraphy-based sleep were more pronounced for two dimensions. Longer SD was significantly associated with higher somatic/vegetative symptom dimension score (Table 2; $\beta = 0.113$, $p = 0.021$) but not with mood/cognition and sleep symptom dimension scores. Lower SE was associated with higher sleep symptom dimension score (Table 2; $\beta = -0.101$, $p = 0.011$) but not with mood/cognition and somatic/vegetative symptom dimension scores.

Patterns of sleep, CR, and PA in individual symptoms

Figure 1 shows the association of individual symptoms with 14-day PA, sleep, and CR patterns. Associations of GMA and RA between daytime and night-time activity level showed more

Table 1. Demographics and psychopathology in the NESDA sample

	All subjects (<i>n</i> = 359)
Demographics	
Age, mean (s.d.)	49.5 (12.6)
Female, <i>n</i> (%)	228 (63.7%)
Education [year], mean (s.d.)	12.9 (3.1)
Psychopathology	
Depressive and/or anxiety disorders	
No lifetime disorders, <i>n</i> (%)	90 (25.0%)
Remitted disorder, <i>n</i> (%)	176 (49.0%)
Current disorder, <i>n</i> (%)	93 (26.%)
IDS Score, mean (s.d.)	14.2 (11.7)
Mood/Cognition symptom score (IDS), mean (s.d.)	8.4 (7.9)
Somatic/vegetative symptom score (IDS), mean (s.d.)	5.1 (3.6)
Sleep symptom score (IDS), mean (s.d.)	3.0 (2.0)
Antidepressant use, <i>n</i> (%)	71 (19.7%)

significant associations with individual symptoms compared to other actigraphy variables.

For GMA and RA 63% (*n* = 19) and 53% (*n* = 16) of all individual symptoms – spread across different dimensions – were significantly associated with lower GMA and lower RA between daytime and night-time activity level. Other symptoms – although not significantly associated – followed a similar, expected negative direction in their associations with GMA and RA. By inspecting the forest plots, no differential association of GMA and RA with symptom dimensions can be confirmed.

When considering actigraphy-based SD, in total, eight of the 30 individual symptoms were significantly associated with longer SD. Although most of somatic/vegetative symptoms followed the same direction in the association with SD, ‘increased weight’ deviated from the overall pattern. Also, as expected ‘early morning awakenings’ in the sleep dimension was associated with objective shorter SD.

Although not captured in the analyses using symptom dimensions, 30% (*n* = 9) of all symptoms were associated with later sleep midpoint. Of these seven symptoms, four belong to the mood/cognition dimension and three belong to the sleep dimension. Most individual symptoms were following the same positive direction in the association with sleep midpoint, with the exception of ‘early morning awakenings’ that showed an – expected – negative direction. This may possibly show that there is more heterogeneity in terms of effect estimates for individual symptoms.

Only two symptoms were associated with lower SE. As expected, all sleep symptoms were following the same negative direction in the association with SE as shown by the sleep dimension.

Discussion

This is the first study to examine 14-day actigraphy estimates of sleep, CR, and PA and their relationship with symptoms and symptom dimensions of depression. As all three symptom dimensions were associated with lower and dampened 14-day PA and

Table 2. Univariate association^a between symptom dimensions and sleep, circadian rhythm, and physical activity (*n* = 359)

Symptom dimensions	Physical activity			Sleep			Circadian rhythm								
	Gross motor activity			Sleep duration			Relative amplitude between daytime and nighttime activity								
	β	S.E.	<i>p</i>	β	S.E.	<i>p</i>	β	S.E.	<i>p</i>						
Mood/Cognition	-0.163	0.043	<0.001	0.081	0.042	0.066	-0.048	0.036	0.200	-0.122	0.036	0.003	0.091	0.047	0.066
Somatic/vegetative	-0.146	0.038	<0.001	0.113	0.044	0.021	-0.043	0.038	0.263	-0.116	0.039	0.008	0.090	0.046	0.066
Sleep	-0.169	0.038	<0.001	0.078	0.039	0.066	-0.101	0.036	0.011	-0.190	0.039	<0.001	0.069	0.043	0.132

Correction for multiple testing by using false discovery rate (FDR).

Significance of bold values = *p* < 0.05.

^aGEE model (outcomes = sleep, circadian rhythm, and physical activity) adjusted for age, sex, and education level.




Fig. 1. Univariate association between individual IDS symptoms and the actigraphy measures: physical activity (i.e. gross motor activity), sleep (i.e. sleep duration, sleep efficiency), and circadian rhythm (i.e. relative amplitude, sleep midpoint) in all sample ($n = 359$). Error bars represent 95% confidence intervals. Adjusted for age, sex, and education.
 Note: ● = mood/cognition, ▲ = somatic/vegetative, ■ = sleep, I = no dimension; black = $p < 0.05$, grey = $p \geq 0.05$.

circadian patterns, it could be concluded that physical inactivity and low circadian amplitude are general indicators of depressive symptom severity. However, sleep disturbances may be more specifically linked to somatic/vegetative and sleep symptom dimensions, as associations with longer SD and lower SE were more pronounced for these symptom dimensions. While exercise and behavioural activation may be used as adjunctive treatments in all patients with depression, sleep interventions may be especially effective in people with more self-reported somatic/vegetative and sleep symptoms.

While few studies have suggested that actigraphy may serve as an objective measure of psychomotor retardation in patients with depression (Krane-Gartiser, Henriksen, Vaaler, Fasmer, & Morken, 2015), our results seem to support that physical inactivity is a general feature of depression (Burton et al., 2013) and symptom severity (Minaeva et al., 2020). This was also reflected in our analyses with individual symptoms, as over 60% of depressive symptoms – across different symptom dimensions – were associated with low GMA with moderate effect sizes. The association between PA and depression has been studied before and appears to be bidirectional. Patients with depression are typically less active and they experience a range of barriers to engaging in PA such as depressive symptoms, higher body mass index, physical co-morbidity, and lower self-efficacy (Vancampfort et al., 2015). Low levels of PA increase the risk of depression (Mammen & Faulkner, 2013) and, PA and exercise (i.e. structured physical activity) can improve depressive symptoms (Schuch et al., 2016). A large meta-analysis in the general population (Biswas et al., 2015) demonstrated that sedentary behaviour is associated with an increased risk of developing cardiovascular disease, type 2 diabetes, cardiovascular and all-cause mortality. As persons with depression are less active than the general population, they

encounter additional risks in developing cardiovascular and chronic diseases and mortality. PA and behavioural activation, predominantly through exercise, may produce an antidepressant effect through multiple biological and psychosocial pathways (Kandola, Ashdown-Franks, Hendrikse, Sabiston, & Stubbs, 2019). Neuroplasticity, neuroendocrine responses, inflammation, and oxidative stress have been suggested to play a role at a biological level. Several psychosocial factors accompany and potentially interact with, these biological changes to influence depression, such as self-esteem (Kandola et al., 2019). While there is less research into the psychosocial benefits of exercise on depression, they are likely to be of parallel importance. As we observed lower PA across all symptom dimensions, PA and behavioural activation may be of help to all patients with depression as it may produce antidepressant effects through multiple biological and psychosocial pathways.

Similarly, dampened circadian amplitude was generally associated with all symptom dimensions, supporting previous findings on the association between dampened rhythms and mood disorders (Lyall et al., 2018). As reviewed by Vadnie and McClung (2017), multiple mechanisms can explain the association between CR disruptions and mood disorders including depression. One of the main theories is that disrupted CRs in the master pacemaker, or suprachiasmatic nucleus may be the cause of mood disturbances. However, also other environmental factors, such as light, or genetic perturbations may influence mood independently from the central clock. Additionally, other studies have suggested that sleep and CR disturbances may be symptoms of mood disorders. As dampened circadian amplitude was associated with all symptom dimensions, CR interventions such as behavioural activation may be administered as adjunctive treatment to all patients with depression.

When studying sleep, it is important to make a clear distinction between subjective and objective sleep measures. The first provides information about a person's perception of his/her sleep quantity and quality. The latter refers to the objective measure of SD and SE with actigraphy. When assessing subjective sleep, persons diagnosed with depression often report sleep disturbances such as insomnia and hypersomnia (Nutt et al., 2008; van Mill et al., 2010). While in our previous work using actigraphy (Difrancesco et al., 2019) we have shown that depressive symptom severity is associated with longer objective SD, in these follow-up analyses the association appeared to be more pronounced in patients with somatic/vegetative symptoms. We found a possible explanation for these findings. Persons with more somatic/vegetative symptoms experience more physical complaints, such as aches and pains, bodily symptoms, than may directly impact on their ability to be physically active and resulting in more sleep disturbances. The sleep dimension itself may be problematic as it combines opposite clinical presentations (i.e. insomnia and hypersomnia) for which the associations with objective sleep is expected in opposite directions. However, as could be expected, lower SE assessed objectively with actigraphy was associated with sleep symptoms assessed with self-ratings. Although in our previous work we did not find an association between objective SE and subjective sleep assessed with the Women's Health Initiative Insomnia Rating Scale (Difrancesco et al., 2019), objective measure of SE seems indicative of self-reported sleep disturbances assessed with the IDS. Sleep interventions may focus not only on patients reporting sleep problems but also on those reporting somatic/vegetative symptoms. Actigraphy may be used to assess the effectiveness of such interventions.

This study has several limitations. First, although we adjusted for multiple testing, we explored several associations when looking at individual symptoms and that may lead to chance findings. However, we aimed to look more at broader patterns than to focus on specific symptoms. Second, the assessment of depression symptoms and actigraphy were not done at the same time, although there was a median number of 23 days between the two assessments. Another limitation is that sleep and CRs are complicated concepts and actigraphy may provide only an indirect and rough estimate of them based on wrist movement, whereas polysomnography is the gold standard. In addition, sleep midpoint was evaluated during the entire period and it can be argued that chronotype is often assessed with sleep midpoint on free days corrected for the sleep debt accumulated during the working days. However, sleep midpoint measured with actigraphy provides a valid approximation of chronotype and it is moderately correlated with it as shown by Santisteban, Brown, and Gruber (2018). Although not assessed in our study, shift work may affect sleep schedules and therefore it may influence the results on the association with sleep and CRs (James, Honn, Gaddameedhi, & Van Dongen, 2017). While data did not show reversed activity/sleep patterns as would be expected with night shifts, early and late day shifts could have occurred and may also have an impact. Individual symptoms may fit in different symptoms dimensions and classifications are not always consistent across studies. We based our classification on those of Wardenaar et al. (2010), in which classification, psychomotor retardation, and low energy level/fatigability, for instance, were part of the mood/cognition dimension, but they could also be fit in the somatic/vegetative dimension. However, by looking at individual symptom patterns we found consistency as most of symptoms follow the same direction in the associations as the

dimension they belong to. Although we did not investigate the role of anxiety, as Wardenaar et al. (2010) have shown that symptom dimensions were consistent across diagnostic groups (i.e. remitted depression, lifetime anxiety, healthy controls), comorbid anxiety does not seem to confound the studied associations. An important strength of the study is the attempt to investigate the heterogeneity of depressive symptoms with actigraphy in a relatively large sample. Another important strength is that on average 14 days of actigraphy data per participant were available.

To conclude, this is the first study to examine the different symptom dimensions of depression and their association with several objective indicators of PA, sleep, CR gathered with wrist-worn actigraphy devices. As lower PA level and dampened circadian amplitude were associated with higher scores on all three dimensions, lower PA level and dampened CRs appeared to be general indicators of depression and depression severity. Disturbances in objective sleep were more pronounced for somatic/vegetative symptoms, suggesting that longer SD may be more closely linked to this dimension. Sleep interventions may focus not only on patients reporting sleep problems but also on those reporting somatic/vegetative symptoms. PA, behavioural activation, and CR interventions may be not limited to groups of patients based on their symptoms but may be promoted in all patients with depression. However, clinical implementation awaits further evidence based on empirical data, if indeed, actigraphy may be used to monitor treatment response of such interventions.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291720004870>

Acknowledgments. The infrastructure for the NESDA study (www.nesda.nl) is funded through the Geestkracht program of the Netherlands Organisation for Health Research and Development (ZonMw, grant number 10-000-1002) and financial contributions by participating universities and mental health care organizations (VU University Medical Center, GGZ inGeest, Leiden University Medical Center, Leiden University, GGZ Rivierduinen, University Medical Center Groningen, University of Groningen, Lentis, GGZ Friesland, GGZ Drenthe, Rob Giel Onderzoekscentrum). The analytical work was financially supported by Innovative Medicines Initiative 2 Joint undertaking under grant agreement No 115902.

Conflict of interest. All authors declare that they have no conflicts of interest.

Data availability statement. According to European law (GDPR) data containing potentially identifying or sensitive patient information are restricted; our data involving clinical participants are not freely available in a public repository. However, data are available upon request via the NESDA Data Access Committee (nesda@ggzingeest.nl).

References

- Avery, D., & Silverman, J. (1984). Psychomotor retardation and agitation in depression: Relationship to age, sex, and response to treatment. *Journal of Affective Disorders*, 7(1), 67–76. [https://doi.org/10.1016/0165-0327\(84\)90066-1](https://doi.org/10.1016/0165-0327(84)90066-1)
- Biswas, A., Oh, P. I., Faulkner, G. E., Bajaj, R. R., Silver, M. A., Mitchell, M. S., & Alter, D. A. (2015). Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: A systematic review and meta-analysis. *Annals of Internal Medicine*, 162(2), 123–132. <https://doi.org/10.7326/M14-1651>
- Burton, C., McKinsty, B., Szentagotai Tatar, A., Serrano-Blanco, A., Pagliari, C., & Wolters, M. (2013). Activity monitoring in patients with depression: A systematic review. *Journal of Affective Disorders*, 145(1), 21–28. <https://doi.org/10.1016/j.jad.2012.07.001>

- da Silva, I., van Hees, V. T., Ramires, V. V., Knuth, A. G., Bielemann, R. M., Ekelund, U., ... Hallal, P. C. (2014). Physical activity levels in three Brazilian birth cohorts as assessed with raw triaxial wrist accelerometry. *International Journal of Epidemiology*, 43(6), 1959–1968. <https://doi.org/10.1093/ije/dyu203>
- Difrancesco, S., Lamers, F., Riese, H., Merikangas, K. R., Beekman, A. F., van Hemert, A. M., ... Penninx, B. W. J. H. (2019). Sleep, circadian rhythm, and physical activity patterns in depressive and anxiety disorders: A 2-week ambulatory assessment study. *Depression and Anxiety*, 36(10), 1–12. <https://doi.org/10.1002/da.22949>
- Droomers, M., Schrijvers, C. T. M., & Mackenbach, J. P. (2001). Educational level and decreases in leisure time physical activity: Predictors from the longitudinal globe study. *Journal of Epidemiology and Community Health*, 55(8), 562–568. <https://doi.org/10.1136/jech.55.8.562>
- Fried, E. I. (2017). Moving forward: How depression heterogeneity hinders progress in treatment and research. *Expert Review of Neurotherapeutics*, 17(5), 423–425. <https://doi.org/10.1080/14737175.2017.1307737>
- Fried, E. I., & Nesse, R. M. (2015). Depression is not a consistent syndrome: An investigation of unique symptom patterns in the STAR*D study. *Journal of Affective Disorders*, 172, 96–102. <https://doi.org/10.1016/j.jad.2014.10.010>
- Gaynes, B. N., Warden, D., Trivedi, M. H., Wisniewski, S. R., Fava, M., & Rush, A. J. (2009). What did STAR*D teach us? Results from a large-scale, practical, clinical trial for patients with depression. *Psychiatric Services*, 60(11), 1439–1445. <https://doi.org/10.1176/ps.2009.60.11.1439>
- Hegeman, J. M., Wardenaar, K. J., Comijs, H. C., de Waal, M. W. M., Kok, R. M., & van der Mast, R. C. (2012). The subscale structure of the inventory of depressive symptomatology self report (IDS-SR) in older persons. *Journal of Psychiatric Research*, 46(10), 1383–1388. <https://doi.org/10.1016/j.jpsychires.2012.07.008>
- Helgadottir, B., Forsell, Y., & Ekblom, Ö. (2015). Physical activity patterns of people affected by depressive and anxiety disorders as measured by accelerometers: A CrossSectional study. *PLoS ONE*, 10(1), Article e0115894. <https://doi.org/10.1371/journal.pone.0115894>
- Hori, H., Koga, N., Hidese, S., Nagashima, A., Kim, Y., Higuchi, T., & Kunugi, H. (2016). 24-H Activity rhythm and sleep in depressed outpatients. *Journal of Psychiatric Research*, 77, 27–34. <https://doi.org/10.1016/j.jpsychires.2016.02.022>
- James, S. M., Honn, K. A., Gaddameedhi, S., & Van Dongen, H. P. A. (2017). Shift work: Disrupted circadian rhythms and sleep – implications for health and well-being. *Current Sleep Medicine Reports*, 3(2), 104–112. <https://doi.org/10.1007/s40675-017-0071-6>
- Kandola, A., Ashdown-Franks, G., Hendrikse, J., Sabiston, C. M., & Stubbs, B. (2019). Physical activity and depression: Towards understanding the anti-depressant mechanisms of physical activity. *Neuroscience and Biobehavioral Reviews*, 107, 525–539. <https://doi.org/10.1016/j.neubiorev.2019.09.040>
- Krane-Gartiser, K., Henriksen, T. E. G., Vaaler, A. E., Fasmer, O. B., & Morken, G. (2015). Actigraphically assessed activity in unipolar depression: A comparison of inpatients with and without motor retardation. *Journal of Clinical Psychiatry*, 76(9), 1181–1187. <https://doi.org/10.4088/JCP.14m09106>
- Lamers, F., Van Oppen, P., Comijs, H. C., Smit, J. H., Spinhoven, P., Van Balkom, A. J. L. M., ... Penninx, B. W. J. H. (2011). Comorbidity patterns of anxiety and depressive disorders in a large cohort study: The Netherlands study of depression and anxiety (NESDA). *Journal of Clinical Psychiatry*, 72(3), 342–348. <https://doi.org/10.4088/JCP.10m06176blu>
- Lyall, L. M., Wyse, C. A., Graham, N., Ferguson, A., Lyall, D. M., Cullen, B., ... Smith, D. J. (2018). Association of disrupted circadian rhythmicity with mood disorders, subjective wellbeing, and cognitive function: A cross-sectional study of 91 105 participants from the UK Biobank. *The Lancet Psychiatry*, 5(6), 507–514. [https://doi.org/10.1016/S2215-0366\(18\)30139-1](https://doi.org/10.1016/S2215-0366(18)30139-1)
- Mammen, G., & Faulkner, G. (2013). Physical activity and the prevention of depression: A systematic review of prospective studies. *American Journal of Preventive Medicine*, 45(5), 649–657. <https://doi.org/10.1016/j.amepre.2013.08.001>
- Mantua, J., & Simonelli, G. (2019). Sleep duration and cognition: Is there an ideal amount? *Sleep*, 42(3), Article zsz010. <https://doi.org/10.1093/sleep/zsz010>
- Minaeva, O., Booi, S. H., Lamers, F., Antypa, N., Schoevers, R. A., Wichers, M., & Riese, H. (2020). Level and timing of physical activity during normal life in depressed and non-depressed individuals. *Translational Psychiatry*, 10, Article 259. <https://doi.org/10.1038/s41398-020-00952-w>
- Murray, C. J. L., Vos, T., Lozano, R., Naghavi, M., Flaxman, A. D., Michaud, C., ... Lopez, A. D. (2012). Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: A systematic analysis for the global burden of disease study 2010. *Lancet (London, England)*, 380(9859), 2197–2223. [https://doi.org/10.1016/S0140-6736\(12\)61689-4](https://doi.org/10.1016/S0140-6736(12)61689-4)
- Nutt, D. J., Wilson, S., & Paterson, L. (2008). Sleep disorders as core symptoms of depression. *Dialogues in Clinical Neuroscience*, 10(3), 329–336.
- Penninx, B. W. J. H., Beekman, A. T. F., Smit, J. H., Zitman, F. G., Nolen, W. A., Spinhoven, P., ... van Dyck, R. (2008). The Netherlands Study of Depression and Anxiety (NESDA): Rationale, objectives and methods. *International Journal of Methods in Psychiatric Research*, 17(3), 121–140. <https://doi.org/10.1002/mpr.256>
- Puetz, T. W. (2006). Physical activity and feelings of energy and fatigue. *Sports Medicine*, 36(9), 767–780. <https://doi.org/10.2165/00007256-200636090-00004>
- Rodney, J., Prior, N., Cooper, B., Theodores, M., Browning, J., Steinberg, B., & Evans, L. (1997). The comorbidity of anxiety and depression. *Australian and New Zealand Journal of Psychiatry*, 31(5), 700–703. <https://doi.org/10.3109/00048679709062683>
- Rush, A. J., Gullion, C. M., Basco, M. R., Jarrett, R. B., & Trivedi, M. H. (1996). The Inventory of Depressive Symptomatology (IDS): Psychometric properties. *Psychological Medicine*, 26(3), 477–486. <https://doi.org/10.1017/s0033291700035558>
- Santisteban, J. A., Brown, T. G., & Gruber, R. (2018). Association between the Munich Chronotype Questionnaire and Wrist Actigraphy. *Sleep Disorders*, 2018, Article 5646848. <https://doi.org/10.1155/2018/5646848>
- Schuch, F. B., Vancampfort, D., Richards, J., Rosenbaum, S., Ward, P. B., & Stubbs, B. (2016). Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. *Journal of Psychiatric Research*, 77, 42–51. <https://doi.org/10.1016/j.jpsychires.2016.02.023>
- Stamatakis, K. A., Kaplan, G. A., & Roberts, R. E. (2007). Short sleep duration across income, education and race/ethnic groups: Population prevalence and growing disparities over 34 years of follow-up. *Annals of Epidemiology*, 17(12), 948–955. <https://doi.org/10.1016/j.annepidem.2007.07.096>
- Stubbs, B., Koyanagi, A., Schuch, F. B., Firth, J., Rosenbaum, S., Veronese, N., ... Vancampfort, D. (2016). Physical activity and depression: A large cross-sectional, population-based study across 36 low- and middle-income countries. *Acta Psychiatrica Scandinavica*, 134(6), 546–556. <https://doi.org/10.1111/acps.12654>
- Vadnie, C. A., & McClung, C. A. (2017). Circadian rhythm disturbances in mood disorders: Insights into the role of the suprachiasmatic nucleus. *Neural Plasticity*, 2017, Article 1504507. <https://doi.org/10.1155/2017/1504507>
- Vancampfort, D., Stubbs, B., Sienaert, P., Wyckaert, S., De Hert, M., Rosenbaum, S., & Probst, M. (2015). Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. *Journal of Affective Disorders*, 210, 139–150. <https://doi.org/10.1016/j.jad.2016.10.050>
- van Mill, J. G., Hoogendijk, W. J. G., Vogelzangs, N., Van Dyck, R., & Penninx, B. W. J. H. (2010). Insomnia and sleep duration in a large cohort of patients with major depressive disorder and anxiety disorders. *Journal of Clinical Psychiatry*, 71(3), 239–246. <https://doi.org/10.4088/JCP.09m05218gry>
- Verduijn, J., Verhoeven, J. E., Milaneschi, Y., Schoevers, R. A., van Hemert, A. M., Beekman, A. T. F., & Penninx, B. W. J. H. (2017). Reconsidering the prognosis of major depressive disorder across diagnostic boundaries: Full recovery is the exception rather than the rule. *BMC Medicine*, 15, Article 215. <https://doi.org/10.1186/s12916-017-0972-8>
- Wardenaar, K. J., Van Veen, T., Giltay, E. J., Den Hollander-Gijsman, M., Penninx, B. W. J. H., & Zitman, F. G. (2010). The structure and dimensionality of the Inventory of Depressive Symptomatology Self Report (IDS-SR) in patients with depressive disorders and healthy controls. *Journal of Affective Disorders*, 125(1–3), 146–154. <https://doi.org/10.1016/j.jad.2009.12.020>
- Wittchen, H. U. (1994). Reliability and validity studies of the WHO – Composite International Diagnostic Interview (CIDI): A critical review. *Journal of Psychiatric Research*, 28(1), 57–84. [https://doi.org/10.1016/0022-3956\(94\)90036-1](https://doi.org/10.1016/0022-3956(94)90036-1)