

University of Groningen

Return to work after total hip or total knee arthroplasty

Kamp, Tamara

DOI:
[10.33612/diss.864171469](https://doi.org/10.33612/diss.864171469)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2024

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Kamp, T. (2024). *Return to work after total hip or total knee arthroplasty*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.864171469>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Influence of social support on return to work after total hip or total knee arthroplasty: a prospective multicentre cohort study

Authors:

T. Kamp
M. Stevens
J. van Beveren
P.C. Rijk
R.W. Brouwer
S. Bulstra
S. Brouwer

BMJ Open, 2022

5

Abstract

Objectives

There is strong evidence that social support is an important determinant of return to work (RTW). Little is known about the role of social support in RTW after total hip or knee arthroplasty (THA/TKA). Objective was to examine the influence of preoperative and postoperative perceived social support on RTW status 6 months postoperatively.

Design

A prospective multicentre cohort study was conducted.

Setting

Orthopaedic departments of four Dutch medical centres; a tertiary university hospital, two large teaching hospitals, and a general hospital.

Participants

Patients planned to undergo THA/TKA, aged 18-63 and employed preoperatively were included.

Main outcome measures

Questionnaires were filled out preoperatively and 3 and 6 months postoperatively, and included questions to assess patients' perceived social support targeting three sources of social support: from home (friends, family), from work (co-workers, supervisors) and from healthcare (occupational physician, general practitioner, other caregivers). Control variables included age, gender, education, type of arthroplasty and comorbidities. RTW was defined as having fully returned to work 6 months postoperatively. Univariate and multivariate logistic regression analyses were conducted.

Results

Enrolled were 190 patients (n=77 THA, n=113 TKA, median age 56 years, 56% female). The majority returned to work (64%). Preoperatively, social support from the occupational physician was associated with RTW (OR 2.53, 95%CI 1.15-5.54). Postoperatively, social support from the occupational physician (OR 3.04, 95%CI 1.43-6.47) and the supervisor (OR 2.56, 95%CI 1.08-6.06) was associated with RTW.

Conclusions

This study underscores the importance of work-related social support originating from the occupational physician and supervisor in facilitating RTW after primary THA/TKA, both preoperatively and postoperatively. Further research is needed to confirm our results and to understand the facilitating role of social support in RTW, as arthroplasty is being performed on a younger population for whom work participation is critical.

Introduction

Adequate social support is known to have positive effects on health status and health behaviours (1), wellbeing and work participation (2,3). Social support has been defined as the assistance and protection given to an individual (1), which can come from a variety of sources such as friends, family, co-workers, organizations and healthcare professionals. There are different dimensions of social support – instrumental, informational, appraisal and emotional, where the former two are known as instrumental support and the latter two as perceived social support (4–7).

There is strong evidence that perceived social support from home, work and occupational healthcare is an important determinant in the return to work (RTW) process and work disability among a variety of working populations (2,3,8–13). Social support within and outside the workplace has shown to contribute to the RTW process (2,8–12). In a recent systematic review about the influence of social support and social integration on RTW outcomes among individuals with work-related injuries, receiving support from family, regular contact and good communication with the employer, and genuine concern and support from co-workers and supervisors were identified as facilitators of RTW (2). Whereas perceived lack of emotional support, especially lack of on-going support from supervisors, was seen as a barrier to the RTW process (2). Regarding healthcare support, positive RTW recommendations from healthcare professionals showed to be associated with a 60% higher RTW rate in a cohort of 325 patients with low back injury (14). Multiple qualitative studies conducted among different patient groups showed the important role of perceived support from healthcare professionals in the RTW process (15–17). Although these studies emphasize the importance of social support from home, work and healthcare, so far little is known about the role of social support in the RTW process among the rapidly growing patient group undergoing a total hip arthroplasty (THA) or total knee arthroplasty (TKA).

The number of THA and TKA procedures performed annually in the Netherlands continues to increase steadily, most rapidly among working-age patients (18). In 2018, 14,768 primary THAs and 12,777 primary TKAs were performed among working-age adults in the Netherlands, a 56% and 32% increase compared to 2010, respectively (19). Similar trends, with the largest increase among working-age patients, are seen in the United States and other Western countries (20,21). This increase is mainly due to increased prosthetic survivorship and the fact that particularly the severity of the osteoarthritis (OA) and patients' preferences, instead of age, have become a major criteria when deciding whether to undergo THA or TKA (22,23). On the one hand the rise in THA and TKA procedures

performed in younger patients and on the other hand the increase in retirement age results in higher numbers of patients expecting to remain in paid employment after surgery (18,24). Previous studies show that 59-85% of patients return to work within 6 months (25–27), so the absolute number of patients who have not returned to work within 6 months is substantial.

Our previous study, which also used data from the “Work participation In Patients with Osteoarthritis” (WIPO) cohort, showed the importance of psychosocial working conditions on time to RTW after THA or TKA (28). However, little research has been conducted among THA and TKA patients on the effect of social support on RTW outcomes. Some qualitative studies have shown that absence of workplace support by the supervisor was associated with a negative experience of returning to work in arthroplasty patients (29). It was also found that a supportive environment at home and at work, as well as supportive care from healthcare professionals might be helpful in facilitating successful RTW, rehabilitation, and postoperative satisfaction (29–31). No quantitative studies have been found so far that examined the effect of different types of social support on RTW among THA and TKA patients. No evidence exists either on the timing of social support, i.e. the effect of social support immediately before or after surgery compared to later postoperatively. The aim of this study was therefore to investigate the influence of perceived social support from different sources (home, work, healthcare) on RTW status 6 months postoperatively in a sample of THA and TKA patients.

Materials and methods

Design and procedure

A prospective multicentre cohort study was conducted among patients who underwent THA or TKA for primary OA. This study was part of the “Work participation In Patients with Osteoarthritis” cohort (WIPO, Trial-ID NTR3497) (28,32–34). Between March 2012 and July 2014 Patients were recruited at the orthopaedic departments of the following Dutch medical centres: (1) University Medical Center Groningen (tertiary university hospital), (2) Martini Hospital Groningen (large teaching hospital), (3) Medical Center Leeuwarden (large teaching hospital) and (4) Röpcke-Zweers Hospital Hardenberg (general hospital), all in the northern Netherlands. The study was approved by the Medical Ethical Committee of University Medical Center Groningen (METc 2012.153). Patients waiting for THA or TKA were contacted by phone and invited to participate. Preoperative questionnaires were filled in approximately one month before surgery. Postoperative follow-up data, for this study, were collected after 3 and 6

months. If applicable, missing answers were added later to the questionnaire after retrieving them by telephone. Informed consent was assumed as being obtained when patients returned finished questionnaires and thereby granting our request to participate in the study. If patients did not want to participate in the study, they were asked to return a blank questionnaire. Patients were informed of this consent method by mail, in an information letter that also communicated the voluntary nature of the study and the anonymous nature of all the data to be processed. The Medical Ethical Committee specifically approved this consent procedure.

Study population

Patients with primary hip and knee OA undergoing THA or TKA, aged 18-63 and employed preoperatively were included. Excluded were patients who in the previous six months received another joint arthroplasty, THA or TKA due to secondary OA, unicompartimental knee arthroplasty, THA or TKA revision and with inadequate understanding of the Dutch language. A dropout was defined as a patient leaving the study preterm by not filling in the 6-month postoperative questionnaire for any reason.

Measures

Dependent variable

Return to work (yes/no) was measured at the 6-month postoperative follow-up. Patients were asked whether they returned to work, with the following answering possibilities: no return to work, partial return to work, full return to work. RTW was defined as participants who answered that they fully returned to work after surgery, no RTW was defined as participants who answered that they did not or partially return to work.

Independent variables

Perceived social support was measured preoperatively (baseline) and 3 months postoperatively using three questionnaires targeting support from home, work, and healthcare.

Social support from home, i.e. friends and family, was assessed with the Groningen Orthopaedic Social Support Scale (GO-SSS). The GO-SSS consists of 12 questions divided into two subscales: perceived social support (seven items) and instrumental social support (five items). This study focused on the perceived social support subscale. On a Likert scale four answers were possible (never or rarely, occasionally, regularly, often). A sum score was computed, where higher scores

indicated more perceived social support. The GO-SSS showed to be a reliable and valid instrument to assess social support for patients following arthroplasty, with a 0.89 Cronbach alpha for the entire questionnaire and 0.86 internal consistency for the perceived social support (PSS) subscale (35).

Social support from work was assessed with a self-constructed scale focusing on perceived social support. The questionnaire consisted of two questions about perceived support from co-workers and the supervisor. Each item is preceded by the question “How much support did you receive during your period of recuperation from...” with responses on a 1–3 point scale (no support, little support, ample support). Dichotomous variables were computed, distinguishing between no perceived support and perceived support (consisting of little or ample support). The two questions were analysed separately.

Social support from healthcare was measured with a self-constructed scale focusing on perceived social support regarding work. The questionnaire included three questions about perceived support from an occupational physician (OP), a general practitioner (GP) and other caregivers. Each item is preceded by the question “How much support regarding work did you receive during your period of recuperation from...” with responses on a 1–3 point scale (no support, little support, ample support). Dichotomous variables were computed, distinguishing between no perceived support and perceived support (consisting of a little or ample support). The three questions were analysed separately.

Covariates

Data about the following sociodemographic characteristics were collected preoperatively: age (years), gender, education (categorized into elementary, secondary and higher), being breadwinner (yes/no). Disease-related information was gathered by inquiring about type of arthroplasty (THA or TKA), body mass index (BMI) divided into normal (<25 kg/m²) and overweight or obese (>25 kg/m²), and comorbidity measured with a 27-item chronic conditions questionnaire (Statistics Netherlands. Health questionnaire 1989) (36). Amount of comorbidities was divided into none, one or two, or more than two. Data about work-related characteristics included questions about self-employment (yes/no), company size (number of employees: 1-9, 10-99, more than 100), contractual hours (h), working hours (h), type of job (executive/administrative/advisory/management/policy), and type of tasks (physical/mental/combo). Executive jobs cover blue collar workers, i.e. requiring manual labour. Physical work demands were measured by

asking whether patients had to perform physical activities like standing, sitting, walking, kneeling or squatting during work (yes/no).

Statistical analysis

Descriptive statistics – mean (SD), n (%) – were used to describe baseline characteristics of the study population. Univariate and multivariate logistic regression analyses were used to study the prognostic factors for RTW 6 months postoperatively. Separate analyses were conducted for perceived social support measured preoperatively and 3 months postoperatively.

The association between each potential prognostic factor and RTW was univariately assessed. All prognostic factors with a p-value ≤ 0.20 in the univariate analyses were included in the multivariate regression analyses (37), after checking for multicollinearity. Variables were omitted by backward selection, depending on their level of statistical significance ($P < 0.05$). Control variables for the analyses included sex, age, education, type of surgery, comorbidities, and work tasks (38–41). Control variables were based on previous literature and were defined a priori. Sensitivity analyses were conducted for THA and TKA groups separately, since previous literature suggests that postoperative recovery and RTW differs between these groups (42,43). Odds ratios were calculated, including 95% confidence intervals (CI). A non-response analysis was performed. Statistical analyses were performed with IBM Statistical Package for the Social Sciences (SPSS) version 25.0 and Mplus version 7.1.

Patient and public involvement statement

Neither patients nor the public were involved in the design, conduct, reporting or dissemination plans of our research.




Fig. 1 Flowchart study enrolment and follow-up

Results

From the 311 patients who had undergone a primary THA or TKA, 190 (n=77 THA, n=113 TKA) were included in the study. Figure 1 is a flowchart showing the total number of patients at baseline and the drop-outs to follow-up. The characteristics of the study sample are presented in table 1 and online supplemental Table 1. Median age was 56 years (interquartile range (IQR) 52-60 years). The sample consisted of 84 (44%) men and 106 (56%) women, 77 (41%) THA patients and 113 (59%) TKA patients. For educational level, 33% had completed elementary school, 44% secondary school and 21% higher education. BMI of 77% was above 25 kg/m² and 46% had two or more comorbidities. Patients worked on average 32 hours. Our cohort had mostly executive jobs (55%; blue collar). A combination of physically and mentally challenging tasks was performed by 39% of patients; the remaining patients were divided equally into performing either physical or mental work tasks. Work demands of the majority included sitting and/or walking, and a quarter of the patients had to perform kneeling or squatting work demands.

The majority of patients returned to work (64%) by 6 months post-surgery. To correct for the drop-out rate during follow-up we conducted a non-response analysis, which showed no significant differences on baseline characteristics or independent variables.

Table 1: Baseline study population characteristics.

Variables	Total (N=190)
Age, median (IQR)	56 (52 – 60)
Male/female, n (%)	84 (44) / 106 (56)
Highest educational level (n (%))	
- Lower (elementary school, vocational education)	62 (33)
- Secondary (high school, intermediate vocational education)	84 (44)
- Higher (higher professional education university)	39 (21)
Wage earner, n (%)	106 (56)
THA/TKA, n (%)	77 (41) / 113 (59)
BMI (kg/m ²), n (%)	
- <25	40 (21)
- >25	147 (77)
Number of comorbidities, n (%)	
- No	19 (10)
- One or two	62 (33)
- More than two	88 (46)
Self-employed, n (%)	22 (12)
Company size (number of employees), n (%)	
- 1-9	28 (15)
- 10-99	50 (26)
- >100	112 (59)
Contractual hours (median, IQR)	32 (21 to 37)
Working hours (median, IQR)	32 (22 to 40)
Job type, n (%)	
- Executive	105 (55)
- Administrative	22 (12)
- Advisory	11 (6)
- Management	27 (14)
- Policy	23 (12)
Work tasks n (%)	

Table 1: *Continued.*

Variables	Total (N=190)
- Physical	57 (30)
- Mental	57 (30)
- Both	74 (39)
Work demands, n (%)	
- Standing	100 (47)
- Sitting	107 (56)
- Walking	104 (55)
- Kneeling or squatting	52 (27)

All numbers are represented as median with interquartile range (IQR), or numbers (n) and percentages (%).

Univariate and multivariate logistic regression analyses

In the *preoperative* univariate analyses, social support from the OP was the only variable below the cut-off value of $p < 0.2$, therefore no multivariate analyses were performed. Preoperative social support from the OP was univariately significantly associated with RTW (OR 2.53, 95%CI 1.15–5.54; table 2). In the *postoperative* univariate analyses social support from the supervisor, the OP, the GP and other caregivers were below the cut-off value of $p < 0.2$ and were therefore used in the multivariate analyses. In the multivariate model perceived social support from the OP (OR 3.04, 95%CI 1.43-6.47) and from the supervisor (OR 2.56, 95%CI 1.08-6.06) showed statistically significant associations with RTW. The odds of an individual having returned to work 6 months post-surgery increased by 3.04 and 2.56 for those patients who perceived social support from the OP and from the supervisor, respectively (table 2).

Table 2: Preoperative and three months postoperative univariate and multivariate logistic regression analyses of perceived social support variables on return to work (RTW) status

Variables	Univariate			Multivariate		
	OR	P	95% CI	OR	P	95% CI
Preoperative						
Support from home	1.04	0.40	0.95 to 1.14			
Support from co-workers (ref=no)	1.26	0.64	0.48 to 3.31			
Support from supervisor (ref=no)	1.57	0.30	0.68 to 3.62			
Support from OP (ref=no)	2.53	0.02*	1.15 to 5.54			
Support from GP (ref=no)	1.46	0.30	0.71 to 2.98			
Support from other caregivers (ref=no)	1.24	0.57	0.59 to 2.63			
Three months postoperative						
Support from home	1.01	0.92	0.92 to 1.10			
Support from co-workers (ref=no)	1.28	0.56	0.56 to 2.93			
Support from supervisor (ref=no)	2.71	0.02†	1.18 to 6.23	2.56	0.03*	1.08 to 6.06
Support from OP (ref=no)	3.17	0.00†	1.51 to 6.66	3.04	0.00*	1.43 to 6.47
Support from GP (ref=no)	2.51	0.02†	1.19 to 5.29			
Support from other caregivers (ref=no)	1.64	0.17†	0.81 to 3.32			

Adjusted for sex, age, education, comorbidities, type of surgery and work tasks; † $p < 0.2$; * $p < 0.05$; OR, odds ratio; CI, confidence intervals; OP, occupational physician; GP, general practitioner.

Sensitivity analyses

Analysing the THA and TKA groups separately, the *preoperative* multivariate model showed no association between social support and RTW in both subgroups (table 3). The *postoperative* multivariate model of THA patients showed that perceived social support from the supervisor was significantly associated with RTW (OR 1.90, 95%CI 1.12–21.53; table 3). The *postoperative* multivariate model of TKA patients showed a significant association between perceived social support from the OP and RTW (OR 5.14, 95%CI 1.84–14.36; table 3).

Table 3: Preoperative and three months postoperative univariate and multivariate logistic regression analyses of perceived social support variables on return to work (RTW) status among subsamples of THA and TKA patients

Variables	Univariate			Multivariate		
	OR	95% CI	P	OR	95% CI	P
Preoperative						
THA (n=77)						
Support from home	1.03	0.88 to 1.20	0.76			
Support from co-workers (ref=no)	2.04	0.35 to 11.90	0.43			
Support from supervisor (ref=no)	2.79	0.55 to 14.07	0.21			
Support from OP (ref=no)	3.33	0.81 to 13.69	0.10†			
Support from GP (ref=no)	1.15	0.34 to 3.90	0.83			
Support from other caregivers (ref=no)	0.67	0.19 – 2.33	0.53			
TKA (n=113)						
Support from home	1.05	0.94 to 1.18	0.38			
Support from co-workers (ref=no)	1.10	0.32 to 3.76	0.88			
Support from supervisor (ref=no)	1.25	0.45 to 3.48	0.67			
Support from OP (ref=no)	2.06	0.76 to 5.57	0.15†			
Support from GP (ref=no)	1.64	0.64 to 4.21	0.31			
Support from other caregivers (ref=no)	1.64	0.60 to 4.49	0.33			
Three months postoperative						
THA (n=77)						
Support from home	1.09	0.93 to 1.27	0.29			

Table 3: Continued.

Variables	Univariate			Multivariate		
	OR	95% CI	P	OR	95% CI	P
Support from co-workers (ref=no)	3.13	0.55 to 17.80	0.20			
Support from supervisor (ref=no)	1.90	1.12 to 21.53	0.04†	1.90	1.12 to 21.53	0.04*
Support from OP (ref=no)	1.85	0.51 to 6.81	0.35			
Support from GP (ref=no)	3.24	0.77 to 13.61	0.11†			
Support from other caregivers (ref=no)	0.65	0.18 to 2.39	0.52			
TKA(n=113)						
Support from home	0.97	0.87 to 1.08	0.60			
Support from co-workers (ref=no)	1.26	0.46 to 3.43	0.66			
Support from supervisor (ref=no)	2.65	0.87 to 8.07	0.09†			
Support from OP (ref=no)	5.14	1.84 to 14.36	0.00†	5.14	1.84 to 14.36	0.00*
Support from GP (ref=no)	2.40	0.94 to 6.11	0.07†			
Support from other caregivers (ref=no)	2.32	0.91 to 5.90	0.08†			

Adjusted for sex, age education, comorbidities, and work tasks; † $p < 0.2$; * $p < 0.05$; OR, odds ratio; CI, confidence intervals; OP, occupational physician; GP, general practitioner.

Discussion

This study aimed to investigate the influence of preoperative and postoperative perceived social support from home, work and healthcare on RTW status 6 months postoperatively in a sample of THA and TKA patients. We found that patients who perceived social support from the OP preoperatively had 2.5 times higher odds of RTW within 6 months postoperatively compared to patients who perceived no support. Patients who perceived social support from the OP and from the supervisor 3 months postoperatively had 3.0 and 2.6 times higher odds of RTW, respectively. These results imply the important role of workplace support in the RTW process, as both the OP and supervisor are linked to the workplace.

In our study the majority of patients (64%) returned to work within 6 months postoperatively, which is in line with previous studies (25–27). Our findings that perceived social support from the OP is important, both preoperatively and postoperatively, is in line with previous quantitative studies on social support from the OP in other populations (13,14,17). In qualitative studies among THA and TKA patients, employers and clinicians also indicated the added value of OPs, especially if there already was contact before surgery (29,44).

Our findings that social support from the supervisor was associated with RTW is also in line with previous studies conducted among other population groups (2,45,46). Supervisors play a considerable role in initiating effective support strategies (47–49): they are expected to communicate the process of RTW with the employee and the OP and implement accommodations, both in agreement with the OP (2,11). In our multivariate analyses, we only found an association between postoperative and not preoperative social support from the supervisor and RTW, leaving questions about optimal timing. An explanation might be that the supervisor is better able to perform specific actions postoperatively to facilitate RTW.

In contrast to previous studies, we did not find an association between social support from home or co-workers and RTW in our study population. A possible explanation for this absence in our study might relate to the duration of sickness absence: other studies that found an association between social support from home or co-workers and RTW were mainly conducted among population groups with long-term absence (>6 months) (3,13), whereas a THA or TKA often leads to a short-term work absence (<3-6 months) for most patients. Disease chronicity and long-term absence may influence the necessity and contributing value of social support from home and co-workers for RTW outcomes.

In our study we did not find an effect of perceived social support from other caregivers (e.g. physiotherapists) on RTW. This might be because we did not

further specify the question and patients could have experienced it as implicit. The role of social support from a physiotherapist on RTW warrants further research, since our particular subsample has frequent contact with these specific healthcare professionals. Value of a physiotherapist is illustrated by Lysaght et al., who reported in their qualitative research that half of the workers experienced support by a physiotherapist (11). More research is needed to evaluate the role of physiotherapists and their contribution to the RTW process.

Our sensitivity analyses showed some differences in factors associated with RTW between THA and TKA patients. Postoperative perceived social support from the supervisor was associated with RTW of THA patients and postoperative perceived social support from the OP was associated with RTW of TKA patients. This dissimilarity in findings may be explained by differences in the rehabilitation process. It is known that for THA patients rehabilitation is easier than for TKA patients (42,43). However, it must be kept in mind that the wide 95% CI indicated our sample size is too small. These results need to be replicated with a larger sample size before definitive conclusions can be drawn.

Finally, our non-response analyses did not show significant differences on baseline characteristics or independent variables. However, it might be that non-response could partly be explained by unfavourable return to work outcomes.

Strengths & limitations

An important strength of this study is its prospective multicentre design with a relatively large number of patients and a follow-up of 6 months. Another strength is the representative sample of patients and therefore the generalizability of the outcomes. We provided multivariate analysis on three different sources of social support, plus investigated both preoperative and postoperative data, in contrast to previous research on social support among other patient groups (2). This study does have some limitations. Due to limited power our study only focused on preoperative and postoperative data separately. The sample sizes of our subgroups (THA and TKA) in the sensitivity analyses lacked power to draw definitive conclusions, and we only focused on the first time workers fully returned to work. Future research should also include sustainable RTW to assess the impact of social support on these RTW trajectories. Finally, another limitation were the self-reported measurements, which are generally susceptible to the effects of reporting bias.

Implications

Changing workforce dynamics and trends towards THA or TKA surgery among working-age employees propel an urgent need to understand the facilitators and

barriers for RTW, besides those of pain and function (33). There are still many uncertainties about the potential influence of psychosocial work factors (including social support), timing of interventions designed to facilitate RTW, and engagement of clinicians and employers as key actors in the RTW process.

To our knowledge, this is the first quantitative study to examine the role of social support among this specific population. The differences in predicting factors between THA and TKA patients might imply a need for group-specific approaches. Further research on social support is needed to confirm our results and to understand the facilitating role of social support on RTW. The optimal timing to implement contact, i.e. social support, the course (change over time) of social support from different sources and their effect on RTW should also be investigated. Therefore, studies among THA and TKA patients specifically focused at social support, and using validated questionnaires to measure social support from different sources (50,51), would be very valuable.

Conclusion

This study showed that, in particular, perceived social support from OPs and supervisors may predict RTW after THA and TKA. Both preoperative and postoperative social support were associated with RTW, which may suggest that perceived work-related social support from OPs and supervisors are important factors over an extended period of time. Some differences in factors were found between THA and TKA patients, where postoperative social support from the supervisor predicted RTW of THA patients and postoperative social support from the OP predicted RTW of TKA patients. Further research on the role of social support in returning to work after THA and TKA is needed, as arthroplasty is being performed on an increasingly younger population for whom work participation is of critical importance.

References

1. Langford CPH, Bowsher J, Maloney JP, Lillis PP. Social support: A conceptual analysis. *J Adv Nurs*. 1997;25(1):95–100.
2. White C, Green RA, Ferguson S, Anderson SL, Howe C, Sun J, et al. The Influence of Social Support and Social Integration Factors on Return to Work Outcomes for Individuals with Work-Related Injuries: A Systematic Review. Vol. 29, *Journal of Occupational Rehabilitation*. Springer New York LLC; 2019. p. 636–59.
3. Englund ACD, Rydström I, Dellve L, Ahlstrom L. Social support outside work and return to work among women on long-term sick leave working within human service organizations. *Appl Nurs Res*. 2016 May 1;30:187–93.
4. House JS. *Work Stress and Social Support* (Addison-Wesley series on occupational stress): James S. House: 9780201031010: Amazon.com: Books. Addison-Wesley Educational Publishers Inc; 1981.
5. Krause N. Understanding the stress process: linking social support with locus of control beliefs. *J Gerontol*. 1987 Nov;42(6):589–93.
6. Weinert C, Brandt PA. Measuring Social Support with the Personal Resource Questionnaire. *West J Nurs Res*. 1987 Nov;9(4):589–602.
7. Veiel HOF. Dimensions of social support: a conceptual framework for research. *Soc Psychiatry*. 1985 Dec;20(4):156–62.
8. Brouwer S, Krol B, Reneman MF, Bültmann U, Franche R-L, van der Klink JLL, et al. Behavioral determinants as predictors of return to work after long-term sickness absence: an application of the theory of planned behavior. *J Occup Rehabil*. 2009 Jun;19(2):166–74.
9. Brouwer S, Reneman MF, Bültmann U, Van Der Klink JLL, Groothoff JW. A prospective study of return to work across health conditions: Perceived work attitude, self-efficacy and perceived social support. *J Occup Rehabil*. 2010 Mar;20(1):104–12.
10. Tjulin Å, MacEachen E, Ekberg K. Exploring workplace actors experiences of the social organization of return-to-work. *J Occup Rehabil*. 2010 Sep;20(3):311–21.
11. Lysaght RM, Larmour-Trode S. An exploration of social support as a factor in the return-to-work process. Vol. 30, *Work*. IOS Press; 2008.
12. Lau B, Shiryayeva O, Ruud T, Victor M. What are they returning to? Psychosocial work environment as a predictor of returning to work among employees in treatment for common mental disorders: A prospective observational pre–post study. Kamperman AM, editor. *PLoS One*. 2019 Apr 24;14(4):e0215354.
13. Snippen NC, De Vries HJ, Van Der Burg-Vermeulen SJ, Hagedoorn M, Brouwer S. Influence of significant others on work participation of individuals with chronic diseases: A systematic review. Vol. 9, *BMJ Open*. BMJ Publishing Group; 2019. p. e021742.
14. Dasinger LK, Krause N, Thompson PJ, Brand RJ, Rudolph L. Doctor proactive communication, return-to-work recommendation, and duration of disability after a workers' compensation low back injury. *J Occup Environ Med*. 2001 Jun;43(6):515–25.
15. van Velzen JM, van Bennekom CAM, van Dormolen M, Sluiter JK, Frings-Dresen MHW. Factors influencing return to work experienced by people with acquired brain injury: a qualitative research study. *Disabil Rehabil*. 2011;33(23–24):2237–46.
16. De Vries G, Koeter MWJ, Nabitz U, Hees HL, Schene AH. Return to work after sick leave due to depression; A conceptual analysis based on perspectives of patients, supervisors and occupational physicians. *J Affect Disord*. 2012 Feb;136(3):1017–26.

17. Kosny A, Franche R-L, Pole J, Krause N, Côté P, Mustard C. Early healthcare provider communication with patients and their workplace following a lost-time claim for an occupational musculoskeletal injury. *J Occup Rehabil*. 2006 Mar;16(1):27–39.
18. Otten R, van Roermund PM, Picavet HSJ. Trends in the number of knee and hip arthroplasties: considerably more knee and hip prostheses due to osteoarthritis in 2030. *Ned Tijdschr Geneesk*. 2010;154:A1534.
19. Reus IMA de, Spekenbrink-Spooren A, van Steenberghe D ir. LN, Denissen GAW, Rijnsburger E, Togh DCR van der. LROI annual report 2019. 2019.
20. Singh JA, Yu S, Chen L, Cleveland JD. Rates of Total Joint Replacement in the United States: Future Projections to 2020–2040 Using the National Inpatient Sample. *J Rheumatol*. 2020;11:2020.
21. Leitner L, Türk S, Heidinger M, Stöckl B, Posch F, Maurer-Ertl W, et al. Trends and Economic Impact of Hip and Knee Arthroplasty in Central Europe: Findings from the Austrian National Database. *Sci Rep*. 2018 Dec 1;8(1):4707.
22. Ferguson RJ, Palmer AJ, Taylor A, Porter ML, Malchau H, Glyn-Jones S. Hip replacement. Vol. 392, *The Lancet*. Lancet Publishing Group; 2018. p. 1662–71.
23. Price AJ, Alvand A, Troelsen A, Katz JN, Hooper G, Gray A, et al. Knee replacement. Vol. 392, *The Lancet*. Lancet Publishing Group; 2018. p. 1672–82.
24. Kurtz SM, Lau E, Ong K, Zhao K, Kelly M, Bozic KJ. Future young patient demand for primary and revision joint replacement: National projections from 2010 to 2030. Springer New York; Oct 10, 2009 p. 2606–12.
25. Kievit AJ, van Geenen RCI, Kuijer PPFM, Pahlplatz TMJ, Blankevoort L, Schafroth MU. Total Knee Arthroplasty and the Unforeseen Impact on Return to Work: A Cross-Sectional Multicenter Survey. *J Arthroplasty*. 2014 Jun;29(6):1163–8.
26. Sankar A, Davis AM, Palaganas MP, Beaton DE, Badley EM, Gignac MA. Return to work and workplace activity limitations following total hip or knee replacement. *Osteoarthritis Cartil*. 2013 Oct;21(10):1485–93.
27. Kuijer PPFM, Kievit AJ, Pahlplatz TMJ, Hooiveld T, Hoozemans MJM, Blankevoort L, et al. Which patients do not return to work after total knee arthroplasty? *Rheumatol Int*. 2016 Sep;36(9):1249–54.
28. Kamp T, Brouwer S, Hylkema TH, van Beveren J, Rijk PC, Brouwer RW, et al. Psychosocial Working Conditions Play an Important Role in the Return-to-Work Process After Total Knee and Hip Arthroplasty. *J Occup Rehabil*. 2021 Sep 28;1–11.
29. Bardgett M, Lally J, Malviya A, Kleim B, Deehan D. Patient-reported factors influencing return to work after joint replacement. *Occup Med (Lond)*. 2016 Apr;66(3):215–21.
30. Marcinkowski K, Wong VG, Dignam D. Getting back to the future: A grounded theory study of the patient perspective of total knee joint arthroplasty. *Orthop Nurs*. 2001;24(3):202–9.
31. Malviya A, Wilson G, Kleim B, Kurtz SM, Deehan D. Factors influencing return to work after hip and knee replacement. *Occup Med (Chic Ill)*. 2014 Sep 1;64(6):402–9.
32. Hylkema TH, Stevens M, Van Beveren J, Rijk PC, van Jonbergen HP, Brouwer RW, et al. Preoperative characteristics of working-age patients undergoing total knee arthroplasty. *PLoS One*. 2017;12(8):e0183550.
33. Hylkema TH, Brouwer S, Stewart RE, van Beveren J, Rijk PC, Brouwer RW, et al. Two-year recovery courses of physical and mental impairments, activity limitations, and participation restrictions after total knee arthroplasty among working-age patients. *Disabil Rehabil*. 2020 May 22;1–10.
34. Hylkema TH, Stevens M, Beveren J van, Rijk PC, Brouwer RW, Bulstra SK, et al. Recovery Courses of Patients Who Return to Work by 3, 6 or 12 Months After Total Knee Arthroplasty. *J Occup Rehabil*. 2021 Sep 1;31(3):627.

35. Van Den Akker-Scheek I, Stevens M, Spriensma A, Van Horn JR. Groningen Orthopaedic Social Support Scale: Validity and reliability. *J Adv Nurs*. 2004 Jul;47(1):57–63.
36. Statistics Netherlands. Health questionnaire 1989. Voorburg/Heerlen. 1989.
37. Hosmer DW, Lemeshow S, Sturdivant RX. *Applied Logistic Regression*. Third edit. Wiley; 2013. 1–510 p.
38. Tilbury C, Schaasberg W, Plevier JWM, Fiocco M, Nelissen RGHH, Vliet Vlieland TPM. Return to work after total hip and knee arthroplasty: a systematic review. *Rheumatology*. 2014 Mar 1;53(3):512–25.
39. Hoorntje A, Janssen KY, Bolder SBT, Koenraadt KLM, Daams JG, Blankevoort L, et al. The Effect of Total Hip Arthroplasty on Sports and Work Participation: A Systematic Review and Meta-Analysis. *Sport Med*. 2018;
40. Van Leemput D, Neiryneck J, Berger P, Vandenneucker H. Return to Work after Primary Total Knee Arthroplasty under the Age of 65 Years: A Systematic Review. *J Knee Surg*. 2021;
41. Mj Pahlplatz T, Schafroth MU, Kuijer PP. Patient-related and work-related factors play an important role in return to work after total knee arthroplasty: a systematic review. *JISAKOS*. 2017;2:127–32.
42. Bourne RB, Chesworth B, Davis A, Mahomed N, Charron K. Comparing patient outcomes after THA and TKA: Is there a difference? In: *Clinical Orthopaedics and Related Research*. Springer New York; 2010. p. 542–6.
43. O'Brien S, Bennett D, Doran E, Beverland DE. Comparison of hip and knee arthroplasty outcomes at early and intermediate follow-up. *Orthopedics*. 2009 Mar;32(3):168.
44. Nouri F, Coole C, Narayanasamy M, Baker P, Khan S, Drummond A. Managing Employees Undergoing Total Hip and Knee Replacement: Experiences of Workplace Representatives. *J Occup Rehabil*. 2018 Aug 21;
45. Netterstrøm B, Eller NH, Borritz M. Prognostic Factors of Returning to Work after Sick Leave due to Work-Related Common Mental Disorders: A One-And Three-Year Follow-Up Study. *Biomed Res Int*. 2015;2015.
46. Islam T, Dahlui M, Majid HA, Nahar AM, Mohd Taib NA, Su TT. Factors associated with return to work of breast cancer survivors: A systematic review. In: *BMC Public Health*. BioMed Central Ltd.; 2014.
47. Franche R-L, Baril R, Shaw W, Nicholas M, Loisel P. Workplace-based return-to-work interventions: optimizing the role of stakeholders in implementation and research. *J Occup Rehabil*. 2005 Dec;15(4):525–42.
48. MacEachen E, Clarke J, Franche RL, Irvin E, Cullen K, Frank J, et al. Systematic review of the qualitative literature on return to work after injury. Vol. 32, *Scandinavian Journal of Work, Environment and Health*. Nordic Association of Occupational Safety and Health; 2006. p. 257–69.
49. Wilkie R, Pransky G. Improving work participation for adults with musculoskeletal conditions. *Best Pract Res Clin Rheumatol*. 2012 Oct;26(5):733–42.
50. Karasek R, Brisson C, Kawakami N, Houtman I, Bongers P, Amick B. The Job Content Questionnaire (JCQ): an instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol*. 1998;3(4):322–55.
51. WÄnnström I, Peterson U, Åsberg M, Nygren Å, Gustavsson JP. Psychometric properties of scales in the General Nordic Questionnaire for Psychological and Social Factors at Work (QPS): confirmatory factor analysis and prediction of certified long-term sickness absence. *Scand J Psychol*. 2009 Jun;50(3):231–44.

Supplementary table 1: Descriptive information on social support

Variables*	<i>Preoperative</i>	<i>Postoperative (3 months)</i>
Total (N=190)		
Support from home, median (IQR)	25 (21 – 27)	25 (22 – 28)
Support from co-workers, n (%)	139 (73)	139 (73)
Support from supervisor, n (%)	119 (63)	115 (61)
Support from OP, n (%)	57 (30)	78 (41)
Support from GP, n (%)	73 (38)	75 (40)
Support from other caregivers, n (%)	73 (38)	66 (35)

**All numbers are represented as median with interquartile range (IQR), or numbers (n) and percentages (%).*

