

University of Groningen

Swimming exercise and clove oil can improve memory by molecular responses modification and reduce dark cells in rat model of Alzheimer's disease

Karaji, Z Gorgin; Fathi, M; Mirnasori, R; van der Zee, E A

Published in:
Experimental Gerontology

DOI:
[10.1016/j.exger.2023.112192](https://doi.org/10.1016/j.exger.2023.112192)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Karaji, Z. G., Fathi, M., Mirnasori, R., & van der Zee, E. A. (2023). Swimming exercise and clove oil can improve memory by molecular responses modification and reduce dark cells in rat model of Alzheimer's disease. *Experimental Gerontology*, 177, 112192. <https://doi.org/10.1016/j.exger.2023.112192>

Copyright


Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Swimming exercise and clove oil can improve memory by molecular responses modification and reduce dark cells in rat model of Alzheimer's disease

Z. Gorgin Karaji^a, M. Fathi^{a,*}, R. Mirnasori^a, E.A. van der Zee^b

^a Dept. of Sport Sciences, Faculty of Human Sciences, Lorestan University, Khorramabad, Iran

^b Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, Groningen, Netherlands

ARTICLE INFO

Section Editor: Zsolt Radak

Keywords:

Alzheimer
Swimming
Memory
 $\alpha 7$ nAChR

ABSTRACT

Alzheimer's disease (AD) is marked by reduced acetylcholine receptor (AChR) density and an increase in nucleotide oligomerization domain (NOD)-like receptors NLR family, pyrin domain containing 1 (NLRP1). We examined the effect of swimming and consumption of clove supplements on memory, dark cells, and $\alpha 7$ nAChR and NLRP1 mRNA and protein expression in the hippocampus of the rat model of AD. Forty-eight rats were divided into six groups: sham (sh), healthy-control (HC), Alzheimer (-control) (AC), -training (AT), -training-supplement (ATS), and -supplement (AS). Alzheimer was induced by injection of amyloid β_{1-42} ($A\beta_{1-42}$). Swimming exercise protocol (30 min) and gavaging clove supplement (0.1 mg/kg) were administered daily for three weeks. The results indicated that in response to AD, $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) mRNA and protein rate ($p = 0.001$) and memory ($p = 0.003$) were significantly decreased. In contrast, NLRP1 mRNA and protein rate ($p = 0.001$) and dark cells ($p = 0.001$) were significantly increased. This is while exercise and clove supplementation improved Alzheimer-induced changes in $\alpha 7$ nAChR, NLRP1, memory, and dark cells ($p < 0/05$). The present study indicated that exercising and consuming clove supplementation could improve memory by increasing $\alpha 7$ nAChR and decreasing NLRP1 and dark cells.

1. Introduction

AD is the common age-related cause of dementia (Jia et al., 2019), recognized by dark cells, cognitive dysfunction, inflammatory responses (Saresella et al., 2016), and memory loss (Jia et al., 2019). AD results in many economic, medicinal, and social problems. Despite the expanded studies in improving remedial drugs, treatment strategies remain insufficient (Esfandiarei et al., 2019). There is growing support reporting that nutrition and exercise can positively affect neurodegenerative diseases (Alkhouli et al., 2019). The Clinical researches demonstrated that physical activity could limit the beginning or delay the progression of age-related human diseases such as AD (Esfandiarei et al., 2019; Alkhouli et al., 2019). Physical activity and a healthy diet are harmless, cheap, and helpful interventions that can improve living quality in patients with AD (Esfandiarei et al., 2019). Exercise training may affect cognitive function (Bernardo et al., 2020), the expression of neurotrophic factors (Jiang et al., 2014), and the levels of inflammatory responses by affecting the cholinergic anti-inflammatory pathway

(Cechella et al., 2014).

The importance of eugenol-rich diets to manage neuronal cells of anti-inflammatory responses was reported (Taher et al., 2015). Previous data have shown that eugenol, as a significant part of clove oil, is well-known to involve glutamatergic and cholinergic systems (Cortés-Rojas et al., 2014; Wie et al., 1997). Eugenol also can promote anti-inflammatory properties (Abbasi et al., 2019). Besides, clove oil has shown links with anti-oxidant properties and neuroprotective effects tightly (Baghshahi et al., 2014). It may improve cognitive performance (Han and Parker, 2017) and reduce AChE activity and oxidative damage. Additionally, clove oil can delay the progression of AD (Kumar et al., 2016).

Inflammatory mediators are central to the neuroinflammation in AD (Saresella et al., 2016). Stimulation of the $\alpha 7$ nAChR, as ligand-gated ion channels (Ni et al., 2013; Parri et al., 2011) and a member of the cholinergic anti-inflammatory pathway (Neumann et al., 2015), can reduce the inflammatory responses (Huang et al., 2019). On the other hand, high levels of the NLRP1 inflammasome were found in the brain

* Corresponding author.

E-mail address: fathi.m@lu.ac.ir (M. Fathi).

<https://doi.org/10.1016/j.exger.2023.112192>

Received 10 December 2022; Received in revised form 24 April 2023; Accepted 26 April 2023

Available online 29 April 2023

0531-5565/© 2023 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

(Tan et al., 2014). NLRP1, as a component of the inflammasome, can activate caspase-1 to induce the inflammatory response and cell death (Tan et al., 2014). Besides, inhibition of the NLRP1 inflammasome could decrease the innate immune response and improve cognitive deficits (Tan et al., 2014). There is growing evidence that NLRP1 inflammasome increase in AD (Tan et al., 2014), while a loss of $\alpha 7$ nAChR has been reported in the brain of Alzheimer's patients (Ni et al., 2013). Activation of $\alpha 7$ nAChRs decreases the inflammation complex level (such as NLRPs) and reduces the pro-inflammatory cytokines secretion (Abbasi et al., 2019; Kelley et al., 2019).

According to the mentioned studies on the effect of physical activity and also clove oil on the physiological and cognitive factors of the brain, the present study aimed to investigate if clove oil treatment and exercise, either alone or together, could be considered a treatment to improve the anti-inflammatory pathway and subsequently reduce AD progression. In this study, we examined the efficacy of clove oil and/or exercise on spatial memory, dark cells, and $\alpha 7$ nAChR and NLRP1 mRNA and protein levels in the brain of the rat model of AD.

2. Results

2.1. Confirmation of the Alzheimer's model

10 days after the induction of AD pathology, obtaining the Alzheimer's model was confirmed by the immunofluorescence study (Fig. 1). There was a significant difference between the mean expression of (amyloid β) A β plaques in HC (256.7 ± 34.80) and AC ($12,949 \pm 441.8$) groups ($p = 0.001$, $t = 28.64$).

Swimming exercise and clove oil supplementation improved spatial learning and spatial memory in rat model of AD in the Morris water maze test.

Memory and learning impairments are consequences of AD. We used the MWM to evaluate the effect of exercise and clove oil on learning and spatial memory in our rat model of AD. A two-way ANOVA was used to investigate the difference between groups and days on escape latency time. These results showed that the difference between days ($F = 28.079$, $p = 0.000$, $df = 3$), and groups ($F = 63.163$, $p = 0.000$, $df = 6$) were significant, but no interaction between groups*day ($F = 0.478$, $p = 0.957$, $df = 18$).

According to Fig. 2A, in four days of the learning test, a significant difference in spatial learning ability was observed among the studied groups ($F = 13.873$, $p = 0.001$). Also, the percentage of spatial learning ability of the AC group was significantly lower than the other groups ($p < 0.05$). There was no difference in pair-wise comparison of Sh, HC, AT,

ATS, and AS groups in four days ($p \geq 0.05$).

The mean time elapsed in the quadrant containing the hidden platform was considered spatial memory (Fig. 2B). A significant difference in spatial memory ability was observed among the study groups ($F = 16.699$, $p = 0.001$). Also, the mean time elapsed in the quadrant containing the hidden platform of the AC group was significantly lower than HC ($p = 0.003$) and, in the ATS group, it was significantly higher than that in the AT group ($p = 0.022$). Although time elapsed in the quadrant containing the hidden platform of the ATS group was higher than the AS group, there was no significant difference between the ATS and AS groups ($p = 0.321$). Moreover, there was no significant difference between the HC group with Sh ($p = 1.00$), AT ($p = 0.682$), ATS ($p = 0.325$), and AS ($p = 1.00$) groups. It means that although AD reduces spatial memory, exercise, clove oil, or both can compensate for the behavioral deficit. According to the results of this study swimming and the consumption of clove oil have a desirable effect on spatial memory in the Rat model of AD.

2.2. Swimming exercise and clove oil supplementation decrease the degree of AD-induced dark cells percentage in the hippocampus

The percentage of dark cells in the CA1 region of the hippocampus in the Rat model of AD in different groups is presented in Fig. 3. Results showed there was no difference between dark cells percentage in the hippocampus in the HC and Sh groups (1.00).

AD induction increased the percentage of dark cells in the CA1 region of the hippocampus of the AC group compared to HC ($p = 0.001$). In contrast, there was no difference between percentage of dark cells in the AT group ($p = 0.491$) and the AS group ($p = 0.491$) compared to the AC group. However, swimming and the consumption of clove oil (in the ATS group) ($p = 0.039$) significantly reduced the percentage of dark cells compared to the AC group. The percentage of dark cells in AT ($p = 0.009$) remained significantly higher than HC group but not in AS ($p = 0.056$) and ATS ($p = 0.221$) groups. Also, there was no difference between the percentage of dark cells in the ATS group compared to AT ($p = 0.169$), and the AS ($p = 0.491$) groups.

2.3. Swimming and clove oil improved the expression of reduced $\alpha 7$ nAChR mRNA and protein levels in the hippocampus of the Rat model of AD

Inflammation is involved in AD. It has been reported that $\alpha 7$ nAChR is affected by an inflammatory response. In this study, we examined if the expression of $\alpha 7$ nAChR mRNA and protein levels are affected by




Fig. 1. Photomicrographs of immunofluorescent staining of A β plaques in the CA1 region of the hippocampus (control and Alzheimer's rats). Green structures indicate the accumulation of A β plaques. Magnification: 100 \times , 100 μ m. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. Effect of swimming exercise and clove oil on spatial learning (A) and spatial memory (B) in the MWM test. The groups consisted of healthy-control (HC), Alzheimer control (AC), Alzheimer-training (AT), Alzheimer-supplement (AS), Alzheimer-training-supplement (ATS), and Sham (Sh). Data are expressed as the mean \pm SD. Significant level: $p < 0.05$. Bar with different letters indicates a significant difference (ANOVA and subsequent Tukey's HSD, $p < 0.05$).

Alzheimer, exercise, and clove oil. Photomicrographs of immunofluorescence staining and the mean expression of $\alpha 7nAChR$ mRNA and rate of protein are shown in Fig. 4. There is a significant difference in the expression of $\alpha 7nAChR$ mRNA and its protein levels observed among the studied groups ($F = 48.766$, $p = 0.001$) and ($F = 172.414$, $p = 0.001$), respectively. There was no difference between the mean expression of $\alpha 7nAChR$ mRNA and protein in the HC and Sh groups ($p = 0.964$, $p = 1.00$). The mean expression of $\alpha 7nAChR$ mRNA and protein levels of the AC group was significantly lower than the HC group ($p = 0.001$). The mean expression of $\alpha 7nAChR$ mRNA and protein levels in group AT ($p = 0.001$), ATS ($p = 0.001$), and AS ($p = 0.001$) was significantly higher than the AC group.

The level of $\alpha 7nAChR$ protein in the hippocampus of AS and AST group rats was higher than the AT group ($p = 0.001$). But this significant increase in the protein level was not associated with a significant increase in the gene expression level because the increase in $\alpha 7nAChR$ mRNA of AS and AST groups to the AT group was not significant ($p = 0.998$, $p = 0.937$). However, differences in the $\alpha 7nAChR$ mRNA and protein levels between HC and the treated groups were significant ($p < 0.05$). Our results illustrate that the swimming, and clove oil improved the expression of reduced $\alpha 7nAChR$ mRNA and protein levels in the rat model of AD.

2.4. Swimming and clove oil supplementation reduced the increased expression of NLRP1 mRNA and protein levels in the hippocampus of the Rat model of AD

NLRP1 induces inflammatory responses and is recognized to affect AD and dark cell percentage. This study examined the effect of AD, exercise, and clove oil on the expression of NLRP1 mRNA and its protein level. Photomicrographs of immunofluorescence staining and the mean expression of NLRP1 mRNA and protein level are shown in Fig. 5. This study showed a significant difference in NLRP1 mRNA and its protein level was observed among the studied groups ($F = 52.025$, $p = 0.001$) and ($F = 188.402$, $p = 0.001$), respectively. This study showed no difference between the HC and Sh groups ($p = 1.00$). $A\beta_{1-42}$ injection caused an increase of NLRP1 mRNA and its protein level in the AC group compared to the HC group ($p = 0.001$). The mean NLRP1 mRNA and its protein level in the ATS ($p = 0.001$) and AS ($p = 0.001$) groups were lower than in the AC group; nevertheless, no difference was observed between AT and AC groups ($p = 0.191$, $p = 0.055$). This showed that clove oil supplement (with or without exercise) could balance the


increased expression of NLRP1 protein level induced by $A\beta_{1-42}$ injection. A comparison between treated groups showed that treatment with clove oil and exercise (ATS group) resulted in a more considerable decrease in NLRP1 protein rate than exercise (AT; $p = 0.001$) and clove oil (AS; $p = 0.006$) alone among the Alzheimer groups. Nonetheless, no reduction was found in the NLRP1 mRNA of the ATS group compared to AT ($p = 0.108$) and AS ($p = 1.000$) groups.

3. Discussion

The present study showed that spatial learning and memory were impaired in the Rat model of AD. In treated $A\beta_{1-42}$ injected rats, the protein rate and mRNA expression of $\alpha 7nAChR$ were comparable to those in the healthy control group. $\alpha 7nAChR$ expression has been reported to be reduced in the brains of Alzheimer's patients (Oz et al., 2013). Because of AD and at high doses of $A\beta$, a cross-link between $A\beta$ and $\alpha 7nAChR$ is established and ultimately leads to blockage and dysfunction of $\alpha 7nAChR$ (Ni et al., 2013). While according to the study by Tan et al., $\alpha 7nAChR$ can restrain the formation of interleukin-1 β and interleukin-18 by inhibiting NLRP1 and preventing inflammation-related injuries (Tan et al., 2014). In the current study, due to the $A\beta_{1-42}$ injections, the increase in mRNA expression and protein rate of NLRP1 was found in the CA1 region of the hippocampus. It seems $A\beta$ can activate inflammation (Masters and O'Neill, 2011). Increasing $A\beta$ activates caspase-1, leads to the secretion of pro-inflammatory cytokines, and finally, increases pyroptosis (Lamkanfi and Dixit, 2014). Therefore, the increase in NLRP1 is probably due to a rise in $A\beta$ and a decrease in $\alpha 7nAChR$.

In addition, the increase in dark cells following AD has been confirmed by some researchers (Tan et al., 2014; Lamkanfi and Dixit, 2014). We have seen an increase in dark cells in the CA1 region of the hippocampus following $A\beta_{1-42}$ injection in the rat model of AD. One of the significant causes of dark cells is the activation of caspases (Song et al., 2018). The activation of caspases can be due to inflammasomes such as NLRP1, which eventually leads to an increase in dark cells (Song et al., 2018). Increased dark cells can cause memory impairment (Lee et al., 2003). Researchers have previously stated that excessive release of AChE, and afterward, the reduction in the level of acetylcholine, is one of the leading causes of impaired spatial memory (Farioli-Vecchioli and Tirone, 2015; Amirahmadi et al., 2022). The results illustrated that the reduced $\alpha 7nAChR$ expression, increased NLRP1 expression, and activated dark cells pathway had contributed to cognitive impairment and

A



B




Fig. 3. Effect of swimming exercise and clove oil on the Percentage of dark cells in the CA1 region of the hippocampus in the rat model of AD in different groups. (A) The dark cells Percentage in the CA1 region of the hippocampus in other groups. Significant level: $p < 0.05$. (B) Purple plaques indicate dark cells. Bar with different letters shows a significant difference (ANOVA and subsequent Tukey's HSD, $p < 0.05$). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

the reduction of spatial memory caused by AD.

An active lifestyle may prevent Alzheimer's (Brasure et al., 2018) and delay the onset of AD. This study discovered that exercise was a positive and influential factor in increasing the protein expression level and mRNA of $\alpha 7nAChR$ in Alzheimer's groups. The findings also emphasized the role of exercise in reducing NLRP1 mRNA and protein levels. In the study of Khakroo Abkenar et al. (2019), high-intensity aerobic exercise increased interleukin-1 β , interleukin-18, and NLRP3 and subsequent neuronal death, while low-intensity exercise has backfired. According to moderate-intensity aerobic exercise, NLRP1 changes to physical exercise are associated with exercise intensity; high-intensity exercise may increase NLRP1 levels and worsen AD symptoms. However, confirmation or rejection of this hypothesis requires further investigation. Due to the direct relationship between A β and NLRP1 in the hippocampus (Tan

et al., 2014), the exercise by reducing A β levels in the rat model of AD (Alkadhi and Dao, 2018) may potentially reduce NLRP1 expression and prevent dark cells and memory impairment.

Furthermore, Farzi et al. have reported that exercise improves spatial memory by reducing AChE activity in the rat model of AD; As a result, decreased AChE activity can be considered as one of the mechanisms to enhance cognition and memory in AD (Farzi et al., 2018). Accordingly, the reduction in dark cells in the ATS group scan is attributed to the effect of exercise on increasing decreased $\alpha 7nAChR$ levels and decreasing increased NLRP1 levels in patient groups. In addition, reducing dark cells after exercise can improve memory.

Researchers have already shown that aerobic exercise reduces NLRP1 and interleukin (Khakroo Abkenar et al., 2019). Eight weeks of resistance training reduces NLRP3 gene expression and serum caspase-1





Fig. 4. Effect of swimming exercise and clove oil on the expression of $\alpha 7nAChR$ mRNA (A) and protein levels (B, C) in the CA1 region of the hippocampus in the rat model of AD in different groups. Green immunostaining indicates $\alpha 7nAChR$ expression. Significant level: $P < 0.05$. (A) Percentage of $\alpha 7nAChR$ protein expression in other groups. (B) $\alpha 7nAChR$ mRNA. (C) Immunofluorescence-stained images (Green dots express $\alpha 7nAChR$ protein). Magnification: 400 \times . Bar with different letters indicates a significant difference (ANOVA and subsequent Tukey's HSD, $p < 0.05$). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)





C

levels (Mejías-Peña et al., 2017). On the other hand, it is known that eugenol decreases interleukin (Abbasi et al., 2019). Exercise seems to improve memory and learning by enhancing the role of eugenol in suppressing inflammation.

Clove oil has antioxidant, anti-inflammatory, and anti-viral properties (Han and Parker, 2017). In this study, clove oil consumption partially compensated for the decreased protein expression level and

mRNA of $\alpha 7nAChR$ in the CA3 region of the hippocampus of the rat model of AD. The effect of clove oil on reducing AChE was confirmed in the study by Dalai et al. (Dalai et al., 2014). They demonstrated eugenol to be helpful in cognitive diseases such as Alzheimer's by affecting AChE (Dalai et al., 2014). Since the protein expression and mRNA of $\alpha 7nAChR$ in the rat model of AD increased after consumption of clove oil and according to previous research (Wie et al., 1997; Dalai et al., 2014),



C

Fig. 5. Effect of swimming exercise and clove oil on the expression of NLRP1 mRNA (A) and protein levels (B, C) in the CA1 region of the hippocampus in the rat model of AD in different groups. Green immunostaining indicates NLRP1 expression. Significant level: $p < 0.05$. (A) Percentage of NLRP1 protein expression in different groups. (B) NLRP1 mRNA. (C) Immunofluorescence-stained images (Green dots express NLRP1 protein). Magnification: 400×. Bar with different letters indicates a significant difference (ANOVA and subsequent Tukey's HSD, $p < 0.05$). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

clove oil may effectively improve the protein expression level and mRNA of $\alpha 7nAChR$ by reducing AChE activity.

According to previous research, eugenol alters the nuclear factor kappa B (NF- κ B) pathway, and it reduces interleukin-6 production (through its anti-inflammatory activity on (COX)-2 and 5-LOX) Also, eugenol improves learning and memory by altering neurotransmitters in the brain in Kunming mice (Wie et al., 1997). Eugenol also effectively suppresses inflammation and subsequently can reduce neuronal death (Taher et al., 2015). In the present study, eugenol may promote learning and memory and reduce neuronal death by affecting the inflammatory pathway.

Consuming clove oil also reduced the protein expression level and mRNA of NLRP1 in the CA1 region of the hippocampus of the rat model of AD. Clove oil has anti-inflammatory activities by activating microglia (Taher et al., 2015). NLRP1 is recognized as one of the most crucial components of inflammation, and clove oil appears to reduce NLRP1 mRNA and protein expression by affecting the NF- κ B pathway and interleukin-6 production (Wie et al., 1997).

As the results showed, clove oil reduced the percentage of dark cells in the CA1 region of the hippocampus (although it was not significant). Previously, it has been found that clove oil effectively suppresses inflammation (Taher et al., 2015). In this study, clove oil stops dark cells by reducing inflammation. The results showed that consuming clove oil improved learning and memory in the ATS group, which improved the cognitive function of the ATS group to the level of the healthy control group. Clove oil can modify cognitive function by reducing AChE activity, reducing oxidative damage, and improving the neuroprotective effect (Song et al., 2018).

The role of clove oil consumption in reducing NLRP1 (as a component of the inflammation pathway) was confirmed. On the other hand, inflammation can reduce cognitive function by intensifying the process of dark cells. Therefore, clove oil seems to improve learning and memory by altering $\alpha 7nAChR$ levels and subsequently affecting the inflammatory pathway.

Generally, it seems that A β is dose-dependent. Under normal conditions, the low-level A β triggers the PI3K/MAPK current by binding to $\alpha 7nAChR$ and forming A β - $\alpha 7nAChR$ complexes (Parri et al., 2011). Under these conditions, $\alpha 7nAChR$ can also inhibit the formation of interleukin-1 β and interleukin-18 by inhibiting NLRP1; in this way, it prevents inflammatory-related injuries (Tan et al., 2014).

In AD, an increased level of A β decreases $\alpha 7nAChR$ efficiency (Ni et al., 2013); up to previous research, we know it reduces $\alpha 7nAChR$ results in the increase of NLRP1 and ultimately leads to inflammation (Abbasi et al., 2019; Ni et al., 2013). Conversely, increasing caspase-1 causes inflammation (Khakroo Abkenar et al., 2019), enhancing dark cells percentage (Khakroo Abkenar et al., 2019).

According to previous research, clove supplementation and physical activity reduce A β levels (Brasure et al., 2018). The combination of training with clove oil supplementation is probably associated with further reductions in A β , caspase-1, interleukin-1 β , and interleukin-18 levels (Khakroo Abkenar et al., 2019). In addition, it will likely decrease dark cells (Khakroo Abkenar et al., 2019)-clove supplementation and physical activity double effect reducing NLRP1 levels. Decreased NLRP1 levels are likely associated with decreased caspase-1 levels and eventually reduced dark cells.

Clove oil supplementation and physical activity can increase reduced $\alpha 7nAChR$ levels and reduce elevated NLRP1 levels in AD by decreasing levels of caspase-1, interleukin-1 β , interleukin-18 (Khakroo Abkenar et al., 2019), AChE, and A β (Brasure et al., 2018). Hence, clove oil can potentially improve the quality of life of Alzheimer's patients. However, confirmation of the details of this pathway requires more future extensive research.

4. Conclusion

In conclusion, the present study showed that endurance training and

clove oil supplementation could enhance spatial memory by modifying $\alpha 7nAChR$ and NLRP1 levels and dark cells. Endurance training and clove oil supplementation caused an increase in $\alpha 7nAChR$ and a decrease in NLRP1 in the hippocampus of the Rat model of AD. Positive changes in these factors improve spatial learning and memory probably.

This study showed that the combination of endurance training and clove oil supplementation caused a significant improvement in studied factors in the Rat Model of AD because the training and supplementation group rats experienced better memory performance than the other rats. Therefore, the results of this study present evidence to investigate new therapeutic options in the rat models of AD.

5. Methods

5.1. Animals

Forty-eight male Wistar rats (6 weeks old) were purchased from the Pastor Institute (Tehran, Iran). The rats were housed (4 rats per cage) at 22 ± 2 °C, with cycles of light/dark 12:12 h, and were fed a pellet diet (Behparvar, Thran, Iran) and water ad libitum. The rats have been cared for following the International Association for the Study of Pain (IASP). The Lorestan University Animal Ethics Committee approved the study in Khorramabad, Iran (LU.ECRA.2018.15). This Committee confirms that all experiments were performed under relevant guidelines and regulations. Further, the present study complied with the Animal Research: Reporting of in vivo Experiments (ARRIVE) guidelines. After one week of adaptation to the new environment, rats were familiarized with the swimming protocol (30 s swimming/2 periods). After that, randomly, the animals were divided into six groups with eight rats in each group, as follows: healthy-control (HC), Alzheimer control (AC), Alzheimer-training (AT), Alzheimer-supplement (AS), Alzheimer-training-supplement (ATS), and Sham (Sh) (Fig. 6). Eight samples (same animals) were used for each immunofluorescence/histological and PCR analysis.

5.2. Hippocampal injections of A β_{1-42} to induce AD pathology

A β_{1-42} was provided from Sigma-Aldrich, USA. Next, A β_{1-42} aggregation was prepared by dissolving that in sterile saline. The solution was incubated to a final 1 mg/ml concentration at 37 °C for seven days. This allowed the peptide to fibrillize associated with toxicity and stored at -20 °C. The A β_{1-42} solution containing insoluble precipitates of both fibril-like structures and different-sized oligomers facilitates marked learning deficits. After that, the rats were anesthetized with ketamine (75 mg/kg, Interchemie, Netherland) and Xylazine (10 mg/kg, Bremer/pharmaGmbH, Germany) and placed onto a stereotaxic frame (RWD68505, Brazil). The skull was positioned in the frame, and a sagittal midline incision was made along the midline. The area surrounding the bregma was cleaned and dried. After that, the incubated A β_{1-42} solution was injected into the CA1 region with a volume of 5 μ l containing 5 μ g A β_{1-42} very slowly (over 2 min) bilaterally. Stereotaxic injection coordinates were as anterior-posterior (AP), -3.3 mm; medial-lateral (ML), ± 1.8 mm; ventral (V), -2.5 mm; by measuring the distances from bregma base on the stereotaxic Atlas of Paxinos and Watson (2006). To prevent A β_{1-42} reflux out of the needle tract and maximize diffusion, the needle remained at the injection site for 1 min after injection (Limón et al., 2012). Injections of A β_{1-42} (Limón et al., 2012) or isotonic saline solution (Panahzadeh et al., 2022) (Vehicle) for AD or Sham groups were infused with a Hamilton syringe. Following surgery, the rats were returned to their cages and frequently monitored during recovery.

5.3. Evaluation of the Alzheimer's model

In addition to 48 rats in the main groups, in the first phase, four animals were euthanized ten days after injection to perform histological




Fig. 6. Experimental design: AD was induced by injection of $A\beta_{1-42}$, and after 10 & 11 days, treatment with clove oil and swimming began for three weeks. MWM was performed on the 17th day of exercise interventions; finally, 48 h after the last intervention, animals were euthanized for more study.

analyses to evaluate the Alzheimer's model 32. The remainder rats were used for the next phase. Immunofluorescence was used for evaluating the Alzheimer's model.

5.4. Supplement treatment

Clove oil was purchased from the Shafa Kurdistan Company (Kurdistan, Iran). The GC-MS method was used to determine the quantity of the active compounds in the clove oil (Hewlett-Packard, HP-6890, USA). The column used was HP-1MS (methyl silicon-cross link). Column (60 m × 0.20 mm, the film thickness 0.25 μm). The oven temperature was as follows: Initial temperature: 160 °C, Final temperature 230 °C, at a rate of 7 °C/min. The carrier gas was Helium (99.9999 %), with a 1 ml/min flow rate. Injector temperatures were 250 °C. For Mass Spectrometry-MS, HP-59-70 (Hewlett-Packard, USA) was used. The gas

Table 1
Chemical composition of clove oil.

Compound	PA (%) ^a	KI ^b
Eugenol	85/5	2350
αCaryophyllene	7/5	2460
Caryophyllene	0/8	2580
Eugenyl acetate	1/3	759
Others	4/9	different

^a Kovats index.

^b Peak area percentage in relation to peak total area.

chromatographic analysis of clove oil is shown in Table 1. The supplement group was treated with 0.1 ml/kg of clove oil daily for 21 days (Kumar et al., 2016).

5.5. Swimming protocol

Rats were placed in a box containing 40 cm water at 37 °C (45 cm long, 25 cm wide, and 45 cm-high plastic containers). 1 ml of soap was added to decrease surface tension and to avoid floating behavior. The swimming protocol had two phases: The adaptation phase and the primary phase. Rats were submitted to swimming, according to Table 1. This swimming protocol was a high-intensity training model, and according to previous studies, blood lactate concentration remained more than 1 mmol/l (Table 2) (Kuphal et al., 2007) (with some minor modifications).

Table 2
Swimming protocol.

Phase	Adaptation phase				Main phase
	Day 1	Day 2	Day 3	Day 4	
Days	Day 1	Day 2	Day 3	Day 4	Day 5–20
Time	2	2	3	2	1
Repetition	30 s	2 min	10 min	15 min	30 min
Interval between swimming	2 h	2 h	5 min	5 min	no interval

5.6. Morris Water Maze test

The Morris Water Maze method (MWM) tested spatial learning and memory. MWM was performed on the 17th day of exercise interventions. MWM consisted of four successive trials and one probe day. This method used a water tank (50 cm in height and 130 cm in diameter). The tank was filled up with water (to a depth of 30 cm). The temperature of the water was 26 ± 2 °C. The MWM tank hypothetically was parted into four quadrants. During the successive trials, a movable hidden escape platform (8×6 cm² area) was located underwater surface in the fourth quadrant of the tank during the swimming time. In successive trials and probes, days were measured using the Noldus Ethovision system, version 5 (USA). Rats were trained for four successive trials with a maximum of 120 s daily to locate the platform. After finding the hidden escape platform, the animal was permitted to stay there for 30 s. Each animal's time to detect and locate the platform was considered escape latency time. If an animal could not find the platform, it was placed by hand on the platform. During the probe day, the platform was removed, and the animals were allowed to search the platform for 60 s. Each rat's time in the target quadrant was noted and analyzed later (Rahman et al., 2019).

5.7. Tissue preparation

After the end of the MWM, animals were anesthetized with ketamine (100 mg/kg) and Xylazine (5 mg/kg) and flowed by transcardial perfusion with saline (100 ml) and 4 % paraformaldehyde (250 ml) in phosphate buffer (0.1 mg) with pH 7.4 (Keshvari et al., 2020). The brain was immediately removed, and the hippocampus was dissected out. After that, half part of the hippocampus was frozen and the other part was fixed in paraffin for further studies.

5.8. RNA extraction and cDNA synthesis

About 50 mg of the hippocampus was homogenized. Then chloroform was added, and the microtube was shaken vigorously (for 15 s). Subsequently, the microtube was centrifuged (at 4 °C for 15 min at 12,000 PRM). The RNA was presented in the aqueous phase. Lastly, the extracted RNA was washed in ethanol and dissolved in 20 µL RNase-free water. After that, its concentration was checked. For cDNA synthesis, 1 µg of total RNA was used. Up to the manufacturer's protocol, synthesize using the Revert Aid First Strand cDNA Synthesis kit (Thermo Scientific, USA). The cDNA was kept at -20 °C for further use (Zhang et al., 2015).

5.9. Real-time-PCR

For $\alpha 7$ nAChR and NLRP1 mRNA expression analyses, quantitative real time-PCR was performed using an ABI Step-One real-time PCR system and software (Applied Biosystems, USA). B-ACTIN was used as the reference gene. The amplification reactions were detected using the Real Q Plus Master Mix Green high ROX (Ampliqon, Denmark). The standard PCR conditions were 95 °C (10 min), 95 °C (15 s), and 60 °C (1 min) for 40 cycles. The expression levels of $\alpha 7$ nAChR and NLRP1 transcripts were analyzed by the $2^{-\Delta\Delta}$ CT method. Results were obtained from ratios between the target gene ($\alpha 7$ nAChR and NLRP1 mRNA) and the B-ACTIN housekeeping mRNA (Saresella et al., 2016). The primer sequences in the qRT-PCR were as follows: B-ACTIN, F→GTGTGATGGTGGGTATGGGT, and R→GGTCATTGTAGAAAGTGTGGTG; $\alpha 7$ nAChR, F→ATTGAAGATGTGGAATGGGAGGT, and R→AGGTTGACGATGTAGAAGGAGGA; NLRP1, F→CAAGAGGGAAAGGTGGACAG, and R→GGAAGTGATGGGGATGAAGTGT.

5.10. Immunofluorescence

For immunofluorescence, the dissected hippocampus was immersion fixed in 10 % formalin for 48 h (Rahman et al., 2021) Blocks of the

hippocampus were embedded in paraffin and then cut into 5 µm thick. Paraffin was removed, and the sections were blocked with goat serum (10 %) and Triton X-100 (3 %) in PBS for 30 min. Subsequently, a primary antibody was added in diluting 1:100 in PBS. Antibodies included $\alpha 7$ nAChR (cat. no. sc-5544; Santa-Cruz Biotechnology Inc., USA) and NLRP1 (cat. no. ab3683; Abcam, UK). Sections with primary antibodies were incubated at 2 to 8 °C overnight. After washing with PBS (4 times for 5 min), the secondary antibody (Goat Anti-Rabbit IgG H&L; cat. no. ab6717; Abcam, UK) was diluted 1:150 in PBS and incubated with sections at 37 °C for 1.30 h in darkness. After washing four times, DAPI was added (Zhan et al., 2015). 500 µm after and before of injection, the section was selected. Each sample was sectioned 5 µm with a 25 µm interval between each section based on neuron size [each neuron is usually 15–25 µm]. For each rat, 20 tissue sections with a thickness of 5 µm were examined; the average of every 20 tissue stained and counted. All of them were calculated in the statistics. The sections were investigated with a fluorescence microscope (LABOMED, USA) and a camera (Delta Pix, Denmark). The images were analyzed using the software Image J 1.5.

5.11. Cresyl violet

We sectioned the CA1 region of the hippocampus at 5-µm thickness, using the Paxinos and Watson atlas (2006) as a reference. After that, hippocampal sections were Nissle-stained for 20 min, with 0.1 % Cresyl violet solution. Subsequently, the sections were washed in water, then in 70 % ethanol (2 steps/min), cleared in Xylenol (2 steps/5 min), and dried in the fume hood. The percentage of dark cells in the CA1 region of the hippocampus was evaluated. After staining, counting each section in three incisions was done with a minimum distance of 50 µm. An area of 1350 µm² was considered for counting neurons in each group. The counting of dark cells was examined using a light microscope (LABOMED, USA) with a magnification of $\times 400$. Image J software v1.8 (NIH, Wayne Rasband, USA) was used (Azad et al., 2011). Apoptotic cells were evaluated based on cell shrinkage, loss of uniformity of Nissl body, cytoplasm, nucleus density, and pyknotic nucleus. Viable cells were considered cells with orderly arranged cells with normal morphology, abundant cytoplasm, and evident nucleus and nucleolus (Pakravan et al., 2022).

5.12. Statistical analysis

Shapiro-Wilk Test investigated the normality of data. The one-way ANOVA was utilized to compare groups with different variables. Tukey's post hoc test was used as the supplementary test. Kruskal-Wallis analysis was used to evaluate the dark cells. To examine the interaction of the day*group on spatial learning, we used a two-way ANOVA test.

The independent *t*-test was utilized for the evaluation of the Alzheimer's model. Data was defined according to the mean \pm SD. The significance level was considered as $p \leq 0.05$. SPSS 20 was used for data analysis.

Sponsor role

The sponsor has no role in the design, methods, subject recruitment, data collection, analysis, and preparation of the paper.

Funding

No funding was obtained for this study.

CRedit authorship contribution statement

Z. GK. And M. F. designed this study; Z. GK. Analyzed the data. Z. GK., M. F., R. M. and E. VZ. Performed research and wrote the paper.

Declaration of competing interest

The authors declare no competing interests.

Data availability

The datasets used and/or analyzed during the current study available from the corresponding author on reasonable request.

Acknowledgments

This study is the result of a doctoral dissertation and has been done in collaboration with the deputy of Research and Technology of Lorestan University.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.exger.2023.112192>.

References

- Abbasi, Y., et al., 2019. Neuroprotective effect of ethanol and modafinil on focal cerebral ischemia in rats. *Metab. Brain Dis.* 34 (3), 805–819. <https://doi.org/10.1007/s11011-018-0378-0>.
- Alkadhri, K.A., Dao, A.T., 2018. Exercise decreases BACE and APP levels in the hippocampus of a rat model of Alzheimer's disease. *Mol. Cell. Neurosci.* 86, 25–29. <https://doi.org/10.1016/j.mcn.2017.11.008>.
- Alkhouli, M.F., et al., 2019. Exercise and resveratrol increase fracture resistance in the 3xTg-AD mouse model of Alzheimer's disease. *BMC Complement. Altern. Med.* 19 (1), 39. <https://doi.org/10.1186/s12906-019-2451-6>.
- Amirahmadi, S., et al., 2022. Minocycline attenuates cholinergic dysfunction and neuroinflammation-mediated cognitive impairment in scopolamine-induced Alzheimer's rat model. *Inflammopharmacology* 30, 2385–2397. <https://doi.org/10.1007/s10787-022-01071-2>.
- Azad, N., et al., 2011. Neuroprotective effects of carnosic acid in an experimental model of Alzheimer's disease in rats. *Cell J.* 13 (1), 39. <https://doi.org/10.3233/JAD-190914>. Yakhteh.
- Baghshahi, H., et al., 2014. Antioxidant effects of clove bud (*Syzygium aromaticum*) extract used with different extenders on ram spermatozoa during cryopreservation. *Cryobiology* 69 (3), 482–487. <https://doi.org/10.1016/j.cryobiol.2014.10.009>.
- Bernardo, T.C., et al., 2020. Physical exercise mitigates behavioral impairments in a rat model of sporadic Alzheimer's disease. *Behav. Brain Res.* 379, 112358 <https://doi.org/10.1016/j.bbr.2019.112358>.
- Brasure, M., et al., 2018. Physical activity interventions in preventing cognitive decline and Alzheimer-type dementia: a systematic review. *Ann. Intern. Med.* 168 (1), 30–38. <https://doi.org/10.7326/M17-1528>.
- Cechella, J.L., et al., 2014. Moderate swimming exercise and caffeine supplementation reduce the levels of inflammatory cytokines without causing oxidative stress in tissues of middle-aged rats. *Amino Acids* 46 (5), 1187–1195. <https://doi.org/10.1155/2014/729827>.
- Cortés-Rojas, D.F., de Souza, C.R.F., Oliveira, W.P., 2014. Clove (*Syzygium aromaticum*): a precious spice. *Asian Pac. J. Trop. Biomed.* 4 (2), 90–96. [https://doi.org/10.1016/S2221-1691\(14\)60215-X](https://doi.org/10.1016/S2221-1691(14)60215-X).
- Dalai, M.K., et al., 2014. Anti-cholinesterase activity of the standardized extract of *Syzygium aromaticum* L. *Pharmacogn. Mag.* 10 (Suppl. 2), S276. <https://doi.org/10.4103/0973-1296.133275>.
- Esfandiari, M., et al., 2019. Beneficial effects of resveratrol and exercise training on cardiac and aortic function and structure in the 3xTg mouse model of Alzheimer's disease. *Drug Des. Devel. Ther.* 13, 1197. <https://doi.org/10.2147/DDDT.S196119>.
- Farioli-Vecchioli, S., Tirone, F., 2015. Control of the cell cycle in adult neurogenesis and its relation with physical exercise. *Brain Plast.* 1 (1), 41–54. <https://doi.org/10.3233/BPL-150013>.
- Farzi, M.A., et al., 2018. Exercise improves recognition memory and acetylcholinesterase activity in the beta amyloid-induced rat model of Alzheimer's disease. *Ann. Neurosci.* 25 (3), 121–125. <https://doi.org/10.1159/000488580>.
- Han, X., Parker, T.L., 2017. Anti-inflammatory activity of clove (*Eugenia caryophyllata*) essential oil in human dermal fibroblasts. *Pharm. Biol.* 55 (1), 1619–1622. <https://doi.org/10.1080/13880209.2017.1314513>.
- Huang, W., et al., 2019. Association of a functional polymorphism in the *CHRFAM7A* gene with inflammatory response mediators and neuropathic pain after spinal cord injury. *J. Neurotrauma* 36 (21), 3026–3033. <https://doi.org/10.1089/neu.2018.6200>.
- Jia, J., et al., 2019. Effects of vitamin D supplementation on cognitive function and blood A β -related biomarkers in older adults with Alzheimer's disease: a randomised, double-blind, placebo-controlled trial. *J. Neurol. Neurosurg. Psychiatry* 90 (12), 1347–1352. <https://doi.org/10.2147/DDDT.S196119>.
- Jiang, P., et al., 2014. The impacts of swimming exercise on hippocampal expression of neurotrophic factors in rats exposed to chronic unpredictable mild stress. *Evid. Based Complement. Alternat. Med.* 2014 <https://doi.org/10.1155/2014/729827>.
- Kelley, N., et al., 2019. The NLRP3 inflammasome: an overview of mechanisms of activation and regulation. *Int. J. Mol. Sci.* 20 (13), 3328. <https://doi.org/10.3390/ijms20133328>.
- Keshvari, M., et al., 2020. Effects of endurance exercise and *Urtica dioica* on the functional, histological and molecular aspects of the hippocampus in STZ-induced diabetic rats. *J. Ethnopharmacol.*, 112801 <https://doi.org/10.1016/j.jep.2020.112801>.
- Khakroo Abkenar, I., Rahmani-Nia, F., Lombardi, G., 2019. The effects of acute and chronic aerobic activity on the signaling pathway of the inflammasome NLRP3 complex in young men. *Medicina* 55 (4), 105. <https://doi.org/10.3390/medicina55040105>.
- Kumar, A., et al., 2016. Possible neuroprotective mechanisms of clove oil against icv-colicidine induced cognitive dysfunction. *Pharmacol. Rep.* 68 (4), 764–772. <https://doi.org/10.1016/j.pharep.2016.03.005>.
- Kuphal, K.E., Fibuch, E.E., Taylor, B.K., 2007. Extended swimming exercise reduces inflammatory and peripheral neuropathic pain in rodents. *J. Pain* 8 (12), 989–997. <https://doi.org/10.1016/j.jpain.2007.08.001>.
- Lamkanfi, M., Dixit, V.M., 2014. Mechanisms and functions of inflammasomes. *Cell* 157 (5), 1013–1022.
- Lee, M.-H., et al., 2003. Treadmill exercise suppresses ischemia-induced increment in apoptosis and cell proliferation in hippocampal dentate gyrus of gerbils. *Life Sci.* 73 (19), 2455–2465. [https://doi.org/10.1016/s0024-3205\(03\)00655-6](https://doi.org/10.1016/s0024-3205(03)00655-6).
- Limón, D., et al., 2012. Neuroprotective effect of the aminoestrogen prolame against impairment of learning and memory skills in rats injected with amyloid- β -25–35 into the hippocampus. *Eur. J. Pharmacol.* 685 (1–3), 74–80. <https://doi.org/10.1016/j.ejphar.2012.04.020>.
- Masters, S.L., O'Neill, L.A., 2011. Disease-associated amyloid and misfolded protein aggregates activate the inflammasome. *Trends Mol. Med.* 17 (5), 276–282. <https://doi.org/10.1016/j.molmed.2011.01.005>.
- Mejías-Peña, Y., et al., 2017. Impact of resistance training on the autophagy-inflammation-apoptosis crosstalk in elderly subjects. *Aging* 9 (2), 408. <https://doi.org/10.18632/aging.101167>. Albany NY.
- Neumann, S., et al., 2015. Innate immunity and inflammation post-stroke: an α 7-nicotinic agonist perspective. *Int. J. Mol. Sci.* 16 (12), 29029–29046. <https://doi.org/10.3390/ijms161226141>.
- Ni, R., Marutle, A., Nordberg, A., 2013. Modulation of α 7 nicotinic acetylcholine receptor and fibrillar amyloid- β interactions in Alzheimer's disease brain. *J. Alzheimers Dis.* 33 (3), 841–851. <https://doi.org/10.3233/JAD-2012-121447>.
- Oz, M., et al., 2013. On the interaction of β -amyloid peptides and α 7-nicotinic acetylcholine receptors in Alzheimer's disease. *Curr. Alzheimer Res.* 10 (6), 618–630. <https://doi.org/10.2174/15672050113109990132>.
- Pakravan, N., Abbasi, A., Hassan, Z.M., 2022. New paradigm in cell therapy using sperm head to restore brain function and structure in animal model of Alzheimer's disease: support for boosting constructive inflammation vs. anti-inflammatory approach. *J. Immunol. Res.* 2022 <https://doi.org/10.1155/2022/8343763>.
- Panahzadeh, F., Mirnasuri, R., Rahmati, M., 2022. Exercise and *Syzygium aromaticum* reverse memory deficits, apoptosis and mitochondrial dysfunction of the hippocampus in Alzheimer's disease. *J. Ethnopharmacol.* 286, 114871 <https://doi.org/10.1016/j.jep.2021.114871>.
- Parri, H.R., Hernandez, C.M., Dineley, K.T., 2011. Research update: Alpha7 nicotinic acetylcholine receptor mechanisms in Alzheimer's disease. *Biochem. Pharmacol.* 82 (8), 931–942. <https://doi.org/10.1016/j.bcp.2011.06.039>.
- Paxinos, G., Watson, C., 2006. *The rat brain in stereotaxic coordinates: hard cover edition. Elsevier.*
- Rahman, S.O., et al., 2019. Neuroprotective role of astaxanthin in hippocampal insulin resistance induced by A β peptides in animal model of Alzheimer's disease. *Biomed. Pharmacother.* 110, 47–58. <https://doi.org/10.1016/j.biopha.2018.11.043>.
- Rahman, M.A., et al., 2021. Alcoholic fixation over formalin fixation: a new, safer option for morphologic and molecular analysis of tissues. *Saudi J. Biol. Sci.* <https://doi.org/10.1016/j.sjbs.2021.08.075>.
- Saresella, M., et al., 2016. The NLRP3 and NLRP1 inflammasomes are activated in Alzheimer's disease. *Mol. Neurodegener.* 11 (1), 23. <https://doi.org/10.1186/s13024-016-0088-1>.
- Song, S.-H., et al., 2018. Treadmill exercise and wheel exercise improve motor function by suppressing apoptotic neuronal cell death in brain inflammation rats. *J. Exerc. Rehab.* 14 (6), 911. <https://doi.org/10.12965/jer.1836508.254>.
- Taher, Y.A., et al., 2015. Experimental evaluation of anti-inflammatory, antinociceptive and antipyretic activities of clove oil in mice. *Libyan J. Med.* 10 (1) <https://doi.org/10.3402/ljmv.v10.28685>.
- Tan, M., et al., 2014. Amyloid- β induces NLRP1-dependent neuronal pyroptosis in models of Alzheimer's disease. *Cell Death Dis.* 5 (8) <https://doi.org/10.1038/cddis.2014.348> e1382-e1382.
- Wie, M.-B., et al., 1997. Eugenol protects neuronal cells from excitotoxic and oxidative injury in primary cortical cultures. *Neurosci. Lett.* 225 (2), 93–96. [https://doi.org/10.1016/s0304-3940\(97\)00195-x](https://doi.org/10.1016/s0304-3940(97)00195-x).
- Zhan, X., et al., 2015. Inflammation combined with ischemia produces myelin injury and plaque-like aggregates of myelin, amyloid- β and A β PP in adult rat brain. *J. Alzheimers Dis.* 46 (2), 507–523. <https://doi.org/10.3233/JAD-143072>.
- Zhang, A., et al., 2015. Mechanisms that lead to the regulation of NLRP3 inflammasome expression and activation in human dental pulp fibroblasts. *Mol. Immunol.* 66 (2), 253–262. <https://doi.org/10.1016/j.molimm.2015.03.009>.