

Journal Pre-proof

Environmental factors regulate soil microbial attributes and their response to drought in rangeland ecosystems

Santiago Toledo, Viviana F. Bondaruk, Laura Yahdjian, Gastón R. Oñatibia, Alejandro Loydi, Juan Alberti, Martín Bruschetti, Jesús Pascual, Guadalupe Peter, Walter D. Agüero, Pedro R. Namur, Lisandro Blanco, Pablo L. Peri



PII: S0048-9697(23)03027-9

DOI: <https://doi.org/10.1016/j.scitotenv.2023.164406>

Reference: STOTEN 164406

To appear in: *Science of the Total Environment*

Received date: 3 March 2023

Revised date: 19 May 2023

Accepted date: 20 May 2023

Please cite this article as: S. Toledo, V.F. Bondaruk, L. Yahdjian, et al., Environmental factors regulate soil microbial attributes and their response to drought in rangeland ecosystems, *Science of the Total Environment* (2023), <https://doi.org/10.1016/j.scitotenv.2023.164406>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ENVIRONMENTAL FACTORS REGULATE SOIL MICROBIAL ATTRIBUTES AND THEIR RESPONSE TO DROUGHT IN RANGELAND ECOSYSTEMS

Toledo, Santiago^{*a}; Bondaruk, Viviana F.^{b,c}; Yahdjian, Laura^{b,c}; Oñatibia, Gastón R.^{b,c}; Loydi, Alejandro^d; Alberti, Juan^e; Bruschetti, Martín^e; Pascual, Jesús^e; Peter, Guadalupe^{f,g}; Agüero, Walter D.^h; Namur, Pedro R.^h; Blanco, Lisandro^h; Peri, Pablo L.ⁱ

- a) CIT-CONICET/Universidad Nacional de la Patagonia Austral, Santa Cruz, Argentina.
- b) Instituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a la Agricultura (IFEVA), CONICET-Universidad de Buenos Aires, Buenos Aires, Argentina.
- c) Facultad de Agronomía, Departamento de Recursos Naturales y Ambiente, Cátedra de Ecología, Universidad de Buenos Aires, Buenos Aires, Argentina.
- d) Centro de Recursos Naturales Renovables de la Zona Semiárida (CERZOS), CONICET. DBByF, Universidad Nacional del Sur, Bahía Blanca, Argentina.
- e) Instituto de Investigaciones Marinas y Costeras (IIMyC), CONICET - Universidad de Mar del Plata, Buenos Aires, Argentina.
- f) Universidad Nacional de Río Negro, Sede Atlántica, CEANPa, Viedma, Argentina.
- g) CONICET, Argentina
- h) Instituto Nacional de Tecnología Agropecuaria (INTA) - EEA La Rioja, La Rioja, Argentina.
- i) Instituto Nacional de Tecnología Agropecuaria (INTA), EEA Río Gallegos, Santa Cruz, Argentina.

E-mails: toledo.santiago@inta.gob.ar; bondaruk@agro.uba.ar; yahdjian@agro.uba.ar; onatibia@agro.uba.ar; alejandro.loydi@gmail.com; juanalberti@gmail.com; mbrusche@mdp.edu.ar; pascualjesu@gmail.com; gpeter@unrn.edu.ar; aguero.walter@inta.gob.ar; namur.pedror@inta.gob.ar; blanco.lisandro@inta.gob.ar; peri.pablo@inta.gob.ar.

***Correspondence: Santiago Toledo (toledo.santiago@inta.gob.ar)**

ABSTRACT

In ecosystems, soil microbial variables characterization are used to determine soil biological health and the response of soils to environmental stress. Although there are strong associations between plants and soil microorganisms, they may respond asynchronously to environmental factors and severe droughts. We aimed to: I) evaluate the spatial variation of soil microbiome such as microbial biomass carbon (MBC) and nitrogen (MBN), soil basal respiration (SBR) and microbial indexes in eight rangeland sites located across an aridity gradient (distributed from arid to mesic climates); II) analyze the relative importance of main environmental factors (climate, soils, and plants) and their relationships with microbial variables in the rangelands; and III) assess the effect of drought on microbial and plant variables in field-based manipulative experiments. First, we found significant changes of microbial variables along a precipitation and temperature gradient. The responses of MBC and MBN were strongly dependent on soil pH, soil nitrogen (N), soil organic carbon (SOC), C:N ratio and vegetation cover. In contrast, SBR was influenced by the aridity index (AI), the mean annual precipitation (MAP), the soil pH and vegetation cover. MBC, MBN and SBR were negatively related with soil pH compared to the other factors (C, N, C:N, vegetation cover, MAP and AI) that had a positive relationship. Second, we found a stronger soil microbial variables response to drought in arid sites compared to humid rangelands. Third, the responses of MBC, MBN, and SBR to drought showed positive relationships with vegetation cover and aboveground biomass, but with different regression slopes, this suggest that plant and microbial communities responded differently to drought. The results from this study improves our understanding about the microbial response to drought in different rangelands, and

may facilitate the development of predictive models on responses of soil microorganisms in carbon cycle under global change scenarios.

KEYWORDS

Aridity index; Reduced precipitation; Soil microbiome; Plant-soil-microorganism; Differential response, Multiple soil classes

1. INTRODUCTION

Rangelands are among the most widely distributed terrestrial biomes, covering around 52.5 million km² (40.5% of the global land area), and located predominantly in arid and semiarid regions (Sala et al., 2017). Rangeland ecosystems provide multiple goods (e.g., livestock production) and services, and play an important role in global carbon (C) cycling, containing more than one third of above and belowground C stocks (Havstad et al. 2007; Yahdjian et al. 2015; Zhao et al., 2020; Peri et al, 2021; Jaman et al., 2022). In these and other terrestrial ecosystems, active soil living microorganisms represents only 0.1 – 2 % of the total soil volume (Blagodatskaya and Kuzyakov, 2013), but are involved in 90% of soil ecosystem functions (Nannipieri et al., 2003). Characterization of soil microbial variables is being increasingly used to determine soil biological health, including response of soils to environmental stress such as severe droughts, anthropogenic disturbances, and as an indicator of ecosystem resilience (Marcos and Olivera, 2016; Li et al., 2018; Fierer et al., 2021). Several studies have investigated the biogeographic patterns of the soil microbial variables across precipitation and temperature gradients (Bachar et al., 2010; Maestre et al., 2015; Delgado-Baquerizo et al., 2016; Yao et al., 2017; Ochoa-Hueso et al., 2018; Wang et al., 2021). However, mechanisms by which edaphic, vegetation, and climatic factors interact to shape local microbial communities are not yet fully understood (Waldrop et al., 2017; Sun et al., 2020a; Deng et al., 2021). This is critical for improving our understanding of the major environmental drivers of soil microbial

communities and their roles in ecosystem functioning (Saccá et al., 2017; Jansson and Hofmockel, 2020), especially in vulnerable rangelands facing climate change. Moreover, the response of the microbial biomass, activity, and community structure to environmental drivers may depend on each site's location across to aridity gradients (Bachar et al., 2010; Stomeo et al., 2013; Waldrop et al., 2017; Wang et al., 2021; Ding and Eldridge, 2022).

Current climate change scenarios are impacting the hydrological cycle by altering precipitation patterns and increasing the frequency of extreme dry events (Trenberth et al., 2014; Bonan and Doney, 2018; IPCC, 2022). In recent years, droughts are increasing in magnitude, frequency and duration, especially in arid and semiarid rangelands (Yahdjian and Sala, 2008; Hoover and Rogers, 2016; Canarini et al., 2017; Zhao et al., 2021), which may lead to land degradation and desertification (Lin et al., 2015; Huang et al., 2016; Middleton, 2018), jeopardizing sustainable livestock production worldwide (Yahdjian and Sala, 2008; Gaitán et al., 2018; Oñatibia et al., 2020; Bondaruk et al., 2022). In addition, the magnitude of change in precipitation (amount, intensity, and frequency) affects processes that control soil organic carbon pools and dynamics in terrestrial biomes as ecosystem responses are largely dependent on the belowground microbial communities (Bardgett et al., 2008; Canarini et al., 2017; Ren et al., 2017; Fuchslueger et al., 2019; Sun et al., 2020b; Deng et al., 2021). Changes in soil microbial variables have subsequent interactive feedbacks on numerous soil functions, such as nutrient cycling and carbon sequestration (Six et al., 2006; Bardgett et al., 2008; Canarini et al., 2017; Deng et al., 2021). Thus, identifying the microbial community responses to change in precipitations can greatly improve our understanding of the ability of ecosystems to

deal with future global climate change (Bardgett et al., 2008; Beier et al., 2012; Liang et al., 2017, 2019).

Given the importance of drought, manipulated precipitation experiments have been increasingly used in ecological studies over the past decade (Beier et al., 2012; Nielsen and Ball, 2015; Homyak et al., 2017; Zhou et al., 2018; Xu et al., 2020).

However, results have been diverse and there is still substantial uncertainty in describing microbial community responses to rainfall reductions at regional scales (Shen et al., 2015; Ochoa-Hueso et al., 2018; Xu et al., 2020; Deng et al., 2021).

Therefore, more studies of soil microbial response to rainfall reductions are urgently needed to improve predictions of global change feedback on the terrestrial carbon cycle.

Ochoa-Hueso et al. (2018) and Ren et al. (2018) found that rainfall reductions modified soil microbial biomass by reducing soil water and altering plant net primary productivity. Rainfall reductions also decrease the mobility of soil solutes and litter inputs, and limit substrate supply by changing rhizodepositions (Sanaullah et al., 2011; Fuchslueger et al., 2009). In addition, reductions in aboveground net primary production due to reduced rainfall can lead to decline in carbon and nitrogen from detritus entering the soil, thus reducing microbial biomass (Sanaullah et al., 2011; Zhang and Xi, 2021; Jaman et al., 2022) through inhibited microbial growth and activity (Bardgett et al., 2008; Dijkstra et al., 2015; Zhao et al., 2021). Therefore, rainfall reductions could have differential responses on soil microbial biomass and activity depending on the aridity of rangelands. Despite an increasing number of studies testing above and belowground ecosystems responses to changes in precipitation (Knapp et al., 2015, 2017; Bondaruk et al., 2022; Jaman et al., 2022), a critical knowledge gap is the combined responses of plants and soil microbes to

changes in precipitation. This gap could lead to a biased understanding of rangeland function and the ecosystem services they provide under climate change as soil microorganisms play a key role in carbon cycling processes, such as litter decomposition and CO₂ emissions (Six et al., 2006; Bardgett et al., 2008; Benner, 2011; Liang et al., 2017). Although, there are strong associations between plants and soil microbes through exchanges at root-soil interfaces, plants and microbes may respond asynchronously to soil resource availability due to contrasting life history strategies (Thakur et al., 2015; Sun et al., 2020a; Zhang and Xi, 2021). Soil microorganisms can adapt to changes in soil moisture more rapidly than plants due to their fast growth, considerable capacity for osmotic adjustment under fluctuating soil moisture and community composition shifts (Schimel et al., 2007; Lau and Lennon, 2011; Fuchslueger et al., 2016; Karłowscy et al., 2018b). The asynchrony between plant and microbial biomass may have significant implications for the competitive balance between plants and soil microbes, as well as for the regulation of biogeochemical cycles (Karłowscy et al., 2018b; Williams and de Vries, 2020). Argentina has an extensive territory covering broad ecological zones, with distinct climates and rangelands (Cyrzabab et al., 2018), accounting for more than two thirds of the total continental area (Anderson et al., 2011). Given their contrasting ecological conditions, these rangelands are expected to respond differently to environmental pressures. Thus, these rangelands are ideal for evaluating ecosystem services and resilience because they vary across climate gradients (Yahdjian and Sala, 2008; Verón et al., 2018; Bondaruk et al., 2022), and are likely to be unevenly affected by forecasted droughts (González et al., 2017). In Argentinean rangelands, soil microbial variables have been rarely studied (Montecchia et al., 2011; Prieto et al., 2011; Olivera et al., 2016; Toledo et al., 2021; Viruel et al., 2022). Therefore, an

improved understanding of soil microbial carbon cycling, as well as its interaction with environmental factors, is urgently needed to accurately represent soil microbial feedbacks in ecosystem models to improve predictions of rangeland responses to projected climate change scenarios (IPCC, 2022).

In this work, we built on our former study to investigate the main environmental drivers of soil microbial communities across an aridity gradient in Argentinean rangelands (Bondaruk et al., 2022). As microorganisms play a key role for rangeland carbon cycling, we added the soil component to the previous study focused on vegetation, and we compared the effects of drought on plants and microbes. In this research we used a coordinated manipulative drought experiment in eight Argentinean rangelands to: I) study the variation patterns of soil microbial community attributes such as microbial biomass (C and N), soil basal respiration and microbial indexes along an aridity gradient (arid, semiarid, sub humid and humid); II) analyzed the relative importance of climate, soil and plant factors on microbial variables and the relationships between microbial biomass (C and N) and soil basal respiration with the main environmental factors occurring along the aridity gradient; and III) we further assessed the effects of drought on microbial and plant variables. More importantly, we also assessed the relationships and differential responses to drought of soil microbial variables and vegetation.

2. MATERIALS AND METHODS

2.1. Study sites and aridity gradient descriptions

The sites are part of an international collaborative research network consisting of coordinated drought experiments (Drought Network). Were selected eight rangeland ecosystems across a natural aridity gradient in Argentina, distributed from 29° 57' to

51°54' S and 58° 09' to 70°24' W (Fig. A.1 supplementary). Based on the meteorological stations data (period 2015-2022), the mean annual temperature ranges from 6.5 to 22 °C, and the mean annual precipitation varies from 137.1 to 1022.3 mm, where 70–80% of rain events occurs during the growing season (September - March). For each study site, we calculated an aridity index (AI) based on mean annual precipitation (MAP) and mean annual temperature (MAT) as: $AI = MAP / (10 + MAT)$ (De Martonne, 1926). Considering the aridity categories defined by Liu et al. (2016), the eight rangelands were sorted into four ecosystem types: arid ($5 < AI < 10$), semiarid ($10 < AI < 20$), semihumid ($20 < AI < 30$), and humid ($30 < AI < 60$) (Fig. A.1 supplementary; Table 1). Also, the rangelands encompass a great variability in plant composition, plant species diversity and richness, vegetation cover, aboveground net primary productivity (ANPP), and soil physico-chemical properties (Table 1). These natural rangelands mainly sustain extensive livestock production.

TABLE 1

FIGURE A.1 supplementary

2. 2. Experimental design

The sites followed a common experimental protocol which allows reliable comparisons among contrasting ecosystems (Knapp et al., 2017) and included experimental rainfall manipulations to understand the ecological drought impact on vegetation (see Bondaruk et al., 2022). The simulation of droughts was induced through a passive well-tested design based on rainout shelters that intercept a fraction of the incoming precipitation (Yahdjian and Sala, 2002). To determine the percentage of precipitation interception, we applied the standardized protocol presented in the precipitation manipulation tool of the Drought Network. This is a software that allows to upload the longest precipitation time series of each site (<https://drought-net.colostate.edu/terrestrial-precipitation-analysis-package>; Lemoine

et al., 2016). Thus, we reduced annual precipitation to $\approx 50\%$ in sites. Intercepted precipitation was collected in gutters and directed away by pipes in order to avoid water infiltrating into the experimental drought plots (see Table A.1 supplementary).

TABLE A.1 supplementary

The rainout shelters were constructed with transparent plastic tiles, placed above the plant canopy (1.20 to 1.60 m height) in order to minimize impacts on micrometeorological conditions. The rainout-shelter design employed in this study has proven to accomplish the desired reduction in water input and the expected effects on soil moisture on vegetation (Yahdjian and Sala, 2002; Gherardi and Sala, 2013; Bondaruk et al., 2022). The experimental plots (3 × 3 m) were installed at the end of the growing season in different years (2015 to 2017) following a completely randomized block design that included paired control plots of the same size. At each experimental site, the experiments were established in a block design and generally replicated three times ($n = 3$), although some sites had four replicates ($n = 4$). All experimental sites were fenced to prevent grazing on experimental plots.

2. 3. Soil microbial community variables

Soil microbial variables were determined after 5 years of the imposition of drought with rainout shelters, during summer (December - March), which is coincident with the peak of the plant growing season. At each experimental site, eighteen soil samples were taken ($n = 2$ treatments × 3 plots × 3 composite samples). The composite soil samples consisted of 5 - 7 subsamples from the first 5 cm depth (without plant residues) taken with a metal cylinder (5.6 cm diameter). The samples were placed in a bag and taken to the laboratory, where they were sieved (2 mm) and conditioned to send to Laboratory INTA Río Gallegos (Santa Cruz, Argentina). They were kept refrigerated at 4 °C in the laboratory until the quantification analysis for microbial variables were determined.

The carbon in the microbial biomass (MBC) was estimated using the “chloroform-fumigation extraction” method (Vance et al., 1987). MBC was calculated as: $(OC_f - OC_{nf})/kEC$; where: OC_f = organic carbon extracted from fumigated samples; OC_{nf} = Carbon extracted from non-fumigated samples; and kEC = fumigation efficiency constant = 0.45. MBC was expressed as $mg\ C\ kg^{-1}$ dry soil.

The nitrogen in the microbial biomass (MBN) was determined by a modification of the fumigation–incubation method (Joergensen and Mueller, 1996). Liquid chloroform (1mL) was added directly to 30 g moist soil samples, stirred and left for 20 h in sealed beakers in desiccators. Chloroform was then removed using a vacuum pump.

Fumigated and non-fumigated samples were incubated at field capacity for 10 days at 25 °C. After incubation, samples were extracted with 2M KCl and analyzed for NH_4-N by the Berthelot reaction. MBN was calculated as the difference in N between fumigated and non-fumigated samples, divided by a correction factor. Non-extractable amount of microbial N was compensated for by a correction factor of $kN = 0.54$. MBN was expressed as $mg\ N\ kg^{-1}$ dry soil.

The soil basal respiration (SBR) was estimated by quantifying the carbon dioxide (CO_2) released in the process of microbial respiration during incubation days 1, 7, 14 and 21. This was done by placing 75 g fresh soil (moistened to 60 % of field capacity) into 1.5-L capacity glass containers with hermetic lids, together with a smaller flask containing 20 mL 0.2 M NaOH to capture the released CO_2 . The CO_2 was determined by titration with 0.1 M HCl, after precipitation of the barium carbonate formed by adding barium chloride ($BaCl_2$) aqueous solution to the NaOH solution, utilizing phenolphthalein as an indicator (Robertson et al., 1999). The SBR was measured until day 21 of incubation because during that period the samples reached

a relatively constant CO₂ production. The mean soil respiration (accumulated SBR/21) during the incubation time was expressed as mg C kg⁻¹ soil day⁻¹.

The soil microbial indexes, metabolic quotient (qCO₂) was calculated as the C-CO₂ evolution (SBR) per unit MBC and per unit time (Anderson and Domsch, 1990). The MBC/MBN ratio was obtained and the contributions of microbial biomass to SOC (MBC/SOC) were also calculated, expressed as percentage (Anderson and Domsch, 1990).

2. 4. Vegetation cover and aboveground biomass

During the experiment, we determined total vegetation cover and aboveground biomass at the peak of the growing season. Cover was determined in 1 m² permanent plots located in the center area of each experimental unit. Aboveground biomass was estimated by clipping to the soil surface all aboveground biomass within two frames of 0.2 × 0.5 m from each experimental unit near permanent plots. Then, we dried the samples (60 °C) for 48h to obtain dry matter plant biomass (total) expressed in g dry matter (DM) m⁻². For the Río Mayo (RM) and Chacra Patagones (ChP) sites, the estimation of aboveground biomass was made non-destructively, through allometric equations that related the relative cover of each plant species present with their aboveground biomass (e.g. Flombaum and Sala, 2007).

2. 5. Data Analysis

Descriptive analyses of means and standard deviations were realized for climatic, vegetation and soil characteristics. The microbial variables (MBC, MBN, MBC:MBN, SBR, qCO₂ and qMC) were analyzed with ANOVA, using the sites (RM, SPV, PA, LC, ChP, N, LCh and SC) and aridity categories (arid, semiarid, semihumid and humid) as factors. Significant differences between means were separated in all cases by Tukey's test with a significance level of p < 0.05. Soil biological attributes from all sites (MBC, MBN, MBC:MBN, SBR, qCO₂ and qMC) were analyzed using principal

component analysis (PCA). We ran random forest (RF) models by means of the *cforest* function in *party* package for R (Hothorn et al., 2008; Strobl et al., 2009), in order to assess relationships between microbial variables (MBC, MBN and SBR) and the climatic, vegetation and soil variables. The variable importance values were determined by using the *varimp* function. To quantify variability in variable importance scores, we developed 100 RF models based on random selections of 80 % of cases, each time using the other 20 % to evaluate model accuracy. RF in *party* package can be used reliably even in situations where the independent variables vary in their scale of measurement. We used main variables obtained from RF model to fit simple linear regressions between climatic, vegetation, soil and soil microbial community variables. The dependent variables were MBC, MBN and SBR and the independent variables were pH, C, N, C:N, vegetation cover, AI and MAP. The differences induced by experimental drought were tested for significance using one-way ANOVA. Significant differences between means were separated by Tukey's test with a significance level of $p < 0.05$. In addition, we calculated the response ratio (RR) to drought for each microbial variable (MBC, MBN, MBC:MBN, SBR, qCO_2 and qMC) in all sites using the meta-analytical methods (Hedges et al., 1999). Values of RR was calculated as: $\ln(RR) = \ln(X_t) / \ln(X_c)$, where X_t represents the drought treatment and X_c the respective mean values of a particular microbial variable in the control treatment. Significant responses ($p < 0.05$) were determined if the bootstrap confidence interval (CI) did not overlap with zero (Koricheva et al., 2013). Simple regressions were used to examine relationships between the microbial variables' RR (MBC, MBN, MBC:MBN, SBR, and qCO_2) and vegetation variables' RR (vegetation cover and aboveground biomass) to drought.

3. RESULTS

3.1. Soil microbial variables along the aridity gradient

Soil microbial variables varied significantly among sites and aridity categories ($p < 0.05$; Table 2). On the one hand, MBC, MBN and SBR ranged from 85.6 to 613.9, 16.9 to 148.3 and 22.7 to 85.9, respectively, and were significantly higher in LCh site (humid) compared with the other sites. On the other hand, metabolic and microbial indexes such as MBC:MBN, qCO_2 , and qMC decreased along the aridity gradient ($p < 0.05$ in all cases), except for one humid site (SC) which had high values of MBC:MBN and qMC .

TABLE 2

Principal components analysis (PCA; Fig. A.2 supplementary) determined that the first two axes explained 79.3 % of the total variance of the samples (57.7% and 21.6% for PC1 and PC2, respectively). In the analysis of axis 1, there was a clear separation between a group containing RM, SPV and LC sites and another with PA, N, LCh and SC sites. This pattern is explained because the first three sites presented higher qCO_2 values, while the other sites exhibited higher MBC, MBN and SBR values. In addition, axis 2 separated PA and SC from N and LCh sites, and also SPV site from LC site (Fig. A.2 supplementary).

FIGURE A.2 supplementary

The Random Forest (RF) model explained 93% of the variance for MBC, and 86 % for MBN (Fig. A.3 supplementary). The most important variables for predicting MBC and MBN across all sites were soil pH, C:N, vegetation cover, soil N contents and SOC (Fig. 1). For SRB, the RF model explained 73% of the variance (Fig. A.3 supplementary), and the main explanatory variables were aridity index, MAP, and soil

pH. In addition, soil pH was the single most important variable in RF models explaining variability for soil biological variables (Fig. 1).

FIGURE 1

FIGURE A.3 supplementary

To better visualize the relationship between soil microbial variables and environmental variables (soil physicochemical characteristics, vegetation, and climate), we analyzed biological correlations by using the most significant variables obtained from RF model. MBC was positively correlated with C:N ratio and vegetation cover ($p < 0.05$; Fig. 2 B and C). Similarly, MBN was positively correlated with N and SOC ($p < 0.05$; Fig. 2 E and F), whereas SBR was positively correlated with MAP and the aridity index ($p < 0.05$; Fig. 2 H and I). In contrast, MBC, MBN, and SBR were negatively correlated with the soil pH ($p < 0.05$; Fig. 2 A, D and G).

FIGURE 2

3.2. Responses of soil microorganisms to drought along the aridity gradient

The effect of drought on soil microbial variables (biomass, activity and microbial indexes) significantly varied among sites ($p < 0.05$; Fig. A.4 supplementary; Fig. 3). However, drought effects on these variables were not significantly related with the aridity index ($p > 0.05$). Drought decreased the MBC by 22 % and 16 % in RM and LCh, respectively, but it increased these variables in the sites SPV (31 %) and PA (8 %). Drought reduced MBN in RM (32 %), PA (8 %) and N (46 %), but increased it in the SPV, LC and LCh sites by 48 %, 40 % and 15 %, respectively. In contrast, drought significantly increased MBC:MBN ratio in RM (13 %), PA (17 %) and N (43 %), but decreased it in SPV (16 %), LC (24 %) and LCh (31 %). The SBR significantly increased under drought by 60 %, 42 % and 11 % in SPV, LC and N, respectively, and it decreased in RM (31 %) and SC (13 %). Finally, the qCO_2

decreased under drought conditions in RM (12 %) and SC (9 %), while it increased in SPV (28 %), LC (25 %), N (13 %) and LCh (15 %) (Fig. A.4 supplementary; Fig. 3).

FIGURE 3

FIGURE A.4 supplementary

3.3. Responses of plants and soil microorganisms to drought

Along the aridity gradient, the vegetation cover increased from 35.7 ± 4.1 % to 100 ± 0 % (Table A.2 supplementary). In addition, the aboveground biomass increased along the aridity gradient from 25.5 ± 3.6 g DM m⁻² in the most arid site (RM) to 769.3 ± 130.1 g DM m⁻² in the most humid site (SC; Table A.2 supplementary).

Drought differently impacted on vegetation cover and aboveground biomass among the sites (Table A.2 supplementary). Vegetation cover and aboveground biomass were significantly reduced by drought in RM (68 and 64 % respectively), and N (37 and 63 %). However, in SPV, vegetation cover and aboveground biomass significantly increased by 30 and 49 %, respectively in response to the drought (Table A.2 supplementary).

TABLE A.2 supplementary

Drought impact on vegetation cover estimated by the response ratio (RR) was positively correlated with the effect of drought on MBC, MBN, and SBR, and negatively correlated with the effect of drought on the MBC:MBN (Fig. 4). Similar results were found between responses to drought of aboveground biomass and microbial variables (MBC, MBN, MBC:MBN, and SBR) (Fig. A.5 supplementary). Finally, we found that the drought effects on microbial variables (MBC, MBN, SBR, and qCO₂) were positively correlated (Fig. A.6 supplementary).

FIGURE 4

FIGURE A.5 supplementary

FIGURE A.6 supplementary

4. DISCUSSION

Soil microbial biomass and activity response to aridity gradient mainly was determined by biotic and soil factors, and at a lower extent by climatic variables.

When manipulating precipitation (drought) at a site scale, microbial responses varied depending on vegetation cover rather than aridity index.

4.1. Soil microbial variables along the aridity gradient

Our findings provided new insights regarding the impact of environmental factors (soil, vegetation and climate) that drive the changes of soil microbial variables along an aridity gradient in rangeland ecosystems. We found that the responses of MBC and MBN were strongly dependent on soil pH, soil N and SOC, C:N ratio and vegetation cover. In contrast, SBR was influenced by the AI, MAP, pH and vegetation cover. Soil pH showed a negative relationship with MBC, MBN, and SBR compared with other factors that had a positive relationship. Regional scale patterns of microbial community (biomass and activity) were related to the biotic, edaphic and climatic factors such as plant composition and productivity, soil nitrogen, carbon, pH, precipitation and temperature (Yao et al., 2017; Ren et al., 2018; Deng et al., 2021; Wang et al., 2021). This is consistent with our study where soil microbial response varied depending on rangeland characteristics. From arid to humid, these sites differ in their biotic and abiotic factors such as aridity index, MAP, MAT, vegetation types and soil physicochemical properties. Several studies suggested that the regional climate determines vegetation types and soil characteristics, and therefore, directly and/or indirectly affects soil microbial variables (Chen et al., 2015; Maestre et al., 2015). In our study we found that the biomass and basal respiration of soil microorganisms increased with the aridity index (arid < semiarid < semihumid < humid). Many studies showed that microbial biomass (C and N) or activity (SBR) were higher with increasing MAP and MAT in rangelands, forests and other biomes

(Bachar et al., 2010; Griffiths et al., 2011; Delgado-Baquerizo et al., 2016; Ochoa-Hueso et al., 2018). In our study, random forest analysis indicated that soil and vegetation factors affected microbial biomass, with pH, C:N ratio, N content, SOC and vegetation cover being the most important factors explaining both MBC and MBN (Figs. 1 and 2). This contrasts with previous studies that have identified climate variables such as MAT (Nielsen and Ball, 2015; Yao et al., 2017; Fuchslueger et al., 2019; Yang et al., 2022) and MAP (Ma et al., 2015) as the most important factors regulating growth and activity of soil microbial communities. However, some other studies found results similar to ours, finding that soil microbial community was mainly explained by soil factors (soil pH, soil N content, SOC and C:N), followed by vegetation factors (vegetation cover, plant productivity and species richness) and finally by climatic factors (Lauber et al., 2008; Fierer et al., 2009; Chen et al., 2016; Takriti et al., 2018). In our study, soil microbial response may be the result of climate-plant-soil-microorganisms' interactions, where rangelands with higher rainfall (humid) and vegetation cover or ANPP (Table 1), may increase organic matter mineralization (confirmed with high SBR values), and favor a higher microbial growth (MBC and MBN). This is consistent with previous study, where microbial biomass (bacterial and fungal) increased with SOC and soil N (Fierer et al., 2009; Wang et al., 2021). In particular, soil pH lead to reduced soil microbial biomass and activity as soil pH tends to become alkaline (from 5.7 to 8.4). Several studies have identified soil pH as a key environmental variable driving soil microbial communities (Fierer and Jackson, 2006; Sinsabaugh et al., 2008; Liu et al., 2022), where soil microbial biomass decreased with increasing soil pH (6–9) at both regional and continental scales (Fierer and Jackson 2006; Lauber et al. 2009; Rousk et al., 2010; Liu et al., 2022). In our rangelands, differences in soil pH (see, Table 1) can arise from many factors,

including vegetation type, soil characteristics, and climatic variables, which agrees with previous studies (Rengel, 2011; Liu et al., 2022). Thus, pH may serve as an integrating variable representing the physicochemical characteristics of a particular soil (Lauber et al., 2009). Regardless of the mechanism, our results demonstrated that pH was a reasonably good predictor of soil microbial variables (MBC, MBN, and SBR) at the regional scale (Fierer and Jackson, 2006; Hermans et al., 2022; Pan et al., 2023).

Moreover, our study showed the SBR was mainly explained by climatic factors (AI and MAP), followed by soil pH and vegetation cover. Zeng et al. (2013) and Chen et al. (2014) found that microbial activity increased with precipitation, soil N content and low pH associated with bacterial-dominated microbial communities. Differences in MBC:MBN ratio indicates shifts of microbial community composition (Cleveland and Liptzin, 2007). Jansson and Hofmockel (2020) reported that the shifts in the fungi:bacteria ratio was correlated with the MBC:MBN. For instance, in our humid sites, we determined lower values of MBC:MBN, which would suggest a dominance of bacteria in the soil. Conversely, increased MBC:MBN in the arid sites may result in fungal dominance (Sun et al., 2020b). Thus, soils with fungal networks are more stable under water limitations (e.g. arid and semiarid ecosystems) than bacterial communities (Fierer and Jackson, 2006; Barnard et al., 2015; Maestre et al., 2015; de Vries et al., 2018). In addition, results showed that the variation of soil microbial biomass and basal respiration along an aridity gradient in rangelands depends on particular microbial populations adapted to resource availability (soil nutrients and plant input) and habitat conditions (climate and other soil properties) (Fierer and Jackson, 2006; Bachar et al., 2010; Yao et al., 2017; Ding and Eldridge, 2022).

4.2. Responses of soil microorganisms to drought along the aridity gradient

In our study, the effects of drought on soil microbial community showed variable responses. While several studies reported that drought increased or decreased the soil microbial abundance and activity (Fuchslueger et al., 2014, 2016; Ochoa-Hueso et al., 2018), other studies determined no effect (Rousk et al., 2013; McHugh and Schwartz, 2015; Canarini et al., 2016; Hoover and Rogers, 2016). In the present work, we found at a regional level (considering eight rangeland sites) that MBC, MBN and MBC:MBN in the natural rangelands of Argentina were not affected by long-term (5 years) drought under field manipulative experiments. Nonetheless, an increase in SBR and qCO_2 was observed under drought plots. The direction and magnitude of soil microbial differential responses to drought might depend on environmental factors in each site (Ren et al., 2017; Schimel et al., 2021). This was demonstrated in our study (Fig. 4), where in the arid site (RM), microbial biomass (C and N) and SBR significantly decreased (20 - 30 %) with drought, while MBC:MBN increased. In addition, our results showed that drought modified the efficiency of microorganisms in these arid rangelands (determined with low qCO_2 values). These changes in microbial community could influence C use efficiency used by microbes for growth (Manzoni et al., 2012, 2014; Dijkstra et al., 2015). Thus, drought could directly inhibit microbial growth due to soil water reduction and indirectly by modifying plant and soil resources, as it was reported in other studies (Bardgett et al., 2008; Sanaullah et al., 2011; Manzoni et al., 2012; Ren et al., 2018). In addition, the reductions in aboveground biomass due to decreased rainfall can cause a decline in microbial biomass and activity through reductions in C and N from detritus entering the soil (Nielsen and Ball, 2015; Schrama and Bardgett, 2016; Canarini et al., 2017; Deng et al., 2021). This was supported in our study, where plants' response to drought was negative in the arid site (Table A.2 supplementary).

In our subhumid and humid sites (AI De Martonne ≥ 22) the responses of biomass and SBR to drought were less consistent. This may indicate that soil microbial community in humid ecosystems are less sensitive to drought (Manzoni et al., 2012; Ren et al., 2017, 2018; Zhou et al., 2018; Xu et al., 2020). Particularly, in the N site (subhumid) the response of microbial biomass to drought was variable; C remained constant, N decreased and C:N increased with drought (Fig. 3). Conversely, we found that drought had negative effects on MBC and MBC:MBN ratio in humid rangelands. Additionally, drought did not generate significant changes on SBR in our sites (Figs. 3 and A.4). The responses of soil microbiome to drought in these rangelands is possibly buffered by complex interactions with environmental factors. Thus, these interactions may enhance the responsiveness of microbial (C and N) turnover in soil under drought (Ren et al., 2017; Deng et al., 2021). Indeed, this response was supported by the regression analysis that indicated a positive relationship between SBR, MCB, M₂N and qCO₂ in our rangelands (Fig. A. 6 supplementary).

The decreased precipitation tended to increase or maintain the microbial biomass in semiarid sites compared to the other three rangeland ecosystems (arid, semihumid and humid). In contrast, drought stimulated a greater SBR in the semiarid sites (most evident in SPV and LC). Several studies have documented that responses to drought in semiarid ecosystems are very different and depend on the environment (Yang et al., 2021; Zhao et al., 2021). In our study, a possible explanation for this response in semiarid sites could be related to changes in the microorganism communities, as it was found in MBC:MBN (Figs. 3 and A.4). In the semiarid rangelands (PA and ChP sites), high microbial MBC:MBN may indicate fungal dominated soil microbiome

(Cleveland and Liptzin, 2007, Sun et al., 2020a), therefore, community structure change drive to better adaptation of soil microorganisms to drought.

Unexpectedly, the microbial biomass exhibited positive trends with drought in the sites SPV and LC (Fig. 3). This may reduce competition for nutrients from plants under drought, leading to higher soil nutrients and increased N and C availability for soil microorganisms (Dijkstra et al., 2015; Fuchslueger et al., 2016; Karlowsky et al., 2018a; Jaman et al., 2022). Drought RR analyses of aboveground biomass at these study sites may confirm these responses (Table A.2 supplementary). Further, higher microbial biomass (C and N) under drought (Figs. 3 and A.4) might have contributed to N and C resorption, which is considered as an essential strategy of nutrient conservation under stressed conditions (Six et al., 2006; Benner, 2011; Karlowsky et al., 2018b; Sun et al., 2020a).

Main results suggested that in humid rangelands, might lead to less soil microbial response to drought. However, in arid sites (AI <7) reduced precipitation exerted a negative impact on the soil microbial community. On the other hand, drought impacts on semiarid sites (AI from 10 to 17) are more variable on microbial attributes but could be very vulnerable if droughts intensify in duration and frequency during the next years due to climate change.

Further studies are needed to better explain the specific mechanisms linking plant biomass to microbial community abundance and activity related to above and below ground substrate availability (litter quantity and quality, root biomass or rhizoposition) (Zhang et al., 2018; Zhang and Xi, 2021; Jaman et al., 2022). The results from this study will improve our understanding of what the main environmental factors that drive the response microbial to drought in different types of rangelands and facilitate the development of predictive models on responses of soil

microorganisms in C cycles and vegetation shifts in rangelands across regional scales under global change scenarios.

4.3. Differential responses of plants and soil microorganisms to drought

Positive relationships between plants and soil microbial (biomass and activity) have widely been reported for several natural ecosystems (Paterson, 2003; van der Heijden et al., 2008; Sanaullah et al., 2011; Fuchslueger et al., 2016), indicating a close coupling between plants and microorganisms (Bardgett et al., 2005; Fierer et al., 2009; Karlowisky et al., 2018b). Our study showed clear evidence that SBR, MBC and MBN positively correlated with vegetation cover (Fig. 4) and aboveground biomass (Fig. A. 5 supplementary) after 5 years of drought treatments in contrasting rangelands. However, the impacts of drought on microorganisms and plants responded differently. Our findings revealed a trend of decoupling of plants and soil microbial community under precipitation reductions.

In the evaluated rangelands drought caused significant decrease in vegetation cover and aboveground biomass in most sites (Table A.2 supplementary; and Fig. 4).

Under drought stress plant growth is limited due to stomata closure (Pirasteh-Anosheh et al., 2016; Jaman et al., 2022) and often results in larger root systems and frequently increased C allocation to the roots (Liu et al., 2004; Karlowisky et al., 2018b) facilitating water and nutrient uptake (Huang and Gao, 2000; Liu and Li, 2005). Drought can also alter the quantity and quality of root exudates due to changes in plant species composition (Canarini et al., 2019), and legacy effects have been shown to cause a reduction of belowground C inputs (Fuchslueger et al., 2016). Although root biomass or plant C input were not the focus of this study, plant community dynamics and associated belowground C inputs are likely an important player in the response of microbial communities to drought. However, the response between plant and microorganisms under drought is bi-directional, where plants can

shape the soil community (de Vries et al., 2019) and in turn the soil microorganisms can have important consequences for the plant diversity, community composition and survival (Teste et al., 2017; Zhang and Xi, 2021). Therefore, it is not possible to conclude to which extent effects on the plant community could determine responses in soil microbial community or reciprocally.

Certainly, in our study we found that responses to drought in rangelands (arid, semiarid, semihumid, and humid) between microbial and vegetation variables (cover or aboveground biomass) differed in their magnitude (Figs 3 and A.5). This suggests that plant and microbial communities may respond differently to drought along aridity gradients. This highlights the importance of plant–microbial interactions under climate change scenarios due to a potential decoupling in the response of both groups.

Recently, two meta-analyses found that the asymmetric responses of plant biomass and soil microbial to precipitation change varied with climate conditions (Liu et al., 2016; Wilcox et al., 2017). In our study, we found that the microbial growth and activity showed a higher percentage change in response to drought in arid or semiarid sites, and the opposite trend in humid sites. The asymmetric responses and different change magnitudes between plants and microorganisms under drought stress are possibly related to the root biomass (e.g. exudation, rhizoposition) that buffered for the growth and activity of microorganisms under drought conditions (Rasse et al., 2005; Dijkstra and Cheng, 2007; Henry et al., 2007; Karlowsky et al., 2018a; Canarini et al., 2019; Williams and de Vries, 2020).

In addition, changing microbial activities in soils experiencing drought may indicate a faster microbial turnover given that drought might also have contributed to higher substrate inputs (labile microbial nutrients), generating a buffer effect that is compensated by the feedback of the microorganisms that die and those that remain

active under drought. In our study, the microbial community exposed to drought showed faster growth and respiration rates and lower qCO_2 values. A recent study found a large shift in the microbial community (bacteria, archaea and fungi), after 10 years of recurrent drought as an adaptation mechanism under water stress (Canarini et al., 2021). Previous studies have shown that microbial communities acclimatized to drought by enhancing osmolyte production, synthesis of capsules and exopolymeric substances to retain water, dormancy and sporulation (Sleator and Hill, 2002; Warren, 2014; Canarini et al., 2021), which may determine a more drought tolerant or resilient microbial community compared with plants responses.

Thus, long-term or more frequent exposure to drought events due to climate change can select a more tolerant soil microbial community that may influence the plant-microorganism interactions, as it was reported in other studies (Canarini et al., 2016; Sihi et al., 2018; Zhou et al., 2018; Zhang and Xi, 2021). The results of the present work can be used for improving the modelling of ecosystem C cycling and climate change impact on rangelands by incorporating microbial community variables into ecosystem process models (Luo et al., 2017; Deng et al., 2021).

5. CONCLUSION

Our findings provided new insights regarding the effect of main environmental factors (soil, vegetation and climate) on soil microbial response (biomass and basal respiration) in rangelands of South America. This study clearly demonstrated that soil microbial variables decreased in the aridity gradient from humid to arid. In this rangeland ecosystems, SOC and soil N contents and vegetation cover are the most important factors that regulate MBC and MBN. In addition, climate (MAP and aridity index) is the main variable which influences SBR. Moreover, the increase of soil pH negatively influences soil microbial variables in this rangeland ecosystems. Our results suggest that plant and microbial variables respond differently to drought along

aridity gradients, and reveal a trend towards a decoupling between plants and soil microorganisms to reduced precipitation. Therefore, our findings improve our understanding of the microorganism's response to the main environmental factors in different rangelands, and contributes to the inclusion of microbial variables in the predictive models of the carbon cycle under global change scenarios.

6. CRediT AUTHORSHIP CONTRIBUTION STATEMENT

Toledo, S.: Investigation; Methodology and performed the soil property analyses; Performed the data analyses; Writing original draft.

Bondaruk, V. F.: Collected the soil samples in the field; Writing-review and editing.

Yahdjian, L.: Conceptualization; Designed the experiments and coordinated the Project (Drought Net Argentina); Writing-review and editing.

Oñatibia, G R.: Collected the soil samples in the field; Writing-review and editing.

Loydi, A.: Collected the soil samples in the field; Writing-review and editing.

Alberti, J.: Collected the soil samples in the field; Writing-review and editing.

Bruschetti, M.: Collected the soil samples in the field; Writing-review and editing.

Pascual, J.: Collected the soil samples in the field; Writing-review and editing.

Peter, G.: Collected the soil samples in the field; Writing-review and editing.

Agüero, W. D.: Collected the soil samples in the field; Writing-review and editing.

Namur, P. R.: Collected the soil samples in the field; Writing-review and editing.

Blanco, L.: Collected the soil samples in the field; Writing-review and editing.

Peri, P. L.: Conceptualization; Investigation, Resources; Writing-review and editing.

T.S.; B.V.F.; Y.L.; O.G.R.; L.A.; A.J.; B.M.; P.J.; P.G.; A.W.D.; N.P.R.; B. L.; P.P.L.:

Drafted the manuscript, and all authors were involved in critical revision and approval of the final version.

7. ACKNOWLEDGEMENTS

We thank the cooperation of Morales Cristian, Ferrer Francisco, and Toledo

Guillermo for their help in the Laboratory.

8. FUNDING

This work was supported by the FONCyT agency, government of Argentina (grant numbers PICT 3659-2016).

9. CONFLICT OF INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

10. DATA AVAILABILITY STATEMENT

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

11. REFERENCES

- Anderson, T.H., Domsch, K.H., 1990. Application of eco-physiological quotients (qCO₂ and qD) on microbial biomasses from soils of different cropping histories. *Soil Biology and Biochemistry*, 22 (2), 251-255. [https://doi.org/10.1016/0038-0717\(90\)90094-G](https://doi.org/10.1016/0038-0717(90)90094-G).
- Anderson, D.L., Bonvissuto, G.L., Brizuela, M.A., Chiossone, G., Cibils, A.F., Cid, M.S., Feldman, I., Fernández-Grecco, R.C., Kunst, C., Oesterheld, M., Oliva, G., Paruelo, J.M., Peinetti, R., Villagra, E.S., 2011. Perspectives on rangeland management education and research in Argentina. *Rangelands*, 33 (1), 2-12. <https://doi.org/10.2111/RANGELANDS-D-10-00016.1>
- Bachar, A., Al-Ashhab, A., Soares, M.I.M., Sklarz, M.Y., Angel, R., Ungar, E.D., Gillor, O., 2010. Soil microbial abundance and diversity along a low precipitation gradient. *Microbial Ecology*, 60 (2), 453-461. <https://doi.org/10.1007/s00248-010-9727-1>.
- Bardgett, R.D., Bowman, W.D., Kaufmann, R., Schmidt, S.K., 2005. A temporal approach to linking aboveground and belowground ecology. *Trends in ecology & evolution*, 20 (11), 634-641. <https://doi.org/10.1016/j.tree.2005.08.005>.

- Bardgett, R.D., Freeman, C., Ostle, N.J., 2008. Microbial contributions to climate change through carbon cycle feedbacks. *The ISME journal*, 2 (8), 805-814. <https://doi.org/10.1038/ismej.2008.58>.
- Barnard, R.L., Osborne, C.A., Firestone, M.K., 2015. Changing precipitation pattern alters soil microbial community response to wet-up under a Mediterranean-type climate. *The ISME journal*, 9 (4), 946-957. <https://doi.org/10.1038/ismej.2014.192>.
- Beier, C., Beierkuhnlein, C., Wohlgemuth, T., Penuelas, J., Emmett, B., Körner, C., Hansen, K., 2012. Precipitation manipulation experiments—challenges and recommendations for the future. *Ecology letters*, 15 (8), 899-911. <https://doi.org/10.1111/j.1461-0248.2012.01793.x>.
- Benner, R., 2011. Biosequestration of carbon by heterotrophic microorganisms. *Nature Reviews Microbiology*, 9 (1), 75-75. <https://doi.org/10.1038/nrmicro2386-c3>.
- Blagodatskaya, E., Kuzyakov, Y., 2013. Active microorganisms in soil: critical review of estimation criteria and approaches. *Soil Biology and Biochemistry*, 67, 192-211. <https://doi.org/10.1016/j.soilbio.2013.08.024>.
- Bonan, G.B., Doney, S.C., 2018. Climate, ecosystems, and planetary futures: The challenge to predict life in Earth system models. *Science*, 359 (6375), eaam8328. DOI: 10.1126/science.aam8328.
- Bondaruk, V.F., Oñatibia, G.R., Fernández, R.J., Agüero, W., Blanco, L., Bruschetti, M., Kröpfl, A., Loydi, A., Pascual, J., Peri, P., Peter, G., Quiroga, R.E., Yahdjian, L., 2022. Forage provision is more affected by droughts in arid and semi-arid than in mesic rangelands. *Journal of Applied Ecology*, 59 (9), 2404-2418. <https://doi.org/10.1111/1365-2664.14243>.

- Brocca, L., Ciabatta, L., Massari, C., Moramarco, T., Hahn, S., Hasenauer, S., Levizzani, V., 2014. Soil as a natural rain gauge: Estimating global rainfall from satellite soil moisture data. *Journal of Geophysical Research: Atmospheres*, 119 (9), 5128-5141. <https://doi.org/10.1002/2014JD021489>.
- Canarini, A., Carrillo, Y., Mariotte, P., Ingram, L., Dijkstra, F.A., 2016. Soil microbial community resistance to drought and links to C stabilization in an Australian grassland. *Soil Biology and Biochemistry*, 103, 171-180. <https://doi.org/10.1016/j.soilbio.2016.08.024>.
- Canarini, A., Kiær, L.P., Dijkstra, F.A., 2017. Soil carbon loss regulated by drought intensity and available substrate: A meta-analysis. *Soil Biology and Biochemistry*, 112, 90-99. <https://doi.org/10.1016/j.soilbio.2017.04.020>.
- Canarini, A., Kaiser, C., Merchant, A., Richter, A., Wanek, W., 2019. Root exudation of primary metabolites: mechanisms and their roles in plant responses to environmental stimuli. *Frontiers in Plant Science*, 10, 157. <https://doi.org/10.3389/fpls.2019.00157>.
- Canarini, A., Schmidt, H., Fuchs-Sueger, L., Martin, V., Herbold, C.W., Zezula, D., Richter, A., 2021. Ecological memory of recurrent drought modifies soil processes via changes in soil microbial community. *Nature communications*, 12, 5308. <https://doi.org/10.1038/s41467-021-25675-4>.
- Chen, S., Zou, J., Hu, Z., Chen, H., Lu, Y., 2014. Global annual soil respiration in relation to climate, soil properties and vegetation characteristics: Summary of available data. *Agricultural and Forest Meteorology*, 198, 335-346. <https://doi.org/10.1016/j.agrformet.2014.08.020>.
- Chen, D., Mi, J., Chu, P., Cheng, J., Zhang, L., Pan, Q., Bai, Y., 2015. Patterns and drivers of soil microbial communities along a precipitation gradient on the

- Mongolian Plateau. *Landscape Ecology*, 30, 1669-1682.
<https://doi.org/10.1007/s10980-014-9996-z>.
- Chen, Y.L., Ding, J.Z., Peng, Y.F., Li, F., Yang, G.B., Liu, L., Yang, Y.H., 2016.
 Patterns and drivers of soil microbial communities in Tibetan alpine and global
 terrestrial ecosystems. *Journal of Biogeography*, 43 (10), 2027-2039. <https://doi.org/10.1111/jbi.12806>.
- De Martonne, E., 1926. Une nouvelle fonction climatologique: l' indice d' aridite.
Meteorologie (Paris), 2, 449-458.
- de Vries, F.T., Williams, A., Stringer, F., Willcocks, R., Waring, R., Langridge, H.,
 Straathof, A.L., 2019. Changes in root-exudate induced respiration reveal a
 novel mechanism through which drought affects ecosystem carbon cycling.
New Phytologist, 224 (1), 132-145. <https://doi.org/10.1111/nph.16001>.
- Delgado-Baquerizo, M., Maestre, F.T., Reich, P.B., Jeffries, T.C., Gaitan, J.J.,
 Encinar, D., Singh, B.K., 2018. Microbial diversity drives multifunctionality in
 terrestrial ecosystems. *Nature communications*, 7, 10541.
<https://doi.org/10.1038/ncomms10541>.
- Deng, L., Peng, C., Kim, D.G., Li, J., Liu, Y., Hai, X., Kuzyakov, Y., 2021. Drought
 effects on soil carbon and nitrogen dynamics in global natural ecosystems.
Earth-Science Reviews, 214, 103501.
<https://doi.org/10.1016/j.earscirev.2020.103501>.
- Dijkstra, F.A., Cheng, W., 2007. Moisture modulates rhizosphere effects on C
 decomposition in two different soil types. *Soil Biology and Biochemistry*, 39,
 2264-2274. <https://doi.org/10.1016/j.soilbio.2007.03.026>.
- Dijkstra, P., Salpas, E., Fairbanks, D., Miller, E.B., Hagerty, S.B., van Groenigen,
 K.J., Schwartz, E., 2015. High carbon use efficiency in soil microbial

- communities is related to balanced growth, not storage compound synthesis. *Soil Biology and Biochemistry*, 89, 35-43.
<https://doi.org/10.1016/j.soilbio.2015.06.021>.
- Ding, J., Eldridge, D.J., 2022. Drivers of soil biodiversity vary with organism type along an extensive aridity gradient. *Applied Soil Ecology*, 170, 104271.
<https://doi.org/10.1016/j.apsoil.2021.104271>.
- Fierer, N., Jackson, R.B., 2006. The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences*, 103 (3), 626-631. <https://doi.org/10.1073/pnas.0507535103>
- Fierer, N., Strickland, M.S., Liptzin, D., Bradford, M.A., Cleveland, C.C., 2009. Global patterns in belowground communities. *Ecology letters*, 12 (11), 1238-1249.
<https://doi.org/10.1111/j.1461-0248.2009.01360.x>.
- Fierer, N., Wood, S.A., de Mesquita, C.F.B., 2021. How microbes can, and cannot, be used to assess soil health. *Soil Biology and Biochemistry*, 153, 108111.
<https://doi.org/10.1016/j.soilbio.2020.108111>.
- Flombaum, P., Sala, O.E., 2007. A non-destructive and rapid method to estimate biomass and aboveground net primary production in arid environments. *Journal of Arid Environments*, 69 (2), 352-358.
<https://doi.org/10.1016/j.jaridenv.2006.09.008>
- Fuchslueger, L., Bahn, M., Fritz, K., Hasibeder, R., Richter, A., 2014. Experimental drought reduces the transfer of recently fixed plant carbon to soil microbes and alters the bacterial community composition in a mountain meadow. *New Phytologist*, 201 (3), 916-927. <https://doi.org/10.1111/nph.12569>.
- Fuchslueger, L., Bahn, M., Hasibeder, R., Kienzl, S., Fritz, K., Schmitt, M., Richter, A., 2016. Drought history affects grassland plant and microbial carbon

- turnover during and after a subsequent drought event. *Journal of Ecology*, 104 (5), 1453-1465. <https://doi.org/10.1111/1365-2745.12593>.
- Fuchslueger, L., Wild, B., Mooshammer, M., Takriti, M., Kienzl, S., Knoltsch, A., Richter, A., 2019. Microbial carbon and nitrogen cycling responses to drought and temperature in differently managed mountain grasslands. *Soil Biology and Biochemistry*, 135, 144-153. <https://doi.org/10.1016/j.soilbio.2019.05.002>.
- Gaitán, J.J., Bran, D.E., Oliva, G.E., Aguiar, M.R., Buono, G.G., Ferrante, D., Nakamatsu, V., Cian, G., Salomone, J.M., Massara, V., García Martínez, G., Maestre, F.T., 2018. Aridity and overgrazing have convergent effects on ecosystem structure and functioning in Patagonian rangelands. *Land Degradation & Development*, 29 (2), 210-218. <https://doi.org/10.1002/ldr.2694>.
- Gherardi, L.A., Sala, O.E., 2013. Automated rainfall manipulation system: A reliable and inexpensive tool for ecologists. *Ecosphere*, 4 (2), 1-10. <https://doi.org/10.1890/ES12-00371.1>.
- González, M.H., Garbarini, E.M., Fofa, A.L., Eslamian, S., 2017. Meteorological drought indices: Rainfall prediction in Argentina. In *Handbook of drought and water scarcity* (pp. 541-570), Edited by Saeid Eslamian, Faezeh A. Eslamian, CRC Press. <https://doi.org/10.1201/9781315404219>.
- Griffiths R.I., Thomson, B.C., James, P., Bell, T., Bailey, M., Whiteley, A.S., 2011. The bacterial biogeography of British soils. *Environ Microbiol*, 13, 1642-1654. <https://doi.org/10.1111/j.1462-2920.2011.02480.x>.
- Havstad, K.M., Peters, D.P.C., Skaggs, R., Brown, J., Bestelmeyer, B., Fredrickson, E., Herrick, J., Wright, J., 2007. Ecological services to and from rangelands of the United States. *Ecology and the Economy*, 64 (2), 261-268. <https://doi.org/10.1016/j.ecolecon.2007.08.005>.

- Hedges, L.V., Gurevitch, J., Curtis, P.S., 1999. The meta-analysis of response ratios in experimental ecology. *Ecology*, 80, 1150-1156.
[https://doi.org/10.1890/0012-9658\(1999\)080\[1150:TMAORR\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1150:TMAORR]2.0.CO;2).
- Henry, A., Doucette, W., Norton, J., Bugbee, B., 2007. Changes in crested wheatgrass root exudation caused by flood, drought, and nutrient stress. *J. Environ. Qual*, 36, 904-912. <https://doi.org/10.2134/jeq2006.0425sc>.
- Hermans, S.M., Buckley, H.L., Case, B.S., Curran-Cournane, F., Taylor, M., Lear, G., 2020. Using soil bacterial communities to predict physico-chemical variables and soil quality. *Microbiome*, 8, 1-13. <https://doi.org/10.1186/s40168-020-00858-1>.
- Homyak, P.M., Allison, S.D., Huxman, T.E., Goulden, M.L., Treseder, K.K., 2017. Effects of drought manipulation on soil nitrogen cycling: A meta-analysis. *Journal of Geophysical Research: Biogeosciences*, 122 (12), 3260-3272.
<https://doi.org/10.1002/2017JG004146>.
- Hoover, D.L., Rogers, B.M., 2016. Not all droughts are created equal: the impacts of interannual drought pattern and magnitude on grassland carbon cycling. *Global Change Biology*, 22 (5), 1809-1820. <https://doi.org/10.1111/gcb.13161>.
- Hothorn, T., Hornik, K., Strobl, C., Zeileis, A., 2008. Party: a Laboratory for recursive part(y)itioning: R package. v 1.3-1. R Packag version 09-0. URL <http://CRAN.R-project.org>.
- Huang, B.R., Gao, H.W., 2000. Root physiological characteristics associated with drought resistance in tall fescue cultivars. *Crop Sci*, 40, 196-203.
<https://doi.org/10.2135/cropsci2000.401196x>.

- Huang, J., Yu, H., Guan, X., Wang, G., Guo, R., 2016. Accelerated dryland expansion under climate change. *Nature Climate Change*, 6, 166-71. <https://doi.org/10.1038/nclimate2837>.
- IPCC, 2022. Central and South America. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 1689-1816. doi:10.1017/9781009325844.014.
- Jaman, M., Wu, H., Yu, Q., Tan, Q., Zhang, Y., Dang, Q.K., Luo, Y., 2022. Contrasting responses of plant above and belowground biomass carbon pools to extreme drought in six grasslands spanning an aridity gradient. *Plant and Soil*, 473 (1), 167-180. <https://doi.org/10.1007/s11104-021-05258-4>.
- Jansson, J.K., Hofmockel, K.S., 2020. Soil microbiomes and climate change. *Nature Reviews Microbiology*, 18 (1), 35-46. <https://doi.org/10.1038/s41579-019-0265-7>.
- Joergensen, R.G., Mueller, T., 1996. The fumigation extraction method to estimate soil microbial biomass: calibration of the kN-factor. *Soil Biology and Biochemistry*, 28, 33-37. [https://doi.org/10.1016/0038-0717\(95\)00102-6](https://doi.org/10.1016/0038-0717(95)00102-6).
- Karlowsky, S., Augusti, A., Ingrisch, J., Akanda, M.K.U., Bahn, M., Gleixner, G., 2018a. Drought-induced accumulation of root exudates supports post-drought recovery of microbes in mountain grassland. *Frontiers in Plant Science*, 9, 1593. <https://doi.org/10.3389/fpls.2018.01593>.

- Karlowsky, S., Augusti, A., Ingrisch, J., Hasibeder, R., Lange, M., Lavorel, S., Gleixner, G., 2018b. Land use in mountain grasslands alters drought response and recovery of carbon allocation and plant-microbial interactions. *Journal of Ecology*, 106 (3), 1230-1243. <https://doi.org/10.1111/13652745.12910>.
- Knapp, A.K., Carroll, C.J.W., Denton, E.M., La Pierre, K.J., Collins, S.L., Smith, M.D., 2015. Differential sensitivity to regional-scale drought in six central US grasslands. *Oecologia*, 177, 949-957. <https://doi.org/10.1007/s00442-015-3233-6>.
- Knapp, A.K., Avolio, M.L., Beier, C., Carroll, C.J.W., Collins, S.L., Dukes, J.S., Fraser, L.H., Griffin-Nolan, R.J., Hoover, D.L., Jentsch, A., Loik, M.E., Phillips, R.P., Post, A.K., Sala, O.E., Slette, I.J., Yehoujian, L., Smith, M.D., 2017. Pushing precipitation to the extremes in distributed experiments: recommendations for simulating wet and dry years. *Glob. Chang. Biol*, 23, 1774-1782. <https://doi.org/10.1111/gcb.13504>.
- Koricheva, J., Gurevitch, J., Menges, K., 2013. *Handbook of Meta-analysis in Ecology and Evolution*. Princeton University Press.
- Lau, J.A., Lennon, J.T., 2011. Evolutionary ecology of plant-microbe interactions: soil microbial structure alters selection on plant traits. *New Phytologist*, 192 (1), 215-224. <https://doi.org/10.1111/j.1469-8137.2011.03790.x>.
- Lauber, C.L., Strickland, M.S., Bradford, M.A., Fierer, N., 2008. The influence of soil properties on the structure of bacterial and fungal communities across land-use types. *Soil Biology and Biochemistry*, 40 (9), 2407-2415. <https://doi.org/10.1016/j.soilbio.2008.05.021>.
- Lemoine, N.P., Sheffield, J., Dukes, J.S., Knapp, A.K., Smith, M.D., 2016. Terrestrial precipitation analysis (TPA): A resource for characterizing long-term

- precipitation regimes and extremes. *Methods in Ecology and Evolution*, 7 (11), 1396-1401. <https://doi.org/10.1111/2041-210X.12582>.
- Li, C., Fultz, L.M., Moore-Kucera, J., Acosta-Martínez, V., Kakarla, M., Weindorf, D.C., 2018. Soil microbial community restoration in Conservation Reserve Program semi-arid grasslands. *Soil Biology and Biochemistry*, 118, 166-177. <https://doi.org/10.1016/j.soilbio.2017.12.001>
- Liang, C., Schimel, J.P., Jastrow, J.D., 2017. The importance of anabolism in microbial control over soil carbon storage. *Nature microbiology*, 2 (8), 1-6. <https://doi.org/10.1038/nmicrobiol.2017.105>.
- Liang, C., Amelung, W., Lehmann, J., Kästner, M., 2019. Quantitative assessment of microbial necromass contribution to soil organic matter. *Global change biology*, 25 (11), 3578-3590. <https://doi.org/10.1111/gcb.14781>.
- Lin, L., Gettelman, A., Feng, S., Fu, C., 2015. Simulated climatology and evolution of aridity in the 21st century. *J. Geophys. Res. Atmos.*, 120, 5795-815. <https://doi.org/10.1002/2014JD022912>.
- Liu, H.S., Li, F.M., Xu, H., 2014. Deficiency of water can enhance root respiration rate of drought-sensitive but not drought-tolerant spring wheat. *Agric. Water Manage*, 64, 41-46. [https://doi.org/10.1016/S0378-3774\(03\)00143-4](https://doi.org/10.1016/S0378-3774(03)00143-4).
- Liu, H.S., Li, F.M., 2005. Root respiration, photosynthesis and grain yield of two spring wheat in response to soil drying. *Plant Growth Regul*, 46, 233-240. <https://doi.org/10.1007/s10725-005-8806-7>.
- Liu, L., Wang, X., Lajeunesse, M.J., Miao, G., Piao, S., Wan, S., Deng, M., 2016. A cross-biome synthesis of soil respiration and its determinants under simulated precipitation changes. *Global change biology*, 22 (4), 1394-1405. <https://doi.org/10.1111/gcb.13156>.

- Liu, K., Liu, Z., Li, X., Shi, X.R., Lock, T.R., Kallenbach, R.L., Yuan, Z.Y., 2022. Precipitation-mediated responses of plant biomass production and allocation to changing soil pH in semiarid grasslands. *Agriculture, Ecosystems & Environment*, 339, 108123. <https://doi.org/10.1016/j.agee.2022.108123>.
- Luo, Z., Feng, W., Luo, Y., Baldock, J., Wang, E., 2017. Soil organic carbon dynamics jointly controlled by climate, carbon inputs, soil properties and soil carbon fractions. *Global Change Biology*, 23 (10), 4430-4439. <https://doi.org/10.1111/gcb.13767>.
- Ma, L., Guo, C., Lü, X., Yuan, S., Wang, R., 2015. Soil moisture and land use are major determinants of soil microbial community composition and biomass at a regional scale in northeastern China. *Biogeochemistry*, 12 (8), 2585-2596. <https://doi.org/10.5194/bg-12-2585-2015>.
- Maestre, F.T., Delgado-Baquerizo, M., Jeffries, T.C., Eldridge, D.J., Ochoa, V., Gozalo, B., Singh, B.K., 2015. Increasing aridity reduces soil microbial diversity and abundance in global drylands. *Proceedings of the National Academy of Sciences*, 112 (51), 15684-15689. <https://doi.org/10.1073/pnas.1516684112>.
- Manzoni, S., Schimek, I.F., Porporato, A., 2012. Responses of soil microbial communities to water stress: results from a meta-analysis. *Ecology*, 93, 930-938. <https://doi.org/10.1890/11-0026.1>.
- Marcos, M.S., Olivera, N.L., 2016. Microbiological and Biochemical Indicators for Assessing Soil Quality in Drylands from Patagonia. In: Olivera, N., Libkind, D., Donati, E. (Eds.) *Biology and Biotechnology of Patagonian Microorganisms*. Springer, Cham. https://doi.org/10.1007/978-3-319-42801-7_6.

- McHugh, T.A., Schwartz, E., 2015. Changes in plant community composition and reduced precipitation have limited effects on the structure of soil bacterial and fungal communities present in a semiarid grassland. *Plant and Soil*, 388, 175-186. <https://doi.org/10.1007/s11104-014-2269-4>.
- Middleton, N., 2018. Rangeland management and climate hazards in drylands: dust storms, desertification and the overgrazing debate. *Natural hazards*, 92 (1), 57-70. <https://doi.org/10.1007/s11069-016-2592-6>.
- Montecchia, M.S., Correa, O.S., Soria, M.A., Frey, S.D., García, A.F., Garland, J.L., 2011. Multivariate approach to characterizing soil microbial communities in pristine and agricultural sites in Northwest Argentina. *Applied soil ecology*, 47 (3), 176-183. <https://doi.org/10.1016/j.apsoil.2010.12.008>.
- Nannipieri, P., Ascher, J., Ceccherini, M., Landi, L., Pietramellara, G., Renella, G., 2003. Microbial diversity and soil functions. *European journal of soil science*, 54 (4), 655-670. <https://doi.org/10.1046/j.1351-0754.2003.0556.x>.
- Nielsen, U.N., Ball, B.A., 2015. Impacts of altered precipitation regimes on soil communities and biogeochemistry in arid and semi-arid ecosystems. *Global change biology*, 21 (1), 1407-1421. <https://doi.org/10.1016/j.apsoil.2010.12.008>.
- Ochoa-Hueso, R., Collins, S.L., Delgado-Baquerizo, M., Hamonts, K., Pockman, W.T., Sinsabaugh, R.L., Power, S.A., 2018. Drought consistently alters the composition of soil fungal and bacterial communities in grasslands from two continents. *Global change biology*, 24 (7), 2818-2827. <https://doi.org/10.1111/gcb.14113>.
- Olivera, N.L., Prieto, L., Bertiller, M.B., Ferrero, M.A., 2016. Sheep grazing and soil bacterial diversity in shrublands of the Patagonian Monte, Argentina. *Journal*

- of Arid Environments, 125, 16-20.
<https://doi.org/10.1016/j.jaridenv.2015.09.012>.
- Oñatibia, G.R., Amengual, G., Boyero, L., Aguiar, M.R., 2020. Aridity exacerbates grazing-induced rangeland degradation: A population approach for dominant grasses. *Journal of Applied Ecology*, 57, 1999-2009.
<https://doi.org/10.1111/1365-2664.13704>.
- Oyarzabal, M., Clavijo, J., Oakley, L., Biganzoli, F., Tognetti, P., Barberis, I., Maturo, H.M., Aragón, R., Campanello, P.I., Prado, D., Oesterheld, M., León, R.J.C., 2018. Unidades de vegetación de la Argentina. *Ecología Austral*, 28 (1), 40-63. <https://doi.org/10.25260/ea.18.28.1.0.399>.
- Pan, J., Peng, Y., Wang, J., Tian, D., Zhang, R., Li, Y., Niu, S., 2023. Controlling factors for soil bacterial and fungal diversity and composition vary with vegetation types in alpine grasslands. *Applied Soil Ecology*, 184, 104777.
<https://doi.org/10.1016/j.apsoil.2022.104777>.
- Pasternak, Z., Al-Ashhab, A., Gatica, J., Gafny, R., Avraham, S., Minz, D., Jurkevitch, E., 2013. Spatial and temporal biogeography of soil microbial communities in arid and semiarid regions. *PLoS One*, 8 (7), e69705.
<https://doi.org/10.1371/journal.pone.0069705>.
- Paterson, E., 2003. Importance of rhizodeposition in the coupling of plant and microbial productivity. *European journal of soil science*, 54 (4), 741-750.
<https://doi.org/10.1046/j.1351-0754.2003.0557.x>.
- Pirasteh-Anosheh, H., Saed-Moucheshi, A., Pakniyat, H., Pessarakli, M., 2016. Stomatal responses to drought stress. *Water stress and crop plants: A sustainable approach*, 1, 24-40. <https://doi.org/10.1002/9781119054450.ch3>.

- Prieto, L.H., Bertiller, M.B., Carrera, A.L., Olivera, N.L., 2011. Soil enzyme and microbial activities in a grazing ecosystem of Patagonian Monte, Argentina. *Geoderma*, 162 (3-4), 281-287. <https://doi.org/10.1016/j.geoderma.2011.02.011>.
- Rasse, D.P., Rumpel, C., Dignac, M.F., 2005. Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation. *Plant and Soil*, 269, 341-356. <https://doi.org/10.1007/s11104-004-0907-y>.
- Ren, C., Zhao, F., Shi, Z., Chen, J., Han, X., Yang, G., Ren, G., 2017. Differential responses of soil microbial biomass and carbon-degrading enzyme activities to altered precipitation. *Soil Biology and Biochemistry*, 115, 1-10. <https://doi.org/10.1016/j.soilbio.2017.08.002>.
- Ren, C., Chen, J., Lu, X., Doughty, R., Zhao, F., Zhong, Z., Ren, G., 2018. Responses of soil total microbial biomass and community compositions to rainfall reductions. *Soil Biology and Biochemistry*, 116, 4-10. <https://doi.org/10.1016/j.soilbio.2017.09.028>.
- Rengel, Z., 2011. Soil pH, Soil Health and Climate Change. In: Singh, B., Cowie, A., Chan, K. (Eds) *Soil Health and Climate Change*. *Soil Biology*, Vol 29. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-20256-8_4.
- Robertson, G.P., Wedin, D., Groffmann, P.M., Blair, J.M., Holland, E.A., Nadelhoffer, K.J., Harris, D., 1999. Soil carbon and nitrogen availability: nitrogen mineralization, nitrification, and soil respiration potentials. In *Standard soil methods for long-term ecological research* (pp. 258-271).
- Rousk, J., Brookes, P.C., Bååth, E., 2010. The microbial PLFA composition as affected by pH in an arable soil. *Soil Biology and Biochemistry*, 42 (3), 516-520. <https://doi.org/10.1016/j.soilbio.2009.11.026>.

- Rousk, J., Smith, A.R., Jones, D.L., 2013. Investigating the long-term legacy of drought and warming on the soil microbial community across five European shrubland ecosystems. *Global Change Biology*, 19, 3872-3884. <https://doi.org/10.1111/gcb.12338>.
- Saccá, M.L., Barra Caracciolo, A., Di Lenola, M., Grenni, P., 2017. Ecosystem Services Provided By Soil Microorganisms. In: Lukac, M., Grenni, P., Gamboni, M. (Eds) *Soil Biological Communities and Ecosystem Resilience. Sustainability in Plant and Crop Protection*. Springer Cham. https://doi.org/10.1007/978-3-319-63336-7_2.
- Sala, O.E., Austin, A.T., 2000. Methods of estimating aboveground net primary productivity. In *Methods in Ecosystem Science* (pp. 31-43). https://doi.org/10.1007/978-1-4612-1224-9_3.
- Sala, O.E., Yahdjian, L., Havstad, K., Acuar, M.R., 2017. Rangeland ecosystem services: Nature's supply and humans' demand. https://doi.org/10.1007/978-3-319-46709-2_14.
- Sanaullah, M., Blagodatskaya, E., Chabbi, A., Rumpel, C., Kuzyakov, Y., 2011. Drought effects on microbial biomass and enzyme activities in the rhizosphere of grasses depend on plant community composition. *Applied Soil Ecology*, 48 (1), 38-44. <https://doi.org/10.1016/j.apsoil.2011.02.004>.
- Schimel, J., Balsler, T.C., Wallenstein, M., 2007. Microbial stress-response physiology and its implications for ecosystem function. *Ecology*, 88 (6), 1386-1394. <https://doi.org/10.1890/06-0219>.
- Schrama, M., Bardgett, R.D., 2016. Grassland invasibility varies with drought effects on soil functioning. *Journal of Ecology*, 104 (5), 1250-1258. <https://doi.org/10.1111/1365-2745.12606>.

- Shen, R.C., Xu, M., Li, R.Q., Zhao, F.X., Sheng, Q.K., 2015. Spatial variability of soil microbial biomass and its relationships with edaphic, vegetational and climatic factors in the Three-River Headwaters region on Qinghai-Tibetan Plateau. *Applied Soil Ecology*, 95, 191-203.
<https://doi.org/10.1016/j.apsoil.2015.06.011>.
- Sihi, D., Davidson, E.A., Chen, M., Savage, K.E., Richardson, A.D., Keenan, T.F., Hollinger, D.Y., 2018. Merging a mechanistic enzymatic model of soil heterotrophic respiration into an ecosystem model in two AmeriFlux sites of northeastern USA. *Agricultural and Forest Meteorology*, 252, 155-166.
<https://doi.org/10.1016/j.agrformet.2018.01.023>
- Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw, C., Zeglin, L.H., 2008. Stoichiometry of soil enzyme activity at global scale. *Ecology letters*, 11 (11), 1252-1264.
<https://doi.org/10.1111/j.1461-0248.2008.01245.x>.
- Six, J., Frey, S.D., Thiet, R.K., Parton, K.M., 2006. Bacterial and fungal contributions to carbon sequestration in agroecosystems. *Soil Science Society of America Journal*, 70 (2), 555-569. <https://doi.org/10.2136/sssaj2004.0347>.
- Sleator, R.D., Hill, C., 2002. Bacterial osmoadaptation: the role of osmolytes in bacterial stress and virulence. *FEMS Microbiol. Rev.*, 26, 49-71.
<https://doi.org/10.1111/j.1574-6976.2002.tb00598.x>.
- Soil Survey Staff, 2010. *Keys to Soil Taxonomy*, 11th ed. USDA-Natural Resources Conservation Service, Washington, DC.
- Stomeo, F., Valverde, A., Pointing, S.B., McKay, C.P., Warren-Rhodes, K.A., Tuffin, M.I., Cowan, D.A., 2013. Hypolithic and soil microbial community assembly

- along an aridity gradient in the Namib Desert. *Extremophiles*, 17 (2), 329-337.
<https://doi.org/10.1007/s00792-013-0519-7>.
- Strobl, C., Malley, J., Tutz, G., 2009. Characteristics of classification and regression trees, bagging and random forests. *Psychol Methods*, 14, 323-348.
<https://doi.org/10.1037/a0016973.An>.
- Sun, Y., Liao, J., Zou, X., Xu, X., Yang, J., Chen, H.Y., Ruan, H., 2020a. Coherent responses of terrestrial C: N stoichiometry to drought across plants, soil, and microorganisms in forests and grasslands. *Agricultural and Forest Meteorology*, 292, 108104. <https://doi.org/10.1016/j.agrformet.2020.108104>.
- Sun, Y., Chen, H.Y., Jin, L., Wang, C., Zhang, R., Ruan, H., Yang, J., 2020b. Drought stress induced increase of fungi: bacteria ratio in a poplar plantation. *Catena*, 193, 104607. <https://doi.org/10.1016/j.catena.2020.104607>.
- Takriti, M., Wild, B., Schneckner, J., Mooshammer, M., Knoltsch, A., Lashchinskiy, N., Richter, A., 2018. Soil organic matter quality exerts a stronger control than stoichiometry on microbial substrate use efficiency along a latitudinal transect. *Soil Biology and Biochemistry*, 121, 212-220.
<https://doi.org/10.1016/j.soilbio.2018.02.022>.
- Teste, F.P., Kardol, P., Turner, B.L., Wardle, D.A., Zemunik, G., Renton, M., Laliberté, E., 2017. Plant-soil feedback and the maintenance of diversity in Mediterranean-climate shrublands. *Science*, 355(6321), 173-176. DOI: 10.1126/science.aai8291.
- Thakur, M.P., Milcu, A., Manning, P., Niklaus, P.A., Roscher, C., Power, S., Eisenhauer, N., 2015. Plant diversity drives soil microbial biomass carbon in grasslands irrespective of global environmental change factors. *Global Change Biology*, 21 (11), 4076-4085. <https://doi.org/10.1111/gcb.13011>.

- Toledo, S., Peri, P.L., Correa, O.S., Gargaglione, V., Gonzalez-Polo, M., 2021. Soil microbial communities respond to an environmental gradient of grazing intensity in South Patagonia Argentina. *Journal Arid Environ*, 184, 104300. <https://doi.org/10.1016/j.jaridenv.2020.104300>.
- Trenberth, K.E., Dai, A., van der Schrier, G., Jones, P.D., Barichivich, J., Briffa, K.R., Sheffield, J., 2014. Global warming and changes in drought. *Nature Clim. Change*, 4, 17-22. <https://doi.org/10.1038/nclimate2067>.
- Van Der Heijden, M.G., Bardgett, R.D., Van Straalen, N.M., 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology letters*, 11 (3), 296-310. <https://doi.org/10.1111/j.1461-0248.2007.01139.x>.
- Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil microbial biomass C. *Soil biology and Biochemistry*, 19 (6), 703-707. [https://doi.org/10.1016/0038-0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6).
- Verón, S.R., Blanco, L.J., Texeira, M.A., Irisarri, J.G.N., Paruelo, J.M., 2018. Desertification and ecosystem services supply: The case of the Arid Chaco of South America. *Journal of Arid Environments*, 159, 66-74. <https://doi.org/10.1016/j.jaridenv.2017.11.001>.
- Viruel, E., Fontana, C.A., Puglisi, E., Nasca, J.A., Banegas, N.R., Cocconcelli, P.S., 2022. Land-use change affects the diversity and functionality of soil bacterial communities in semi-arid Chaco region, Argentina. *Applied Soil Ecology*, 172, 104362. <https://doi.org/10.1016/j.apsoil.2021.104362>.
- Waldrop, M.P., Holloway, J.M., Smith, D.B., Goldhaber, M.B., Drenovsky, R.E., Scow, K.M., Grace, J.B., 2017. The interacting roles of climate, soils, and plant

- production on soil microbial communities at a continental scale. *Ecology*, 98 (7), 1957-1967. <https://doi.org/10.1002/ecy.1883>.
- Wang, S., Zuo, X., Awada, T., Medima-Roldan, E., Feng, K., Yue, P., Cheng, H., 2021. Changes of soil bacterial and fungal community structure along a natural aridity gradient in desert grassland ecosystems, Inner Mongolia. *Catena*, 205, 105470. <https://doi.org/10.1016/j.catena.2021.105470>.
- Warren, C.R., 2014. Response of osmolytes in soil to drying and rewetting. *Soil Biology and Biochemistry*, 70, 22-32. <https://doi.org/10.1016/j.soilbio.2013.12.008>.
- Wilcox, K.R., Shi, Z., Gherardi, L.A., Lemoine, N.P., Koerner, S.E., Hoover, D.L., Luo, Y., 2017. Asymmetric responses of primary productivity to precipitation extremes: A synthesis of grassland precipitation manipulation experiments. *Global Change Biology*, 23, 4376-4385. <https://doi.org/10.1111/gcb.13706>.
- Williams, A., de Vries, F.T., 2020. Plant root exudation under drought: implications for ecosystem functioning. *New Phytologist*, 225 (5), 1899-1905. <https://doi.org/10.1111/nph.16223>.
- Xu, S., Geng, W., Sayer, E.J., Zhou, G., Zhou, P., Liu, C., 2020. Soil microbial biomass and community responses to experimental precipitation change: A meta-analysis. *Soil Ecology Letters*, 2 (2), 93-103. <https://doi.org/10.1007/s42832-020-0033-7>.
- Yahdjian, L., Sala, O.E., 2002. A rainout shelter design for intercepting different amounts of rainfall. *Oecologia*, 133 (2), 95-101. <https://doi.org/10.1007/s00442-002-1024-3>.

- Yahdjian, L., Sala, O.E., 2008. Climate change impacts on South American rangelands. *Rangelands*, 30 (3), 34-39. [https://doi.org/10.2111/1551-501X\(2008\)30\[34:CCIOSA\]2.0.CO;2](https://doi.org/10.2111/1551-501X(2008)30[34:CCIOSA]2.0.CO;2).
- Yahdjian, L., Sala, O.E., Havstad, K.M., 2015. Rangeland ecosystem services: Shifting focus from supply to reconciling supply and demand. *Front. Ecol. Environ.*, 13, 44-51. <https://doi.org/10.1890/140156>.
- Yang, Y., Li, T., Wang, Y., Cheng, H., Chang, S.X., Liang, C., An, S., 2021. Negative effects of multiple global change factors on soil microbial diversity. *Soil Biology and Biochemistry*, 156, 108229. <https://doi.org/10.1016/j.soilbio.2021.108229>.
- Yao, M., Rui, J., Niu, H., Heděnc, P., Li, J., He, Z., Li, X., 2017. The differentiation of soil bacterial communities along a precipitation and temperature gradient in the eastern Inner Mongolia steppe. *Catena*, 152, 47-56. <https://doi.org/10.1016/j.catena.2017.01.007>.
- Zhang, L., Xie, Z., Zhao, R., Zhang, Y., 2018. Plant, microbial community and soil property responses to an experimental precipitation gradient in a desert grassland. *Applied Soil Ecology*, 127, 87-95. <https://doi.org/10.1016/j.apsoil.2018.02.005>.
- Zhang, C., Xi, N., 2021. Precipitation changes regulate plant and soil microbial biomass via plasticity in plant biomass allocation in grasslands: a meta-analysis. *Frontiers in plant science*, 12, 614968. <https://doi.org/10.3389/fpls.2021.614968>.
- Zhao, Y., Liu, Z., Wu, J., 2020. Grassland ecosystem services: a systematic review of research advances and future directions. *Landscape Ecology*, 35 (4), 793-814. <https://doi.org/10.1007/s10980-020-00980-3>.

- Zhao, M., Guo, S., Wang, R., 2021. Diverse soil respiration responses to extreme precipitation patterns in arid and semiarid ecosystems. *Applied Soil Ecology*, 163, 103928. <https://doi.org/10.1016/j.apsoil.2021.103928>.
- Zhou, X., Chen, C., Wang, Y., Xu, Z., Duan, J., Hao, Y., Smaill, S., 2013. Soil extractable carbon and nitrogen, microbial biomass and microbial metabolic activity in response to warming and increased precipitation in a semiarid Inner Mongolian grassland. *Geoderma*, 206, 24-31. <https://doi.org/10.1016/j.geoderma.2013.04.020>.
- Zhou, Z., Wang, C., Luo, Y., 2018. Response of soil microbial communities to altered precipitation: A global synthesis. *Global Ecology and Biogeography*, 27 (9), 1121-1136. <https://doi.org/10.1111/geb.12761>.

FIGURES AND TABLES

Figure 1. Relative importance values of independent environmental variables used to characterize the studied rangeland sites in Argentina, resulting from the random forest analysis. Environmental variables representing climate were: mean annual precipitation and temperature (MAP and MAT), and aridity index (AI); the soil: soil pH, soil organic carbon (SOC), nitrogen (N) content, phosphorus (P) content, and C:N ratio; and the vegetation: vegetation cover, aboveground net primary production (ANPP), and plant species richness, for the eight sites and microbial variables microbial biomass carbon (MBC); microbial biomass nitrogen (MBN); and soil biological respiration (SBR).

Figure 2. Relationships between microbial variables and the most important soil, vegetation, and climatic variables selected by the random forest model/analysis, in eight rangelands of Argentina. Microbial biomass carbon (MBC) as a function of A)

Soil pH, B) Soil C:N ratio, and C) Vegetation cover. Microbial biomass nitrogen (MBN) as a function of D) soil pH, E) soil N content, and F) SOC. Soil biological respiration (SBR) as a function of G) soil pH, H) mean annual precipitation (MAP), and I) aridity index (AI). Experimental sites were (from arid to humid): Rio Mayo (RM), San Pablo Valdes (SPV), Los Cerrillos (LC), Potrok Aike (PA), Chacra de Patagones (ChP), Naposta (N), San Claudio (SC) and Las Chilcas (LCh). Sites are depicted as arid (circles), semiarid (squares), semihumid (triangles), and humid (rhombuses).

Figure 3. Drought effects (estimated by the Response Ratio, RR) on soil microbial variables (biomass, activity, and microbial indexes) in eight rangelands of Argentina. The lines/whiskers indicated the 95 % confidence interval (CI). Significant treatment effects were found when the error bars did not overlap with zero. Sites were Rio Mayo (RM), San Pablo Valdes (SPV), Potrok Aike (PA), Chacra de Patagones (ChP), Los Cerrillos (LC), Naposta (N), Las Chilcas (LCh), and San Claudio (SC). Circles represent arid sites, squares semiarid sites, triangles semihumid sites, and rhombuses humid sites.

Figure 4. Relationships between drought response ratio of total vegetation cover and microbial variables: A) microbial biomass carbon (MBC), B) microbial biomass nitrogen (MBN), C) MBC:MBN ratio, and D) soil biological respiration (SBR). Experimental sites are (from arid to humid): Rio Mayo (RM), San Pablo Valdes (SPV), Los Cerrillos (LC), Potrok Aike (PA), Chacra de Patagones (ChP), Naposta (N), Las Chilcas (LCh), and San Claudio (SC). Sites are depicted as arid (circles), semiarid (squares), semihumid (triangles), and humid (rhombuses).




Fig. 1

Journal Pre-proof




Fig. 2




Fig. 3




Fig. 4

Table 1. Characterization of the study sites in an aridity gradient in Argentina. Geographic location, description of climatic variables, vegetation, and soil physicochemical characteristics of the eight sites are shown from the most arid to the most humid. Values are mean and standard deviation (\pm SD). Values of soil variables followed by different lowercase letters in each file indicate significant differences ($p < 0.05$) among sites.

	Río Mayo (RM)	San Pablo Valdes (SPV)	Potro k Aike (PA)	Los Cerrillos (LC)	Chacra Patagones (ChP)	Napostá ^(a) (N)	Las Chilcas (LCh)	San Claudio (SC)
Coordinates								
Latitude (S)	45°23' 51.72"	42°39' 12.04"	51°54' 57.60"	29°57' 01.08"	40°43'2 3.16"	38°25' 26.04"	36° 09'47.5 2"	35°54' 55.44"
Longitude (W)	70°18' 21.24"	64°10' 17.08"	70°24' 26.64"	65°52' 24.60"	62°53'5 6.04"	62°17' 16.44"	58° 09'20.1 6"	61°08' 54.24"
Climatic								
MAP (mm yr ⁻¹)	137.1± 44.7	249.3± 83.1	186.6± 32.7	372.9± 71.6	144.2± 57.6	579.4 ±63.4	998.7± 152.9	1022.3 ±233.5
MAT (°C)	9.6±0.3	13.9±0.3	6.5±0.3	22±1	17.4±2	15.4± 0.6	15.4±0.4	15.1±0.3
Aridity Index De Martonne (MAP/MAT+10)	7	10.4	11.3	11.7	16.2	22.9	39.3	40.8
Aridity categories	Arid	Semiarid	Semiarid	Semiarid	Semiarid	Semi Humid	Humid	Humid
Vegetation								
Vegetation types	Grass steppe: grasses and dwarf shrubs	Grass steppe: grasses and dwarf shrubs	Grass steppe: grasses and dwarf shrubs	Grass-shrub steppe	Grassland	Grassland	Grassland	Grassland
ANPP (g m ⁻² yr ⁻¹)	29.01c	160.8cd	45.56e	268.99c	103.80d	347.72c	583.45b	1219.01a
Vegetation cover (%)	55.8c	60.1b	74.7b	94a	71.7b	94.7a	97.8a	96.3a
Species richness (α diversity)	6	11	19	12	5	6	16	4
Dominant species	<i>Mulinum spinosum</i> ; <i>Pappostipa speciosa</i> ; <i>Poa ligularis</i>	<i>Nassella tenuis</i> ; <i>Poa lanuginosa</i> ; <i>Sporobolus rigens</i> ; <i>Vulpia bromoides</i>	<i>Carex andina</i> ; <i>Nardophyllum bryoides</i> ; <i>Poa spiciformis</i> ; <i>Pappostipa speciosa</i>	<i>Larrea divaricata</i> ; <i>Pappophorum krapovickasii</i> ; <i>Aristida mendocina</i>	<i>Nassella longiglumis</i> ; <i>Poa ligularis</i>	<i>Bromus catharticus</i> ; <i>Nassella nessiana</i> ; <i>Nassella trichotoma</i>	<i>Festuca arundinacea</i> ; <i>Leersia hexandra</i> ; <i>Panicum gouninii</i> ; <i>Paspalum paludivagum</i>	<i>Festuca arundinacea</i> ; <i>Sorghum halepense</i>
Soil								
Soil Taxonomy	Aridisol	Entisol	Aridisol	Aridisol	Entisols	Mollis	Alfisols	Molliso

USDA ^(b)	s	s	s	s		<i>o/s</i>		<i>ls</i>
Textural class	Sandy	<i>Sandy</i>	Sandy loam	Sandy loam	Sandy clay loam	Sandy loam	Silt loam	Loam
pH	7.4c	8.0b	6.2d	8.4a	7.6bc	6.0de	5.7e	6.1de
SOC (g Kg ⁻¹)	5.35c	7.37c	24.74b	4.03c	25.37b	35.40 a	40.40a	21.71b
N (g Kg ⁻¹)	0.52e	1.13d	3.03b	0.55e	2.83b	3.34a b	3.57a	1.70c
P extractable (ppm)	10.1b	3.5c	28.0a	6.5bc	8.3b	25.0a	24.6a	11.1b
C:N	10.3b	6.5d	8.2c	7.3d	8.9c	10.6b	11.3a	12.8a

(a) Napostá: Campo Experimental Napostá; convenio UNS y MDA-PBA; **(b)** Soil Survey Staff, 2010

Journal Pre-proof

Table 2. Soil biological variables and microbial indexes for the eight sites and sites grouped by aridity in rangelands of Argentina. Mean values and standard deviation (\pm SD) are shown. Microbial biomass carbon and nitrogen: MBC and MBN (mg kg^{-1} dry soil), MBC:MBN ratio, soil biological respiration: SBR ($\text{mg C-CO}_2 \text{ kg}^{-1}$ dry soil d^{-1}), metabolic quotient ($q\text{CO}_2$) ($\mu\text{gC-CO}_2 \text{ mg}^{-1} \text{MBC d}^{-1}$) and microbial carbon quotient: $q\text{MC}$ (%). Values followed by different letters in each column indicate significant differences ($p < 0.05$) among aridity categories (uppercase letters) and sites (lowercase letters).

Microbial variables	MBC	MBN	SBR	MBC:MBN	$q\text{CO}_2$	$q\text{MC}$
Aridity categories ^(a)						
Arid	122 \pm 35D	22 \pm 8C	29 \pm 7D	6 \pm 2A	0.24 \pm 0.03A	2.4 \pm 0.7A
Semiarid	252 \pm 143C	43 \pm 22B	40 \pm 12C	6 \pm 3A	0.21 \pm 0.12A	2.2 \pm 0.6A
Semihumid	425 \pm 43B	148 \pm 37A	50 \pm 9B	3 \pm 1B	0.12 \pm 0.02B	1.3 \pm 0.1B
Humid	557 \pm 56A	116 \pm 49A	74 \pm 10A	6 \pm 2A	0.13 \pm 0.03B	2.0 \pm 0.7A
All sites ^(b)						
RM	122 \pm 35e	22 \pm 8c	29 \pm 7ef	6 \pm 2bcd	0.24 \pm 0.03a	2.4 \pm 0.7ab
SPV	137 \pm 45e	17 \pm 10c	33 \pm 16def	10 \pm 4a	0.25 \pm 0.09a	1.8 \pm 0.6cde
PA	437 \pm 36c	70 \pm 8b	40 \pm 5cde	6 \pm 1bc	0.09 \pm 0.01c	1.9 \pm 0.2cd
LC	86 \pm 37e	24 \pm 17c	23 \pm 5f	4 \pm 3cd	0.27 \pm 0.12a	2.1 \pm 0.9bc
ChP	295 \pm 35d	58 \pm 11b	45 \pm 9cd	5 \pm 1bcd	0.15 \pm 0.03b	2.3 \pm 0.3abc
N	425 \pm 43c	148 \pm 37a	50 \pm 9c	3 \pm 1d	0.12 \pm 0.02bc	1.3 \pm 0.1e
LCh	614 \pm 63a	148 \pm 27a	86 \pm 11a	4 \pm 1cd	0.14 \pm 0.03b	1.5 \pm 0.1de
SC	557 \pm 56b	72 \pm 9b	66 \pm 10b	8 \pm 1ab	0.12 \pm 0.02bc	2.7 \pm 0.2a

^(a) Sites are depicted as arid (RM), semiarid (SPV, PA, LC, and ChP), semihumid (N), and humid (LCh and SC). ^(b) Experimental sites are: Rio Mayo (RM), San Pablo Valdes (SPV), Los Cerrillos (LC), Potrok Aike (PA), Chacra de Patagones (ChP), Naposta (N), San Claudio (SC) and Las Chilcas (LCh).

CRedit authorship contribution statement

Toledo, S.: Investigation; Methodology and performed the soil property analyses; Performed the data analyses; Writing original draft.

Bondaruk, V. F.: Collected the soil samples in the field; Writing-review and editing.

Yahdjian, L.: Conceptualization; Designed the experiments and coordinated the Project (Drought Net Argentina); Writing-review and editing.

Oñatibia, G R.: Collected the soil samples in the field; Writing-review and editing.

Loydi, A.: Collected the soil samples in the field; Writing-review and editing.

Alberti, J.: Collected the soil samples in the field; Writing-review and editing.

Bruschetti, M.: Collected the soil samples in the field; Writing-review and editing.

Pascual, J.: Collected the soil samples in the field; Writing-review and editing.

Peter, G.: Collected the soil samples in the field; Writing-review and editing.

Agüero, W. D.: Collected the soil samples in the field; Writing-review and editing.

Namur, P. R.: Collected the soil samples in the field; Writing-review and editing.

Blanco, L.: Collected the soil samples in the field; Writing-review and editing.

Peri, P. L.: Conceptualization; Investigation, Resources; Writing-review and editing.


T.S.; B.V.F.; Y.L.; O.G.R.; L.A.; A.J.; B.M.; P.J.; P.G.; A.W.D.; N.P.R.; B. L.; P.P.L.: Drafted the manuscript, and all authors were involved in critical revision and approval of the final version.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof



Graphical abstract

HIGHLIGHTS

Microbial biomass and respiration decreased along an aridity gradient, from humid to arid

Soil C, N, and vegetation cover positively regulate microbial biomass (C and N)

Soil pH negatively affected microbial biomass (C and N) and soil basal respiration

Differential responses of microbial biomass and soil basal respiration depended upon aridity index

Soil microbial variables showed a close relationships with vegetation cover under drought

Journal Pre-proof