




Case Report



Safety of metamizole analgesic therapy in patients with dyspepsia syndrome: A case report



Khoirul Anam ¹, Rina Widiyawati ¹, Tita Sugesti ²,
 M. Hari Pristantiningtyas ³, Herya Putra Dharma ⁴, Muhammad Muchlis ⁵,
 Jainuri Erik Pratama ⁶, Adji Prayitno Setiadi ¹, Marisca Evalina
 Gondokesumo ^{1*}

- ¹ Faculty of Pharmacy, University of Surabaya, Surabaya, 60293 Indonesia
- ² Kidney Disease Ward, Mardi Waluyo Regional Public Hospital, Blitar, 66137 Indonesia
- ³ Pharmacy, Mardi Waluyo Regional Public Hospital, Blitar, 66137 Indonesia
- ⁴ Department of Medical Services and Support, Mardi Waluyo Regional Public Hospital, Blitar, 66137 Indonesia
- ⁵ Mardi Waluyo Regional Public Hospital, Blitar, 66137 Indonesia
- ⁶ Pharmacy Department, Dr. Kariadi Central General Hospital, Semarang, 50244 Indonesia

Abstract: The selection of drugs in patients who have experienced dyspepsia syndrome needs to be studied for its use. Non-steroidal anti-inflammatory drugs (NSAIDs) are a class of drugs that are effective in controlling various inflammatory conditions and treating inflammatory symptoms such as pain but are limited in their use in patients with dyspepsia. We report case of patients with metamizole NSAIDs during three days of hospitalization with a diagnosis of dyspepsia. Other side effects such as cardiovascular, neurological disorders and agranulocytosis can also occur with the use of metamizole. There were no side effects that exacerbated dyspepsia symptoms in patients or other side effects during the three days of using metamizol. It can be concluded that short-term metamizole therapy is safe for use in patients with dyspeptic syndrome. Concomitant use with gastric acid blocking agents such as H2 receptor antagonists may also reduce the effects of gastrointestinal disturbances.

Keywords: Metamizole, Gastrointestinal, Adverse effect, Dyspepsia

INTRODUCTION

Dyspepsia is a clinical syndrome consisting of symptoms of pain or burning that primarily occur in the epigastric or upper abdominal area, accompanied by symptoms of early satiety, bloating, nausea, and vomiting.¹⁻³ About 40% of individuals with dyspepsia complaints seek medical consultation, while 15% of dyspepsia patients are referred for further examination and management.⁴ Dyspepsia is categorized into organic/structural dyspepsia and non-organic/functional dyspepsia. Organic/structural dyspepsia is caused by abnormalities such as peptic ulcers, gastritis, stomach cancer, and gastroesophageal reflux disease (GERD), whereas functional dyspepsia shows no abnormalities on physical examination and endoscopy.⁵ Functional dyspepsia is associated with abnormal gastrointestinal motility, visceral hypersensitivity, genetic

Corresponding author.

E-mail address: a6.apt1@gmail.com (Khoirul Anam)

DOI: [10.29238/teknolabjournal.v12i2.443](https://doi.org/10.29238/teknolabjournal.v12i2.443)

Received 11 March 2023; Received in revised form 28 June 2023; Accepted 30 June 2023

© 2023 The Authors. Published by [Poltekkes Kemenkes Yogyakarta](#), Indonesia.

This is an open-access article under the [CC BY-SA license](#).

factors, H. pylori infection, post-infection factors, psychosocial factors, as well as environmental and dietary factors.^{4,6}

Pharmacological therapy for dyspepsia is based on acid-inhibiting drugs, namely proton pump inhibitors (PPIs) and histamine-2 receptor antagonists (H2RAs).^{7,8} Prokinetic agents are also administered to accelerate disrupted gastrointestinal motility due to changes in visceral sensitivity.^{6,9} The British Society of Gastroenterology (BSG) recommends PPIs as the first-line therapy for dyspepsia with strong recommendations and high-quality evidence. H2 receptor antagonists are also used as first-line treatment, but with weaker recommendations and low-quality evidence.¹⁰ PPIs like omeprazole and lansoprazole, H2RAs like cimetidine, famotidine, ranitidine, and prokinetic agent domperidone are commonly used for dyspepsia therapy in Indonesia (Nabila et al., 2022; Sakaguchi et al., 2012).^{11,12} Restricting foods and drugs that can alter gastrointestinal motility is also essential.

The selection of medications for patients who have experienced dyspepsia syndrome requires careful assessment of their use. Non-steroidal anti-inflammatory drugs (NSAIDs) are one effective class of drugs for controlling various inflammatory conditions and alleviating inflammatory symptoms like pain.¹³ NSAIDs are widely used; however, the risk of adverse effects on the gastrointestinal tract, especially the gastric mucosa, can occur, warranting limitation or dose reduction of NSAIDs in dyspepsia patients.¹⁴ A meta-analysis demonstrated a significantly higher prevalence of dyspepsia events in NSAID users (OR 1.59, 95% CI 1.27-1.99).¹⁵ Adverse effects such as nausea, vomiting, and gastric irritation can occur with parenteral NSAID administration.¹⁶ This can exacerbate dyspeptic symptoms if administered to dyspeptic patients. A case report is presented involving the use of metamizole, a parenteral NSAID, in a patient with dyspepsia syndrome.

CASE REPORT

A 66-year-old woman was admitted to the hospital with complaints of nausea and breathlessness while walking. The patient was diagnosed with dyspepsia, anemia, and hypertension. She had a history of lung infection and had undergone a blood transfusion one month ago. Using the Visual Analogue Scale (VAS), the patient assessed her pain as a score of 4. Vital signs were blood pressure 117/90 mmHg, heart rate 122 beats per minute, respiratory rate 28 breaths per minute, and body temperature 37.40°C. Laboratory results revealed hemoglobin level of 5.43 g/dL and leukocyte count of 19,800/Cmm. The patient received fluid resuscitation therapy with 0.9% NaCl at a rate of 19 drops per minute, ranitidine injection 2x50 mg, metoclopramide injection 3x10 mg, metamizole injection 3x1 g, and packed red blood cell transfusion. Monitoring on the second and third days of therapy showed a decrease in body temperature to a range of 36-36.70°C. By the third day, the complaints of breathlessness and bloating had improved, and the anemia condition showed improvement with a post-treatment hemoglobin increase of 11.7 g/dL. Pain improvement was also observed with a decrease in VAS scores on the second day (score of 3) and on the third day (score of 2), with no signs of gastrointestinal bleeding. Upon discharge, the patient was prescribed omeprazole 2x20 mg, amlodipine 10 mg, spironolactone 50 mg, nucral syrup 3x15 mL, and rindobion 1x1.

DISCUSSION

Metamizole sodium is classified as a commonly used parenteral NSAID in Indonesia. Metamizole is a pyrazolone derivative with a chemical structure related to amidopyrine. This medication exerts analgesic-antipyretic, spasmolytic, and weak anti-inflammatory effects.^{17,18} As a non-selective NSAID, metamizole inhibits the cyclooxygenase (COX) enzymes, namely COX-1 and COX-2, which are

involved in prostaglandin (PG) precursor synthesis.¹⁹ Inhibition of COX-1 results in the suppression of PGE2's gastroprotective function, thus increasing the risk of gastrointestinal disorders like dyspepsia.^{18,20} An infrequently but dangerously occurring adverse effect associated with metamizole usage is agranulocytosis. Agranulocytosis is defined as a neutrophil count lower than $0.5 \times 10^9 \text{ L}^{-1}$ ($<500 \mu\text{L}^{-1}$). In specific patients, metamizole can induce agranulocytosis with an average risk occurrence after one week of usage.^{21,22}

Cardiovascular adverse effects of metamizole usage have been reported in systematic reviews and meta-analyses of metamizole's side effects. Administration of parenteral metamizole causes hypotension compared to paracetamol (RR 3.48, 95% CI 1.07-11.27). Meanwhile, compared to other NSAIDs, side effects such as headache, dizziness, and vertigo are most commonly observed (RR 0.75, 95% CI 0.57-0.99).¹⁸

This case report demonstrates the absence of emerging adverse effects during metamizole usage. Metamizole exhibits good gastric tolerance.¹⁶ Although metamizole strongly inhibits COX-1, very few reports indicate occurrences of duodenal ulcer-related adverse effects. Research conducted by Konijnenbelt et al (2017) revealed that metamizole is the NSAID with the lowest upper gastrointestinal tract side effects compared to other NSAIDs.²³ In line with Konijnenbelt's findings, a meta-analysis conducted by Kotter et al (2015) indicated no difference in side effect occurrences between metamizole and placebo, paracetamol, aspirin, or other NSAIDs. A total of 79 trials involving 4000 patients administered short-term metamizole showed no difference in side effect occurrences and no reports of agranulocytosis.¹⁸ The administration of gastrointestinal protective agents in this case, such as ranitidine, an H2 receptor antagonist, besides alleviating the patient's complaints of nausea and vomiting, can be implemented to prevent gastrointestinal side effects.²⁴

CONCLUSION

As a mild analgesic, metamizole exhibits good gastric tolerance. Short-term metamizole therapy is safe for use in patients with dyspepsia syndrome. Administering it concurrently with gastric acid inhibitors like H2 receptor antagonists can also reduce gastrointestinal disturbances.

AUTHORS' CONTRIBUTIONS

Khoirul Anam and Rina Widiyawati collected research data and wrote this journal. Tita Sugesti, M. Hari Pristantiningtyas, Herya Putra Dharma, and Muhammad Muchlis selected the hospital cases that could be used as case reports, and also guided the writing of this journal. Jainuri Erik Pratama, Adji Prayitno Setiadi, and Marisca Evalina Gondokesumo reviewed and supervised the journal. All authors have read and approved the final version of the journal.

ACKNOWLEDGEMENT

Thanks to Mardi Waluyo Regional Public Hospital, Blitar, East Java, Indonesia where the research data in this journal was collected.

FUNDING INFORMATION

None.

DATA AVAILABILITY STATEMENT

The utilized data to contribute in this journal are available from the author on reasonable request.

DISCLOSURE STATEMENT

The views and opinions expressed in this journal are those of the authors after reviewing various literatures and do not necessarily reflect the official policy or position of any affiliated agency of the authors.

REFERENCE

1. Ford AC, Moayyedi P. Dyspepsia. *BMJ*. 2013;347(aug29 1):f5059-f5059. doi:10.1136/bmj.f5059
2. Madisch A, Andresen V, Enck P, Labenz J, Frieeling T, Schemann M. The diagnosis and treatment of functional dyspepsia. *Dtsch Arztebl Int*. Published online March 30, 2018. doi:10.3238/arztebl.2018.0222
3. Harmon RC, Peura DA. Review: Evaluation and management of dyspepsia. *Therap Adv Gastroenterol*. 2010;3(2):87-98. doi:10.1177/1756283X09356590
4. Black CJ, Houghton LA, Ford AC. Insights into the evaluation and management of dyspepsia: recent developments and new guidelines. *Therap Adv Gastroenterol*. 2018;11:175628481880559. doi:10.1177/1756284818805597
5. Oustamanolakis P, Tack J. Dyspepsia: Organic versus functional. *J Clin Gastroenterol*. 2012;46(3):175-190. doi:10.1097/MCG.0b013e318241b335
6. Yarandi SS, Christie J. Functional dyspepsia in review: Pathophysiology and challenges in the diagnosis and management due to coexisting gastroesophageal reflux disease and irritable bowel syndrome. *Gastroenterol Res Pract*. 2013;2013:1-8. doi:10.1155/2013/351086
7. Pinto-Sanchez MI, Yuan Y, Bercik P, Moayyedi P. Proton pump inhibitors for functional dyspepsia. *Cochrane Database of Systematic Reviews*. Published online March 8, 2017. doi:10.1002/14651858.CD011194.pub2
8. Chaves J, Pita I, Libânio D, Pimentel-Nunes P. Pharmacological Treatment of Functional Dyspepsia: An Old Story Revisited or a New Story to Be Told? A Clinical Review. *GE Port J Gastroenterol*. 2023;30(2):86-97. doi:10.1159/000526674
9. Quigley EMM. Prokinetics in the management of functional gastrointestinal disorders. *J Neurogastroenterol Motil*. 2015;21(3):330-336. doi:10.5056/jnm15094
10. Black CJ, Paine PA, Agrawal A, et al. British society of gastroenterology guidelines on the management of functional dyspepsia. *Gut*. 2022;71(9):1697-1723. doi:10.1136/gutjnl-2022-327737
11. Sakaguchi M. Comparison of PPIs and H2-receptor antagonists plus prokinetics for dysmotility-like dyspepsia. *World J Gastroenterol*. 2012;18(13):1517. doi:10.3748/wjg.v18.i13.1517
12. Nabila H, Dewi NMAR, Lestarini IA. Evaluasi pola penggunaan obat dispepsia berdasarkan indikator WHO (World Health Organization) pada pasien rawat jalan Puskesmas Karang Taliwang tahun 2019. *Sasambo Journal of Pharmacy*. 2022;3(1):11-17. doi:10.29303/sjp.v3i1.139
13. Ida Ghlichloo, Valerie Gerriets. Nonsteroidal Anti-Inflammatory Drugs (NSAIDs) - StatPearls - NCBI Bookshelf. StatPearls [Internet]. Published May 1, 2023. Accessed December 3, 2023. <https://www.ncbi.nlm.nih.gov/books/NBK547742/>
14. Savarino V, Marabotto E, Zentilin P, et al. Proton pump inhibitors: use and misuse in the clinical setting. *Expert Rev Clin Pharmacol*. 2018;11(11):1123-1134. doi:10.1080/17512433.2018.1531703
15. Ford AC, Marwaha A, Sood R, Moayyedi P. Global prevalence of, and risk factors for, uninvestigated dyspepsia: a meta-analysis. *Gut*. 2015;64(7):1049-1057. doi:10.1136/gutjnl-2014-307843
16. Nikolova I, Petkova V, Tencheva J, Benbasat N, Voinikov J, Danchev N. Metamizole: A review profile of a Well-Known "Forgotten" Drug. Part II:

- Clinical profile. *Biotechnology & Biotechnological Equipment*. 2013;27(2):3605-3619. doi:10.5504/BBEQ.2012.0135
17. Harahap MuhW, Ahmad MuhR. Perbandingan antara Dexametason dan Metamizol intravena terhadap kadar neutrofil pasca seksio sesarea. *Jurnal Anestesi Obstetri Indonesia*. 2022;5(3). doi:10.47507/obstetri.v5i3.108
 18. Kötter T, da Costa BR, Fässler M, et al. Metamizole-Associated adverse events: A systematic review and meta-analysis. *PLoS One*. 2015;10(4):e0122918. doi:10.1371/journal.pone.0122918
 19. Jasiocka A, Maślanka T, Jaroszewski JJ. Pharmacological characteristics of metamizole. *Pol J Vet Sci*. 2014;17(1):207-214. doi:10.2478/pjvs-2014-0030
 20. Coxib and traditional NSAID Trialists' (CNT) Collaboration. Vascular and upper gastrointestinal effects of non-steroidal anti-inflammatory drugs: meta-analyses of individual participant data from randomised trials. *The Lancet*. 2013;382(9894):769-779. doi:10.1016/S0140-6736(13)60900-9
 21. Nikolova I, Tencheva J, Voinikov J, Petkova V, Benbasat N, Danchev N. Metamizole: A review profile of a Well-Known "Forgotten" drug. Part I: Pharmaceutical and nonclinical profile. *Biotechnology & Biotechnological Equipment*. 2012;26(6):3329-3337. doi:10.5504/BBEQ.2012.0089
 22. Miljkovic M, Dragojevic-Simic V, Rancic N, et al. Metamizole utilization and expenditure during 6-Year period: Serbia vs. Croatia. *Front Public Health*. 2018;6. doi:10.3389/fpubh.2018.00213
 23. Konijnenbelt-Peters J, van der Heijden C, Ekhart C, Bos J, Bruhn J, Kramers C. Metamizole (dipyrone) as an alternative agent in postoperative analgesia in patients with contraindications for nonsteroidal anti-inflammatory drugs. *Pain Practice*. 2017;17(3):402-408. doi:10.1111/papr.12467
 24. Kumar S, Thakur PK, Sowmya K, Priyanka S. Evaluation of prescribing pattern of NSAIDs in south Indian teaching hospital. *Journal of Chitwan Medical College*. 2016;6(18):54-58. doi:10.3126/jcmc.v6i4.16716



JTL

www.teknolabjournal.com



Jurnal Teknologi Laboratorium

About this journal

Browse this journal



Editorial Team

Editor-in-Chief

Abdul Rohman (Pharmacist)

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia)

Journal Manager

Budi Setiawan (Parasitologist)

(Department of Medical Laboratory Technology, Health Polytechnic of Yogyakarta, Indonesia)

Editorial Board

Dora Dayu Rahma Turista (Microbiologist)

(Department of Biology, University of Mulawarman, Indonesia)

Dipak Kumar Yadav (Physician)

(Kathmandu University, Nepal)

Christian Ebere Enyoh (Chemical)

(Imo State University, Owerri, Nigeria)

Gunes Bolatli (Physician)

(Department of Anatomy, Faculty of Medicine, Siirt University, Siirt, Turkey)

Anis Nurwidayati (Biologist)

(Vector-Borne Disease Research Unit Donggala, National Institute of Health Research and Development, Ministry of Health, Donggala, Central Sulawesi Province, Indonesia)

Sirisha Kanugala (Chemist)

(Department of Biotechnology Senior research fellow- CSIR-Indian Institute of Chemical Technology)

Anna Yuliana (Laboratory Technologist)

(STIKes Bakti Tunas Husada, Tasikmalaya, Indonesia)



Jurnal Teknologi Laboratorium

About this journal



Browse this journal



Vol. 12 No. 2 (2023): In Progress



POLKESYO

Poltekkes Kemenkes Yogyakarta

DOI: <https://doi.org/10.29238/teknolabjournal.v12i2>

Published : 2023-09-17

► Histology

Decreasing alpha-synuclein aggregation on rotenone-induced adult Zebrafish as Parkinson's Diseases model by ethanol extraction of Keluwih (*Artocarpus camansi*) leaves >

Marisca Evalina Gondokesumo, Faisal Akhmal Muslikh, Ni Putu Diah Nopitasari, Putu Dea Angelita Putri

 DOI : <https://doi.org/10.29238/teknolabjournal.v12i2.408>

 Pages: 53-60

Published : 2023-09-17

336	6
Views	Downloads

Immunology

Overview of trygleseride level on coffee consumption >

Amellya Octifani, Tarisa Suci Novianti , Farida Anwari, Arif Rahman Nurdianto, Fery Setiawan, Rizal Fauzi Nurdianto

 DOI : <https://doi.org/10.29238/teknolabjournal.v12i2.427>

 Pages: 81-88

Published : 2023-12-12

126	13
Views	Downloads

Serology

Bioactivity of Nelumbo nucifera extract on sperm recovery due to 2-methoxyethanol exposure: In vivo and in silico study >

Angella Ananda Syaputra, Putri Ayu Ika Setiyowati, Badriatul Musyarofah, Yunita Ainul Kasanah, Amelia Kartika Reza Rahmawati, Helga Syasya Qotrunnada

 DOI : <https://doi.org/10.29238/teknolabjournal.v12i2.450>

 Pages: 89-100

Published : 2023-12-29

10	0
Views	Downloads

Bacteriology

In vitro antibacterial activity of patchouli batik (Pogostemon Cablin) plant extract varieties from Southeast Sulawesi and South Sulawesi >

Tuty Yuniarty, Ratih Feraritra Danu Atmaja, Julianti Isma Sari Usman, Theosobia Grace Orno

 DOI : <https://doi.org/10.29238/teknolabjournal.v12i2.439>

Published : 2023-12-30

21
Views

Clinical condition of patients with Obesity Hypoventilation Syndrome (OHS): case report >

Made Oka Heryana, Heru Widjono

DOI : <https://doi.org/10.29238/teknolabjournal.v12i2.440>

Pages: 71-80

Published : 2023-12-25

158	19
Views	Downloads

Therapy of resistant hypertension in patients with chronic kidney disease complications of anemia in hemodialysis: a case report >

Yenry Sumarlim, Emilia Gan, Desantika Wuryana, M. Hari Pristantiningtyas, Herya Putra Dharma, Muhammad Muchlis, Jainuri Erik Pratama, Adji Prayitno Setiadi, Marisca Evalina Gondokesumo

DOI : <https://doi.org/10.29238/teknolabjournal.v12i2.444>

Pages: 61-65

Published : 2023-11-04

189	14
Views	Downloads

Safety of metamizole analgesic therapy in patients with dyspepsia syndrome: a case report >

Khoirul Anam, Rina Widiyawat, Tita Sugesti, M. Hari Pristantiningtyas, Herya Putra Dharma, Muhammad Muchlis, Jainuri Erik Pratama, Adji Prayitno Setiadi, Marisca Evalina Gondokesumo

DOI : <https://doi.org/10.29238/teknolabjournal.v12i2.443>

Pages: 66-70

Published : 2023-11-04

143	9
Views	Downloads



Tweets from @JurnalTeknolog1



**Nothing to
see here - yet**

When they Tweet, their Tweets
will show up here.

Keywords



JURNAL TEKNOLOGI LABORATORIUM Published by Poltekkes Kemenkes Yogyakarta

is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

Number of Visitors

00379133

[View My Stats](#)



JURNAL TEKNOLOGI LABORATORIUM

JURUSAN ANALIS KESEHATAN POLTEKKES KEMENKES YOGYAKARTA

P-ISSN : 23385634 < > E-ISSN : 25800191 Subject Area : Science, Education



0

Impact Factor



812

Google Citations



Sinta 2

Current Accreditation

[Google Scholar](#) [Garuda](#) [Website](#) [Editor URL](#)

History Accreditation

2017

2019

2020

2021

2022

2023

Garuda

[Google Scholar](#)

[Does gender affect immune response in HIV patients?](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress 14-21](#)

2021 [DOI: 10.29238/teknolabjournal.v10i1.261](#) [Accred : Sinta 2](#)

[Absolute leukocytes count and NLR as a diagnostic and prognostic biomarkers for severity of COVID-19 infection](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress 03-05](#)

2021 [DOI: 10.29238/teknolabjournal.v10i1.263](#) [Accred : Sinta 2](#)

Turkey

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress 22-30](#)

2021 [DOI: 10.29238/teknolabjournal.v10i1.266](#) [Accred : Sinta 2](#)

[A simple method for isolation of rest of trypsinized stem cells with magnetic beads](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress 01-02](#)

2021 [DOI: 10.29238/teknolabjournal.v10i1.268](#) [Accred : Sinta 2](#)

[Conserved B-cell epitope identification of envelope glycoprotein \(GP120\) HIV-1 to develop multi-strain vaccine candidate through](#)

[bioinformatics approach](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress 06-13](#)

📅 2021 [DOI: 10.29238/teknolabjournal.v10i1.274](#) [Accred : Sinta 2](#)

[Molecular identification of pathogenic bacteria causing foodborne disease in Caulerpa racemosa](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress 31-39](#)

📅 2021 [DOI: 10.29238/teknolabjournal.v10i1.276](#) [Accred : Sinta 2](#)

[Air-Dried and Wet Fixation on Fine Needle Aspiration Biopsy \(FNAB\) Specimen](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress](#)

📅 2021 [DOI: 10.29238/teknolabjournal.v10i1.279](#) [Accred : Sinta 2](#)

[Digital communication in health promotion in handling tuberculosis sputum](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress](#)

📅 2021 [DOI: 10.29238/teknolabjournal.v10i1.285](#) [Accred : Sinta 2](#)

[Multi drugs resistance to Diabetes Mellitus patients with tuberculosis in Manado City](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 10 No 1 \(2021\): inpress 40-45](#)

📅 2021 [DOI: 10.29238/teknolabjournal.v10i1.286](#) [Accred : Sinta 2](#)

[Konfirmasi Index Eritrosit Dengan Sediaan Apus Darah Tepi \(Sadt\) Pada Penduduk Tambaklorok Semarang Yang Terpajan Pb](#)

POLTEKKES KEMENKES YOGYAKARTA [Jurnal Teknologi Laboratorium Vol 9 No 2 \(2020\): \(In Progress\)](#)

📅 2020 [DOI: 10.29238/teknolabjournal.v9i2.213](#) [Accred : Sinta 2](#)

[View more ...](#)