



Different methods and times to estimate heat production in sheep fed with sunflower meal

Luciana Castro Geraseev^{1*} Sarah Silva Santos¹
Amália Saturnino Chaves² Fernanda Samarini Machado³

¹Instituto de Ciências Agrárias da Universidade Federal de Minas Gerais (UFMG), 1000, Montes Claros, MG, Brasil. E-mail: lgeraseev@gmail.com.

*Corresponding author.

²Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora, MG, Brasil.

³Embrapa Gado de Leite, Juiz de Fora, MG, Brasil.

ABSTRACT: The objective of this study was to assess the oxygen pulse and heart rate method (O₂P-HR) using a respiration chamber at different measurement times for estimate the heat production (HP) of lambs fed increasing levels of sunflower meal in their diet. Twenty-four lambs were assigned to four experimental diets (0, 100, 200, and 300 g of sunflower meal kg⁻¹ DM). Heat production was estimated using the O₂P-HR (HP_{O₂P}) method and a respirometry chamber (HP_{RC}). Measurements were obtained by simultaneously measuring heart rate (HR) and oxygen consumption over 3, 6, 9, 12, 15, 18, 21 and 24 h. A flow-through respirometry chamber for small ruminants was used to determine oxygen consumption (VO₂) and carbon dioxide and methane production. Data on dietary treatment, measurement times and their interactions were analyzed as repeated measures using mixed model procedures and Restricted Maximum Likelihood (REML) estimation. The Pearson's correlation coefficient was used to compare techniques. There was no effect of the different levels of sunflower meal inclusion on VO₂ and heat production. The HP_{O₂P} (126.16 kcal/ BW^{0.75}/day) was 2% higher than that of the HP_{RC} (124.61 kcal/ BW^{0.75}/day), and the correlation coefficients was 0.628. The coefficient of variation was greater for the HP_{O₂P} (21.33%) than for HP_{RC} (11.44%). HR (beats/min), VO₂ (mL/min/BW^{0.75}) and O₂P-HR (mL/beat) required measurement times of 24, 15 and 9 hours, respectively. A measurement time of 24 h was necessary to ensure a more accurate estimate of the heat production using the O₂P-HR method.

Key words: bioenergetic, energy requirements, indirect calorimetry, lamb, ovine.

Diferentes métodos e tempos de medição para estimar a produção de calor em ovinos alimentados com farelo de girassol

RESUMO: O objetivo com este estudo foi avaliar o método do pulso de oxigênio (O₂P-FC) usando câmara respirométrica em diferentes tempos de medição para estimar a produção de calor de cordeiros alimentados com níveis crescentes de farelo de girassol na dieta. Vinte e quatro cordeiros foram distribuídos em quatro dietas experimentais (0, 10, 20 e 30% de farelo de girassol). A produção de calor foi estimada pelo método de O₂P-FC (PC_{O₂P}) e por câmara respirométrica (PC_{CR}). As estimativas foram obtidas medindo-se simultaneamente a frequência cardíaca (FC) e o consumo de oxigênio (VO₂) durante 3, 6, 9, 12, 15, 18, 21 e 24 horas. Uma câmara de respirométrica para pequenos ruminantes foi usada para determinar o VO₂ e a produção de dióxido de carbono e metano. Os dados referentes a dieta experimental, tempos de medição e suas interações foram analisados como medidas repetidas usando os procedimentos de modelo misto e estimativa de máxima verossimilhança restrita. A correlação de Pearson foi usada para comparar as duas técnicas de estimativa da produção de calor. Não houve efeito dos diferentes níveis de inclusão de farelo de girassol sobre o consumo de oxigênio e produção de calor dos animais. A PC_{O₂P} (126,16 kcal/ PV^{0.75}/dia) foi 2% maior que a PC_{CR} (124,61 kcal/ PV^{0.75}/dia), e o coeficiente de correlação foi de 62,8%. O coeficiente de variação foi maior para PC_{O₂P} (21,33%) comparado com PC_{CR} (11,44%). A FC (batimentos/min), VO₂ (mL/min/PC^{0.75}) e o O₂P-FC (mL/batimento) requerem tempos de medição de 24, 15 e 9 horas, respectivamente. É necessário a mensuração por 24 horas para garantir uma estimativa mais precisa da produção de calor usando o método de O₂P-FC.

Palavras-chave: bioenergética, calorimetria indireta, cordeiros, exigências de energia, ovinos.

1 INTRODUCTION

2
3 Energy is the primary nutrient limiting
4 ruminants. It is derived from the oxidation of dietary
5 nutrients and is essential for maintenance of vital
6 processes. This nutrient is dissipated by animals during
7 ingestion and metabolism of food: first, energy is

consecutively lost in feces, urine and fermentative gases,
and subsequently lost as heat increases. The remaining
energy is primarily directed towards maintenance
(approximately 70% of the net energy available) and
production processes (TEIXEIRA et al., 2017).

The standard method for measuring energy
expenditure in ruminants involves the use of open

1 circuit respirometry chambers. In this method, the
2 products resulting from the animal's metabolism,
3 such as gas exchanges with the environment
4 (oxygen consumption, carbon dioxide and methane
5 production), combined with urinary nitrogen
6 excretion, are quantified (SILVA, 2011; OSS et
7 al., 2016). A respirometry chamber is an accurate
8 technique, however it is used under laboratory
9 conditions and is extremely expensive. It also requires
10 significant expertise and infrastructure, which makes it
11 impractical for small rural properties (RODRIGUEZ
12 et al., 2007; MACHADO et al., 2016).

13 Measuring the heat production of animals
14 can provide insights into how efficiently they utilize
15 the nutrients in their diet, thereby helping optimize feed
16 efficiency, since heat production is closely linked to the
17 metabolic processes and energy utilization of animals.

18 Researchers are seeking to estimate heat
19 production in ruminants by using heart rate adjusted
20 for oxygen consumption per beat as there is a linear
21 relationship between heart rate (HR) and oxygen
22 consumption (VO_2) in homeothermic animals,
23 thereby indicating that it is possible to estimate heat
24 production through HR measurements (TALMON
25 et al., 2023). The primary goal is to improve and
26 develop techniques capable of measuring the energy
27 requirements of animals in a shorter time frame,
28 with cheaper equipment, and without changing the
29 behavior and normal conditions of animal husbandry
30 (BROSH, 2007; CHAVES et al., 2015).

31 The O_2P -HR technique can be used as an
32 alternative method to determine heat production.
33 However, there are some problems associated with
34 the ideal time to measure oxygen consumption,
35 heart rate, and O_2P -HR, especially considering the
36 intraday changes that interact directly with animals.
37 According to OSS et al. (2016), O_2P -HR is an
38 alternative technique, but has a greater between-
39 animal coefficient of variation, which has a negative
40 effect on the power of the experiments. Further
41 studies should be performed to investigate ways
42 to minimize the errors associated with the O_2P -HR
43 method to increase the precision and statistical power
44 of experiments using this technique.

45 The objective of this study was to evaluate
46 the O_2P -HR method at different measurement times
47 to estimate heat production in crossbred lambs fed
48 diets containing increasing levels of sunflower meal.

49 MATERIALS AND METHODS

50
51
52 Twenty-four crossbred (Santa Inês x
53 Dorper) intact male lambs with a mean age of

4 months were arranged into three blocks, four
treatments, and two replicates per block using a
randomized block design. A 10-day adaptation period
was allowed before the data collection.

5 The lambs received four isoproteic
6 experimental diets formulated according to the NRC
7 (2007) recommendations for lambs on maintenance
8 levels. The diets containing a roughage:concentrate
9 ratio of 40:60 on a dry matter basis (DM). Corn
10 silage was supplied as the roughage source and the
11 concentrate was formulated by replacing soybean
12 meal with increasing levels of sunflower meal (0,
13 100, 200 and 300 g kg^{-1} DM) (Table 1). The diet was
14 provided in two daily meals at 8 a.m. and 4 p.m.

15 The chemical composition of the diets
16 andorts was determined by analyzing the dry matter
17 (DM), ash, crude protein (CP), neutral detergent
18 fiber (NDF), acid detergent fiber (ADF), ether
19 extract (EE), non-fibrous carbohydrates (NFC) and
20 total carbohydrate (TC) content according to the
21 procedures of INCT-CA (DETMANN et al., 2012).

22 Before starting the experiment, the lambs
23 were weighed, drenched and vaccinated against
24 clostridial diseases. During the experiment, lambs
25 were housed in individual metabolic cages provided
26 with feed and water troughs, which allowed the
27 collection of urine and fecal samples. After the
28 adaptation period DM intake was measured, and
29 urine was collected for nitrogen determination for
30 five days. Heat production was estimated by using a
31 respirometry chamber after adaptation to the diet.

32 The O_2 consumption (VO_2) and CO_2
33 production data were recorded using a Sable System
34 (Sable Systems International, Las Vegas, NV, USA).
35 The lambs were individually placed in a respirometry
36 chamber for 24 h and the same dietary treatment
37 offered during the adaptation period was administered
38 to each lamb once in the morning.

39 Ambient air flowed through the chamber at a
40 controlled flow rate based on lamb's weight (0.6 liters/
41 kg of body weight/minute), and it was mixed with the
42 exhaled air. Samples were taken every 5 min for 24 h
43 to determine O_2 , CO_2 , and CH_4 concentrations. All data
44 were recorded using an automated data acquisition
45 program (Expedata; Sable Systems International).

46 The maximum allowable concentration of
47 CO_2 in the chamber was 1.0%. Oxygen consumption
48 and CO_2 production were calculated by comparing
49 the composition and volume of the air that flowed
50 through the respirometry chamber with the air
51 released. The temperature was kept at 22 °C by
52 using an air conditioner placed inside the chamber to
53 provide thermal comfort to lambs.

Table 1 - Nutritional composition of the experimental diets.

Item (g kg ⁻¹)	-----Sunflower meal level (g kg ⁻¹ DM)-----			
	0	100	200	300
Corn silage	400	400	400	400
Soybean meal	264	196	118	18
Corn grain	315	281	256	261.6
Sunflower meal	0	100	200	300
Vitamin-Mineral Premix ¹	10.5	15.0	22.0	36.6
Dicalcium phosphate	10.5	8.0	4.0	0
	-----Nutrients-----			
DM (g/kg ⁻¹ NM)	643.4	641.7	653.3	658.8
MM (g/ kg ⁻¹ DM)	45.0	43.2	45.7	46.1
CP (g/ kg ⁻¹ DM)	207.9	203.8	195.6	189.5
NDF(g/ kg ⁻¹ DM)	337.9	364.5	391.2	419.8
ADF(g/ kg ⁻¹ DM)	176.3	212.5	227.2	302.8
EE (g/ kg ⁻¹ DM)	65.5	62.9	60.9	60.1
NFC(g/ kg ⁻¹ DM)	343.7	325.6	306.6	284.5
TC (g/ kg ⁻¹ DM)	681.6	690.1	697.8	704.3

DM = Dry matter, NM = Natural Matter, CP = Crude Protein, NDF = Neutral detergent Fiber, ADF = Acid detergent fiber, EE = Ether extract, NFC = Non-fibrous carbohydrate, TC = Total carbohydrate.

¹Composition of Vitamin-Mineral Premix: Calcium (Max.) 150 g, Calcium (Min.) 130 g, Phosphorus (Min.) 65 g, Sodium (Min.) 130 g, Fluorine (Max.) 50 mg, Sulfur (Min.) 12 g, Magnesium (Min.) 10 g, Iron (Min.) 1000 mg, Manganese (Min.) 3000 mg, Cobalt (Min.) 80 mg, Zinc (Min.) 5000 mg, Iodine (Min.) 60 mg, Selenium (Min) 10 mg, Vitamin A (Min.) 50000 IU, Vitamin E (Min.) 312 IU.

1 Heat production was estimated using the
2 respirometry chamber technique (HP_{RC}) according to
3 Brouwer's equation (1965) as follows:

$$4 \text{ HP}_{\text{RC}} (\text{Kj}) = 16.18 \times \text{VO}_2 (\text{L}) + 5.02 \times \text{VCO}_2 (\text{L}) - 5.88$$

$$5 \times \text{UN} (\text{g}) - 2.17 \times \text{VCH}_4 (\text{L})$$

6 HP_{RC} = Estimation of heat production using the respirometry chamber technique

7 VO₂ = Oxygen consumption

8 VCO₂ = Carbon dioxide production

9 UN = Urinary nitrogen

10 VCH₄ = Methane production

11 Estimation of heat production using the
12 O₂ pulse methodology (HP_{O₂P}) was based on a protocol adapted from BROSH et al. (1998). After
13 the adaptation period, the lambs were monitored
14 for four days to record the mean heart rate using
15 a POLAR® RS800 transmitter. The transmitters
16 were attached to the girth of the lambs using elastic strips. Data were recorded at 60 s intervals and
17 subsequently transferred to a computer using an infrared sensor.

18 After determining the mean heart rate
19 (HR during the four days of measurement), data on
20 heartbeat (HR-RC) and oxygen consumption (VO₂)
21 were collected simultaneously for 24 h using a
22 respirometry chamber, as described above. These data

1 were used to calibrate the O₂ volume per heartbeat.
2 The oxygen pulse and heart rate were calculated as
3 VO₂ per heartbeat.

4 Daily heat production was obtained by
5 multiplying the total O₂ consumption by the constant
6 4.89 kcal/L of O₂ (NICOL & YOUNG, 1990). The
7 results were expressed as metabolic weight (kcal/kg
8 BW^{0.75}/day). Heat production was estimated using the
9 following equation:

$$10 \text{ HP}_{\text{O}_2\text{P}} = \frac{\text{kcal}}{\text{day} \times \text{kg BW}^{0.75}} = (\text{HR} - \text{RC} \times 2 \text{ O}_2\text{P} \times 4.89) \times 1440 / (\text{kg BW}^{0.75})$$

11 HP_{O₂P} = Estimation of heat production using the oxygen pulse and heart rate method

12 HR-RC: Mean heartbeat (beat/min)

13 O₂P: Oxygen consumption per heartbeat (L/beat).

14 Data analysis

15 The dietary treatments, measurement
16 times and their interactions were analyzed as repeated
17 measures (each treatment was analyzed at eight
18 measurement times: 3, 6, 9, 12, 15, 18, 21 and 24 h)
19 because the observations were interdependent. Data
20 were analyzed using the Proc MIXED procedure in
21 SAS (SAS 9.0 Inst. Inc.) and Restricted Maximum
22 Likelihood (REML) estimation according to the
23 following model:
24
25
26

$$y_{ijt} = \mu + \alpha_i + dj(i) + \gamma t + (\alpha\gamma)it + (b + \beta_j) + e_{ijt}$$

Where, y_{ijt} = the expected outcome for the dependent variable Y observed at the measurement time t for the lamb j fed the diet i ; μ is the overall mean; α_i is the fixed effect of diet; $dj(i)$ is the random effect of lamb j nested within diet i ; γt is the fixed effect of measurement time; $(\alpha\gamma)it$ is the interaction between diet and measurement time; b is the regression coefficient; β_j is the slope deviation (diet i) of the regression coefficient b ; e_{ijt} is the random error associated with lamb j fed diet i at the measurement time t , $e_{ijt} \sim \text{NID}((0, \sigma_e^2))$ (data is approximately normally distributed with mean of 0 and variance of σ_e^2); and the values of $dj(i)$ and e_{ijt} are assumed to be independent.

Five variance-covariance matrix structures were tested as follows: variance components (VC - variances are equal and observations are independent, i.e., there is no correlation between observations over time); compound symmetry (CS - equality of variances and covariances); first-order autoregressive model (AR (1) - equality of variances and covariances with higher correlation between adjacent measures); first-order ante-dependence (ANTE (1) - the magnitude of the covariance depends on the values of both correlation and standard deviations associated with them); unstructured (UN - each variance and covariance is estimated exclusively from the data) (SAS, 2004). The best model for each set of variables was selected based on the lowest corrected Akaike Information Criterion (AICc) value.

The variance-covariance matrix structure of the best fit for the measurement time was selected based on the lowest corrected Akaike Information Criterion (AICc) value (LITTELL et al., 2006). The ANTE (1) model provided the best fit for HR, O_2P (mL/beat/ $\text{BW}^{0.75}$), HPO_2P (kcal/day) and HPO_2P (kcal/day/ $\text{BW}^{0.75}$), thereby modeling

the covariance structure and thus generating valid tests. The AR (1) model provided the best fit for the VO_2 (L/day), VO_2 (mL/min/ $\text{BW}^{0.75}$) and O_2P (L/beat), whereas the ANTE (1) model did not converge. The UN model did not converge for VO_2 (L/day) or VO_2 (mL/min/ $\text{BW}^{0.75}$). For the other parameters, problems were encountered when the Hessian matrix was applied, which demonstrates that the UN structure was inappropriate.

After defining the best model for each set of variables, the result of the fixed effect analysis (measurement time) was used as a criterion to test the significance of the treatment effect ($\alpha=0.05$). The parameters were subjected to regression analysis (PROC REG) using SAS (2004) when the diet was significant. Differences between groups means (each measurement time vs 24 h) were determined by calculating the minimum significant difference for $p = 0.05$ using the Tukey's test' when measurement time was significant. To express the accuracy and repeatability of the test, the coefficient of variation was calculated using the PROC UNIVARIATE procedure (SAS, 2004). Pearson's correlation was used to compare techniques using PROC CORR (SAS, 2004). Statistical significance was set at $P < 0.05$.

RESULTS AND DISCUSSION

The inclusion of sunflower meal did not change ($P > 0.05$) the DM intake, as the animals were fed at a maintenance level, with averages of 612 g/day and 50.48 g/ $\text{BW}^{0.75}$ /day. There was no effect on CP intake owing to the lack an effect on DM intake and the isonitrogenous profile of the diets (Table 2). However, there was a significant difference ($P < 0.05$) in NDF intake among the treatments, with a linearly increasing effect observed. This behavior can be

Table 2 - Means, coefficient of variation (CV) for dry matter intake (DMI), crude protein intake (CPI) and neutral detergent fiber intake (NDFI) of lambs fed with different levels of sunflower meal inclusion.

Variables	-----Sunflower meal level (g kg ⁻¹ DM)-----				CV (%)	-----P-----	
	0	100	200	300		Linear	Quadratic
DMI (g/day)	637.63	596.57	601.7	613.80	10.70	0.593	0.5440
DMI (g/ $\text{BW}^{0.75}$ /day)	50.94	50.16	50.28	50.54	2.63	0.6576	0.5850
CPI (g/ $\text{BW}^{0.75}$ /day)	10.9	9.85	10.04	10.04	2.72	0.8721	0.5929
NDFI (g/ $\text{BW}^{0.75}$ /day) ¹	17.49	19.58	22.47	25.14	14.17	<.0001	<.0001

¹ $y = 17.29 + 0.0258x$; $R^2 = 0.967$.

1 explained by the increased fiber concentration in the
2 diets resulting from the inclusion of SFM.

3 The difference in NDF intake was not
4 sufficient to change the VO_2 or heat production
5 ($P > 0.05$). This behavior was possibly due to the
6 animals' energy intake being close to maintenance
7 levels. Typically, animals with higher digestible
8 energy intake consume more oxygen during
9 metabolic processes.

10 The HR-RC, normal HR, VO_2 , HP_{O_2P} and
11 HP_{RC} did not differ among sunflower meal inclusion
12 levels ($P > 0.05$), possibly because the lambs were fed
13 near maintenance. Moreover, the interaction between
14 SFM level and time was not significant ($P > 0.05$).

15 The HR of lambs during the four days of
16 measurement varied by only 2.6% in comparison with
17 the HR-RC (Table 3), which suggests the absence of
18 stress or lack of exercise during the evaluation period.
19 LANDAU et al. (2006) observed similar results for
20 daily HR and HR-RC (81.1 ± 5.1 and 79.2 ± 5.1 ,
21 respectively). According to the authors, this response
22 was associated with a lower environmental effect on
23 grazing during the measurement of the O_2 pulse.

24 The VO_2 (Table 3) was similar to that
25 reported by MACHADO et al. (2015) when evaluating
26 heat production in sheep fed sorghum silages at
27 different maturation stages ($18.37 \text{ mL/min/kg BW}^{0.75}$).
28 The similarity in the results is likely related to feeding
29 near maintenance and absence of stress during the
30 execution of both studies. This is also corroborated by
31 the heart rate data as variations of less than 20% were
32 observed between normal heart rates.

33 The O_2P values (Table 3) corroborate those
34 reported by ARIELI et al. (2002) ($0.250 \text{ mL/beat/kg/}$
35 $BW^{0.75}$) in sheep fed high or low energy diets (75%
36 and 25% concentrate, respectively). Under conditions
37 where animals are not subjected to stress, physical

1 activity or if the variation in heart rate is less than
2 20%, O_2P remains constant, and the data are reliable.

3 The coefficient of variation was greater
4 for HP_{O_2P} (21.33%) than for HP_{RC} (11.44%). The
5 repeatability of individual animals over time needs
6 to be high to reliably detect the differences in the
7 heat production of animals through respiration
8 trials. OSS et al. (2016) compared the O_2P method
9 with measurements using a respirometry chamber
10 in crossbred steers (Holstein \times Gyr), which was also
11 confirmed by a greater between-animal coefficient of
12 variation (16.6%) compared to RC (7.7%). According
13 to the authors, the O_2P -HR method had a higher
14 coefficient of variation, and the sample size (n) must
15 be increased to determine the differences in HP
16 between treatments more accurately.

17 Despite the differences in the coefficients
18 of variation, the correlation between HP_{O_2P} and HP_{RC}
19 was 0.628 (Figure 1), thereby validating the efficiency
20 of the O_2P -HR method in predicting heat production.
21 The HP_{O_2P} was 2% higher than the HP_{RC} . In a study
22 evaluating the efficiency of the O_2P -HR method as
23 a tool for determining energy expenditure in sheep,
24 ARIELI et al. (2002) reported that heat production
25 using the O_2P -HR technique was 6.7% higher than
26 that using the comparative slaughter method.


27 Table 4 presents the differences between
28 the overall means observed at each measurement time
29 and after 24 h. A significant difference was detected
30 between measurement times of up to 6 h vs. 24 h for
31 all variables studied. This response is probably due to
32 the initial adjustment phase of gas collection, which
33 is essential for equilibrium between gas production
34 and consumption inside the respirometry chamber.

35 There was an increase in HR and VO_2
36 during the first hours, followed by a gradual reduction
37 after feeding. This response is likely related to

Table 3 - Mean, standard deviation, minimum and maximum of heart rate, oxygen consumption and heat production of sheep.

Item	Mean	SD	Min.	Max.
HR ¹ , bpm	79.52	9.28	65.70	96.80
HR-RC ² , bpm ³	81.65	11.50	61.84	105.71
VO_2 ⁴ , ml/min/kg $BW^{0.75}$	18.09	2.52	12.21	29.42
O_2P ⁵ , ml/beat/kg $BW^{0.75}$	0.225	0.039	0.143	0.287
HP_{O_2P} ⁶ , kcal/day/kg $BW^{0.75}$	126.16	26.91	68.72	164.92
HP_{RC} ⁷ , kcal/day/kg $BW^{0.75}$	124.61	14.26	95.94	149.03

¹Mean heart rate during the four days of measurement. ²Mean heart rate collected for 24 hours using a respirometry chamber. ³Beat per minute. ⁴Oxygen consumption. ⁵Oxygen consumption per heart beat. ⁶Heat production using the O_2 pulse methodology. ⁷Heat production using the respirometry chamber methodology.



1 reduced stress levels after the initial period and a more
 2 extended post-feeding period, as the diet was provided
 3 once a day at the beginning of the measurement. The
 4 feeding time and the physical activities of chewing
 5 and swallowing were the leading causes of the
 6 increase in HR and VO₂ during the first few hours,
 7 as they showed a gradual reduction after feeding.
 8 According to TALMON et al. (2023) eating was the
 9 activity that most increased HP, VO₂, and HR.

10 Measurement time had a significant effect
 11 on HR (Table 4) throughout the 24 h period ($P <$

0.0001). Nevertheless, the variations observed after
 the 9 h period were lower than 15%. According to
 BROSH (2007), variations in a normal heartbeat
 are acceptable for the determination of O₂P, thereby
 ensuring the reliability of our database.

Based on the statistical analysis, we
 observed that HR measurements should be performed
 for 24 h for more complete data collection, avoiding
 intraday variations such as feeding time and diet
 quality (metabolic activity increases during digestion
 and absorption), lower heart rate at night (when

Table 4 - Variation in heart rate, oxygen consumption, oxygen volume per heart beat, and heat production using the O₂ pulse methodology in sheep (n = 23), expressed as the difference between the mean measurement at each time studied and that obtained during 24 hours.

Variable	Measurement time							
	3h	6h	9h	12h	15h	18h	21h	24h
HR, beat/min	26.52***	17.08***	12.88***	8.34***	5.17***	2.64***	0.97***	81.65 ± 2.40
VO ₂ , l/day	5.40***	4.13**	3.26**	2.11*	1.02	-0.16	-0.04	30.53 ± 8.92
VO ₂ , ml/min/BW ^{0.75}	3.19***	2.43**	1.94*	1.24*	0.60	0.06	-0.02	18.09 ± 0.53
O ₂ P, ml/beat	-0.30**	-0.20*	-0.14	-0.10	-0.09	-0.09	-0.04	2.6±0.11
O ₂ P, ml/beat/BW ^{0.75}	-0.03*	-0.02*	-0.01*	-0.008*	-0.007*	-0.007*	-0.002*	0.225±0.008
HP _{O₂P} , Kcal/day	-174.9*	-107.3*	-79.51*	-55.98*	-52.63*	-52.63*	-20.48*	1485.2±74.32
HP _{O₂P} , Kcal/day/BW ^{0.75}	-14.51*	-8.75*	-6.48*	-4.60*	-4.23*	-4.23*	-1.63*	126.17±5.61

*** $P < 0.0001$; ** $P < 0.001$; * $P < 0.05$. HR=Mean heart rate during the four days of measurement. VO₂=Oxygen consumption. O₂P=Oxygen consumption. per heart beat. HP_{O₂P}=Heat production using the O₂ pulse methodology.

1 animals are at rest) and excitement resulting from
2 the presence of people. BARKAI et al. (2002),
3 ARIELI et al. (2002), AHARONI et al. (2003),
4 and LANDAU et al. (2006) used the methodology
5 of BROSH et al. (1998) with HR and VO_2
6 measurements for 15-20 min throughout the day
7 and obtained results similar to those found in the
8 literature. However, according to PUCHALA et al.
9 (2007), the HR and energy expenditure of goats
10 consuming different quality diets varied within 24
11 h, thereby corroborating our observations.

12 The measurement time (up to 12 hours)
13 affected VO_2 (L/day) and VO_2 (mL/min/ $\text{BW}^{0.75}$).
14 From 15 h onwards, the parameters were similar
15 to those obtained after 24 h (Table 4). Oxygen
16 consumption may have varied at the beginning of
17 the gas collection phase owing to the start of feeding
18 and the initial stress associated with the chamber,
19 which resulted in increased O_2 consumption.
20 VAN MILGEN et al. (1997) observed that
21 oxygen consumption varied according to animal
22 behavior when assessing O_2 consumption and CO_2
23 production during the resting state, feeding and
24 physical activity in pigs. BARKAI et al. (2002)
25 and LANDAU et al. (2006) estimated oxygen
26 consumption for 15-20 min at different times of the
27 day using the methodology of BROSH et al. (1998)
28 and observed no variation in oxygen consumption.
29 However, based on our observations, more accurate
30 measurements of the oxygen consumption require
31 longer measurement times.

32 Although it was possible to measure O_2P
33 (mL/beat) for 9 h, the effect of time on O_2P (mL/beat/
34 $\text{BW}^{0.75}$) and $\text{HP}_{\text{O}_2\text{P}}$ (Kcal/day and Kcal/day/ $\text{BW}^{0.75}$)
35 over the entire measurement period, demonstrated
36 that these parameters should be measured for 24
37 h when using the O_2P methodology (Table 4).
38 This may be associated with variations in HR or
39 processes involving digestion and diet quality. The
40 roughage:concentrate ratio (40:60) explains the 24 h
41 variations in $\text{HP}_{\text{O}_2\text{P}}$ because the degradation of non-
42 fibrous carbohydrates is fast, whereas the digestion of
43 fibrous carbohydrates occurs more slowly owing to
44 the long lag time.

46 CONCLUSION

47
48 The O_2P -HR method is highly correlated
49 with the respirometry chamber methodology for
50 estimating heat production in sheep; however, O_2P -
51 HR should be measured for 24 h to ensure greater
52 accuracy. Sunflower meal inclusion levels did not
53 affect heat production in the animals.

ACKNOWLEDGEMENTS

The authors would like to thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), the Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001, for financial assistance.

DECLARATION OF CONFLICT OF INTEREST

The authors declare no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

AUTHORS' CONTRIBUTIONS

Conceptualization: LCG and ASC. Data acquisition: SSS, ASC and LCG. Design of methodology and data analysis: LCG and ASC. LCG, SSS, ASC and FSM prepared the draft of the manuscript. All authors critically revised the manuscript and approved of the final version.

BIOETHICS AND BIOSSECURITY COMMITTEE APPROVAL

The experimental procedures involving animals were approved by the Ethics Committee on Animal Use of the Universidade Federal de Minas Gerais (UFMG) under protocol No. 189/15.

REFERENCES

- AHARONI, Y. et al. The variability of the ratio of oxygen consumption to heart rate in cattle and sheep at different hours of the day and under different heat load conditions. *Livestock Production Science*, v.79, p.107-117, 2003. Available from: <[http://dx.doi.org/10.1016/S0301-6226\(02\)00147-1](http://dx.doi.org/10.1016/S0301-6226(02)00147-1)>. Accessed: Nov. 09, 2022. doi: 10.1016/S0301-6226(02)00147-1.
- ARIELI, A. et al. Assessment of energy expenditure by daily heart rate measurement: validation with energy accretion in sheep. *Livestock Production Science*, v.78, p.99-105, 2002. Available from: <[https://doi.org/10.1016/S0301-6226\(02\)00094-5](https://doi.org/10.1016/S0301-6226(02)00094-5)>. Accessed: Nov. 09, 2022. doi: 10.1016/S0301-6226(02)00094-5.
- BARKAI, D. et al. Estimation of energy intake from heart rate and energy expenditure in sheep under confinement or grazing condition. *Livestock Production Science*, v.73, p.237-246, 2002. Available from: <[https://doi.org/10.1016/S0301-6226\(01\)00251-2](https://doi.org/10.1016/S0301-6226(01)00251-2)>. Accessed: Nov. 09, 2022. doi: 10.1016/S0301-6226(01)00251-2.
- BROSH, A. et al. Effects of solar radiation, dietary energy, and time of feeding on thermoregulatory responses and energy balance in cattle in a hot environment. *Journal of Animal Science*, v.76, p.2671-2677, 1998. Available from: <<http://dx.doi.org/10.2527/1998.76102671x>>. Accessed: Nov. 09, 2022. doi: 10.2527/1998.76102671x.
- BROSH, A. Heart rate measurements as an index of energy expenditure and energy balance in ruminants: A review. *Journal of*

- 1 **Animal Science**, v.85, p.1213-1227, 2007. Available from: <<http://dx.doi.org/10.2527/jas.2006-298>>. Accessed: Nov. 09, 2022. doi: 10.2527/jas.2006-298. 1
2
3
4
5 BROUWER, E. Report of Sub-committee on Constants and Factors. 2
6 In **Proceedings of 3rd Symposium on Energy Metabolism**. 3
7 EEAP Publication 11. Academic Press, London, 1965. 4
8
9 DETMANN, E. et al. Métodos para análise de alimentos. Visconde 5
10 do Rio Branco, MG: Suprema, p. 214, 2012. 6
11
12 LANDAU, S. et al. Energy expenditure in Awassi sheep grazing 7
13 wheat stubble in the northern Negev Desert of Israel. **Livestock** 8
14 **Science**, v.105, p.265-271, 2006. Available from: <<https://doi.org/10.1016/j.livsci.2006.05.024>>. Accessed: Nov. 09, 2022. doi: 9
15 10.1016/j.livsci.2006.05.024. 10
16
17 LITTELL, R. C. et al. **SAS® for Mixed Models**. SAS Institute Inc., 11
18 Cary, 2006. 12
19
20 MACHADO, F.S. et al. Energy partitioning and methane emission 13
21 by sheep fed sorghum silages at different maturation stages. **Arq.** 14
22 **Bras. Med. Vet. Zootec.**, v.67, n.3, p.790-800, 2015. Available 15
23 from: <<https://doi.org/10.1590/1678-4162-7177>>. Accessed: Nov. 16
24 27, 2023. doi: 10.1590/1678-4162-7177. 17
25
26 MACHADO, F. S. et al. Technical note: A facility for respiration 18
27 measurements in cattle. **J. Dairy Sci.** v.99, p.4899-4906, 2016. 19
28 Available from: <<http://dx.doi.org/10.3168/jds.2015-10298>>. 20
29 Accessed: Nov. 09, 2022. doi: 10.3168/jds.2015-10298. 21
30
31 NATIONAL RESEARCH COUNCIL (NRC). Nutrient 22
32 requirements of small ruminants: sheep, goats, cervids, and new 23
33 world camelids. **National Academic Press**, 2007. Available from: 24
34 <<https://doi.org/10.17226/11654>>. Accessed: Nov. 27, 2024. doi: 25
35 10.17226/11654. 26
36
37 NICOL, A. M.; YOUNG, B. A. Short-term thermal and metabolic 27
38 responses of sheep to ruminal cooling: effects of level of cooling 28
39 and physiological state. **Canadian Journal of Animal Science**, 29
40 v.70, p.833-843, 1990. Available from: <<https://doi.org/10.4141/cjas90-102>>. Accessed: Nov. 09, 2022. doi: 10.4141/cjas90-102. 30
31
32 PUCHALA, R. et al. The relationship between heart rate and 31
33 energy expenditure in Alpine, Angora, Boer and Spanish goat 32
34 wethers consuming different quality diets at level of intake 33
35 near maintenance or fasting. **Small Ruminant Research**, v.70, 34
36 p.183-193, 2007. Available from: <<http://dx.doi.org/10.1016/j.smallrumres.2006.03.002>>. Accessed: Nov. 09, 2022. doi: 35
37 10.1016/j.smallrumres.2006.03.002. 36
38
39 OSS, D.B. et al. Assessment of the oxygen pulse and heart rate 37
40 method using respiration chambers and comparative slaughter 38
41 for measuring heat production of cattle. **Journal of Dairy** 39
42 **Science**, v.99, p.8885-8890, 2016. Available from: <<http://dx.doi.org/10.3168/jds.2016-11157>>. Accessed: Nov. 09, 2022. doi: 40
43 10.3168/jds.2016-11157. 41
44
45 SAS - Statistical Analysis System. **User's guide to statistics**. 42
46 Versão 9.1. Cary: SAS Institute. 2004. Available from: <https://support.sas.com/documentation/onlinedoc/91pdf/sasdoc_91/stat_ug_7313.pdf>. Accessed: Nov. 27, 2023. 43
47
48 TALMON, D. et al. Effect of animal activity and air temperature 44
49 on heat production, heart rate, and oxygen pulse in lactating 45
50 Holstein cows. **Journal of Dairy Science**, v.106, p.1475-1487, 46
51 2023. Available from: <<https://doi.org/10.3168/jds.2022-22257>>. 47
52 Accessed: Jan. 16, 2023. doi: 10.3168/jds.2022-22257. 48
53
54 TEIXEIRA, I.A.M.A. et al. Body composition, protein and energy 49
55 efficiencies, and requirements for growth of F1 Boer × Saanen goat 50
56 kids. **Journal of Animal Science**, v.95, p.2121-2132, 2017. Available 51
57 from: <<https://doi-org.ez27.periodicos.capes.gov.br/10.2527/jas.2016.1252>>. Accessed: Nov. 27, 2023. doi: 10.2527/jas.2016.1252. 52
58
59 VAN MILGEN, J. et al. Dynamic aspects of oxygen consumption 53
60 and carbon dioxide production in swine. **British Journal of** 54
61 **Nutrition**, v.78, n.3, p.397-410, 1997. Available from: <<https://doi.org/10.1079/bjn19970159>>. Accessed: Nov. 27, 2023. doi: 55
62 10.1079/bjn19970159. 56
63
64