

## **Anesthesia in Cocaine addicts: the risks of chemical interactions**

### **Anestesia em dependentes de Cocaína: os riscos das interações medicamentosas**

DOI:10.34119/bjhrv5n4-122

Recebimento dos originais: 14/04/2022

Aceitação para publicação: 30/06/2022

#### **Erika Amorim Melo Moreira**

Doutoranda em Ciências Farmacêuticas

Institution: Universidade Vila Velha (UVV)

Address: Avenida Comissário José Dantas de Melo, 21, Boa Vista II, Vila Velha - ES,  
CEP: 29102-920

E-mail: erika.amorim@uvv.br

#### **André Carvalho Gonçalves**

Graduando em Medicina

Institution: Universidade Vila Velha (UVV)

Address: Avenida Comissário José Dantas de Melo, 21, Boa Vista II, Vila Velha - ES,  
CEP: 29102-920

E-mail: carvalho.andre@uvvnet.com.br

#### **Elisardo Corral Vasquez**

Pós-Doutorado em Fisiologia

Institution: Universidade Vila Velha (UVV)

Address: Avenida Comissário José Dantas de Melo, 21, Boa Vista II, Vila Velha - ES,  
CEP: 29102-920

E-mail: elisardo.vasquez@uvv.br

#### **Kelly Ribeiro Mouro Barboza**

Doutoranda em Ciências Farmacêuticas

Institution: Universidade Vila Velha (UVV)

Address: Avenida Comissário José Dantas de Melo, 21, Boa Vista II, Vila Velha - ES,  
CEP: 29102-920

Email: kelly.barboza@uvv.br

#### **Pedro Henrique Cassaro Lirio**

Graduação em Farmácia

Institution: Universidade Federal do Espírito Santo (UFES)

Address: Av. Fernando Ferrari, 514, Vitória - ES, Brasil, CEP: 29075-910

E-mail: Pedro.lirio@edu.ufes.br

#### **João Magnago Santos**

Graduando em Medicina

Institution: Universidade Vila Velha (UVV)

Address: Avenida Comissário José Dantas de Melo, 21, Boa Vista II, Vila Velha - ES,  
CEP: 29102-920

E-mail: joomagnago@hotmail.com

## ABSTRACT

The interaction between cocaine and anesthetic drugs has been known since 1884, when it was first used for the treatment of morphine addiction and as local anesthetic. Due to the large number of users of the substance, the interaction with anesthetic drugs has been frequent and it has been seen that many drugs have their effects altered in the presence of cocaine. Through bibliographical survey in PubMed and CrossRef, aiming to highlight the risks of this interaction and conduct in cases of elective surgeries in these patients, we conducted this study and verified that so many drugs have to be avoided due to their interaction effects, but in the absence of clinical signs of intoxication, screening becomes ineffective.

**Keywords:** anesthesiology, Cocaine, drug interaction, psychotropic drugs, elective surgery.

## RESUMO

A interação entre cocaína e drogas anestésicas é conhecida desde 1884, quando foi usada pela primeira vez para o tratamento do vício em morfina e como anestésico local. Devido ao grande número de usuários da substância, a interação com drogas anestésicas tem sido frequente e tem sido observado que muitas drogas têm seus efeitos alterados na presença de cocaína. Através de pesquisa bibliográfica no PubMed e CrossRef, visando destacar os riscos desta interação e conduta em casos de cirurgias eletivas nestes pacientes, realizamos este estudo e verificamos que tantas drogas têm que ser evitadas devido aos seus efeitos de interação, mas na ausência de sinais clínicos de intoxicação, o rastreamento se torna ineficaz.

**Palavras-chave:** anestesiologia, Cocaína, interação com drogas, drogas psicotrópicas, cirurgia eletiva.

## 1 INTRODUCTION

The discovery of cocaine by the German chemist Albert Niemann, in 1860, led to profound changes in the way surgeons promote invasive interventions in patients as suggested by Ball and Westhorpe (2003). However, in the following two decades it was observed that besides inducing addiction in consumers, this drug also had undesirable chemical interactions, as reviewed elsewhere (Ball and Westhorpe, 2003, Alraies *et al.*, 2011), including the potentiation of chemical substances that was used by anesthesiologists during surgery procedures. At that time, Sigmund Freud tested in a friend who was addicted to morphine the hypothesis that cocaine could also interact with this opioid and interrupt that addiction, what happened successfully, but gradually was observed that the patients developed addiction to cocaine (Niemann, 1860).

Along years, the number of patients using illicit drugs, mainly cocaine, has increased in the world, becoming common the presence of effects that, acutely or chronically could interfere with anesthetic management according to Alraies *et al.* (2011); however, the early diagnosis of cocaine poisoning continues being a challenge, as it is basically clinical, not correlating with toxicological tests, because its inactive metabolite takes about 6-14 days for complete

elimination, making screening difficult as discussed by several authors (Weiss and Gawin, 1988, Lange and Hillis, 2001). According to different studies (Weiss and Gawin, 1988, Ritz, Cone and Kuhar, 1990) the effects of cocaine and its metabolites can generate synergistic interaction with anesthetics, as they act in synapses, involving neurotransmitters, a fact that can potentiate the sedation process or even decrease the metabolization of anesthetic drugs. In this context, it is important to evaluate and monitor these patients in surgeries to minimize the adverse effects of this drug interaction as suggested by Alraies *et al.* (2011).

## 2 MATERIALS AND METHODS

Here, we summarize the state of the art of literature exposing the risks of drug interaction between cocaine and anesthetic drugs, such as their harm to the patient and difficulties in perioperative management through this review. The search for publications was carried out in the electronic bibliographic database, such as PubMed. The descriptors used in the search were: anesthetic drugs, cocaine, drug interaction. It was used as inclusion criteria articles published as systematic or narrative review. Duplicate books, dissertations and studies were excluded.

## 3 RESULTS


### 3.1 MECHANISM OF ACTION

Cocaine and its derivatives came from the plant *Erythroxylum coca*. Belonging to the class of alkaloids, this substance has psychostimulant properties (Tsuchiya, 2017). Due to its diversity of compounds, cocaine can be used by many routes, such as oral, nasal, pulmonary or intravenous route (Corrêa *et al.*, 2014). As for its pharmacokinetics, cocaine is a benzoylmethylenoin, with a plasma half-life of about 30-60 min, suffering hydrolysis by plasma and hepatic esterases, producing inactive metabolites (Ball and Westhorpe, 2003), such as ecgonine methyl ester, which by degrading forms benzoylgonin, a urinary metabolite and Norcaide, metabolite produced through demethylation (Luft and Mendes, 2007).

Cocaine is considered a psychostimulant drug with mechanism of action due to the combination of increased production of catecholamines (dopamine, norepinephrine) and serotonin; and inhibition of reuptake of these substances with emphasis on the increasing availability of dopamine, a mechanism that explains the stimulant and addictive effects of the drug, in addition to the double effect on  $\alpha$  and  $\beta$ -adrenergic receptors, a fact that corroborates the presence of cardiopathies in these patients [Fig 1] (Alraies *et al.*, 2011). As second effect, cocaine has a class I antiarrhythmic effect, blocking sodium and potassium channels, preventing the spread of action potential (Fozzard *et al.*, 2005, Tsuchiya and Mizogami, 2013). Although

it has paradoxically opposite mechanisms of action, its psychostimulant effect is corroborated by the greater intensity of action in the pre-synaptic terminals to the detriment of the antiarrhythmic effect (Lange and Hillis, 2001).

Figure 1. Demonstration of the different sites of action of Cocaine. Adapted from Alraies, MC., Alraiyes AH. and Michota F., 2011. Should surgery be cancelled when surreptitious cocaine use is discovered before elective non-cardiac surgery ?. Middle East Journal of Anaesthesiology, 21(3), pp. 445-446.



### 3.2 INTERACTION OF COCAINE AND ANESTHETIC DRUGS

Cocaine, because its paradoxical effects, provides great concern in case of surgery. Anesthetic drugs, such as ketamine, increase the circulation of catecholamines, and the additive interaction to cocaine, can lead to overstimulation of  $\alpha$  and  $\beta$  receptors, which may culminate in cardiovascular events (Vadivelu *et al.*, 2014) as well as halothane, a halogenated anesthetic that sensitizes the catecholamine conduction system in the myocardium, and when used in cocaine addicts, can generate prolongation of QT interval and arrhythmias (Alraies *et al.*, 2011, Tsuchiya, 2017). We should also avoid other drugs that can increase QT interval, such as methadone and ondansetron [Table 1] (Ehret *et al.*, 2006, Roden, 2004, Dolenska, 2009)

In general anesthesia, the challenge in cocaine users is the control of arterial hypertension, since it is not indicated to administer  $\beta$ -blockers alone (Moran *et al.*, 2015), because they potentiate  $\alpha$ -agonism, giving priority to other vasodilators (Kuczkowski, 2003,

Kuczowski, 2004, Gazoni *et al.*, 2006). An excellent option would be pressure control with Dexmedetomidine, an  $\alpha_2$ -adrenergic agonist (Moran *et al.*, 2015). Despite the risk of hypertensive events, general anesthesia is preferred over local anesthesia due to the possibility of relative hypovolemia with lack of response to vasopressor's action, in addition to combative behavior, altered pain perception and cocaine-induced thrombocytopenia (Hill, Ogunnaiké and Johnson, 2006, Luft and Mendes, 2007), besides the additive effect to local anesthetics, with risk of toxicity, since the safe dose is unpredictable (Corrêa, *et al.*, 2014) and also because seizures by cocaine use do not decrease with anticonvulsant drugs or inhalation anesthetics (Luft and Mendes, 2007).

Regarding the use of muscle relaxers, succinylcholine, a depolarizing relaxant, presents a prolonged response in chronic users, due to the reduction of pseudocholinesterase, responsible for its metabolization, but in cases of acute intoxication, this fact occurs by the competition of cocaine with succinylcholine in metabolism, increasing the serum concentration of both [ table 1] (Fleming *et al.*, 1990; Alraies *et al.*, 2011).

Table 1. Relationship Between Classes of Drugs used in Anesthesia and their cardiovascular risk in cocaine users

Drugs		Cardiovascular adverse effect
Anesthetic	Ketamine	Release of catecholamine sensitizing the myocardium to catecholamines and contributing to the increased incidence of arrhythmias in these patients (ventricular tachycardia, ventricular extrasystole and ventricular fibrillation)
	Halogenated	Arterial hypertension
$\beta$ -blockers	Labetalol	Increased incidence of ventricular arrhythmias
	Propranolol	Arterial hypertension Complication of blood pressure control during surgery
$\alpha_2$ -adrenergic agonist	Dexmedetomidine	Hypertensive events Possibility of relative hypovolemia with lack of response to vasopressor's action
Vasoactive medications	Sodium nitroprusside	Higher incidence of ventricular arrhythmias
Opioids	Methadone	Prolongation of QT interval and arrhythmias
Selective serotonin 5-HT <sub>3</sub> -receptor antagonist which antiemetic activity	Ondansetron	Prolongation of QT interval and arrhythmias
Muscle relaxers	Succinylcholine	Increasing the serum concentration of both (cocaine and succinylcholine) and risk of the arrhythmias

#### 4 DISCUSSION

Surgeries in cocaine users have increased frequently, but despite all the important risks of drug interaction, studies have shown that since these patients show no signs of substance poisoning, the risk of a perioperative complication does not increase when compared to a non-user individual (Alraies *et al.*, 2011). Given that toxicological tests can present a result that does not match intoxication by the substance, as the elimination half-life of its inactive metabolite is up to 14 days, the only exclusion criterion for an elective surgery would be clinical symptoms,

such as excitation, weight loss, anxiety, requests for sedatives or opioid and digestive problems, to prevent cocaine interaction problems in the body (Vadivelu, *et al.*, 2014).

Cocaine has two important actions: psychostimulant and antiarrhythmic. The antiarrhythmic's action, with double effect in  $\alpha$  and  $\beta$ -adrenergic receptors, leads to an important interaction with ketamine due to the release of catecholamines, as well as halogenates does, sensitizing the myocardium to catecholamines and contributing to an increase in the incidence of arrhythmias in these patients. Because cocaine interacts with catecholamines and nitric oxide, it is important the awareness about interactions with commonly used drugs during anesthesia, such as beta-blockers, sodium nitroprusside and consequently complicating the control of blood pressure during the surgery [Table 1] (Bijker *et al.*, 2007). Despite that, Moon *et al.* (2019), conducted a retrospective cohort study, in which it was found that cocaine addicts do not need increased anesthetics or had a higher incidence of ventricular arrhythmias, with only higher blood pressure scores requiring vasodilators. These data emphasize the need for follow-up of these patients to minimize the effects arising from the drug-drug interaction presented in these cases.

However, from the possibility of drug interaction, it leads us to reflect on whether important interaction effects with other anesthetic have gone unnoticed over the years.

Patients presenting acute or chronic intoxication by drug abuse are an additional concerning for the anesthesiologist and surgeons during and after an operation, which sometimes it is possible to delay the scheduled surgery but in other occasions the patient needs an urgent or emergent intervention, and both surgeon and anesthesiologist must be able to deal with the undesirable changes that may occur in these patients. Preoperative targeted questions to the patient regarding drug use can help surgeon and anesthesiologist to know in the screening for drug addiction and can allows them to better take care of the patient during and in the postoperative intervention.

Evidence shows that there are important interactions that can complicate the perioperative period, however these patients do not present higher risks of complications than the general population as a more recent study (Moon *et al.*, 2020) has shown, corroborating the possibility of performing non-cardiac elective surgeries in these patients, if they do not show signs of intoxication, avoiding unnecessary postponement of surgical procedures (Ball and Westhorpe, 2003, Gazoni *et al.*, 2006).

#### **DECLARATION OF INTEREST STATEMENT**

The authors declare no conflicts of interest.

## REFERENCES

- Alraies, MC., Alraiyes AH. and Michota F., 2011. Should surgery be cancelled when surreptitious cocaine use is discovered before elective non-cardiac surgery ?. *Middle East Journal of Anaesthesiology*, 21(3), pp. 445-446.
- Ball, C. and Westhorpe, R., 2003. Local Anaesthesia—Freud, Koller and Cocaine. *Anaesthesia and Intensive Care*, 31(3), pp.247-247.
- Biebuyck, J., Fleming, J., Byck, R. and Barash, P., 1990. Pharmacology and Therapeutic Applications of Cocaine. *Anesthesiology*, 73(3), pp.518-531.
- Bijker, J., van Klei, W., Kappen, T., van Wolfswinkel, L., Moons, K. and Kalkman, C., 2007. Incidence of Intraoperative Hypotension as a Function of the Chosen Definition. *Anesthesiology*, 107(2), pp.213-220.
- Corrêa, C., Oliveira, L., Assis, J. and Barros, R., 2014. Anesthesia in patients who are users of crack and cocaine. *Revista Médica de Minas Gerais*, 24.
- Dolenska, S., 2009. Intraoperative cardiac arrest in acquired long QT syndrome. *British Journal of Anaesthesia*, 102(4), pp.503-505.
- Ehret, G., Voide, C., Gex-Fabry, M., Chabert, J., Shah, D., Broers, B., Piguet, V., Musset, T., Gaspoz, J., Perrier, A., Dayer, P. and Desmeules, J., 2006. Drug-Induced Long QT Syndrome in Injection Drug Users Receiving Methadone. *Archives of Internal Medicine*, 166(12), p.1280.
- Fozzard, H., Lee, P. and Lipkind, G., 2005. Mechanism of Local Anesthetic Drug Action on Voltage-Gated Sodium Channels. *Current Pharmaceutical Design*, 11(21), pp.2671-2686.
- Gazoni, F., Truffa, A., Kawamura, C., Guimarães, H., Lopes, R., Sandre, L. and Lopes, A., 2006. Complicações cardiovasculares em usuário de cocaína: relato de caso. *Revista Brasileira de Terapia Intensiva*, 18(4), pp.427-432.
- Hill, G., Ogunnaike, B. and Johnson, E., 2006. General anaesthesia for the cocaine abusing patient. Is it safe?. *British Journal of Anaesthesia*, 97(5), pp.654-657.
- Kuczkowski, K., 2003. Caesarean section in a cocaine-intoxicated parturient: regional vs. general anaesthesia?. *Anaesthesia*, 58(10), pp.1042-1043.
- Kuczkowski, K., 2004. The cocaine abusing parturient: a review of anesthetic considerations. *Canadian Journal of Anesthesia/Journal canadien d'anesthésie*, 51(2), pp.145-154.
- Lange, R. and Hillis, L., 2001. Cardiovascular Complications of Cocaine Use. *New England Journal of Medicine*, 345(5), pp.351-358.
- Luft, A. and Mendes, F., 2007. Anestesia no paciente usuário de cocaína. *Revista Brasileira de Anestesiologia*, 57(3), pp.307-314.



- Moon, T., Gonzales, M., Sun, J., Kim, A., Fox, P., Minhajuddin, A., Park, T. and Ogunnaike, B., 2019. Recent cocaine use and the incidence of hemodynamic events during general anesthesia: A retrospective cohort study. *Journal of clinical anesthesia*, 55, pp.146-150.
- Moon, T., Pak, T., Kim, A., Gonzales, M., Volnov, Y., Wright, E., Vu, K., Lu, R., Sharifi, A., Minhajuddin, A., Chen, J., Fox, P., Gasanova, I., Fox, A., Stewart, J. and Ogunnaike, B., 2020. A Positive Cocaine Urine Toxicology Test and the Effect on Intraoperative Hemodynamics Under General Anesthesia. *Anesthesia & Analgesia*, 132(2), pp.308-316.
- Moran, S., Isa, J. and Steinemann, S., 2015. Perioperative Management in the Patient with Substance Abuse. *Surgical Clinics of North America*, 95(2), pp.417-428.
- Niemann, A., 1860. Ueber eine neue organische Base in den Cocablättern. *Archiv der Pharmazie*, 153(2), pp.129-155.
- Ritz, M., Cone, E. and Kuhar, M., 1990. Cocaine inhibition of ligand binding at dopamine, norepinephrine and serotonin transporters: A structure-activity study. *Life Sciences*, 46(9), pp.635-645.
- Roden, D., 2004. Drug-Induced Prolongation of the QT Interval. *New England Journal of Medicine*, 350(10), pp.1013-1022.
- Tsuchiya, H., 2017. Anesthetic Agents of Plant Origin: A Review of Phytochemicals with Anesthetic Activity. *Molecules*, 22(8), p.1369.
- Tsuchiya, H. and Mizogami, M., 2013. Interaction of Local Anesthetics with Biomembranes Consisting of Phospholipids and Cholesterol: Mechanistic and Clinical Implications for Anesthetic and Cardiotoxic Effects. *Anesthesiology Research and Practice*, 2013, pp.1-18.
- Vadivelu, N., Mitra, S., Kaye, A. and Urman, R., 2014. Perioperative analgesia and challenges in the drug-addicted and drug-dependent patient. *Best Practice & Research Clinical Anaesthesiology*, 28(1), pp.91-101.
- Weiss, R. and Gawin, F., 1988. Protracted elimination of cocaine metabolites in long-term, high-dose cocaine abusers. *The American Journal of Medicine*, 85(6), pp.879-880.