



Universiteit
Leiden
The Netherlands

Circadian functioning of Locus Cœruleus of the nocturnal rat and diurnal rodent *Arvicanthis*

Caputo, R.; Poirel, V.J.; Paiva, I.; Boutillier, A.L.; Challet, E.; Meijer, J.H.; Raison, S.

Citation


Caputo, R., Poirel, V. J., Paiva, I., Boutillier, A. L., Challet, E., Meijer, J. H., & Raison, S. (2023). Circadian functioning of Locus Cœruleus of the nocturnal rat and diurnal rodent *Arvicanthis*. *Neuroscience Letters*, 799. doi:10.1016/j.neulet.2023.137091

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](#)

Downloaded from: <https://hdl.handle.net/1887/3714007>

Note: To cite this publication please use the final published version (if applicable).



Research article

Circadian functioning of Locus Cœruleus of the nocturnal rat and diurnal rodent *Arvicanthis*

Rosanna Caputo^{a,b}, Vincent-Joseph Poirel^a, Isabel Paiva^c, Anne-Laurence Boutillier^c, Etienne Challet^a, Johanna H. Meijer^b, Sylvie Raison^{a,*}

^a Institute of Cellular and Integrative Neurosciences, CNRS and University of Strasbourg, 8 Allée du Général Rouvillois, Strasbourg 67000, France

^b Department of Cell and Chemical Biology, Leiden University Medical Center, Leiden 2333 ZC, the Netherlands

^c Laboratoire de Neurosciences Cognitives et Adaptatives (LNCA), UMR 7364, Université de Strasbourg-CNRS, 12 rue Goethe, Strasbourg 67000, France



ARTICLE INFO

Keywords:

Circadian
Clock genes
Diurnal
Locus Cœruleus
Tyrosine hydroxylase

ABSTRACT

The noradrenergic Locus Cœruleus is one of the major arousal structures involved in inducing wakefulness. While brain noradrenaline (NA) amounts display 24-h variations, the origin of NA rhythm is currently unknown. In this study, we tested the hypothesis that NA rhythm could result from its rhythmic synthesis. Therefore, we investigated the 24-h expression profile of NA rate-limiting enzyme, tyrosine hydroxylase (*th*), in the Locus Cœruleus (LC) of the nocturnal rat and the diurnal rodent *Arvicanthis*, under 12 h:12 h light/dark (LD) and constant darkness (DD) conditions. In both species, *th* mRNA levels vary significantly over 24-h. In nocturnal rats, *th* mRNA profiles show a unimodal rhythm, with peak values in late day in LD, and in the middle of the subjective day in DD. In contrast, *th* mRNA rhythm in *Arvicanthis* is characterized by a bimodal profile, with higher levels at the beginning of the day and of the night in LD, and in the middle of the subjective day and night in DD. The rhythmic pattern of *th* expression may be dependent on a LC clock machinery. Therefore, we investigated the expression of three clock genes, namely *bmal1*, *per1*, and *per2*, and found that their mRNAs display significant variations between day and nighttime points in both species, but in opposite directions. These data show that NA rhythm may be related to circadian expression of *th* gene in both species, but differs between nocturnal and diurnal rodents. Furthermore, the phase opposition of clock gene expression in the rat compared to *Arvicanthis* suggests that the clock machinery might be one of the mechanisms involved in *th* rhythmic expression.

1. Introduction

The circadian system, orchestrated by the central pacemaker located in the suprachiasmatic nucleus (SCN), regulates the rhythm of many behavioral and physiological functions, including the sleep/wake cycle. Besides being under the control of the SCN, the distribution of wake and rest episodes is regulated by the arousal system [1,2]. The noradrenergic Locus Cœruleus (LC) is one of the major arousal structures involved in inducing wakefulness [3], and contributes to modulate circadian parameters of the sleep/wake cycle [4]. Evidence shows that the functioning of the LC itself is rhythmic. Circadian variations have been reported on the firing activity of LC neurons, which is higher during the active than the inactive phase [5]. Moreover, noradrenaline (NA) levels

in the brain show 24-h oscillations [6,7]. However, the origin of NA rhythm is so far unknown. In the LC, NA levels directly depend on the activity of tyrosine hydroxylase (TH), the rate-limiting enzyme for NA synthesis [8].

In this study, we tested the hypothesis that NA rhythm originates from its synthesis, and is therefore dependent on *th* expression. For that purpose, we investigated the 24-h profile of *th* mRNA levels. Considering the NA role in promoting wakefulness and the correlation of LC activity with the animal's arousal state [9,10], we hypothesize that NA synthesis rhythm differs between nocturnal and diurnal animals. Hence, we compared *th* expression in the nocturnal rat and in the diurnal rodent *Arvicanthis ansorgei*.

To go further, we raised the question about the origin of *th*

Abbreviations: Bmal1, Brain and Muscle ARNT-Like 1; CT, Circadian Time; DD, Constant Darkness; GRE, Glucocorticoid Response Element; LC, Locus Cœruleus; LD, Light/Dark; NA, Noradrenaline; Per1/2, Period 1/2; SCN, Suprachiasmatic Nuclei; Th, Tyrosine Hydroxylase; TTFL, Transcriptional-Translational Feedback Loops; ZT, Zeitgeber Time.

* Corresponding author at: INCI 8, Allée du Général Rouvillois, Strasbourg FR 67000, France.

E-mail address: raison@unistra.fr (S. Raison).

<https://doi.org/10.1016/j.neulet.2023.137091>

Received 7 November 2022; Received in revised form 17 January 2023; Accepted 18 January 2023

Available online 20 January 2023

0304-3940/© 2023 Elsevier B.V. All rights reserved.

rhythmicity. The circadian system may drive *th* rhythmic transcription through different mechanisms, i.e. through direct/indirect neuronal projections to the LC [5], through hormonal outputs, such as glucocorticoids [11], or *th* rhythmic expression may be under the control of clock genes expressed in the LC. Indeed, at a molecular level, circadian rhythms are generated by a self-sustained clock machinery, characterized by transcriptional-translational feedback loops (TTFL) [12]. The core of this TTFL consists in positive and negative loops involving specific clock genes and proteins (e.g. *Bmal1*, *Clock*, *Per1/2*, *Cry1/2*) which are responsible not only for their own transcription/repression, but even for the transcription/repression of other genes. In the mouse midbrain, CLOCK, which heterodimerizes with BMAL1, represses *th* expression by directly binding *Th* promoter at sequences named E-boxes [13]. E-box sequences have been identified in *Th* gene promoter of the rat [14,15].

In this study, we investigated whether a functional clock machinery, which may be involved in *th* rhythmic transcription, is present in the rat and *Arvicanthis* LC. For that purpose, we measured the daily expression of three clock genes: *bmal1*, *per1* and *per2*.

2. Materials & methods

2.1. Animals & housing conditions

All experiments were performed in accordance with the guidelines of the European Commission Directive 2010/63/EU, and the French Ministry of Higher Education and Research. Adult male Sudanian grass rats (*Arvicanthis ansorgei*) weighting 150–250 g (Chronobiotron, UAR 3415, CNRS Strasbourg, France), and adult male Wistar rats weighting 200–400 g (Charles River Laboratories, Saint-Germain-Nuelles, France) were used. All the animals were individually housed in Plexiglas cages, entrained to 12 h:12 h light/dark (LD) cycle (light: 150–200 lx, dark: red dim light, <5 lx, lights on and lights off defining Zeitgeber (ZT) 0 and ZT12, respectively), and provided with food and water *ad libitum*.

2.2. Experimental protocol

For *th* quantification, 84 rats and 84 *Arvicanthis* were entrained to 12 h:12 h LD cycle. Two days before the sacrifice, 42 rats and 42 *Arvicanthis* were exposed to constant darkness (DD), while the others were maintained in LD. Animals were killed by decapitation, after CO2 sedation at seven time points along the 24 h (every 4 h). For LD and DD rats sacrifice starts at ZT6 and CT6 respectively while it was at ZT2 and CT2 for *Arvicanthis* (Supplementary Fig. 1). The last time point was taken at ZT6/CT6 of the second experimental day for rats, indicated as ZT6' and CT6', respectively. For *Arvicanthis*, the last time point was ZT2/CT2 of the second experimental day, indicated as ZT2' and CT2', respectively. Brains were quickly removed, frozen in cold isopentane (-30 °C) and stored at -80 °C. Serial coronal 20 µm-sections were cut along the LC (Interaural from -1.4 mm to -0.2 mm, Paxinos & Watson, 1986) with a cryostat (Leica Instruments GmbH, Nussloch, Germany), and collected on sterile slides. Slides were stored at -20 °C until use.

For RT-qPCR quantification of clock genes, rats and *Arvicanthis* were sacrificed at ZT2, ZT10 and ZT18 (6 animals per group). Brains were quickly removed and frozen in cold isopentane (-30 °C), and subsequently stored at -80 °C. The right and left LC were dissected from four consecutive coronal sections (300 µm thickness, 2 mm diameter), using the 4th ventricle as reference. LC tissue was stored at -80 °C until use.

2.3. Radioactive in situ hybridization

Sense (GGGGAGCTGAAGGCTTATGGTGCAGGGCTGCTGTCTTCC-TACGGAG) and antisense (CTCCGTAGGAAGACAGCAGCCCTGCACCA-TAAGCCTCAGTCC) oligoprobes (Sigma Aldrich) for *th* were designed based on the rat *th* mRNA sequence. The sequence identity of the probes with the *Arvicanthis th* sequence were analyzed on the *Arvicanthis niloticus* (taxid:61156) genome using Nucleotide BLAST (<https://blast.ncbi.nlm.nih.gov/scd-rproxy.u-strasbg.fr/Blast.cgi>).

Probes (46-nucleotides) were 3' end-labelled with [³⁵S]-dATP (46.25 GBq/mol, PerkinElmer, Waltham MA) using the Terminal Deoxynucleotidyl Transferase (40U, Thermo Scientific). Specificity of the antisense probe was tested by a saturation test and by specific signals after hybridization within the rat and *Arvicanthis* LC, and within the *Arvicanthis* Substantia Nigra (Supplementary Fig. 2). Hybridization with oligoprobes on brain slices was performed as described by Caputo et al. 2022 [16], with the following modifications. Hybridization was carried out by depositing 80 µl of oligoprobes (0.1 pMol) in a solution containing 50 % deionized formamide, 4X sodium saline citrate (SSC), 1X Denhardt's solution, 0.25 mg/ml yeast totalRNA, 10 mg/ml salmon sperm DNA, 10 % dextran sulfate, and 300 mM dithiothreitol. After hybridization, the sections were washed in 1X SSC for 5 min at room temperature (RT) and stringency washes were performed in 0.1X SSC for 30 min at 50 °C and then in 0.05X SSC for 30 min at 52 °C. Sections were finally dehydrated, air dried at RT and then exposed to an autoradiographic film (Kodak BioMax; Kodak, Rochester, NY), with a ¹⁴C standard.

2.4. Quantitative analyses of *th* mRNA

Quantitative analyses of the autoradiograms were performed by using NIH ImageJ software. Six 20-µm-thick coronal sections (distance between two sections = 80 µm) along the caudo-rostral extension of the rat and *Arvicanthis* LC (see Fig. 1B and D), were analyzed. For each section, surface of hybridization and total optical density (OD) were measured in the right and left LC and the specific signal intensity was calculated by subtracting the non-specific OD, measured in surrounding area, where the signal was not specific. OD was normalized to relative levels of mRNA, using a ¹⁴C radioactive scale (KBq/g).

2.5. Real-time quantitative polymerase chain reaction

Total RNA was extracted from rat and *Arvicanthis* LC samples using RNeasy Lipid Tissue Mini Kit (74804, QIAGEN) following the manufacturer's recommendations. RNA concentration and purity were measured using a NanoDrop ND-1000 V 3.5 Spectrophotometer (NanoDrop Technologies, Wilmington, DE). Integrity of the RNA was assessed by using the 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). Synthesis of cDNA was performed using, 300 ng of rat total RNA and 400 ng of *Arvicanthis* total RNA, using Thermo Scientific™ Maxima™ H Minus cDNA Synthesis Master Mix, with dsDNase (15696019, Thermo Fisher Scientific) following the manufacturer's recommendations. RT-qPCR was performed using PowerUp™ SYBR™ Green Master Mix (A25776, Thermo Fisher Scientific) and run on the Applied Biosystems Real-Time PCR Instruments and on the CFX96 Touch Real-Time PCR Detection System (Bio-Rad). Real-time PCR amplification was performed after 120 s of denature at 95 °C, and then run for 40 cycles at 95 °C for 3 s, 60 °C for 30 s. The amplified clock genes were *bmal1*, *per1*, *per2* (Supplementary Table 1). Data were analyzed by using the 2^{-ΔΔCT} method [17], and normalized to that of two housekeeping genes: *gapdh* and *36b4* for rat data, two primers pairs for *gapdh*, amplifying different segments of the gene, for *Arvicanthis* data. Changes in clock gene expression were referenced to the mean expression level of the three ZTs.

2.6. Statistical analyses

As no significant differences were found between the right and left LC, the following analyses were performed on the left LC. To test the variation of *th* mRNA expression within the caudo-rostral extension of the rat and *Arvicanthis* LC among the seven time points, two-way analysis of variance (ANOVA II) was performed, with factors "time", "level" and their interaction. The effect of light condition (LD vs DD) was tested by ANOVA III, with factors "time", "level", "light condition" and their interactions. Differences between the two species (rat vs *Arvicanthis*)




Fig. 1. *In situ* hybridization of *th* antisense oligoprobe. In the rat (A) and *Arvicantis* (C) left (L) and right (R) Locus Cœruleus. Six successive 20- μ m-thick sections (distance = 80 μ m) are considered along the caudo-rostral extension of the rat (B) and *Arvicantis* (D) LC. Scale bars: 500 μ m.

were tested by ANOVA III, with factors “time”, “level”, “species” and their interactions. *Post-hoc* analyses were performed to examine the significant difference among time points. Individuals’ mean data were fitted by a non-linear regression using Cosinor analysis (SigmaPlot software, Jandel Scientific, Chicago, IL, USA). To fit either a unimodal (24 h) or a bimodal (24 h + 12 h, or 12 h) rhythm, the following equation was used: $y = A + (B * \cos(2\pi(\times - C)/24) + (D * \cos(2\pi(\times - E)/12))$; where A is the mean level (mesor), B the amplitude of the 24 h rhythm, C the acrophase of the 24 h rhythm, D amplitude of the 12 h rhythm, and E the acrophases of the 12 h rhythm.

For RT-qPCR data, differences among time points were analyzed using an ANOVA I test. Differences between species were tested using an ANOVA II, with factors “time”, “species”, and their interactions, followed by *Post-hoc* analyses. For all statistical procedures, the level of significance was set at $p \leq 0.05$. All data are presented as mean \pm SEM, unless otherwise stated. Sigma Plot (v14) and SPSS (v22) software were used for statistical analyses.

3. Results

3.1. Anatomical extension of the rat and *Arvicantis* Locus Cœruleus

Oligoprobes were designed according to the available rat *th* mRNA sequence. As the *Arvicantis ansorgei* genome is not available, *th* similarity was tested on the *Arvicantis niloticus* sequence. BLAST analysis revealed 97.8 % identity with *A. niloticus* predicted *th* mRNA sequence. Hybridization with *th* antisense oligoprobe allowed clear anatomical identification of left and right LC in rat (Fig. 1A) and *Arvicantis* (Fig. 1C). In the rat LC, as previously described by other authors [18], the left and right LC nuclei show a trapezoidal morphology and are positioned at the edges of the fourth ventricle. The left and right LC nuclei of *Arvicantis* are also located around the fourth ventricle, but display a crescent-shaped morphology. For both species, six successive

anatomical levels of the LC were considered for *th* quantification (Fig. 1B and D).

3.2. *Th* mRNA profile within the Locus Cœruleus in LD and DD

In both LD and DD conditions, *th* mRNA expression was quantified every-four hours for seven time points. For easier visual comparison, one time point in each graph is double-plotted in Fig. 2, which corresponds to ZT2 and CT2 for rat, and ZT6 and CT6 for *Arvicantis*. The levels of *th* expression were significantly different among the anatomical levels of LC both in the rat (LD $p < 0.001$, DD $p < 0.001$) and in *Arvicantis* (LD $p < 0.001$, DD $p < 0.001$).

In the rat, a significant effect of time on *th* expression is observed in both LD (Fig. 2A; $p = 0.012$) and DD (Fig. 2B, $p < 0.001$). Cosinor analyses showed a significant 24-h fit of *th* mRNA expression in both light conditions. The acrophases of the LD and DD rhythms are observed at ZT11.5 and CT6.4, respectively. The LD and DD *th* profiles are significantly different ($p < 0.001$). *Post-hoc* analyses show that the LD and DD profiles significantly differ at ZT2 vs CT2 ($p < 0.001$), ZT10 vs CT10 ($p = 0.027$), ZT14 vs CT14 ($p < 0.001$), ZT18 vs CT18 ($p < 0.001$), ZT22 vs CT22 ($p = 0.019$), and at ZT6' vs CT6' ($p < 0.001$).

In *Arvicantis*, the effect of time is significant in both LD (Fig. 2C, $p < 0.001$) and DD (Fig. 2D, $p < 0.001$) conditions. Cosinor analyses differed from the rat, and displayed a significant 12-h fit with peaks at ZT3.5 and ZT15.5 in LD, and at CT7.5 and CT19.5 in DD. The *th* profiles are significantly different between the LD and DD conditions ($p < 0.001$). *Post-hoc* analyses show that the LD and DD profiles significantly differ at ZT2 vs CT2 ($p = 0.002$), ZT14 vs CT14 ($p = 0.002$), ZT22 vs CT22 ($p = 0.005$), and at ZT2' vs CT2' ($p < 0.001$).

Analyses comparing the rat and *Arvicantis* *th* profiles show that the two patterns are significantly different both in LD ($p = 0.008$), and in DD ($p < 0.001$).




Fig. 2. *Th* mRNA rhythm in the rat and *Arvicantis* Locus Cereuleus. *In situ* hybridization of *th* mRNA levels within the rat (A and B) and *Arvicantis* (C and D) LC housed in 12:12 light/dark (LD, A and C) and constant darkness (DD, B and D) condition. Points are connected by a smoothing line. Dotted lines represent the significant Cosinor fit. White and black horizontal bars represent light and dark phases, respectively. Time is expressed as Zeitgeber Time (ZT) in LD and Circadian Time (CT) in DD. Rat groups were all $n \geq 5$, except: ZT18 ($n = 4$), ZT6' ($n = 3$) and CT6' ($n = 4$). *Arvicantis* groups were all $n \geq 5$, except: ZT10 ($n = 4$) and ZT2' ($n = 4$). The effect of time is significant in both species, in both lighting conditions. Data are presented as mean \pm SEM. Representative autoradiograms of *th* hybridization throughout the LD and DD cycle in the rat and *Arvicantis* left LC (E). Scale bar: 500 μ m.

3.3. Clock genes mRNA expression within the Locus Cereuleus in LD

To investigate whether rhythms in LC are different in nocturnal and diurnal mammals, we measured the expression of three genes central in the functioning of the circadian clock machinery, namely *Bmal1*, *Per1* and *Per2*, at ZT2, ZT10 and ZT18. Differences among the time points were measured for all the three clock genes, both in the rat and *Arvicantis* (Fig. 3). In the rat, the expression levels of *bmal1* significantly decrease from ZT2 to ZT10 ($p = 0.001$) and to ZT18 ($p < 0.001$). Coherently, *per1* and *per2* expressions significantly increase from ZT2 to ZT18 (both $p < 0.001$), from ZT2 to ZT10 (*per1* $p = 0.002$, *per2* $p = 0.021$), and from ZT10 to ZT18 only for *per2* ($p = 0.003$). In *Arvicantis*, clock genes expression varied in opposite ways compared to the rat. *Bmal1* expression varied significantly over the time points ($p = 0.05$) with higher levels at ZT18, while *per1* and *per2* levels decreased from ZT2 to ZT18 ($p = 0.007$ and $p = 0.018$, respectively), and from ZT10 to ZT18 for *per1* ($p = 0.017$). The clock gene mRNA variation over time was significantly different between rats and *Arvicantis*, for all three genes (*bmal1*: $p < 0.001$; *per1*: $p < 0.001$; *per2*: $p < 0.001$).

4. Discussion

4.1. Technical considerations and morphology of rat and Arvicantis LC

In situ hybridization quantification of *th* expression in the LC was performed using radioactive antisense oligoprobe designed on the rat *th* sequence. The high percentage of similarity (97.8 %) of rat *th* oligoprobe with *Arvicantis niloticus* genome, a species belonging to the same genus, made it suitable to detect *th* in the LC of *Arvicantis ansorgei*. The morphology and neurochemical organization of the LC in rats have been extensively described [18,19]. Though its neurotransmitter identity is quite complex, all neurons in the LC contain NA [19]. Therefore, the use of an antisense oligoprobe labelling *th*, noradrenaline rate-limiting enzyme, allowed an accurate morphological characterization of the LC both in the rat, and for the first time in *Arvicantis*. Differences between the two species relate mainly to the LC shape, which is more half-moon curved in the medial part of the *Arvicantis* LC, and extension, which is shorter in *Arvicantis* than in rats of about 200 μ m.

4.2. Rhythmic th mRNA expression in the rat and Arvicantis LC

In this study, we have shown that *th* mRNA levels in the LC are




Fig. 3. Clock gene mRNA levels in the rat and *Arvicanthis* Locus Cereuleus. RT-qPCR analyses of *bmal1*, *per1* and *per2* mRNA levels in the rat (blue) and *Arvicanthis* (red) LC housed in 12:12 light/dark condition. Time is expressed as Zeitgeber Time (ZT). The effect of time is significant for all genes. *Post-hoc* tests show differences among the time points: $p \leq 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***). Interaction between time and species is significant for all genes ($p \leq 0.05$). Experimental groups were all $n \geq 5$; data are presented as mean + SEM.

rhythmic in both a diurnal species, the *Arvicanthis*, and a nocturnal species, the rat, as previously described in the midbrain of another nocturnal mammal, the mouse [13,20].

The LC is a wake-promoting structure that participates in the circadian organization of the sleep/wake cycle. It is reasonable to hypothesize that temporal differences may exist between nocturnal and diurnal species, as they are awake in opposite temporal windows. Accordingly, we showed that the rhythm of *th* expression reflects the animals' behavioral activity pattern. In LD condition, *th* levels in LC show a unimodal rhythm with a peak occurring at the end of the day in the rat. By contrast, in *Arvicanthis*, whose activity pattern is characterized by a bimodal profile (as shown in Caputo et al. 2022 [16]) *th* levels display a bimodal profile, with a peak at the beginning of the day and a peak at the beginning of the night. Even when animals are kept in DD for two days, *th* expression held unimodal in rats and bimodal in *Arvicanthis*, coherently with their locomotor activity profile [21,22], showing that in both species *th* rhythmicity is endogenous. Coherently, our group demonstrated that in another arousal structure, the raphe nuclei, the 24-h mRNA pattern of tryptophan hydroxylase 2, serotonin rate-limiting enzyme, is unimodal in rats [23], and bimodal in *Arvicanthis* [16].

Although *th* mRNA is rhythmic in both lighting conditions within the same species, differences are observed between *th* LD and DD profiles. In the rat, *th* mRNA peak in DD occurs in the middle of the subjective day, 5 h before the peak observed in LD. In *Arvicanthis*, the two peaks observed in DD occur in the middle of the subjective day and in the middle of the subjective night, 4 h delayed compared to the respective peaks in LD. Altogether, these data suggest that although *th* rhythm is endogenous, light may affect *th* expression in both nocturnal and diurnal species, though with some differences. Coherently, in a previous study, we have shown that in the *Arvicanthis* raphe nuclei, tryptophan hydroxylase 2 rhythmic profile in DD differs from the rhythm in LD condition [16]. The importance of light on the noradrenergic system has been shown in rats, in which long term exposure to DD decreases the NA

neurotransmission in the frontal cortex, suggesting that light is necessary for physiological NA transmission [4,24]. In *Arvicanthis*, the effects of light on the LC system are scarce. It has been shown that 1 h light pulse applied 2 h after light-off increases cFOS expression in *Arvicanthis* LC, but not in mice [25], suggesting that light may have a different impact in diurnal and nocturnal rodents. How and to which extent light affects *th* rhythm requires further investigation.

Th rhythmicity suggests that the oscillating amount of NA observed in several brain regions [7,26], may be partly regulated at the level of its synthesis. NA levels are directly dependent on TH amount and activity. Variation of 24-h TH protein levels have been described in the ventral tegmental area (VTA) and nucleus accumbens of rats [27], and mice [20]. Natali and colleagues reported a circadian rhythm of TH activity in the mouse LC [28], while Cahill and Ehret reported a bimodal TH activity in the rat brain stem [29]. Future studies should investigate the rhythm and temporal relationship between *th* mRNA oscillation and TH protein levels/activity in the *Arvicanthis* and rat LC. Considering the role of NA in the circadian regulation of the sleep/wake cycle, different rhythms in NA synthesis may contribute to determine the opposite behavioral phenotype in diurnal and nocturnal animals.

4.3. Clock gene expression in the rat and *Arvicanthis* LC

To investigate possible mechanisms leading to different rhythmic expression of *th* in rats and *Arvicanthis*, we explored the rhythmicity of clock genes in the LC. Here we showed that the clock genes *Bmal1*, *Per1* and *Per2* are expressed in the LC and vary over time in LD condition. As expected, the expression profile of *Per1* and *Per2*, that are part of the negative feedback loop of the clock machinery, are opposite to the expression of *Bmal1*, in both species. Interestingly, clock genes expression in *Arvicanthis* is opposite to that in rats. These data suggest that clock machinery and circadian rhythms of the LC may be in anti-phase in nocturnal compared to diurnal species.

Our data in rats are coherent with a study in another nocturnal species, the hamster, showing that LC *per1* expression in TH cells is higher at CT12.5 compared to CT3 [30]. The presence of a rhythmic clock machinery leads to the possibility that *th* rhythm in the LC may be regulated by clock genes. In the mouse midbrain, *Th* expression can be regulated both by CLOCK [13] and by REV-ERB α [20], another clock protein. Furthermore, the rhythmic expression of *th* in the VTA is abolished in mice with deficient clocks [31]. E-box sequences, recognized by CLOCK:BMAL1 heterodimer, have been described in the rat *Th* promoter [14,15]. The presence of E-boxes in *Arvicanthis Th* gene needs to be verified.

Although clock genes may be involved in *th* rhythmic transcription, other mechanisms may contribute to *th* 24-h rhythm. In rats, a functional indirect SCN projection to the LC has been described to be responsible of LC rhythmic firing activity [5]. It is possible that signals resulting from indirect SCN projection may also affect *th* rhythm in *Arvicanthis*. Furthermore, circadian signals may be relayed indirectly to the LC through SCN hormonal outputs, such as glucocorticoids. *Th* gene promoter contains glucocorticoid response elements (GRE) [11], at least in rats. Therefore, corticosterone rhythm may be another factor involved in *th* mRNA oscillation.

In *Arvicanthis*, *th* expression is bimodal. Based on the three time points investigated in this study, it is not possible to establish whether clock genes expression may be bimodal as well. As for rats, other mechanism may contribute to *th* rhythmic mRNA expression. For example, corticosterone rhythm, which is bimodal in *Arvicanthis* [16,32], may contribute to *th* bimodal expression.

4.4. Conclusions

In conclusion, we found endogenous rhythms of *th* mRNA in the LC of both a diurnal and a nocturnal rodent, and we showed that *th* is influenced by light condition. Different temporal patterns of *th* levels and clock gene expression were observed in the nocturnal rat compared to the diurnal *Arvicanthis*, indicating that the rhythmic functioning of the LC is opposite in these species. Furthermore, daily changes in clock genes suggest that a functional clock machinery is present in the *Arvicanthis* and rat LC and might be involved in mechanisms regulating rhythmic functions within the LC, including *Th* transcription. These data highlight that LC rhythms differ between nocturnal and diurnal species and may contribute to their opposite temporal organization.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

We thank Dominique Ciocca and Nicolas Lethenet (Chronobiotron platform, UAR3415, CNRS, University of Strasbourg) for their help with animal experiments and for the *Arvicanthis ansorgei* breeding.

Funding

This work was supported by IDEX-Unistra [grant number: ANR-10-IDEX-0002-02].

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.neulet.2023.137091>.

[org/10.1016/j.neulet.2023.137091](https://doi.org/10.1016/j.neulet.2023.137091).

References

- [1] C.B. Saper, P.M. Fuller, N.P. Pedersen, J. Lu, T.E. Scammell, Sleep state switching, *Neuron* 68 (6) (2010) 1023–1042.
- [2] D.J. Dijk, S.N. Archer, Circadian and homeostatic regulation of human sleep and cognitive performance and its modulation by PERIOD3, *Sleep Med. Clin.* 4 (2) (2009) 111–125.
- [3] Jones, B.E., *Arousal systems*. Front Biosci, 2003. 8: p. s438–51.
- [4] M.M. González, G. Aston-Jones, Circadian regulation of arousal: role of the noradrenergic locus coeruleus system and light exposure, *Sleep* 29 (10) (2006) 1327–1336.
- [5] G. Aston-Jones, S. Chen, Y. Zhu, M.L. Oshinsky, A neural circuit for circadian regulation of arousal, *Nat. Neurosci.* 4 (7) (2001) 732–738.
- [6] J. Semba, M. Toru, N. Mataga, Twenty-four hour rhythms of norepinephrine and serotonin in nucleus suprachiasmaticus, raphe nuclei, and locus coeruleus in the rat, *Brain Res.* 397 (2) (1984) 211–218.
- [7] H. Agren, M. Koulou, J.M. Saavedra, W.Z. Potter, M. Linnoila, Circadian covariation of norepinephrine and serotonin in the locus coeruleus and dorsal raphe nucleus in the rat, *Brain Res.* 397 (2) (1986) 353–358.
- [8] M. Levitt, et al., Elucidation of the rate-limiting step in norepinephrine biosynthesis in the perfused guinea-pig heart, *J. Pharmacol. Exp. Ther.* 148 (1965) 1–8.
- [9] G. Aston-Jones, F.E. Bloom, Activity of norepinephrine-containing locus coeruleus neurons in behaving rats anticipates fluctuations in the sleep-waking cycle, *J. Neurosci.* 1 (8) (1981) 876–886.
- [10] J. Rajkowski, et al., State-related activity, reactivity of locus coeruleus neurons in behaving monkeys, *Adv. Pharmacol.* 42 (1998) 740–744.
- [11] T. Hagerty, W.W. Morgan, N. Elango, R. Strong, Identification of a glucocorticoid-responsive element in the promoter region of the mouse tyrosine hydroxylase gene, *J. Neurochem.* 76 (3) (2001) 825–834.
- [12] Ko, C.H. and J.S. Takahashi, *Molecular components of the mammalian circadian clock*. Hum Mol Genet, 2006. 15 Spec No 2: p. R271–7.
- [13] Sidor, M.M., et al., *Daytime spikes in dopaminergic activity drive rapid mood-cycling in mice*. Mol Psychiatry, 2015. 20(11): p. 1479–80.
- [14] B.P. Fung, S.O. Yoon, D.M. Chikaraishi, Sequences that direct rat tyrosine hydroxylase gene expression, *J. Neurochem.* 58 (6) (1992) 2044–2052.
- [15] S.O. Yoon, D.M. Chikaraishi, Isolation of two E-box binding factors that interact with the rat tyrosine hydroxylase enhancer, *J. Biol. Chem.* 269 (28) (1994) 18453–18462.
- [16] R. Caputo, V.-J. Poirel, E. Challet, J.H. Meijer, S. Raison, Bimodal serotonin synthesis in the diurnal rodent, *Arvicanthis ansorgei*, *FASEB J.* 36 (4) (2022) e22255.
- [17] K.J. Livak, T.D. Schmittgen, Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method, *Methods* 25 (4) (2001) 402–408.
- [18] R. Grzanna, M.E. Molliver, The locus coeruleus in the rat: an immunohistochemical delineation, *Neuroscience* 5 (1) (1980) 21–40.
- [19] L.A. Schwarz, L. Luo, Organization of the locus coeruleus-norepinephrine system, *Curr. Biol.* 25 (21) (2015) R1051–R1056.
- [20] S. Chung, E. Lee, S. Yun, H. Choe, S.-B. Park, H. Son, K.-S. Kim, D. Dluzen, I. Lee, O. Hwang, G. Son, K. Kim, Impact of circadian nuclear receptor REV-ERB α on midbrain dopamine production and mood regulation, *Cell* 157 (4) (2014) 858–868.
- [21] E. Challet, B. Pitrosky, B. Sicard, A. Malan, P. Pévet, Circadian organization in a diurnal rodent, *Arvicanthis ansorgei* Thomas 1910: chronotypes, responses to constant lighting conditions, and photoperiodic changes, *J. Biol. Rhythms* 17 (1) (2002) 52–64.
- [22] M. Cuesta, D. Clesse, P. Pévet, E. Challet, From daily behavior to hormonal and neurotransmitters rhythms: comparison between diurnal and nocturnal rat species, *Horm. Behav.* 55 (2) (2009) 338–347.
- [23] Z.S. Malek, H. Dardente, P. Pévet, S. Raison, Tissue-specific expression of tryptophan hydroxylase mRNAs in the rat midbrain: anatomical evidence and daily profiles, *Eur. J. Neurosci.* 22 (4) (2005) 895–901.
- [24] M.M. Gonzalez, G. Aston-Jones, Light deprivation damages monoamine neurons and produces a depressive behavioral phenotype in rats, *PNAS* 105 (12) (2008) 4898–4903.
- [25] D.D. Shuboni, S.L. Cramm, L. Yan, C. Ramanathan, B.L. Cavanaugh, A.A. Nunez, L. Smale, Acute effects of light on the brain and behavior of diurnal *Arvicanthis niloticus* and nocturnal *Mus musculus*, *Physiol. Behav.* 138 (2015) 75–86.
- [26] F.R. Cagampang, H. Okamura, S. Inouye, Circadian rhythms of norepinephrine in the rat suprachiasmatic nucleus, *Neurosci. Lett.* 173 (1–2) (1994) 185–188.
- [27] I.C. Webb, R.M. Baltazar, X.u. Wang, K.K. Pitchers, L.M. Coolen, M.N. Lehman, Diurnal variations in natural and drug reward, mesolimbic tyrosine hydroxylase, and clock gene expression in the male rat, *J. Biol. Rhythms* 24 (6) (2009) 465–476.
- [28] J.-P. Natali, A. McRae-Degueurce, P. Keane, G. Debilly, J.-F. Pujol, Genetic studies of daily variations of first-step enzymes of monoamines metabolism in the brain of inbred strains of mice and hybrids. II. Daily variations of tyrosine hydroxylase activity in the locus coeruleus, *Brain Res.* 191 (1) (1980) 205–213.
- [29] A.L. Cahill, C.F. Ehret, Circadian variations in the activity of tyrosine hydroxylase, tyrosine aminotransferase, and tryptophan hydroxylase: relationship to catecholamine metabolism, *J. Neurochem.* 37 (5) (1981) 1109–1115.

- [30] C.E. Mahoney, J. McKinley Brewer, E.L. Bittman, S. Yamazaki, Central control of circadian phase in arousal-promoting neurons, *PLoS One* 8 (6) (2013) e67173.
- [31] C.E. Koch, K. Begemann, J.T. Kiehn, L. Griewahn, J. Mauer, M. E. Hess, A. Moser, S. M. Schmid, J.C. Brüning, H. Oster, Circadian regulation of hedonic appetite in mice by clocks in dopaminergic neurons of the VTA, *Nat. Commun.* 11 (1) (2020).
- [32] L.A.W. Verhagen, P. Pévet, M. Saboureau, B. Sicard, B. Nesme, B. Claustrat, R. M. Buijs, A. Kalsbeek, Temporal organization of the 24-h corticosterone rhythm in the diurnal murid rodent *Arvicanthis ansorgei* Thomas 1910, *Brain Res.* 995 (2) (2004) 197–204.