



Universiteit
Leiden
The Netherlands

SARS-CoV-2 susceptibility and COVID-19 illness course and outcome in people with pre-existing neurodegenerative disorders: systematic review with frequentist and Bayesian meta-analyses

Smadi, M.; Kaburis, M.; Schnapper, Y.; Reina, G.; Molero, P.; Molendijk, M.L.

Citation

Smadi, M., Kaburis, M., Schnapper, Y., Reina, G., Molero, P., & Molendijk, M. L. (2023). SARS-CoV-2 susceptibility and COVID-19 illness course and outcome in people with pre-existing neurodegenerative disorders: systematic review with frequentist and Bayesian meta-analyses. *British Journal Of Psychiatry*, 223, 348-361. doi:10.1192/bjp.2023.43

Version: Publisher's Version

License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)

Downloaded from: <https://hdl.handle.net/1887/3674000>

Note: To cite this publication please use the final published version (if applicable).

Review

SARS-CoV-2 susceptibility and COVID-19 illness course and outcome in people with pre-existing neurodegenerative disorders: systematic review with frequentist and Bayesian meta-analyses

Muhannad Smadi, Melina Kaburis, Youval Schnapper, Gabriel Reina, Patricio Molero and Marc L. Molendijk

Background

People with neurodegenerative disease and mild cognitive impairment (MCI) may have an elevated risk of acquiring severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and may be disproportionately affected by coronavirus disease 2019 (COVID-19) once infected.

Aims

To review all eligible studies and quantify the strength of associations between various pre-existing neurodegenerative disorders and both SARS-CoV-2 susceptibility and COVID-19 illness course and outcome.

Method

Pre-registered systematic review with frequentist and Bayesian meta-analyses. Systematic searches were executed in PubMed, Web of Science and preprint servers. The final search date was 9 January 2023. Odds ratios (ORs) were used as measures of effect.

Results

In total, 136 primary studies (total sample size $n = 97\,643\,494$), reporting on 268 effect-size estimates, met the inclusion criteria. The odds for a positive SARS-CoV-2 test result were increased for people with pre-existing dementia (OR = 1.83, 95% CI 1.16–2.87), Alzheimer's disease (OR = 2.86, 95% CI 1.44–5.66) and Parkinson's disease (OR = 1.65, 95% CI 1.34–2.04). People with

pre-existing dementia were more likely to experience a relatively severe COVID-19 course, once infected (OR = 1.43, 95% CI 1.00–2.03). People with pre-existing dementia or Alzheimer's disease were at increased risk for COVID-19-related hospital admission (pooled OR range: 1.60–3.72). Intensive care unit admission rates were relatively low for people with dementia (OR = 0.54, 95% CI 0.40–0.74). All neurodegenerative disorders, including MCI, were at higher risk for COVID-19-related mortality (pooled OR range: 1.56–2.27).

Conclusions

Our findings confirm that, in general, people with neurodegenerative disease and MCI are at a disproportionately high risk of contracting COVID-19 and have a poor outcome once infected.

Keywords:

Alzheimer's disease; coronavirus disease 2019; dementia; mild cognitive impairment; Parkinson's disease.

Copyright and usage

© The Author(s), 2023. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

The novel 'coronavirus disease 2019' (COVID-19) is a widespread public health threat that is caused by a highly transmissible respiratory pathogen, 'severe acute respiratory syndrome (SARS-CoV-2)'.^{1,2} Although much has returned to normal in our everyday lives, the virus continues to spread and infect millions and to be lethal to thousands of people on a daily basis, across the globe.³ Early in the pandemic, it became clear that there are individual differences in COVID-19 infection susceptibility and severity.⁴ More than half of all COVID-19 casualties and intensive care unit (ICU) admissions were older adults.⁵

Both age and age-related comorbidities are known to be strong risk factors for the development of dementia.^{6–10} The dementias (i.e. Alzheimer's disease; Parkinson's disease with dementia; and mild cognitive impairment (MCI)) are a leading cause of impairment, dependence and mortality, especially among the elderly.^{10,11} People with dementia, including MCI, are more likely to have comorbid conditions that confer a vulnerability for other medical conditions, including COVID-19.¹² In addition, studies have suggested that individuals who have comorbid conditions are more likely to experience severe illness and require hospital admission due to COVID-19 infection.^{9,13} Previous data suggest that a dysregulated immune response in people with dementia can put them at further risk for COVID-19, leading to poor outcome, including death.^{14–16} Furthermore, people with dementia have been particularly susceptible to the stressors brought on by the pandemic and the social restrictions to help deter the spread of the virus.¹⁷ In

particular, social distancing may worsen stress in people with dementia owing to a disruption in routines developed to compensate for their memory loss.¹

Age-related comorbidities, immune dysregulation and exposure to stressors, as well as a reduced ability to comprehend the risks of infection and follow strict protocols to mitigate the spread of the virus, have all been related to infection risk and disease course.^{1,17–19} Consequently, people with dementia may be more susceptible to SARS-CoV-2 infection and a relatively poor course and outcome of this disease once infected.²⁰ A meta-analysis conducted early in the pandemic found a higher risk of death due to COVID-19 in those with dementia compared with those without dementia.² However, the authors reported substantial heterogeneity which remained unexplained, and there was evidence of publication bias. A later meta-analysis showed that the risk for mortality was higher in people with pre-existing dementia.²¹ A limitation of both meta-analyses is the small number of studies synthesised and the likelihood of duplicate data, as both included nationwide data from Italy and Korea multiple times, which may invalidate results.^{22–25} Therefore, we considered conducting an updated meta-analysis. Another reason for such an update is the rapidly evolving situation and recent influx of publications. In addition, past meta-analyses focused solely on dementia and not its precursor, MCI.

The current meta-analysis aims to quantify all eligible cohort studies reporting on infection risk for COVID-19 and course of

disease due to COVID-19 as a function of dementia status. We hypothesise that individuals with pre-existing dementia or MCI are more likely to become infected with SARS-CoV-2 and to experience worse COVID-19 severity and outcome (i.e. COVID-19-related hospital admission, ICU admission or mortality).

Method

The searches and methodology of this systematic review and meta-analysis are reported in accordance with the guidelines set out by Meta-analyses of Observational Studies in Epidemiology (MOOSE)²⁶ and Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).²⁷ A review protocol was drafted and pre-registered with the PROSPERO database (registration number CRD42022299941) and with the Open Science Framework (OSF).

Search and selection strategy

Systematic searches were executed in PubMed and Web of Science. These were supplemented with a non-systematic search in Google Scholar. A grey literature search on the preprint servers PsyArXiv and MedArXiv was also executed. The final search date was 9 January 2023. The search string and terms used per database are presented in the supplementary material available online at <https://dx.doi.org/10.1192/bjp.2023.43>. Eligibility of article inclusion was assessed independently by four members of the research team masked to each other's assessments, based on (a) title and abstract of potential papers, followed by (b) full-text assessment. A final decision on eligibility was made by four members of the review team (M.S., M.L.M., Y.S. and M.K.) based on the set eligibility criteria.

Eligibility criteria

Articles were included when they (a) reported SARS-CoV-2 infection rates (determined by any of the diagnostic methods, including blood, saliva analysis, polymerase chain reaction (PCR) and antibody testing) and the effect of infection on illness course of COVID-19, including mortality in people with pre-existing dementia (any type, including MCI) compared with controls; and (b) were written in English, Dutch, Spanish, Arabic, Hebrew, German, Italian or French. Articles were excluded if (a) no relevant outcome data could be extracted, (b) no original data were reported (e.g. reviews) or (c) they were case studies. When articles used data that we suspected might be overlapping, we included the article that was most informative for our purposes (see Article selection and overlapping data-sets in the supplementary material).

Exposure and outcome variables

Exposure variables were pre-existing dementias, including the precursor condition MCI, as defined by DSM-IV, DSM-5,^{28,29} ICD-10³⁰ or other validated assessment tools, compared with reference groups of people without a dementia. Outcome variables of interest included (a) SARS-CoV-2 infection risk (risk of getting infected with COVID-19), presented as the percentage of SARS-CoV-2 positive tests in the populations under study and (b) the course of COVID-19, further specified as (i) indicators of severity of the disease (e.g. symptomatic versus non-symptomatic, requiring respiratory assistance or not), (ii) hospital admission rates, (iii) ICU admission rates and (iv) COVID-19-related mortality rates.

Data extraction

The following data were extracted from eligible articles: average age (as mean or median in years), gender distribution at follow-up, country in which the study was performed; clinical data (i.e. method of diagnostic assessment, type of disorder), validity of assessment, the covariates that were used in statistical analyses, differences in outcome in covariate adjusted and unadjusted models, whether time-varying covariates were used, the analytical strategy that was used; and raw numbers or effect-size estimates and corresponding 95% confidence intervals (95% CIs) on outcome data. Data extraction was performed independently and masked, by at least two members of the review team (M.S., M.L.M., Y.S. and/or M.K.).

Measures of effect

We extracted ORs and corresponding 95% CIs as measures of effect. Where reported, we extracted data from analyses that controlled for the largest number of potential confounders or that came from (propensity-) matched samples. When results were reported as hazard ratios or risk ratios and raw data were not available, we interpreted these as an OR when the incidence of the reported outcome was <20%. Hazard ratios and risk ratios based on data reporting on an incidence of outcome >20% were transformed.^{31–33}

Assessment of methodological quality

The methodological quality of input studies was scored by three members of the review team (M.S., M.L.M. and Y.S.), who were masked to each other's assessment, using the Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies recommended by the US National Institutes of Health.³⁴

Statistical analysis

All analyses were performed in JASP version 0.17.1 for Apple Silicon (JASP Team, University of Amsterdam, Netherlands; <https://jasp-stats.org/download>). To check the robustness of results, analyses were also performed in IBM SPSS Statistics version 28 for Macintosh and STATA version 17 for Macintosh. Random-effects frequentist meta-analyses were used to pool the data on SARS-CoV-2 infection risk, COVID-19 course, hospital admissions, ICU admissions and mortality rates in relation to the types of pre-existing dementia. Statistical significance was set at $P < 0.05$. Heterogeneity among studies was quantified using the I^2 measure and assessed for statistical significance using the Q^2 statistic.³⁵ Meta-analyses were repeated using a Bayesian approach to verify robustness of results over different analytical approaches. When heterogeneity in outcome was present, subgroup and meta-regression analyses were performed with the aim of identifying study or population characteristics that might explain the heterogeneity. Potential moderators included the percentage of females, average age and methodological quality scores per sample. Subgroup analysis by geographical region was also performed if heterogeneity in outcome was present. Publication bias was assessed by means of Kendall's tau.³⁵

Results

Of the 5548 candidate articles that we retrieved, 136 met the eligibility criteria (Fig. 1). Supplementary Tables 1 and 2 list all the articles that were included for full-text assessment as well as reasons for final inclusion and exclusion.

Tables 1 and 2 provide demographic and clinical information on the samples in the studies included, stratified by SARS-CoV-2 susceptibility and COVID-19 course and outcome, respectively. The median age was 70.1 years (range 35–89.5 years), the percentage




Fig. 1 Flowchart on identification, screening and inclusion of eligible publications. ICU, intensive care unit.

of females was 53% (range 31–82%) and the median sample size per analysis was 94 624 (range 46–62 250 998). The methodological quality of the majority of input studies was high (Supplementary Tables 16 and 17). Supplementary Box 2 lists studies in which data-sets were (suspected to be) used more than once and the choices that we subsequently made to ensure that data on which we performed our analyses were independent. Supplementary Tables 3(a) and 3(b) provide further information on potential overlap and actions taken per analysis. It should be noted that when nationwide data were available for analysis alongside data gathered more locally, we ran analyses once with the nationwide data included and the local data excluded and once with the local data included and the nationwide data excluded. Therefore, we occasionally reported on fewer data-sets per analysis relative to the numbers provided in the flowchart.

SARS-CoV-2 infection risk

The odds for a SARS-CoV-2 positive test result were increased for people with documented pre-existing dementia (OR = 1.83, 95% CI 1.16–2.87; Table 3). These results were evident in multivariable analyses controlling for potential confounding factors such as age, gender and other comorbidities, but not in crude analyses (Supplementary Table 5). Alzheimer's disease, Parkinson's disease and mixed dementia were all associated with an increase in SARS-CoV-2 susceptibility (Fig. 2). When replacing nationwide data with local data, an increase in SARS-CoV-2 susceptibility remained evident in people with Alzheimer's disease, but not in those with the other disorders (Supplementary Table 4). Between-study heterogeneity in outcome was evident in all analyses (Table 3 and Supplementary Table 5). A small positive association between percentage of females and odds for infection was found in people with

dementia (Supplementary Table 7). Methodological quality was not associated with between-study heterogeneity (Supplementary Table 6). The odds for infection risk for all categories of neurodegenerative disorder were not evident in the data gathered in Asia, except for dementia (Supplementary Table 9).

COVID-19 course and outcome

People with pre-existing dementia were more likely to experience a severe COVID-19 course, once infected, relative to people in control conditions (OR = 2.66, 95% CI 1.16–6.12; Supplementary Table 5). This was also evident, although with attenuated effect size, in studies utilising multivariable analyses (OR = 1.43, 95% CI 1.00–2.03; Table 4). People with pre-existing dementia were at lower risk for ICU admission (OR = 0.54, 95% CI 0.40–0.74), but at higher risk for COVID-19-related hospital admission (OR = 1.60, 95% CI 1.09–2.35) and mortality (OR = 1.58, 95% CI 1.39–1.79; Fig. 3) in studies utilising multivariable analyses (Table 4). People with Alzheimer's disease were at higher risk for COVID-19-related hospital admission (OR = 3.72, 95% CI 2.35–5.90), but people with MCI, Parkinson's disease or mixed dementia were not. Based on a single study, it was found that people with Alzheimer's disease or Parkinson's disease were at higher risk for COVID-19-related ICU admissions (pooled OR range: 1.55–1.65; Table 4). All patient groups were at higher risk for COVID-19-related mortality (pooled OR range: 1.56–2.27; Table 4). When replacing nationwide data with local data, higher odds for COVID-19-related mortality remained evident for people with dementia or Parkinson's disease (Supplementary Table 4). Between-study heterogeneity in outcome was observed in most analyses (see Table 4 for two exceptions). Average age was positively associated with odds for COVID-19-related hospital admission in people with Parkinson's disease

Table 1. Characteristics of the studies included and samples reporting on SARS-CoV-2 infection risk.

Study	N	Age	Female (%)	Predictor	Country
Ajayi et al. 2020 ³⁶	657	55 ^{AVG.}	40	Dem	UK
Beobide Telleria et al. 2022 ³⁷	436	87 ^{MED.}	72	Dem	Spain
Castilla et al. 2021 ³⁸	643 757	44 ^{MED.}	37	Dem	Spain
de Malherbe et al. 2022 ³⁹	881	89 ^{AVG.}	79	AD	France
Del Ser et al. 2021 ⁴⁰	913	82 ^{AVG.}	65	AD	Spain
Emmerson et al. 2022 ⁴¹	9571	82 ^{AVG.}	68	Dem	Wales
Karapetyan et al. 2021 ⁴²	99 811	44 ^{AVG.}	60	Dem	Germany
Kim et al. 2022b ⁴³	129 120	45 ^{MED.}	36	Dem	Korea
Orlando et al. 2021 ⁴⁴	3497	48 ^{AVG.}	54	Mix, PD	Italy
Pan et al. 2021 ⁴⁵	12 384	78 ^{AVG.}	57	CI	USA
Proffili et al. 2020 ⁴⁶	1840	65 ^{MED.}	48	Dem	Italy
Scherbaum et al. 2021 ⁴⁷	30 872	78 ^{MED.}	58	PD	Germany
Smith et al. 2021 ⁴⁸	124 167	n.a.	n.a.	Dem	USA
Seon et al. 2021 ⁴⁹	123 480	53 ^{MED.}	61	Dem	Korea
Soldevila et al. 2022 ⁵⁰	8021	86 ^{AVG.}	74	Dem	Spain
Tahira et al. 2021 ⁵¹	12 863	75 ^{MED.}	49	AD, PD, Dem	UK
Wang et al. 2021a ⁵²	61 916 260	47 ^{MED.}	54	Dem	USA
Wang et al. 2021b ⁵³	446	65 ^{MED.}	66	AD, Dem	Korea
Wang et al. 2021c ⁵⁴	60 446	67 ^{AVG.}	67	Dem, AD, PD, Mix	UK
Wong & Lovier 2022 ⁵⁵	3257	74 ^{AVG.}	58	Dem	USA
Worcel et al. 2021 ⁵⁶	173	80 ^{AVG.}	61	AD, Dem	France
Yu et al. 2021 ⁵⁷	13 338	70 ^{AVG.}	51	AD, PD	UK
Zenesini et al. 2022 ⁵⁸	10 172	76 ^{AVG.}	41	PD	Italy
Zhou et al. 2021a ⁵⁹	3884	57 ^{AVG.}	55	AD, Dem	UK

a. Studies are divided by outcome: 'infection risk' and 'course and outcome'. The latter includes only participants with positive COVID-19 infection. AVG., average (mean); MED., median; AD, Alzheimer's disease; n.a., not applicable; CI, cognitive impairment; Dem, dementia; Mix, mixed dementia; PD, Parkinson's disease.

(Supplementary Table 8). In crude analyses, average age was positively associated with odds for mortality in people with dementia or Alzheimer's disease (Supplementary Table 7). The percentage of females was positively associated with odds for mortality in people with Alzheimer's disease in studies utilising crude analyses (Supplementary Table 7) and in people with MCI in studies utilising multivariable analyses (Supplementary Table 8). Methodological quality was not associated with any of the outcomes (Supplementary Tables 7 and 8). The odds of experiencing severe COVID-19, hospital admission, ICU admission and mortality for all categories of neurodegenerative disorders differed by continent (Supplementary Tables 9 and 10).

Bayesian meta-analysis

Supplementary Tables 11–14 present the odds ratios and 95% confidence intervals based on Bayesian analysis. For ease of comparison, frequentist results are also reported in these tables. Overall, the Bayesian analyses yielded largely similar results to the frequentist approach across all neurodegenerative disorders and the evidence for the alternative hypotheses, in case of significant findings, ranges from moderately strong (Bayes factor between 3 and 10) to extremely strong (Bayes factor >100).

Discussion

This systematic review with meta-analysis, which synthesised 136 primary studies, corroborates that individuals with pre-existing neurodegenerative disorders (i.e. dementia, Alzheimer's disease, Parkinson's disease, MCI or mixed dementia) have an increased susceptibility for SARS-CoV-2 infection and, in general, have higher morbidity and mortality rates for COVID-19. A notable observation is the lower risk for ICU admission in people with dementia. Large sample sizes and convergence of findings using both the frequentist and the Bayesian methods suggest robustness of the findings.

Susceptibility for SARS-CoV-2 infection and neurodegenerative disorders

The odds of infection with SARS-CoV-2 are about 1.5 to 3.0 times higher in individuals with pre-existing neurodegenerative disorders. Age and gender are known risk factors for various chronic diseases.¹⁷² In fact, older individuals are more susceptible to SARS-CoV-2¹⁷³ because of age-related changes in the immune system, which deteriorate immune response and efficiency.^{18,174} The living conditions of individuals with a neurodegenerative disorder may be a risk factor, since long-term care facilities (e.g. nursing homes) are predominantly tenanted by the elderly, 48–50.4% of whom have Alzheimer's disease or other dementias.^{10,175} The combination of age and age-related comorbidities (e.g. neurodegenerative disorders) with proximity and exposure of vulnerable individuals in communal housing, through shared (overcrowded) spaces, may translate to increased susceptibility to COVID-19.¹⁷ Nevertheless, associations that were controlled for age also yielded significant findings. Additionally, factors such as poor health behaviour (e.g. decreased physical activity) and non-adherence to public health measures may play a significant role in explaining the increased risk for infection with SARS-CoV-2 in individuals with neurodegenerative disorders. This may be attributed to the inability to comprehend the severity of contracting the virus and thus the necessity of complying with the protocols, owing to memory loss and cognitive impairment in individuals with dementia or MCI.^{2,17} A further explanation may be an unwillingness to adhere to the measures owing to apathy,² which is evident in individuals with dementia.¹⁷⁶ There were no data on whether there was preferential testing among people with dementia that might have led to an increased likelihood of having a diagnosis.

In some instances, we were unable to run subgroup analyses for some of the disorder types by continent (e.g. Asia) owing to the lack of data. Nevertheless, there were some differences in SARS-CoV-2 susceptibility among the geographic continents (i.e. Asia, America, Europe). This might be explained by the financial opportunities of each country to implement safety measures. In addition,

Table 2. Characteristics of the studies included and samples reporting on COVID-19 course and outcome.

Study	N	Age	% Female	Predictor	Outcome	Country
Ajayi et al. 2020 ³⁶	39	73 <small>AVG.</small>	44%	Dem	[5]	UK
Ajayi et al. 2021 ⁶⁰	75	55 <small>AVG.</small>	47%	Dem	[5]	UK
Alqahtani et al. 2021 ⁶¹	101	75 <small>AVG.</small>	58%	Dem	[5]	KSA
An et al. 2021 ⁶²	5596	59 <small>AVG.</small>	53%	Dem	[2], [5]	Korea
Atkins et al. 2020 ⁶³	507	73 <small>AVG.</small>	39%	Dem	[4], [5]	UK
Bae et al. 2021 ⁶⁴	1232	61 <small>AVG.</small>	39%	Dem	[5]	Korea
Baker et al. 2021 ⁶⁵	100	67 <small>AVG.</small>	45%	Dem	[2], [3], [5]	USA
Banoei et al. 2021 ⁶⁶	250	69 <small>MED.</small>	56%	Dem	[5]	USA
Becerra-Muñoz et al. 2021 ⁶⁷	1520	76 <small>MED.</small>	40%	Dem, PD	[5]	Multiple
Bennett et al. 2021 ⁶⁸	1 926 526	47 <small>AVG.</small>	55%	Dem	[2]	USA
Beobide Telleria et al. 2022 ³⁷	173	87 <small>MED.</small>	72%	Dem	[5]	Spain
Bhargava et al. 2021 ⁶⁹	656	64 <small>AVG.</small>	48%	Dem	[5]	USA
Bianchetti et al. 2020 ⁷⁰	627	76 <small>AVG.</small>	53%	Dem	[5]	Italy
Bielza et al. 2021 ⁷¹	630	87 <small>MED.</small>	65%	Dem	[5], [2]	Spain
Booij et al. 2022 ⁷²	134	85 <small>AVG.</small>	60%	Dem	[5]	Netherlands
Boye et al. 2021 ⁷³	4298	70 <small>AVG.</small>	50%	Dem	[4], [5]	USA
Bucholc et al. 2022 ⁷⁴	6036	72 <small>MED.</small>	47%	Dem	[5]	N. Ireland
Busetto et al. 2020 ⁷⁵	92	71 <small>AVG.</small>	62%	Dem	[3], [5]	Italy
Caliskan and Saylan, 2020 ⁷⁶	565	50 <small>AVG.</small>	n.a.	Dem	[3], [5]	Turkey
Carrillo-Garcia et al. 2021 ⁷⁷	165	89 <small>MED.</small>	69%	Dem	[5]	Spain
Castilla et al. 2021 ³⁸	35 387	44 <small>MED.</small>	37%	Dem	[4], [5]	Spain
Chang et al. 2020 ⁷⁸	710 980	71 <small>MED.</small>	50%	Mix	[4]	USA
Chatterjee et al. 2021 ⁷⁹	2337	65 <small>MED.</small>	80%	Dem	[5]	Netherlands
Chen et al. 2022 ⁸⁰	1 271 033	53 <small>AVG.</small>	60%	Dem	[5]	USA
Choi et al. 2021 ⁸¹	7590	40 <small>MED.</small>	60%	Dem	[3], [5]	Korea
Chojnicki et al. 2021 ⁸²	322	78 <small>AVG.</small>	62%	CI	[5]	Poland
Cisterna-Garcia et al. 2022 ⁸³	86 867	54 <small>MED.</small>	53%	Dem	[3], [4], [5]	Spain
COVIDSurg, 2021 ⁸⁴	1063	90 <small>MED.</small>	66%	Dem	[5]	Multiple
Covino et al. 2020 ⁸⁵	69	84 <small>MED.</small>	46%	Dem	[2], [5]	Italy
Covino et al. 2021a ⁸⁶	729	85 <small>MED.</small>	53%	Dem	[5]	Italy
Covino et al. 2021b ⁸⁷	729	85 <small>MED.</small>	53%	Dem	[5]	Italy
Cummins et al. 2021 ⁸⁸	1781	35 <small>MED.</small>	45%	Dem	[3], [4], [5]	UK
de Marcaida et al. 2020 ⁸⁹	36	75 <small>MED.</small>	36%	PD	[5]	USA
De Smet et al. 2020 ⁹⁰	81	85 <small>MED.</small>	59%	Dem	[5]	Belgium
Del Ser et al. 2021 ⁴⁰	62	82 <small>AVG.</small>	39%	AD	[2]	Spain
Descamps et al. 2022 ⁹¹	90 950	85 <small>MED.</small>	55%	Dem	[3], [5]	France
Ellis et al. 2022 ⁹²	1071	84 <small>MED.</small>	52%	Dem, MCI	[4], [5]	Australia
Escribà-Salvans et al. 2022 ⁹³	78	85 <small>AVG.</small>	76%	CI	[2]	Spain
Esme et al. 2021a ⁹⁴	16 942	71 <small>AVG.</small>	51%	Dem	[5]	Turkey
Esme et al. 2021b ⁹⁴	3172	71 <small>AVG.</small>	51%	Dem	[5]	Turkey
España et al. 2021a ⁹⁵	9121	84 <small>AVG.</small>	70%	Dem	[5]	Spain
España et al. 2021b ⁹⁵	2140	84 <small>AVG.</small>	70%	Dem	[5]	Spain
Esteban et al. 2021 ⁹⁶	113	87 <small>MED.</small>	82%	Dem	[2]	Argentina
Fasano et al. 2020 ⁹⁷	1486	71 <small>AVG.</small>	48%	PD	[4], [5]	Italy
Fathi et al. 2021 ⁹⁸	3732	78 <small>AVG.</small>	74%	AD, PD	[5]	Iran
Filardo et al. 2020 ⁹⁹	337	58 <small>MED.</small>	33%	Dem	[5]	USA
Filipe et al. 2021 ¹⁰⁰	3550	68 <small>MED.</small>	41%	Dem	[3], [5]	Switzerland
Fumagalli et al. 2021 ¹⁰¹	221	82 <small>AVG.</small>	39%	Dem	[5]	Italy
Gale and Boland, 2021 ¹⁰²	1181	59 <small>AVG.</small>	40%	Dem	[5]	UK
Ge et al. 2021 ¹⁰³	167 500	43 <small>MED.</small>	52%	Dem	[2], [5]	Canada
Genet et al. 2020 ¹⁰⁴	201	86 <small>AVG.</small>	67%	Dem	[5]	France
Geriatric et al. 2021 ¹⁰⁵	5711	74 <small>MED.</small>	45%	Dem	[3], [5]	Multiple
Ghaffari et al. 2021 ¹⁰⁶	361	62 <small>AVG.</small>	41%	Dem	[2]	Iran
Gómez Antúnez et al. 2020 ¹⁰⁷	746	69 <small>MED.</small>	43%	Dem, Mix	[5]	Spain
Harrison et al. 2020 ¹⁰⁸	31 461	50 <small>MED.</small>	55%	Dem	[5]	USA
Hasani Azad et al. 2021 ¹⁰⁹	2351	47 <small>AVG.</small>	48%	Dem	[3]	Iran
Hatamabadi et al. 2022 ¹¹⁰	5318	60 <small>MED.</small>	43%	AD, PD	[5]	Iran
Hippisley-Cox et al. 2021 ¹¹¹	6 952 440	53 <small>AVG.</small>	52%	Dem, PD	[4], [5]	UK
Hwang et al. 2020 ¹¹²	103	68 <small>AVG.</small>	50%	AD	[5]	Korea
Izurieta et al. 2021 ¹¹³	24 367 476	75 <small>MED.</small>	56%	CI, PD	[4], [5]	USA
Kang and Kong, 2021 ¹¹⁴	4141	50 <small>MED.</small>	58%	Dem	[2], [5]	Korea
Karapetyan et al. 2021 ⁴²	46 071	44 <small>AVG.</small>	61%	Dem	[2]	Germany
Ken-Dror et al. 2020 ¹¹⁵	429	70 <small>AVG.</small>	44%	Dem	[5]	UK
Kim et al. 2020 ¹¹⁶	2959	40 <small>MED.</small>	39%	Dem	[2]	Korea
Kim et al. 2021 ¹¹⁷	2254	58 <small>MED.</small>	64%	Dem	[5]	Korea
Kim et al. 2022a. (65-74) ¹¹⁸	17 890	69 <small>MED.</small>	51%	Dem	[2], [4], [5]	USA
Kim et al. 2022a. (75-84) ¹¹⁸	9500	79 <small>MED.</small>	53%	Dem	[2], [4], [5]	USA
Kim et al. 2022a. (85+) ¹¹⁸	4380	89 <small>MED.</small>	59%	Dem	[2], [4], [5]	USA
Kim et al. 2022b. ⁴³	19 912	45 <small>MED.</small>	36%	Dem	[2]	Korea
Kong et al. 2021 ¹¹⁹	5307	50 <small>AVG.</small>	59%	Dem	[2], [5]	Korea
Kostev et al. 2022 ¹²⁰	28 311	73 <small>AVG.</small>	49%	Dem, VD, AD	[5]	Germany
Kyoung et al. 2021 ¹²¹	1697	47 <small>MED.</small>	58%	Dem	[5]	Korea
Lazcano et al. 2021 ¹²²	91 629	55 <small>AVG.</small>	58%	Dem	[5]	Spain
Li et al. 2020 ¹²³	42	75 <small>MED.</small>	68%	AD	[5]	China
Livingston et al. 2020 ¹²⁴	131	75 <small>AVG.</small>	52%	Dem	[5]	UK

(Continued)

Table 2. (Continued)

Study	N	Age	% Female	Predictor	Outcome	Country
Lozano-Montoya et al. 2021 ¹²⁵	300	86 ^{AVG.}	63%	Dem	[5]	Spain
Lu et al. 2021 ¹²⁶	608 251	74 ^{MED.}	56%	PD, CI	[4], [5]	USA
Lucijanić et al. 2022 ¹²⁷	2586	70 ^{MED.}	44%	Dem	[5]	Croatia
Magallon-Botaya et al. 2021 ¹²⁸	6286	61 ^{MED.}	56%	Dem	[3], [4], [5]	Spain
Maguire et al. 2020 ¹²⁹	224	n.a.	45%	CI	[5]	UK
Mahmoud et al. 2021 ¹³⁰	126	83 ^{AVG.}	43%	Dem	[5]	Italy
Maniero et al. 2022 ¹³¹	102	83 ^{MED.}	45%	Dem, PD	[5]	UK
Martinot et al. 2021 ¹³²	600	71 ^{MED.}	42%	Dem	[3], [5]	France
Meis-Pinheiro et al. 2021 ¹³³	249	87 ^{AVG.}	73%	Dem	[5]	Spain
Menditto et al. 2021 ¹³⁴	283	57 ^{AVG.}	49%	CI	[4]	Italy
Miyashita et al. 2020a ¹³⁵	1557	70 ^{MED.}	47%	Dem	[3], [4], [5]	USA
Miyashita et al. 2020b ¹³⁵	514	70 ^{MED.}	47%	Dem	[3], [4], [5]	USA
Molani et al. 2022 ¹³⁶	4943	60 ^{AVG.}	44%	Dem	[2]	USA
Moon et al. 2021 ¹³⁷	5626	35 ^{MED.}	58%	Dem	[5]	Korea
Munblit et al. 2021 ¹³⁸	3382	56 ^{MED.}	50%	Dem	[5]	Russia
Nojiri et al. 2022 ¹³⁹	5980	79 ^{MED.}	42%	Dem, PD	[2], [5]	Japan
Oh et al. 2022 ¹⁴⁰	5077	n.a.	59%	Dem	[2], [5]	Korea
Ouattara et al. 2021a ¹⁴¹	25 765	66 ^{MED.}	46%	Dem	[5]	France
Ouattara et al. 2021b ¹⁴¹	72 601	66 ^{MED.}	46%	CI	[5]	France
Pan et al. 2021 ¹⁴⁵	13 874	78 ^{AVG.}	57%	CI	[5]	USA
Panagiotou et al. 2021a ¹⁴²	3202	79 ^{MED.}	61%	Dem	[5]	USA
Panagiotou et al. 2021b ¹⁴²	3570	79 ^{MED.}	61%	Dem	[5]	USA
Panagiotou et al. 2021c ¹⁴²	2486	79 ^{MED.}	61%	Dem	[5]	USA
Patel et al. 2022 ¹⁴³	264	78 ^{AVG.}	59%	Dem, PD, AD, VD	[5]	USA
Pisaturo et al. 2021 ¹⁴⁴	69	63 ^{MED.}	39%	Dem, PD	[2], [5]	Italy
Raheja et al. 2021 ¹⁴⁵	355	84 ^{AVG.}	46%	Dem	[5]	USA
Ramos-Rincón et al. 2021a ¹⁴⁶	2772	86 ^{MED.}	51%	Dem	[5]	Spain
Ramos-Rincon et al. 2021b ¹⁴⁷	6189	86 ^{MED.}	51%	Dem	[5]	Spain
Rebora et al. 2021 ¹⁴⁸	516	78 ^{MED.}	38%	Dem	[5]	Italy
Roig-Marín & Roig-Rico 2021 ¹⁴⁹	300	82 ^{AVG.}	49%	Dem	[5]	Spain
Romagnolo et al. 2021a ¹⁵⁰	344	62 ^{AVG.}	41%	Mix	[2], [3], [4], [5]	Italy
Romagnolo et al. 2021b ¹⁵¹	46	62 ^{AVG.}	40%	Mix	[2]	Italy
Rossi et al. 2020 ¹⁵²	2653	63 ^{AVG.}	50%	Dem	[4], [5]	Italy
Russo et al. 2021a ¹⁵³	19 854	72 ^{AVG.}	45%	AD/Dem, PD	[5]	Italy
Russo et al. 2021b ¹⁵³	11 118	83 ^{AVG.}	53%	PD	[5]	Italy
Rutten et al. 2021 ¹⁵⁴	1294	84 ^{AVG.}	64%	Dem, PD	[5]	Netherlands
Salari et al. 2021c ¹⁵⁵	12 909	77 ^{AVG.}	n.a.	PD	[5]	Iran
Samuels et al. 2021 ¹⁵⁶	1692	57 ^{AVG.}	53%	Dem	[3]	USA
Scherbaum et al. 2021 ¹⁴⁷	30 872	78 ^{MED.}	58%	PD	[5]	Germany
Secnik et al. 2023 (≤70) ¹⁵⁷	359	66 ^{AVG.}	n.a.	Dem	[5]	Sweden
Secnik et al. 2023 (71–80) ¹⁵⁷	1521	75 ^{AVG.}	n.a.	Dem	[5]	Sweden
Secnik et al. 2023 (81–90) ¹⁵⁷	2091	85 ^{AVG.}	n.a.	Dem	[5]	Sweden
Secnik et al. 2023 (>90) ¹⁵⁷	1151	93 ^{AVG.}	n.a.	Dem	[5]	Sweden
Seon et al. 2021 ¹⁴⁹	7713	53 ^{MED.}	61%	Dem	[5]	Korea
Shin et al. 2021 ¹⁵⁸	5771	62 ^{MED.}	59%	Dem	[2], [5]	Korea
Soldevila et al. 2022 ⁵⁰	2225	86 ^{AVG.}	27%	Dem	[5]	Spain
Song et al. 2021 ¹⁵⁹	5621	45 ^{MED.}	59%	Dem	[5]	Korea
Stawinski et al. 2021 ¹⁶⁰	186	67 ^{AVG.}	47%	Dem	[5]	USA
Tahira et al. 2021 ⁵¹	12 863	75 ^{MED.}	49%	AD, PD, Dem	[4], [5]	UK
Tsai et al. 2020 ¹⁶¹	627	47 ^{AVG.}	53%	Dem	[5]	USA
Tyson et al. 2021 ¹⁶²	150	78 ^{AVG.}	50%	Dem, PD	[5]	USA
Vekaria et al. 2022 ¹⁶³	10 473	79 ^{MED.}	50%	Dem	[3], [5]	Finland
Venturini et al. 2021 ¹⁶⁴	175	75 ^{MED.}	44%	Dem	[3], [5]	Italy
Vignatelli et al. 2021 ¹⁶⁵	9470	77 ^{AVG.}	55%	Dem	[4], [5]	Italy
Wan et al. 2020 ¹⁶⁶	30	64 ^{AVG.}	67%	Dem	[5]	China
Wang et al. 2021c ⁵⁴	14 874	65 ^{AVG.}	53%	Dem, AD, VD	[5]	UK
Worcel et al. 2021 ⁵⁶	63	85 ^{AVG.}	61%	AD, Dem	[5]	France
Yakar et al. 2021 ¹⁶⁷	1626	68 ^{AVG.}	48%	AD, PD, VD, FTD, Dem	[5]	UK
Yu et al. 2021 ⁵⁷	249	71 ^{MED.}	31%	PD, Dem	[5]	Turkey
Zakaria et al. 2021 ¹⁶⁸	1064	65 ^{AVG.}	51%	Dem	[3], [4], [5]	USA
Zenesini et al. 2022a ⁵⁸	736	78 ^{AVG.}	33%	Dem, PD	[4]	Italy
Zenesini et al. 2022b ⁵⁸	331	81 ^{AVG.}	41%	Dem, PD	[5]	Italy
Zerbo et al. 2021 ¹⁶⁹	219 001	45 ^{MED.}	51%	AD, PD	[5]	USA
Zhang et al. 2021 ¹⁷⁰	387 841	51 ^{MED.}	55%	Mix	[5]	USA
Zhou et al. 2021b ¹⁷¹	4442	45 ^{MED.}	49%	Mix	[2]	China

a. Studies are divided by outcome: 'infection risk' and 'course and outcome'. The latter includes only participants with positive COVID-19 infection. AVG., average (mean); MED., median; AD, Alzheimer's disease; n.a., not applicable; CI, cognitive impairment; Dem, dementia; DLB, dementia with Lewy bodies; FTD, frontotemporal dementia; MCI, mild cognitive impairment; Mix, mixed dementia; PD, Parkinson's disease; VD, vascular dementia; KSA, Kingdom of Saudi Arabia; N. Ireland, Northern Ireland; 1, infection risk; 2, severity; 3, intensive care unit admission; 4, hospital admission; 5, mortality.

Disorder	OR (95% CI)	Studies, <i>k</i>	Participants, <i>n</i>	<i>I</i> ²	Egger's <i>t</i>
Dementia	1.83 (1.16–2.87)**	7	62 859 255	99.3***	−0.078
Alzheimer's disease ^a	2.86 (1.44–5.66)**	4	61 978 500	98.1***	−0.518
Parkinson's disease	1.65 (1.34–2.04)***	6	278 245	86.8***	0.397
Mild cognitive impairment	1.51 (1.35–1.70)***	1	12 384	n.a.	n.a.
Mixed dementia	2.48 (1.17–5.27)*	3	76 327	93.5***	3.745

n.a., not applicable.
a. Estimates come from analyses including nationwide data, at the expense of local data, hence the number of studies (*k*) is relatively low.
P* < 0.05, *P* < 0.01, ****P* < 0.001.

owing to better financial resources and advanced medical technology, such as laboratories, (self-) diagnostic kits, and public and private funded testing stations in high-income countries, more cases have been detected in high-income countries compared with low-income countries.¹⁷⁷ Similarly, nursing homes, more common in high-income countries, tend to have a larger elderly population than in low-income countries, which may also explain differences among continents.¹⁸

Severity, course and outcome of SARS-CoV-2 and neurodegenerative disorders

Individuals with most types of pre-existing neurodegenerative disorder are disproportionately affected by COVID-19 once infected. These effects were evident over disease types and outcome, suggesting an approximately twofold increase in risk of more severe illness, and a relatively poor course and outcome, for people with pre-existing dementia, and about a fourfold increase in risk of hospital admission in people with Alzheimer's disease. It has conclusively been shown that age is a risk factor for severe COVID-19.¹⁷⁸ However, analyses controlled for age yielded similar findings. A possible explanation for the observed findings may be deleterious interactions between COVID-19 and some specific clinical presentations and comorbidities inherent in neurodegenerative disorders. The atypical manifestation of COVID-19 symptoms in the elderly may lead to a delay in detection and diagnosis of the virus, accelerating the risk of developing severe complications and therefore resulting in a higher risk of hospital admission and ICU

admission.^{179,180} In addition, dementia is associated with oropharyngeal dysphagia,¹⁸¹ a serious comorbidity or complication that independently increases the risk of pneumonia, malnutrition and mortality.¹⁸² Furthermore, Parkinson's disease, dementia and dysphagia are well-known independent and substantial risk factors for pneumonia,¹⁸³ which is a common cause of death in advanced dementia.¹⁸⁴ A notable exception that was observed is that people with pre-existing dementia are less likely to be admitted to an ICU because of COVID-19. We are not aware of any studies showing that widespread vaccination altered this association. This finding might best be explained by the triage criteria (e.g. age, frailty and likelihood of benefit) commonly used in disaster situations to maximise the number of survivors.^{185,186} In countries such as Belgium and the UK, it is advised against admitting to an ICU individuals aged 65 years or older presenting a Clinical Frailty Scale (CFS) score ≥ 5, who have been diagnosed with COVID-19 or are suspected of having contracted the virus.^{90,187} Another significant finding is that individuals with pre-existing neurodegenerative disorders are at higher risk for mortality. This finding is in line with previous studies, demonstrating that in 2020, recorded deaths from Alzheimer's disease and from dementia were respectively 13% and 17% higher than expected, compared with 5 years earlier.^{10,188} It should be noted that we conducted separate meta-analyses focused on unadjusted effect estimates and on adjusted effect estimates, and all associations yielded significant findings when adjusted for age and gender.

The results of these meta-analyses raise the question of whether the increased susceptibility of people with neurodegenerative




Fig. 2 Forest plot of pooled effect estimates for SARS-CoV-2 infection risk across all disorders.

Table 4 Neurodegenerative disorders and COVID-19 severity, hospital admission, intensive care unit admission and mortality from multivariable analyses

Disorder	OR (95% CI)	Studies, <i>k</i>	Participants, <i>n</i>	<i>I</i> ²	Egger's <i>t</i>
<i>COVID-19 severity</i>					
Dementia ^a	1.43 (1.00–2.03)*	6	2 058 163	94.2***	1.504
Alzheimer's disease	0.73 (0.20–2.67)	1	913	n.a.	n.a.
Mild cognitive impairment	0.95 (0.59–1.53)	1	78	n.a.	n.a.
Mixed dementia	1.40 (1.12–1.76)**	4	182 781	18.5	1.189
<i>COVID-19 hospital admission</i>					
Dementia ^a	1.60 (1.09–2.35)*	6	94 624	95.2***	1.002
Alzheimer's disease	3.72 (2.35–5.90)***	2	231 864	58.6	n.a.
Parkinson's disease ^a	1.06 (0.67–1.70)	4	24 382 561	88.4*	–1.069
Mild cognitive impairment	1.64 (0.82–3.28)	2	24 368 547	87.1**	n.a.
Mixed dementia	1.01 (0.99–1.03)	1	670 496	n.a.	n.a.
<i>COVID-19 intensive care unit admission</i>					
Dementia ^a	0.54 (0.40–0.74)***	12	47 015	80.4**	–2.020
Alzheimer's disease	1.65 (1.13–2.42)**	1	219 001	n.a.	n.a.
Parkinson's disease	1.55 (1.01–2.38)*	1	219 001	n.a.	n.a.
<i>COVID-19 mortality</i>					
Dementia ^a	1.58 (1.39–1.79)***	29	756 194	86.9***	3.625***
Alzheimer's disease	1.96 (1.34–2.86)***	7	673 716	93.8***	1.405
Parkinson's disease ^a	1.56 (1.25–1.94)***	9	7 942 537	81.9***	0.133
Mild cognitive impairment ^a	1.68 (1.11–2.54)**	5	10 651	89.4***	–1.117
Mixed dementia	2.27 (1.49–3.46)***	6	432 426	73.8**	3.608*

n.a., not applicable.
 a. Estimates come from analyses including nationwide data, at the expense of local data, hence the number of studies (*k*) is relatively low.
 P* < 0.05, *P* < 0.01, ****P* < 0.001.




Fig. 3 Forest plot of pooled effect estimates for COVID-19 mortality in people with dementia.

diseases to COVID-19 and poorer outcome may be attributable (at least partly) to biological factors and support a research agenda on this topic. We propose two main putative pathophysiological underpinnings that deserve further investigation: (a) a dysfunction of first barrier mucosal defences, leading to a higher infection rate and (b) a deteriorated, slower immune response to SARS-CoV-2, particularly an impaired T-cell immunity, essential to reduce the severity of the infection and facilitate the recovery of infected individuals. The first hypothesis may be approached by the investigation of the possible role in SARS-CoV-2 infection rates of oropharyngeal dysphagia, common in dementia,¹⁸¹ and the effect of reduced salivary lactoferrin levels in Alzheimer's disease, which will reduce the defence mechanisms against SARS-CoV-2 and increase COVID-19 susceptibility.¹⁸⁹ The second one would require an investigation of the clinical significance of some changes in peripheral blood cell profiles involved at least in Alzheimer's disease and related to inflammation and immune dysfunction, such as the CD4/CD8 ratio,¹⁹⁰ which in turn may contribute to severe COVID-19.¹⁹¹ In addition, individuals homozygous for apolipoprotein E (APOE) $\epsilon 4$ have shown a higher risk of COVID-19-related hospital admission, which could be explained by the changes associated with APOE $\epsilon 4$ that lead to extensive central nervous system inflammation, neurodegeneration and aggressive inflammatory response due to increased blood-brain barrier permeability, exacerbated microglia-mediated neuroinflammation and increased cytokine production in response to inflammatory stimuli.¹⁹²

Strength and limitations

This meta-analysis followed MOOSE²⁶ and PRISMA²⁷ guidelines. In addition, a review protocol was pre-registered with the PROSPERO database. In support of the open science movement to promote transparency, expand access and broaden the range of research output, all the extracted data are openly available at the Open Science Framework (OSF). A further key strength of the study is the inclusion of independent data, which is a crucial assumption in meta-analysis.¹⁹³ Given that most of the conducted research on the topic of interest is based on freely accessible electronic data-sets, overlapping data-sets were anticipated. Hence, we carefully followed an inclusion protocol, ensuring that no duplicate data were used in each meta-analysis. To achieve this, we ran analyses for both local and nationwide data, and reported similarities or differences among the analyses. An additional strength of the study is the composition of the data-sets. We reported on several outcomes stratified by type of neurodegenerative disease. Last, our study makes use of both frequentist and Bayesian methods. Using this enhanced methodology in our meta-analysis enabled the collection of more exhaustive and reliable data regarding the association between various types of pre-existing neurodegenerative disease and SARS-CoV-2 susceptibility, course and outcome. In this meta-analysis both the frequentist and the Bayesian method showed comparable results, which suggests consistency and overall robustness of the findings.

Aside from these strengths, several limitations need to be considered. First, most of the meta-analyses revealed high between-study heterogeneity which remained unexplained, and publication bias was found in associations between dementia and COVID-19 severity and mortality. Three of the analyses had a small number of studies (fewer than 10) and were difficult to interpret and thought to be unreliable. Accounting for this by means of trim-and-fill methods did not result in different estimates. For the analyses on the associations between dementia and mortality and between mixed dementia and mortality, the trim-and-fill yielded slightly smaller yet significant effect-size estimates (funnel plots and trim-and-fill analyses can be found in the supplementary

material). The number of studies for certain outcomes and disease types was relatively small, which may have resulted in high levels of between-study heterogeneity.¹⁹⁴ However, this could also be attributed to the scarce reporting on potential sources of heterogeneity (such as diagnostic criteria, time frame of diagnostic assessment, type of analysis used) in the majority of studies. Consequently, inadequate reporting limited the ability to examine the effect of these sources by running subgroup analyses, sensitivity analyses or meta-regression. Moreover, primary studies that specified subgroups of neurodegenerative disorders reported effect estimates for each category using the entire sample, rather than by subsample. We were therefore unable to pool all effect estimates for some disease types. The Cochrane handbook advises that results derived from meta-regression should be interpreted only when there are >10 studies available per analysis.¹⁹⁵ Sometimes we reported results based on fewer studies. Last, we cannot attribute causality to the relationships we reported on as all studies were observational and most retrospective.

Implications

Our findings underline the importance of vaccine priority and health surveillance in people with pre-existing neurodegenerative disease, in the current and possibly a next pandemic.

Muhannad Smadi , Institute of Psychology, Department of Clinical Psychology, Leiden University, Leiden, The Netherlands; **Melina Kaburis** , Institute of Psychology, Department of Clinical Psychology, Leiden University, Leiden, The Netherlands; **Youval Schnapper**, Institute of Psychology, Department of Clinical Psychology, Leiden University, Leiden, The Netherlands; **Gabriel Reina** , Navarra Institute for Health Research (IdiSNA), Pamplona, Spain; and Clinica Universidad de Navarra, Department of Microbiology, Pamplona, Spain; **Patricio Molero**, Navarra Institute for Health Research (IdiSNA), Pamplona, Spain; and Clinica Universidad de Navarra, Department of Psychiatry and Medical Psychology, Pamplona, Spain; **Marc L. Molendijk** , Institute of Psychology, Department of Clinical Psychology, Leiden University, Leiden, The Netherlands; and Leiden Institute for Brain and Cognition, Leiden University Medical Centre, Leiden, The Netherlands

Correspondence: Marc L. Molendijk. Email: m.l.molendijk@fsw.leidenuniv.nl

First received 2 Nov 2022, final revision 22 Mar 2023, accepted 22 Mar 2023

Supplementary material

Supplementary material for this article is available at <https://doi.org/10.1192/bjp.2023.43>.

Data availability

The data that support the findings of this study are openly available on the Open Science Framework (OSF) at https://osf.io/fz5j4/?view_only=95054452816442eaa1e05d5d42be4b2e

Acknowledgements

We thank Soldevila et al (2022), Secnik et al (2023) and Taquet et al (2021), who, on request, responded on our queries and/or provided additional data.

Author contributions

M.S. and M.L.M. had full access to all data in the study and take responsibility for the integrity of the data and the accuracy of data analyses. All authors were responsible for the study concept and design. M.S., Y.S., M.K. and M.L.M. contributed to the collecting and processing of the data. M.S. and M.L.M. ran all statistical analyses. M.S., M.K., G.R., P.M. and M.L.M. drafted the manuscript. All authors interpreted and discussed the findings. All authors critically revised the manuscript. All authors agreed on the final manuscript and the decision to submit it for publication.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Declaration of interest

Without relevance to this work, P.M. reports research grants from the Ministry of Education (Spain), the Government of Navarra (Spain), the Spanish Foundation of Psychiatry and Mental Health, and AstraZeneca; he is a clinical consultant for MedAvante-ProPhase and has received lecture honoraria from and/or has been a consultant for AB-Biotics, Adept Field Solutions, Guidepoint, Janssen, Novumed, Roland Berger and Scienta.

References

- John A, Ali K, Marsh H, Reddy PH. Can healthy lifestyle reduce disease progression of Alzheimer's during a global pandemic of COVID-19. *Ageing Res Rev* 2021; **70**: 101406.
- Liu N, Sun J, Wang X, Zhao M, Huang Q, Li H. The impact of dementia on the clinical outcome of COVID-19: a systematic review and meta-analysis. *J Alzheimers Dis* 2020; **78**: 1775–82.
- El-Sadr WM, Vasan A, El-Mohandes A. Facing the new Covid-19 reality. *N Engl J Med* 2023; **388**: 385–7.
- Chojnicki M, Neumann-Podczaska A, Seostianin M, Tomczak Z, Tariq H, Chudek J, et al. Long-term survival of older patients hospitalized for COVID-19: do clinical characteristics upon admission matter?. *Int J Environ Res Public Health* 2021; **18**: 10671.
- CDC COVID-19 Response Team. Severe outcomes among patients with coronavirus disease 2019 (COVID-19) - United States, February 12-March 16, 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 343–6.
- Bulut C, Kato Y. Epidemiology of COVID-19. *Turk J Med Sci* 2020; **50**: 563–70.
- Bunn F, Burn AM, Goodman C, Rait G, Norton S, Robinson L, et al. Comorbidity and dementia: a scoping review of the literature. *BMC Med* 2014; **12**: 192.
- Holder K, Reddy PH. The COVID-19 effect on the immune system and mitochondrial dynamics in diabetes, obesity, and dementia. *Neuroscientist* 2021; **27**: 331–9.
- Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020; **369**: m1966.
- Alzheimer's Association. 2022 Alzheimer's disease facts and figures. *Alzheimers Dement* 2022; **18**: 700–89.
- Wu KM, Zhang YR, Huang YY, Dong Q, Tan L, Yu JT. The role of the immune system in Alzheimer's disease. *Ageing Res Rev* 2021; **70**: 101409.
- Chung SJ, Chang Y, Jeon J, Shin JI, Song TJ, Kim J. Association of Alzheimer's disease with COVID-19 susceptibility and severe complications: a nationwide cohort study. *J Alzheimers Dis* 2022; **87**: 701–10.
- Suleyman G, Fadel RA, Malette KM, Hammond C, Abdulla H, Entz A, et al. Clinical characteristics and morbidity associated with coronavirus disease 2019 in a series of patients in metropolitan Detroit. *JAMA Netw Open* 2020; **3**: e2012270.
- Lutshumba J, Nikolajczyk BS, Bachstetter AD. Dysregulation of systemic immunity in aging and dementia. *Front Cell Neurosci* 2021; **15**: 652111.
- Shi M, Li C, Tian X, Chu F, Zhu J. Can control infections slow down the progression of Alzheimer's disease? talking about the role of infections in Alzheimer's disease. *Front Aging Neurosci* 2021; **13**: 685863.
- Shi M, Chu F, Tian X, Aeqin Q, Zhu F, Zhu J. Role of adaptive immune and impacts of risk factors on adaptive immune in Alzheimer's disease: are immunotherapies effective or off-target. *Neuroscientist* 2022; **28**: 254–70.
- Bianchetti A, Rozzini R, Bianchetti L, Coccia F, Guerini F, Trabucchi M. Dementia clinical care in relation to COVID-19. *Curr Treat Options Neurol* 2022; **24**: 1–15.
- Azarpazhooh MR, Amir A, Morovatdar N, Steinwender S, Rezaei Ardani A, Yassi N, et al. Correlations between COVID-19 and burden of dementia: an ecological study and review of literature. *J Neurol Sci* 2020; **416**: 117013.
- Shea YF, Wan WH, Chan MMK, DeKosky ST. Time-to-change: dementia care in COVID-19. *Psychogeriatrics* 2020; **20**: 792–3.
- Butler MJ, Barrientos RM. The impact of nutrition on COVID-19 susceptibility and long-term consequences. *Brain Behav Immun* 2020; **87**: 53–4.
- Alves VP, Casemiro FG, Araujo BG, Lima MAS, Oliveira RS, Fernandes FTS, et al. Factors associated with mortality among elderly people in the COVID-19 pandemic (SARS-CoV-2): a systematic review and meta-analysis. *Int J Environ Res Public Health* 2021; **18**: 8008.
- Broad WJ. The publishing game: getting more for less. *Science* 1981; **211**: 1137–9.
- Glasziou PP, Sanders S, Hoffmann T. Waste in covid-19 research. *BMJ* 2020; **369**: m1847.
- Treskova-Schwarzbach M, Haas L, Reda S, Pilic A, Borodova A, Karimi K, et al. Pre-existing health conditions and severe COVID-19 outcomes: an umbrella review approach and meta-analysis of global evidence. *BMC Med* 2021; **19**: 212.
- von Elm E, Poglía G, Walder B, Tramèr MR. Different patterns of duplicate publication: an analysis of articles used in systematic reviews. *JAMA* 2004; **291**: 974–80.
- Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. meta-analysis of observational studies in epidemiology (MOOSE) group. *JAMA* 2000; **283**: 2008–12.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders* (4th edn) (DSM-IV). APA, 1994.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders* (5th edn) (DSM-5). APA, 2013.
- World Health Organization. *ICD-10: International Statistical Classification of Diseases and Related Health Problems*. WHO, 1992.
- Davies HT, Crombie IK, Tavakoli M. When can odds ratios mislead?. *BMJ* 1998; **316**: 989–91.
- Grant RL. Converting an odds ratio to a range of plausible relative risks for better communication of research findings. *BMJ* 2014; **348**: f7450.
- Zhang J, Yu KF. What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. *JAMA* 1998; **280**: 1690–1.
- National Institutes of Health. *Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies*. NIH, 2021 (<https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>).
- Sterne JA, Bradburn MJ, Egger M. *Meta-Analysis in StataTM, Systematic Reviews in Health Care: Meta-Analysis in Context*. John Wiley & Sons, 2001: 347–69.
- Ajayi B, Trompeter A, Arnander M, Sedgwick P, Lui DF. 40 days and 40 nights: clinical characteristics of major trauma and orthopaedic injury comparing the incubation and lockdown phases of COVID-19 infection. *Bone Joint Open* 2020; **1**: 330–8.
- Beobide Telleria I, Ferro Uriguen A, Laso Lucas E, Sannino Menicucci C, Enriquez Barroso M, López de Munain Arregui A. Risk factors associated with COVID-19 infection and mortality in nursing homes. *Aten Primaria* 2022; **54**: 102463.
- Castilla J, Guevara M, Miqueleiz A, Baigorria F, Ibero-Esparza C, Navascues A, et al. Risk factors of infection, hospitalization and death from SARS-CoV-2: a population-based cohort study. *J Clin Med* 2021; **10**(12): 2608.
- de Malherbe A, Verdun S, Brenière V, Luquel L, Jourdan M, Harboun M. COVID-19 prevalence in UNIV group nursing homes and multilevel geriatric hospitals: epidemiological study of immunological status with rapid serological tests for diagnostic guidance and follow up. *J Nutr Health Aging* 2022; **26**: 477–84.
- Del Ser T, Fernández-Blázquez MA, Valentí M, Zea-Sevilla MA, Frades B, Alfayate E, et al. Residence, clinical features, and genetic risk factors associated with symptoms of COVID-19 in a cohort of older people in Madrid. *Gerontology* 2021; **67**: 281–9.
- Emmerson C, Hollinghurst J, North L, Fry R, Akbari A, Humphreys C, et al. The impact of dementia, frailty and care home characteristics on SARS-CoV-2 incidence in a national cohort of Welsh care home residents during a period of high community prevalence. *Age Ageing* 2022; **51**(12): afac250.
- Karapetyan S, Schneider A, Linde K, Donnachie E, Hapfelmeier A. SARS-CoV-2 infection and cardiovascular or pulmonary complications in ambulatory care: a risk assessment based on routine data. *PLoS ONE* 2021; **16**: e0258914.
- Kim J, Park SH, Kim JM. Effect of comorbidities on the infection rate and severity of COVID-19: nationwide cohort study with propensity score matching. *JMIR Public Health Surveill* 2022; **8**: e35025.
- Orlando V, Rea F, Savare L, Guarino I, Mucherino S, Perrella A, et al. Development and validation of a clinical risk score to predict the risk of SARS-CoV-2 infection from administrative data: a population-based cohort study from Italy. *PLoS One* 2021; **16**(1): e0237202.
- Pan AP, Meeks J, Potter T, Masdeu JC, Seshadri S, Smith ML, et al. SARS-CoV-2 susceptibility and COVID-19 mortality among older adults with cognitive impairment: cross-sectional analysis from hospital records in a diverse US metropolitan area. *Front Neurol* 2021; **12**: 692662.
- Profili F, Ballo P, Balzi D, Bellini B, Bartalocci S, Zuppiroli A, et al. Chronic diseases and risk of symptomatic COVID-19: results of a case-population study on a sample of patients in the local health unit 'Toscana Centro' (Tuscan Region, Central Italy). *Epidemiol Prev* 2020; **44**: 308–14.

- 47 Scherbaum R, Kwon EH, Richter D, Bartig D, Gold R, Krogias C, et al. Clinical profiles and mortality of COVID-19 in patients with Parkinson's disease in Germany. *Mov Disord* 2021; **36**: 1049–57.
- 48 Smith J, Aboumradi M, Reyes C, Satram S, Young-Xu Y. Predictors of incident severe acute respiratory syndrome coronavirus 2 positivity in a veteran population. *Mil Med* [Epub ahead of print] 20 Oct 2021. Available from: <https://doi.org/10.1093/milmed/usab428>.
- 49 Seon JY, Kim S, Hong M, Lim MK, Oh IH. Risk of COVID-19 diagnosis and death in patients with mental illness: a cohort study. *Epidemiol Psychiatr Sci* 2021; **30**: e68.
- 50 Soldevila L, Prat N, Mas MA, Massot M, Miralles R, Bonet-Simó JM, et al. The interplay between infection risk factors of SARS-CoV-2 and mortality: a cross-sectional study from a cohort of long-term care nursing home residents. *BMC Geriatr* 2022; **22**: 123.
- 51 Tahira AC, Verjovski-Almeida S, Ferreira ST. Dementia is an age-independent risk factor for severity and death in COVID-19 inpatients. *Alzheimers Dement* 2021; **11**: 1818–31.
- 52 Wang Q, Davis PB, Gurney ME, Xu R. COVID-19 and dementia: analyses of risk, disparity, and outcomes from electronic health records in the US. *Alzheimers Dement* 2021; **17**: 1297–306.
- 53 Wang SM, Park SH, Kim NY, Kang DW, Na HR, Um YH, et al. Association between dementia and clinical outcome after COVID-19: a nationwide cohort study with propensity score matched control in South Korea. *Psychiatry Investig* 2021; **18**: 523–9.
- 54 Wang Y, Yang Y, Ren L, Shao Y, Tao W, Dai XJ. Preexisting mental disorders increase the risk of COVID-19 infection and associated mortality. *Front Public Health* 2021; **9**: 684112.
- 55 Wong R, Lovier MA. Relationship between dementia, COVID-19 risk, and adherence to COVID-19 mitigation behaviors among older adults in the United States. *Int J Geriatr Psychiatry* 2022; **37**(6): doi:10.1002/gps.5735.
- 56 Worcel A, Ali BM, Ramos-Pascual S, Stirling P, Chary FG. Low mortality from COVID-19 at a nursing facility in France following a combined preventive and active treatment protocol. *Ann Palliat Med* 2021; **10**: 11288–300.
- 57 Yu YZ, Travaglio M, Popovic R, Leal NS, Martins LM. Alzheimer's and Parkinson's diseases predict different COVID-19 outcomes: a UK Biobank study. *Geriatrics* 2021; **6**(1): 10.
- 58 Zenesini C, Vignatelli L, Belotti LMB, Baccari F, Calandra-Buonaura G, Cortelli P, et al. Risk of SARS-CoV-2 infection, hospitalization and death for COVID-19 in people with Parkinson's disease or parkinsonism over a 15-month period: a cohort study. *Eur J Neurol* [Epub ahead of print] 16 Jul 2022. Available from: <https://doi.org/10.1111/ene.15505>.
- 59 Zhou J, Liu C, Sun Y, Huang W, Ye K. Cognitive disorders associated with hospitalization of COVID-19: results from an observational cohort study. *Brain Behav Immun Health* 2021; **9**: 383–92.
- 60 Ajayi B, Trompeter AJ, Umarji S, Saha P, Arnander M, Lui DF. Catching the second wave: clinical characteristics and nosocomial infection rates in major trauma and orthopaedic patients during the COVID-19 pandemic. *Bone Jt Open* 2021; **2**: 661–70.
- 61 Alqahtani AM, Alshahrani FM, Khalaf MME, Rezk SR. Outcome of COVID-19 among homecare patients and its relation to chronic diseases. *World Family Med* 2021; **19**: 111–7.
- 62 An C, Oh HC, Chang JH, Oh SJ, Lee JM, Han CH, et al. Development and validation of a prognostic model for early triage of patients diagnosed with COVID-19. *Sci Rep* 2021; **11**: 21923.
- 63 Atkins JL, Masoli JAH, Delgado J, Pilling LC, Kuo CL, Kuchel GA, et al. Preexisting comorbidities predicting COVID-19 and mortality in the UK Biobank community cohort. *J Gerontol A Biol Sci Med Sci* 2020; **75**: 2224–30.
- 64 Bae S, Kim Y, Hwang S, Kwon KT, Chang HH, Kim SW. New scoring system for predicting mortality in patients with COVID-19. *Yonsei Med J* 2021; **62**: 806–13.
- 65 Baker SM, Leedy DJ, Klafner JA, Zhang Y, Secrest KM, Osborn TR, et al. Clinical presentation, complications, and outcomes of hospitalized COVID-19 patients in an academic center with a centralized palliative care consult service. *Health Sci Rep* 2021; **4**: e423.
- 66 Banoei MM, Dinparastisaleh R, Zadeh AV, Mirsaedi M. Machine-learning-based COVID-19 mortality prediction model and identification of patients at low and high risk of dying. *Crit Care* 2021; **25**: 328.
- 67 Becerra-Muñoz VM, Núñez-Gil IJ, Eid CM, García Aguado M, Romero R, Huang J, et al. Clinical profile and predictors of in-hospital mortality among older patients hospitalised for COVID-19. *Age Ageing* 2021; **50**: 326–34.
- 68 Bennett TD, Moffitt RA, Hajagos JG, Amor B, Anand A, Bissell MM, et al. Clinical characterization and prediction of clinical severity of SARS-CoV-2 infection among US adults using data from the US national COVID cohort collaborative. *JAMA Netw Open* 2021; **4**(7): e2116901.
- 69 Bhargava A, Szpunar SM, Sharma M, Fukushima EA, Hoshi S, Levine M, et al. Clinical features and risk factors for in-hospital mortality from COVID-19 infection at a tertiary care medical center, at the onset of the US COVID-19 pandemic. *J Intensive Care Med* 2021; **36**: 711–8.
- 70 Bianchetti A, Rozzini R, Guerini F, Boffelli S, Ranieri P, Minelli G, et al. Clinical presentation of COVID-19 in dementia patients. *J Nutr Health Aging* 2020; **24**: 560–2.
- 71 Bielza R, Sanz J, Zambrana F, Arias E, Malmierca E, Portillo L, et al. Clinical characteristics, frailty, and mortality of residents with COVID-19 in nursing homes of a region of Madrid. *J Am Med Dir Assoc* 2021; **22**: 245–252.e2.
- 72 Booij JA, van de Haterd J, Huttjes SN, van Deijck R, Koopmans R. Short- and long-term mortality and mortality risk factors among nursing home patients after COVID-19 infection. *J Am Med Dir Assoc* 2022; **23**: 1274–8.
- 73 Boye KS, Tokar Erdemir E, Zimmerman N, Reddy A, Benneyworth BD, Dabora MC, et al. Risk factors associated with COVID-19 hospitalization and mortality: a large claims-based analysis among people with type 2 diabetes mellitus in the United States. *Diabetes Ther* 2021; **12**: 2223–39.
- 74 Bucholtz M, Bradley D, Bennett D, Patterson L, Spiers R, Gibson D, et al. Identifying pre-existing conditions and multimorbidity patterns associated with in-hospital mortality in patients with COVID-19. *Sci Rep* 2022; **12**: 17313.
- 75 Busetto L, Bettini S, Fabris R, Serra R, Dal Pra C, Maffei P, et al. Obesity and COVID-19: an Italian snapshot. *Obesity* 2020; **28**: 1600–5.
- 76 Caliskan T, Saylan B. Smoking and comorbidities are associated with COVID-19 severity and mortality in 565 patients treated in Turkey: a retrospective observational study. *Rev Assoc Med Bras* 2020; **66**: 1679–84.
- 77 Carrillo-García P, Garmendia-Prieto B, Cristofori G, Montoya IL, Hidalgo JJ, Feijoo MQ, et al. Health status in survivors older than 70 years after hospitalization with COVID-19: observational follow-up study at 3 months. *Eur Geriatr Med* 2021; **12**: 1091–4.
- 78 Chang MH, Moonesinghe R, Truman BI. COVID-19 hospitalization by race and ethnicity: association with chronic conditions among Medicare beneficiaries, January 1–September 30, 2020. *J Racial Ethnic Health Disparities* 2022; **9**(1): 325–34.
- 79 Chatterjee A, Wu GY, Primakov S, Oberije C, Woodruff H, Kubben P, et al. Can predicting COVID-19 mortality in a European cohort using only demographic and comorbidity data surpass age-based prediction: an externally validated study. *PLoS One* 2021; **16**(4): e0249920.
- 80 Chen UI, Xu H, Krause TM, Greenberg R, Dong X, Jiang X. Factors associated with COVID-19 death in the United States: cohort study. *JMIR Public Health Surveill* 2022; **8**: e29343.
- 81 Choi YJ, Park JY, Lee HS, Suh J, Song JY, Byun MK, et al. Variable effects of underlying diseases on the prognosis of patients with COVID-19. *PLoS One* 2021; **16**(7): e0254258.
- 82 Chojnicki M, Neumann-Podczaska A, Seostianin M, Tomczak Z, Tariq H, Chudek J, et al. Long-term survival of older patients hospitalized for COVID-19: Do clinical characteristics upon admission matter?. *Int J Environ Res Public Health* 2021; **18**(20): 10671.
- 83 Cisterna-García A, Guillén-Teruel A, Caracena M, Pérez E, Jiménez F, Francisco-Verdú FJ, et al. A predictive model for hospitalization and survival to COVID-19 in a retrospective population-based study. *Sci Rep* 2022; **12**: 18126.
- 84 COVIDSurg Collaborative. Outcomes after perioperative SARS-CoV-2 infection in patients with proximal femoral fractures: an international cohort study. *BMJ Open* 2021; **11**: e050830.
- 85 Covino M, De Matteis G, Santoro M, Sabia L, Simeoni B, Candelli M, et al. Clinical characteristics and prognostic factors in COVID-19 patients aged ≥ 80 years. *Geriatr Gerontol Int* 2020; **20**: 704–8.
- 86 Covino M, De Matteis G, Polla DAD, Santoro M, Burzo ML, Torelli E, et al. Predictors of in-hospital mortality and death risk stratification among COVID-19 patients aged ≥ 80 years old. *Arch Gerontol Geriatr* 2021; **95**: 104383.
- 87 Covino M, Russo A, Salini S, De Matteis G, Simeoni B, Della Polla D, et al. Frailty assessment in the emergency department for risk stratification of COVID-19 patients aged ≥ 80 years. *J Am Med Dir Assoc* 2021; **22**: 1845–1852.e1.
- 88 Cummins L, Ebyarimpa I, Cheetham N, Tzortziou Brown V, Brennan K, Panovska-Griffiths J. Factors associated with COVID-19 related hospitalisation, critical care admission and mortality using linked primary and secondary care data. *Influenza Other Respi Viruses* 2021; **15**: 577–88.
- 89 de Marcaida JA, Lahrmann J, Machado D, Bluth L, Dagostine M, Moro-de Casillas M, et al. Clinical characteristics of coronavirus disease 2019 (COVID-19) among patients at a movement disorders center. *Geriatrics* 2020; **5**(3): 54.
- 90 De Smet R, Mellaerts B, Vandewinckele H, Lybeert P, Frans E, Ombelet S, et al. Frailty and mortality in hospitalized older adults with COVID-19: retrospective observational study. *J Am Med Dir Assoc* 2020; **21**: 928–32.e1.
- 91 Descamps A, Frenkiel J, Zarca K, Laidi C, Godin O, Launay O, et al. Association between mental disorders and COVID-19 outcomes among inpatients in France: a retrospective nationwide population-based study. *J Psychiatry Res* 2022; **155**: 194–201.

- 92 Ellis RJ, Moffatt CR, Aaron LT, Beaverson G, Chaw K, Curtis C, et al. Factors associated with hospitalisations and deaths of residential aged care residents with COVID-19 during the Omicron (BA.1) wave in Queensland. *Med J Aust* 2023; **218**: 174–9.
- 93 Escribà-Salvans A, Rierola-Fochs S, Farrés-Godayol P, Molas-Tuneu M, Bezerra de Souza DL, Skelton DA, et al. Risk factors for developing symptomatic COVID-19 in older residents of nursing homes: A hypothesis-generating observational study. *MedRxiv* (Preprint) 2022. Available from: <https://doi.org/10.1101/2022.01.18.22269433>.
- 94 Esme M, Koca M, Dikmeer A, Balci C, Ata N, Dogu BB, et al. Older adults with coronavirus disease 2019: a nationwide study in Turkey. *J Gerontol A Biol Sci Med Sci* 2021; **76**: e68–75.
- 95 España PP, Bilbao A, García-Gutiérrez S, Lafuente I, Anton-Ladislao A, Villanueva A, et al. Predictors of mortality of COVID-19 in the general population and nursing homes. *Intern Emerg Med* 2021; **16**: 1487–96.
- 96 Esteban I, Bergero G, Alves C, Bronstein M, Ziegler V, Wood C, et al. Asymptomatic COVID-19 in the elderly: dementia and viral clearance as risk factors for disease progression. *Am J Respir Crit Care Med* 2021; **203**: A3826.
- 97 Fasano A, Cereda E, Barichella M, Cassani E, Ferri V, Zecchinelli AL, et al. COVID-19 in Parkinson's disease patients living in Lombardy, Italy. *Mov Disord* 2020; **35**: 1089–93.
- 98 Fathi M, Taghizadeh F, Mojtahedi H, Zargar Balaye Jame S, Markazi Moghaddam N. The effects of Alzheimer's and Parkinson's disease on 28-day mortality of COVID-19. *Rev Neurol (Paris)* 2022; **178**: 129–36.
- 99 Filardo TD, Khan MR, Krawczyk N, Galitzer H, Karmen-Tuohy S, Coffee M, et al. Comorbidity and clinical factors associated with COVID-19 critical illness and mortality at a large public hospital in New York City in the early phase of the pandemic (March–April 2020). *PLoS One* 2020; **15**(11): e0242760.
- 100 Filipe MS, Maroussia R, Brian F, Amaury T, Anne I, Alexia C, et al. Risk factors for severe outcomes for COVID-19 patients hospitalised in Switzerland during the first pandemic wave, February to August 2020: prospective observational cohort study. *Swiss Med Wkly* 2021; **151**: w20547.
- 101 Fumagalli C, Ungar A, Rozzini R, Vannini M, Coccia F, Cesaroni G, et al. Predicting mortality risk in older hospitalized persons with COVID-19: a comparison of the COVID-19 mortality risk score with frailty and disability. *J Am Med Dir Assoc* 2021; **22**: 1588–92.e1.
- 102 Gale TM, Boland B. COVID-19 deaths in a secondary mental health service. *Compr Psychiatry* 2021; **111**: 152277.
- 103 Ge E, Li Y, Wu S, Candido E, Wei X. Association of pre-existing comorbidities with mortality and disease severity among 167,500 individuals with COVID-19 in Canada: a population-based cohort study. *PLoS One* 2021; **16**: e0258154.
- 104 Genet B, Vidal JS, Cohen A, Bouly C, Beunardeau M, Marine Harlé L, et al. COVID-19 in-hospital mortality and use of renin-angiotensin system blockers in geriatrics patients. *J Am Med Dir Assoc* 2020; **21**: 1539–45.
- 105 Geriatric Medicine Research Collaborative, COVID Collaborative, Welch C. Age and frailty are independently associated with increased COVID-19 mortality and increased care needs in survivors: results of an international multi-centre study. *Age Aging* 2021; **50**: 617–30.
- 106 Ghaffari M, Ansari H, Beladimoghdam N, Aghamiri SH, Haghghi M, Nabavi M, et al. Neurological features and outcome in COVID-19: dementia can predict severe disease. *J Neurovirol* 2021; **27**: 86–93.
- 107 Gómez Antúnez M, Muiño Míguez A, Bendala Estrada AD, Maestro de la Calle G, Monge Monge D, Boixeda R, et al. Clinical characteristics and prognosis of COPD patients hospitalized with SARS-CoV-2. *Int J Chron Obstruct Pulmon Dis* 2020; **15**: 3433–45.
- 108 Harrison SL, Fazio-Eynullayeva E, Lane DA, Underhill P, Lip GYH. Comorbidities associated with mortality in 31,461 adults with COVID-19 in the United States: a federated electronic medical record analysis. *PLoS Med* 2020; **17**(9): e1003321.
- 109 Hasani Azad M, Khorrami F, Kazemi Jahromi M, Alishan Karami N, Shahi M, Davari Dolatabadi N, et al. clinical and epidemiological characteristics of hospitalized COVID-19 patients in Hormozgan, Iran: a retrospective, multicenter study. *Arch Iran Med* 2021; **24**: 434–44.
- 110 Hatamabadi H, Sabaghian T, Sadeghi A, Heidari K, Safavi-Naini SAA, Looha MA, et al. Epidemiology of COVID-19 in Tehran, Iran: a cohort study of clinical profile, risk factors, and outcomes. *Biomed Res Int* 2022; **2022**: 2350063.
- 111 Hippisley-Cox J, Coupland CA, Mehta N, Keogh RH, Diaz-Ordaz K, Khunti K, et al. Risk prediction of COVID-19 related death and hospital admission in adults after COVID-19 vaccination: national prospective cohort study. *BMJ* 2021; **374**: n2244.
- 112 Hwang JM, Kim JH, Park JS, Chang MC, Park D. Neurological diseases as mortality predictive factors for patients with COVID-19: a retrospective cohort study. *Neuro Sci* 2020; **41**: 2317–24.
- 113 Izurieta HS, Graham DJ, Jiao Y, Hu M, Lu Y, Wu Y, et al. Natural history of coronavirus disease 2019: risk factors for hospitalizations and deaths among >26 million US Medicare beneficiaries. *J Infect Dis* 2021; **223**: 945–56.
- 114 Kang IS, Kong KA. Body mass index and severity/fatality from coronavirus disease 2019: a nationwide epidemiological study in Korea. *PLoS ONE* 2021; **16**(6): e0253640.
- 115 Ken-Dror G, Wade C, Sharma S, Law J, Russo C, Sharma A, et al. COVID-19 outcomes in UK centre within highest health and wealth band: a prospective cohort study. *BMJ Open* 2020; **10**: e042090.
- 116 Kim SR, Nam SH, Kim YR. Risk factors on the progression to clinical outcomes of COVID-19 patients in South Korea: using national data. *Int J Environ Res Public Health* 2020; **17**(23): 8847.
- 117 Kim SW, Kim SM, Kim YK, Kim JY, Lee YM, Kim BO, et al. Clinical characteristics and outcomes of COVID-19 cohort patients in Daegu Metropolitan City outbreak in 2020. *J Korean Med Sci* 2021; **36**: e12.
- 118 Kim J, Blaum C, Ferris R, Arcila-Mesa M, Do H, Pulgarin C, et al. Factors associated with hospital admission and severe outcomes for older patients with COVID-19. *J Am Geriatr Soc* 2022; **70**: 1906–17.
- 119 Kong KA, Jung S, Yu M, Park J, Kang IS. Association between cardiovascular risk factors and the severity of coronavirus disease 2019: nationwide epidemiological study in Korea. *Front Cardiovasc Med* 2021; **8**: 732518.
- 120 Kostev K, Gessler N, Wohlmuth P, Arnold D, Bein B, Bohlken J, et al. Is dementia associated with COVID-19 mortality? A multicenter retrospective cohort study conducted in 50 hospitals in Germany. *J Alzheimers Dis* 2023; **91**: 719–26.
- 121 Kyoung DS, Lee J, Nam H, Park MH. Dementia and COVID-19 mortality in South Korea. *Dement Neurocogn Disord* 2021; **20**: 38–40.
- 122 Lazcano U, Cuadrado-Godia E, Grau M, Subirana I, Martínez-Carbonell E, Boher-Massaguer M, et al. Increased COVID-19 mortality in people with previous cerebrovascular disease: a population-based cohort study. *Stroke* 2022; **53**: 1276–84.
- 123 Li JW, Long X, Huang HQ, Tang JN, Zhu CL, Hu SP, et al. Resilience of Alzheimer's disease to COVID-19. *J Alzheimers Dis* 2020; **77**: 67–73.
- 124 Livingston G, Rostampour H, Gallagher P, Kalafatis C, Shastri A, Huzzey L, et al. Prevalence, management, and outcomes of SARS-CoV-2 infections in older people and those with dementia in mental health wards in London, UK: a retrospective observational study. *Lancet Psychiatry* 2020; **7**: 1054–63.
- 125 Lozano-Montoya I, Quezada-Feijoo M, Jaramillo-Hidalgo J, Garmendia-Prieto B, Lisette-Carrillo P, Gómez-Pavón FJ. Mortality risk factors in a Spanish cohort of oldest-old patients hospitalized with COVID-19 in an acute geriatric unit: the OCTA-COVID study. *Eur Geriatr Med* 2021; **12**: 1169–80.
- 126 Lu Y, Jiao Y, Graham DJ, Wu Y, Wang J, Menis M, et al. Risk factors for COVID-19 deaths among elderly nursing home Medicare beneficiaries in the pre-vaccine period. *J Infect Dis* 2022; **225**: 567–77.
- 127 Lucijanić M, Piskač Živković N, Zelenika M, Barišić-Jaman M, Jurin I, Matijaca A, et al. Survival after hospital discharge in patients hospitalized for acute coronavirus disease 2019: data on 2586 patients from a tertiary center registry. *Croat Med J* 2022; **63**: 335–42.
- 128 Magallon-Botaya R, Olivan-Blazquez B, Ramirez-Cervantes KL, Mendez-Lopez-de-la-Manzanara F, Aguilar-Palacio I, Casajua-Closas M, et al. Geographic factors associated with poorer outcomes in patients diagnosed with COVID-19 in primary health care. *Int J Environ Res Public Health* 2021; **18**(7): 3842.
- 129 Maguire D, Woods M, Richards C, Dolan R, Veitch JW, Sim WMJ, et al. Prognostic factors in patients admitted to an urban teaching hospital with COVID-19 infection. *J Transl Med* 2020; **18**(1): 354.
- 130 Mahmoud M, Carmisciano L, Tagliafico L, Muzyka M, Rosa G, Signori A, et al. Patterns of comorbidity and in-hospital mortality in older patients with COVID-19 infection. *Front Med (Lausanne)* 2021; **8**: 726837.
- 131 Maniero C, Patel D, Pavithran A, Naran P, Ng FL, Prowle J, et al. A retrospective cohort study of risk factors and outcomes in older patients admitted to an inner-city geriatric unit in London during first peak of COVID-19 pandemic. *Ir J Med Sci* 2022; **191**: 1037–45.
- 132 Martinot M, Eyrie M, Gravier S, Bonijoly T, Kayser D, Ion C, et al. Predictors of mortality, ICU hospitalization, and extrapulmonary complications in COVID-19 patients. *Infect Dis Now* 2021; **51**: 518–25.
- 133 Meis-Pinheiro U, Lopez-Segui F, Walsh S, Ussi A, Santauegenia S, Garcia-Navarro JA, et al. Clinical characteristics of COVID-19 in older adults: a retrospective study in long-term nursing homes in Catalonia. *PLoS One* 2021; **16**(7): e0255141.
- 134 Menditto VG, Fulgenzi F, Bonifazi M, Gnudi U, Gennarini S, Mei F, et al. Predictors of readmission requiring hospitalization after discharge from emergency departments in patients with COVID-19. *Am J Emerg Med* 2021; **46**: 146–9.
- 135 Miyashita S, Yamada T, Mikami T, Miyashita H, Chopra N, Rizk D. Impact of dementia on clinical outcomes in elderly patients with coronavirus 2019 (COVID-19): an experience in New York. *Geriatrics & Gerontology International* 2020; **20**: 732–4.
- 136 Molani S, Hernandez PV, Roper RT, Duvvuri VR, Baumgartner AM, Goldman JD, et al. Risk factors for severe COVID-19 differ by age for hospitalized adults. *Sci Rep* 2022; **12**: 6568.

- 137 Moon HJ, Kim K, Kang EK, Yang HJ, Lee E. Prediction of COVID-19-related mortality and 30-day and 60-day survival probabilities using a nomogram. *J Korean Med Sci* 2021; **36**: e248.
- 138 Munblit D, Nekliudov NA, Bugaeva P, Blyuss O, Kislova M, Listovskaya E, et al. Stop COVID Cohort: an observational study of 3480 patients admitted to the Sechenov University Hospital network in Moscow city for suspected coronavirus disease 2019 (COVID-19) infection. *Clin Infect Dis* 2021; **73**: 1–11.
- 139 Nojiri S, Irie Y, Kanamori R, Naito T, Nishizaki Y. Mortality prediction of COVID-19 in hospitalized patients using the 2020 diagnosis procedure combination administrative database of Japan. *Intern Med* 2023; **62**: 201–13.
- 140 Oh H, Kim R, Chung W. Sex-specific association between underlying diseases and the severity and mortality due to COVID-19 infection: a retrospective observational cohort analysis of clinical epidemiological information collected by the Korea Disease Control and Prevention Agency. *Healthcare (Basel)* 2022; **10**(10): 1846.
- 141 Ouattara E, Bruandet A, Borde A, Lenne X, Binder-Foucard F, Le-Bourhis-Zaimi M, et al. Risk factors of mortality among patients hospitalised with COVID-19 in a critical care or hospital care unit: analysis of the French national medico-administrative database. *BMJ Open Respir Res* 2021; **8**(1): e001002.
- 142 Panagiotou OA, Kosar CM, White EM, Bantis LE, Yang XF, Santostefano CM, et al. Risk factors associated with all-cause 30-day mortality in nursing home residents with COVID-19. *JAMA Int Med* 2021; **181**: 439–48.
- 143 Patel RA, Stebbins GT, Kishen EB, Barton B. COVID-19 outcomes in hospitalized patients with neurodegenerative disease: a retrospective cohort study. *Neurol Clin Pract* 2022; **12**: 43–51.
- 144 Pisaturo M, Calò F, Russo A, Camaioni C, Giaccone A, Pinchera B, et al. Dementia as risk factor for severe coronavirus disease 2019: a case-control study. *Front Aging Neurosci* 2021; **13**: 698184.
- 145 Raheja H, Chukwuika N, Agarwal C, Sharma D, Munoz-Martinez A, Fogel J, et al. Should COVID-19 patients >75 years be ventilated? An outcome study. *QJM* 2021; **114**: 182–9.
- 146 Ramos-Rincón JM, Bernabeu-Whittel M, Fiteni-Mera I, López-Sampalo A, López-Ríos C, García-Andreu MDM, et al. Clinical features and risk factors for mortality among long-term care facility residents hospitalized due to COVID-19 in Spain. *J Gerontol A Biol Sci Med Sci* 2022; **77**: e138–47.
- 147 Ramos-Rincon JM, Buonaiuto V, Ricci M, Martín-Carmona J, Paredes-Ruiz D, Calderón-Moreno M, et al. Clinical characteristics and risk factors for mortality in very old patients hospitalized with COVID-19 in Spain. *J Gerontol A Biol Sci Med Sci* 2021; **76**: e28–37.
- 148 Reborá P, Rozzini R, Bianchetti A, Blangiardo P, Marchegiani A, Piazzoli A, et al. Delirium in patients with SARS-CoV-2 infection: a multicenter study. *J Am Geriatr Soc* 2021; **69**: 293–9.
- 149 Roig-Marín N, Roig-Rico P. Elderly people with Dementia admitted for COVID-19: how different are they?. *Exp Aging Res* 2022; **48**: 177–90.
- 150 Romagnolo A, Balestrino R, Imbalzano G, Ciccone G, Riccardini F, Artusi CA, et al. Neurological comorbidity and severity of COVID-19. *J Neurol* 2021; **268**: 762–9.
- 151 Romagnolo A, Imbalzano G, Artusi CA, Balestrino R, Ledda C, De Rosa FG, et al. Neurological comorbidities and COVID-19-related case fatality: a cohort study. *J Neurol Sci* 2021; **428**: 117610.
- 152 Rossi PG, Marino M, Formisano D, Venturelli F, Vicentini M, Grilli R, et al. Characteristics and outcomes of a cohort of COVID-19 patients in the province of Reggio Emilia, Italy. *PLoS One* 2020; **15**(8): e0238281.
- 153 Russo AG, Decarli A, Valsecchi MG. Strategy to identify priority groups for COVID-19 vaccination: a population based cohort study. *Vaccine* 2021; **39**: 2517–25.
- 154 Rutten JJS, van Kooten J, van Loon AM, van Buul LW, Joling KJ, Smalbrugge M, et al. Dementia and Parkinson's disease: risk factors for 30-day mortality in nursing home residents with COVID-19. *J Alzheimers Dis* 2021; **84**: 1173–81.
- 155 Salari M, Etemadifar M, Ashrafi F, Ommi D, Aminzade Z, Tehrani Fateh S. Parkinson's disease patients may have higher rates of Covid-19 mortality in Iran. *Parkinsonism Relat Disord* 2021; **89**: 90–2.
- 156 Samuels S, Niu J, Sareli C, Eckardt P. The epidemiology and predictors of outcomes among confirmed COVID-19 cases in a large community healthcare system in South Florida. *J Community Health* 2021; **46**: 822–31.
- 157 Secnik J, Eriksdotter M, Xu H, Annetorp M, Rytarowski A, Johnell K, et al. Dementia and psychotropic medications are associated with significantly higher mortality in geriatric patients hospitalized with COVID-19: data from the StockholmGeroCovid project. *Alzheimers Res Ther* 2023; **15**: 5.
- 158 Shin EK, Choi HY, Hayes N. The anatomy of COVID-19 comorbidity networks among hospitalized Korean patients. *Epidemiol Health* 2021; **43**: e2021035.
- 159 Song J, Park DW, Cha JH, Seok H, Kim JY, Park J, et al. Clinical course and risk factors of fatal adverse outcomes in COVID-19 patients in Korea: a nationwide retrospective cohort study. *Sci Rep* 2021; **11**(1): 10066.
- 160 Stawinski PM, Dziadkowiec KN, Al-Abbasi B, Suarez L, Simms L, Dewaswala N, et al. Model of end-stage liver disease (MELD) score as a predictor of in-hospital mortality in patients with COVID-19: a novel approach to a classic scoring system. *Cureus* 2021; **13**: e15179.
- 161 Tsai A, Diawara O, Nahass RG, Brunetti L. Impact of tocilizumab administration on mortality in severe COVID-19. *Sci Rep* 2020; **10**(1): 19131.
- 162 Tyson B, Erdodi L, Shahein A, Kamrun S, Eckles M, Agarwal P. Predictors of survival in older adults hospitalized with COVID-19. *Neurol Sci* 2021; **42**: 3953–8.
- 163 Vekaria PH, Syed A, Anderson J, Cornett B, Bourbia A, Flynn MG, et al. Association of dementia and patient outcomes among COVID-19 patients: a multi-center retrospective case-control study. *Front Med (Lausanne)* 2022; **9**: 1050747.
- 164 Venturini S, Orso D, Cugini F, Crapis M, Fossati S, Callegari A, et al. Classification and analysis of outcome predictors in non-critically ill COVID-19 patients. *Intern Med J* 2021; **51**: 506–14.
- 165 Vignatelli L, Zenesini C, Belotti LMB, Baldin E, Bonavina G, Calandra-Buonaura G, et al. Risk of hospitalization and death for COVID-19 in people with Parkinson's disease or Parkinsonism. *Mov Disord* 2021; **36**: 1–10.
- 166 Wan Y, Wu J, Ni LH, Luo QQ, Yuan C, Fan F, et al. Prognosis analysis of patients with mental disorders with COVID-19: a single-center retrospective study. *Aging-Us* 2020; **12**: 11238–44.
- 167 Yakar MN, Ergan B, Ergün B, Küçük M, Cantürk A, Ergon MC, et al. Clinical characteristics and risk factors for 28-day mortality in critically ill patients with COVID-19: a retrospective cohort study. *Turk J Med Sci* 2021; **51**: 2285–95.
- 168 Zakaria A, Piper M, Douda L, Jackson NM, Flynn JC, Misra DP, et al. Determinants of all-cause in-hospital mortality among patients who presented with COVID-19 to a community teaching hospital in Michigan. *Heliyon* 2021; **7**(12): e08566.
- 169 Zerbo O, Lewis N, Fireman B, Goddard K, Skarbinski J, Sejvar JJ, et al. Population-based assessment of risks for severe COVID-19 disease outcomes. *Influenza Other Respir Viruses* 2022; **16**: 159–65.
- 170 Zhang Q, Schultz JL, Aldridge GM, Simmering JE, Kim Y, Ogilvie AC, et al. COVID-19 case fatality and Alzheimer's disease. *J Alzheimers Dis* 2021; **84**: 1447–52.
- 171 Zhou JD, Lee S, Wang XS, Li Y, Wu WKK, Liu T, et al. Development of a multi-variable prediction model for severe COVID-19 disease: a population-based study from Hong Kong. *NPJ Digital Medicine* 2021; **4**(1): 66.
- 172 Bellou V, Tzoulaki I, van Smeden M, Moons KGM, Evangelou E, Belbasis L. Prognostic factors for adverse outcomes in patients with COVID-19: a field-wide systematic review and meta-analysis. *Eur Respir J* 2022; **59**: 2002964.
- 173 Wang M, Baker JS, Quan W, Shen S, Fekete G, Gu Y. A preventive role of exercise across the coronavirus 2 (SARS-CoV-2) Pandemic. *Front Physiol* 2020; **11**: 572718.
- 174 Nikolich-Zugich J, Knox KS, Rios CT, Natt B, Bhattacharya D, Fain MJ. SARS-CoV-2 and COVID-19 in older adults: what we may expect regarding pathogenesis, immune responses, and outcomes. *Geroscience* 2020; **42**: 505–14.
- 175 Harris-Kojetin L, Sengupta M, Park-Lee E, Valverde R, Caffrey C, Rome V, et al. Long-term care providers and services users in the United States: data from the national study of long-term care providers, 2013–2014. *Vital Health Stat 3* 2016; (38): x–xii.
- 176 Cagnin A, Di Lorenzo R, Marra C, Bonanni L, Cupidi C, Laganà V, et al. Behavioral and psychological effects of coronavirus disease-19 quarantine in patients with dementia. *Stat Med* 2020; **11**: 578015.
- 177 Bayati M. Why is COVID-19 more concentrated in countries with high economic status?. *Iran J Public Health* 2021; **50**: 1926–9.
- 178 Statsenko Y, Al Zahmi F, Habuza T, Almansoori TM, Smetanina D, Simiyi GL, et al. Impact of age and sex on COVID-19 severity assessed from radiologic and clinical findings. *Front Cell Infect Microbiol* 2021; **11**: 777070.
- 179 Dadrás O, SeyedAlinaghi S, Karimi A, Shamsabadi A, Qaderi K, Ramezani M, et al. COVID-19 mortality and its predictors in the elderly: a systematic review. *Health Sci Rep* 2022; **5**: e657.
- 180 Putri C, Hariyanto TI, Hananto JE, Christian K, Situmeang RFV, Kurniawan A. Parkinson's disease may worsen outcomes from coronavirus disease 2019 (COVID-19) pneumonia in hospitalized patients: a systematic review, meta-analysis, and meta-regression. *Parkinsonism Relat Disord* 2021; **87**: 155–61.
- 181 Rajati F, Ahmadi N, Naghibzadeh ZA, Kazemina M. The global prevalence of oropharyngeal dysphagia in different populations: a systematic review and meta-analysis. *J Transl Med* 2022; **20**: 175.
- 182 Banda KJ, Chu H, Chen R, Kang XL, Jen HJ, Liu D, et al. Prevalence of oropharyngeal dysphagia and risk of pneumonia, malnutrition, and mortality in adults aged 60 years and older: a meta-analysis. *Gerontology* 2022; **68**: 841–53.
- 183 Torres A, Peetermans WE, Vieggi G, Blasi F. Risk factors for community-acquired pneumonia in adults in Europe: a literature review. *Thorax* 2013; **68**: 1057–65.

- 184 Mitchell SL, Teno JM, Kiely DK, Shaffer ML, Jones RN, Prigerson HG, et al. The clinical course of advanced dementia. *N Engl J Med* 2009; **361**: 1529–38.
- 185 Antommaria AHM, Gibb TS, McGuire AL, Wolpe PR, Wynia MK, Applewhite MK, et al. Ventilator triage policies during the COVID-19 pandemic at U.S. hospitals associated with members of the association of bioethics program directors. *Ann Intern Med* 2020; **173**: 188–94.
- 186 Bledsoe TA, Jokela JA, Deep NN, Snyder Sulmasy L. Universal do-not-resuscitate orders, social worth, and life-years: opposing discriminatory approaches to the allocation of resources during the COVID-19 pandemic and other health system catastrophes. *Ann Intern Med* 2020; **173**: 230–2.
- 187 National Institute for Health and Care Excellence. *COVID-19 rapid guideline: Critical care in adults (NICE Guideline NG159)*. NICE, 2022.
- 188 Centers for Disease Control and Prevention. *About Underlying Cause of Death, 1999–2019*. CDC WONDER, 2020 (<https://wonder.cdc.gov/ucd-icd10.html>).
- 189 Bartolomé F, Rosa L, Valenti P, Lopera F, Hernández-Gallego J, Cantero JL, et al. Lactoferrin as immune-enhancement strategy for SARS-CoV-2 infection in Alzheimer's disease patients. *Front Immunol* 2022; **13**: 878201.
- 190 Huang LT, Zhang CP, Wang YB, Wang JH. Association of peripheral blood cell profile with Alzheimer's disease: a meta-analysis. *Front Aging Neurosci* 2022; **14**: 888946.
- 191 De Zuani M, Lazničková P, Tomašková V, Dvončová M, Forte G, Stokin GB, et al. High CD4-to-CD8 ratio identifies an at-risk population susceptible to lethal COVID-19. *Scand J Immunol* 2022; **95**: e13125.
- 192 Numbers K, Brodaty H. The effects of the COVID-19 pandemic on people with dementia. *Nat Rev Neurol* 2021; **17**: 69–70.
- 193 Cheung MW. A guide to conducting a meta-analysis with non-independent effect sizes. *Neuropsychol Rev* 2019; **29**: 387–96.
- 194 von Hippel PT. The heterogeneity statistic I^2 can be biased in small meta-analyses. *BMC Med Res Methodol* 2015; **15**: 35.
- 195 Higgins JPT, Thomas J. *Cochrane Handbook for Systematic Reviews of Interventions*. John Wiley & Sons, 2019.

