



---

Year: 2023

---

## Seladelpar treatment reduces interleukin-31 and pruritus in patients with primary biliary cholangitis

Kremer, Andreas E ; Mayo, Marlyn J ; Hirschfield, Gideon M ; Levy, Cynthia ; Bowlus, Christopher L ; Jones, David E ; Johnson, Jeff D ; McWherter, Charles A ; Choi, Yun-Jung

**Abstract:** BACKGROUND AND AIMS: Pruritus is a debilitating symptom for many people living with primary biliary cholangitis (PBC). In studies with seladelpar, a selective PPAR-delta agonist, PBC patients experienced significant improvement in pruritus and reduction of serum bile acids. Interleukin-31 (IL-31) is a cytokine known to mediate pruritus and blocking IL-31 signaling provides relief in pruritic skin diseases. This study examined the connection between seladelpar's anti-pruritic effects, IL-31 and bile acid levels in PBC patients. APPROACH AND RESULTS: IL-31 levels were quantified in serum samples from the ENHANCE study of PBC patients receiving daily oral doses of placebo (n=55), seladelpar 5 mg (n=53) or 10 mg (n=53) for 3 months and for healthy volunteers (n=55). IL-31 levels were compared with pruritus using a numerical rating scale (NRS, 0-10) and with bile acid levels. Baseline IL-31 levels closely correlated with pruritus NRS ( $r=0.54$ ,  $p<0.0001$ ), and total ( $r=0.54$ ,  $p<0.0001$ ) and conjugated bile acids (up to  $0.64$ ,  $p<0.0001$ ). Decreases in IL-31 were observed with seladelpar 5 mg (-30%,  $p=0.0003$ ) and 10 mg (-52%,  $p<0.0001$ ) versus placebo (+31%). Patients with clinically meaningful improvement in pruritus (NRS $\geq 2$  decrease) demonstrated greater dose-dependent reductions in IL-31 compared to those without pruritus improvement (NRS $< 2$  decrease). Strong correlations were observed for the changes between levels of IL-31 and total bile acids ( $r=0.63$ ,  $p<0.0001$ ) in the seladelpar 10 mg group. CONCLUSIONS: Seladelpar decreased serum IL-31 and bile acids in PBC patients. The reductions of IL-31 and bile acids correlated closely with each other and pruritus improvement suggesting a mechanism to explain seladelpar's anti-pruritic effects.

DOI: <https://doi.org/10.1097/HEP.0000000000000728>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-254272>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Kremer, Andreas E; Mayo, Marlyn J; Hirschfield, Gideon M; Levy, Cynthia; Bowlus, Christopher L; Jones, David E; Johnson, Jeff D; McWherter, Charles A; Choi, Yun-Jung (2023). Seladelpar treatment reduces interleukin-31 and pruritus in patients with primary biliary cholangitis. *Hepatology*:Epub ahead of print.

DOI: <https://doi.org/10.1097/HEP.0000000000000728>

## Seladelpar treatment reduces interleukin-31 and pruritus in patients with primary biliary cholangitis

**Short title:** Seladelpar reduces IL-31 and cholestatic itch

**Authors:**

Andreas E. Kremer<sup>1</sup>, Marlyn J. Mayo<sup>2</sup>, Gideon M. Hirschfield<sup>3</sup>, Cynthia Levy<sup>4,5</sup>, Christopher L. Bowlus<sup>6</sup>, David E. Jones<sup>7</sup>, Jeff D. Johnson<sup>8</sup>, Charles A. McWherter<sup>8</sup>, and Yun-Jung Choi<sup>8</sup>

**Author Affiliations:**

1. Department of Gastroenterology and Hepatology, University Hospital Zürich, University of Zürich, Zürich, Switzerland.
2. Division of Digestive and Liver Diseases, University of Texas SW Medical Center, Dallas, Texas, USA.
3. Toronto Centre for Liver Disease, Toronto General Hospital, University Health Network, Toronto, Ontario, Canada.
4. Division of Digestive Health and Liver Diseases, University of Miami Miller School of Medicine. Miami, Florida, USA.
5. Schiff Center for Liver Diseases, University of Miami, Miami, FL, USA
6. Division of Gastroenterology and Hepatology, Department of Internal Medicine, University of California Davis, Sacramento, California, USA.
7. Clinical and Translation Research Institute, Newcastle University, Newcastle upon Tyne, UK.
8. CymaBay Therapeutics, Inc., Newark, California, USA.

**Keywords:** PPAR-delta agonist; cholestatic pruritus; itch; bile acids; pruritus NRS

**Correspondence:**

Names: Andreas E. Kremer and Yun-Jung Choi

Andreas E. Kremer, M.D., Ph.D., MHBA

Department of Gastroenterology and Hepatology

University Hospital Zürich, Rämistrasse 100, CH-8091

Zürich, Switzerland.

Email: andreas.kremer@usz.ch

Tel: +41-44-255-85 48

Yun-Jung Choi, Ph.D.

CymaBay Therapeutics, Inc.

Newark, California, USA.

Email: ychoi@cymabay.com

Tel: +1-510-954-2755

Graphical abstract

GA1

**AUTHOR CONTRIBUTIONS:**

Yun-Jung Choi, Jeff D. Johnson, Charles A. McWherter, and Andreas E. Kremer contributed to the study concept and study design. Yun-Jung Choi was responsible for data collection and data analysis. Andreas E. Kremer, Marlyn J. Mayo, Gideon M. Hirschfield, Cynthia Levy, Christopher L. Bowlus, and David E. Jones were investigators in the ENHANCE study. All

authors interpreted data, reviewed, and revised drafts of the manuscript, and approved the final manuscript for submission.

#### **AUTHOR E-MAIL AND ORCID ID:**

<b>Name</b>	<b>E-mail</b>	<b>ORCID ID</b>
Andreas E. Kremer	andreas.kremer@usz.ch	0000-0002-9263-948X
Marlyn J. Mayo	marlyn.mayo@utsouthwestern.edu	0000-0002-4874-7010
Gideon Hirschfield	gideon.hirschfield@uhn.ca	0000-0002-6736-2255
Cynthia Levy	clevy@med.miami.edu	0000-0001-5498-6037
Christopher L. Bowlus	clbowlus@ucdavis.edu	0000-0002-3906-6811
David E. J. Jones	david.jones@newcastle.ac.uk	NA
Jeff D. Johnson	jeffjohnson3142@gmail.com	0009-0001-9384-9577
Charles A. McWherter	cmcwherter@cymabay.com	0000-0002-7613-3008
Yun-Jung Choi	ychoi@cymabay.com	0000-0002-5153-9714

#### **CONFLICTS OF INTEREST**

Andreas E. Kremer consults, advises, is on the speakers' bureau, and received grants from Gilead and Intercept. He consults, advises, is on the speakers' bureau for AbbVie, Advanz, Bayer, CymaBay, Eisai, Falk, GlaxoSmithKline, MSD, Novartis, and Roche. He consults and advises Beiersdorf, Escient, FMC, Guidepoint, Medscape, Mirum, Myr, and Viofor. He is on the speakers' bureau for AOP Orphan, Bristol Myers Squibb, CMS, Janssen, Lilly, Newbridge, Viofor, and Zambon. Marlyn J. Mayo consults and is on the speakers' bureau for Intra-Sana Laboratories. She consults and received grants from TARGET. She advises and received grants from CymaBay, GlaxoSmithKline, and Ipsen. She advises and received grants from Mallinckrodt. She received grants from GENFIT, Intercept, and Mirium. Gideon M. Hirschfield consults for and is on the speakers' bureau for GlaxoSmithKline, Intercept, and Ipsen. He consults for Advanz, CymaBay, Escient, Gilead, Kowa, Mirum, and Pliant. Cynthia Levy consults, advises, and received grants from CymaBay, GlaxoSmithKline, Intercept, and Ipsen.

She consults and received grants from Calliditas and Mirium. She consults for Escient and Kowa Research Institute. She received grants from Cara, Escient, Gilead, HighTide, TARGET, and Zydus. She has other interests with AASLD. Christopher L. Bowlus consults and received grants from CymaBay, GlaxoSmithKline, Ipsen, and Mirum. He consults for BiomX, Eli Lilly, Invea Therapeutics, Shire, and Trevi. He received grants from Bristol Myers Squibb, Boston Scientific, Calliditas, Chemomab, COUR Pharmaceuticals, Eli Lilly, GENFIT, Gilead, Hanmi, Intercept, Novartis, Novo Nordisk, Pliant, Takeda, TARGET, and Viking. David E. Jones consults, is on the speakers' bureau, and received grants from Intercept. He consults and is on the speakers' bureau for Ipsen. He consults for CymaBay, Kowa, and Umecrine. He is on the speakers' bureau for Falk and GlaxoSmithKline. Jeff D. Johnson consults for CymaBay. Charles A. McWherter is employed by and owns stock in CymaBay. Yun-Jung Choi is employed by CymaBay.

## DATA AVAILABILITY

Data are available from the corresponding authors upon reasonable request.

**Trial registration:** ClinicalTrials.gov number: NCT03602560.

**Abbreviations:** ALP, alkaline phosphatase; CA, cholic acid; CDCA, chenodeoxycholic acid; DCA, deoxycholic acid; DRG, dorsal root ganglia; FXR, Farnesoid X receptor; GCA, glycocholic acid; GCDCA, glycochenodeoxycholic acid; GDCA, glycodeoxycholic acid; GLCA, glycolithocholic acid; GPCR, G protein-coupled receptor; GUDCA, glyoursodeoxycholic acid; IBAT, ileal bile acid transporter; IL-31, Interleukin-31; IL-31RA, Interleukin-31 receptor A; LCA, lithocholic acid; LPA, lysophosphatidic acid; NRS, numerical rating scale; PBC, primary biliary cholangitis; PSC, primary sclerosing cholangitis; PPAR, peroxisome proliferator-activated receptor; TCA, taurocholic acid; TCDCA, taurochenodeoxycholic acid; TDCA, taurodeoxycholic acid; TLCA, tauroolithocholic acid; Th2, T helper 2; TUDCA, taoursodeoxycholic acid; UDCA, ursodeoxycholic acid; ULN, upper limit of normal.

**Funding information:** CymaBay Therapeutics, Inc.

## Abstract

**Background and Aims:** Pruritus is a debilitating symptom for many people living with primary biliary cholangitis (PBC). In studies with seladelpar, a selective PPAR-delta agonist, PBC patients experienced significant improvement in pruritus and reduction of serum bile acids. Interleukin-31 (IL-31) is a cytokine known to mediate pruritus and blocking IL-31 signaling provides relief in pruritic skin diseases. This study examined the connection between seladelpar's anti-pruritic effects, IL-31 and bile acid levels in PBC patients.

**Approach and Results:** IL-31 levels were quantified in serum samples from the ENHANCE study of PBC patients receiving daily oral doses of placebo (n=55), seladelpar 5mg (n=53) or 10mg (n=53) for 3 months and for healthy volunteers (n=55). IL-31 levels were compared with pruritus using a numerical rating scale (NRS, 0-10) and with bile acid levels. Baseline IL-31 levels closely correlated with pruritus NRS ( $r=0.54$ ,  $p<0.0001$ ), and total ( $r=0.54$ ,  $p<0.0001$ ) and conjugated bile acids (up to  $0.64$ ,  $p<0.0001$ ). Decreases in IL-31 were observed with seladelpar 5mg (-30%,  $p=0.0003$ ) and 10mg (-52%,  $p<0.0001$ ) versus placebo (+31%). Patients with clinically meaningful improvement in pruritus (NRS $\geq$ 2 decrease) demonstrated greater dose-dependent reductions in IL-31 compared to those without pruritus improvement (NRS<2 decrease). Strong correlations were observed for the changes between levels of IL-31 and total bile acids ( $r=0.63$ ,  $p<0.0001$ ) in the seladelpar 10mg group.

**Conclusions:** Seladelpar decreased serum IL-31 and bile acids in PBC patients. The reductions of IL-31 and bile acids correlated closely with each other and pruritus improvement suggesting a mechanism to explain seladelpar's anti-pruritic effects.

## INTRODUCTION

Primary biliary cholangitis (PBC) is a rare, slowly progressive, female-predominant liver disease characterized by immune-mediated destruction of small intrahepatic bile ducts followed by the development of chronic cholestasis.<sup>[1]</sup> If untreated, PBC patients have a higher risk of cirrhosis and its complications, liver failure and death.<sup>[2]</sup> Along with fatigue, pruritus is one of the most common presenting symptoms in patients with PBC.<sup>[3,4]</sup> Treatments for cholestatic pruritus are currently limited, and active pruritus can cause sleep deprivation, exacerbate fatigue, and impact quality of life for patients living with PBC. The currently approved first line (ursodeoxycholic acid; UDCA) and second line therapies (obeticholic acid) improve biochemical markers of PBC associated with disease progression but fail to address key symptoms, including pruritus, associated with the disease. In the absence of efficacious therapy acceptable to patients, pruritus remains a high unmet need for patients with PBC.<sup>[5]</sup>

Seladelpar is a potent and selective peroxisome proliferator-activated receptor (PPAR)-delta agonist in development for the treatment of patients with PBC. In previous studies with PBC patients having insufficient response to UDCA, seladelpar 10 mg demonstrated significant improvements in biochemical markers of cholestasis as well as significant reductions in patient-reported pruritus measurements in those patients with moderate-to-severe pruritus.<sup>[6-8]</sup> Seladelpar treatment also improved sleep disturbance and fatigue measured by the 5-D itch and PBC-40 questionnaires.<sup>[6]</sup>

A variety of pruritogenic molecules have been suggested to play a role in the pruritus experienced by patients with PBC but the precise mechanism remains elusive. Individual bile acids, or sets of them, may be components of an ensemble of pruritogenic or pruritus-sensitizing signaling molecules, although serum bile acids are not always correlated with itch.<sup>[9]</sup>

Nevertheless, bile acids can initiate signaling through G protein-coupled receptors (GPCRs) such as MRGPRX4, which has been localized to human dorsal root ganglia (DRG) associated sensory neurons as well as a variety of other cells.<sup>[10,11]</sup> Autotaxin, a lysophosphatidic acid forming enzyme, is associated with severity of cholestatic itch and is lowered by certain but not all anti-pruritic treatments.<sup>[9,12,13]</sup>

Interleukin-31 (IL-31) is a pruritogenic cytokine produced by immune cells such as monocytes, mast cells, macrophages, fibroblasts, eosinophils, basophils and, especially, T helper 2 (Th2)

cells.<sup>[14-16]</sup> IL-31 has a role in neurite outgrowth and branching, and in transmitting pruritic signals through a heterodimeric receptor composed of IL-31 receptor A (IL-31RA) and oncostatin M receptor  $\beta$  on keratinocytes and DRG neurons.<sup>[17-19]</sup> Higher levels of IL-31 or IL-31RA have been reported in the tissue and serum of patients with pruritic skin diseases such as atopic dermatitis.<sup>[16]</sup> Recent studies with nemolizumab, a humanized antibody against IL-31RA, demonstrated significant decreases of pruritus and improvement of sleep disturbance and quality of life in patients with atopic dermatitis,<sup>[20-22]</sup> but its effects on itch in PBC have not been reported. However, elevated levels of IL-31 have been reported in patients with cholestatic diseases such as PBC, primary sclerosing cholangitis (PSC), and intrahepatic cholestasis of pregnancy.<sup>[23-25]</sup> Furthermore, Xu and colleagues<sup>[23]</sup> demonstrated that farnesoid X receptor (FXR) agonists, which are known to cause itch when treating patients, resulted in increases in levels of IL-31 in humans and in mice with humanized livers.

In this study, we evaluated the association of levels of IL-31, type 2 cytokines, and bile acids with pruritus experienced by PBC patients being treated with seladelpar or placebo. IL-31 and bile acid levels were also assessed in comparison with age-, sex- and BMI-matched healthy volunteers. This report demonstrates for the first time that IL-31, a known pruritogenic mediator, can be reduced in patients with PBC by a therapeutic intervention that concomitantly improves pruritus. Our findings both shed light on potential mediators for cholestatic itch and provide important information on novel therapeutic approaches.

## METHODS

### Study design, patients, and treatment

The ENHANCE study [ClinicalTrials.gov (NCT03602560) and EU Clinical Trials Registry (EudraCT2018-001171-20)]<sup>[8]</sup> was a multicenter, randomized, double-blind, placebo-controlled phase 3 study of seladelpar in patients with PBC with an inadequate response or intolerance to UDCA. Patients (18-72 years old) with a diagnosis of PBC, elevated alkaline phosphatase (ALP) level (ALP $\geq$ 1.67 times the upper limit of normal [ULN]) and total bilirubin $\leq$ 2 times ULN were eligible to participate in the study. Enrolled patients were stratified by ALP level (<350U/L or  $\geq$ 350U/L) and pruritus numerical rating scale (NRS) (<4 or  $\geq$ 4) and randomly assigned to a treatment group to receive once-daily dosing of oral placebo, seladelpar 5 mg or 10 mg. The



study protocol was approved by appropriate local and national institutional review boards or independent ethics committees, and the trial was conducted in accordance with the Declaration of Helsinki and Good Clinical Practice guidelines. All patients provided written informed consent prior to study procedures. The detailed methodology and results of the ENHANCE study have been described in full elsewhere.<sup>[8]</sup> This analysis evaluated 161 PBC patients with available serum samples (placebo, n=55; seladelpar 5 mg, n=53; seladelpar 10 mg, n=53) at baseline and study endpoint (Month 3) to investigate the treatment effects of seladelpar on IL-31 and its association with pruritus NRS, bile acids and other laboratory measures.

To compare baseline IL-31 and bile acid levels in PBC patients with healthy volunteers, serum samples were collected from healthy volunteers (n=55) who were prospectively recruited by matching age, sex and BMI to the PBC patients in the ENHANCE study (BioCollections Worldwide, Inc.).

To confirm the findings in the ENHANCE study, we evaluated the effect of seladelpar on IL-31, bile acids and pruritus in a separate population enrolled in an open-label phase 2 study with PBC patients (NCT02955602; EudraCT # 2016-002996-91).<sup>[7]</sup>

### **Pruritus numerical rating scale (NRS)**

Patients reported a NRS each evening for the previous 24 hours on a scale of 0 (no itching) to 10 (worst imaginable itching) using a daily electronic diary. Pruritus NRS at baseline and study endpoint (Month 3) were determined by the means of all daily recorded scores during the run-in period (up to 14 days) and the week prior to the Month 3 visit, respectively. Changes in IL-31 levels associated with clinically meaningful improvement in pruritus NRS ( $\geq 2$  decrease) as defined by Reich and colleagues<sup>[26]</sup> were evaluated.

### **Cytokine measurements**

Serum IL-31 and other known pruritic cytokines such as IL-4, IL-13, and IL-33 were quantified by Simoa<sup>TM</sup> ultrasensitive immunoassays using Simoa HD-1 Analyzer and single molecule array technology (Rules-Based Medicine, Austin, TX, USA). Capture antibody-conjugated paramagnetic beads were incubated with standards, serum samples or controls and biotinylated detection antibodies. The beads were then washed and incubated with streptavidin  $\beta$ -

galactosidase. After the final wash, the beads were loaded into the Simoa Disc with the enzyme substrate resorufin  $\beta$ -galactopyranoside. The concentration of IL-31, IL-4, IL-13, or IL-33 in each sample was calculated by interpolating from a standard curve. The lower limits of quantitation for IL-31, IL-4, IL-13 and IL-33 were 0.03 pg/mL, 0.022 pg/mL, 0.36 pg/mL, and 0.19 pg/mL, respectively.

### **Autotaxin measurement**

Serum autotaxin levels were measured using human ENPP-2/Autotaxin Quantikine ELISA kit (R&D systems, Inc., USA) according to the manufacturer's instructions.

### **Bile acid measurements**

Serum bile acids in samples at baseline and Month 3 in ENHANCE and in those from healthy volunteers were analyzed quantitatively by liquid chromatography tandem mass spectrometry (Metabolon, Durham, NC, USA). Fifteen primary, secondary bile acids and glycine- and taurine-conjugates were measured: cholic acid (CA), glycocholic acid (GCA), taurocholic acid (TCA), chenodeoxycholic acid (CDCA), glycochenodeoxycholic acid (GCDCA), taurochenodeoxycholic acid (TCDCA), deoxycholic acid (DCA), glycodeoxycholic acid (GDCA), taurodeoxycholic acid (TDCA), lithocholic acid (LCA), glycolithocholic acid (GLCA), tauroolithocholic acid (TLCA), ursodeoxycholic acid (UDCA), glyoursodeoxycholic acid (GUDCA), and taoursodeoxycholic acid (TUDCA).

### **Statistical analyses**

JMP statistical software (version 17; SAS Institute, Cary, NC, USA) was used for statistical analysis. Graphs were created in PRISM (version 9; GraphPad, San Diego, CA, USA). Descriptive statistics were used to summarize clinical data. Paired t-tests were used to detect differences for a given patient between baseline and Month 3 for each treatment group. One-way analysis of variance (ANOVA) followed by a post-hoc Student's t-test was used to evaluate the difference between treatment groups. Pearson's correlation coefficients (r) using multivariate analysis were used to evaluate correlations between IL-31 and pruritus NRS, autotaxin, and bile acids.

## **RESULTS**

## Baseline characteristics

Of the 161 PBC patients evaluated at study endpoint (Month 3), 55 received placebo, 53 received seladelpar 5 mg, and 53 received seladelpar 10 mg. Demographics and baseline characteristics are summarized in Table 1 and were generally well-balanced between treatment groups. Patients were mostly females (94%) living with PBC for 9 years, a mean (SD) age of 56 (9) years, 90% AMA-positive, 95% on UDCA, BMI of 29 (6) kg/m<sup>2</sup>, and 28% with moderate to severe pruritus (NRS $\geq$ 4, mean (SD), NRS=6.1 (1.4)). The number of patients with cirrhosis was limited in this study (11%, n=17) with few suffering from moderate to severe pruritus (Placebo, n=3; 5 mg, n=2; 10 mg, n=1). The 55 healthy volunteers were matched to PBC patients in this study: a female proportion of 95% (52 females and 3 males), a mean age of 56 (5) years and BMI of 29 (3) kg/m<sup>2</sup>.

PBC patients had 31-fold higher serum concentrations of IL-31 compared to those of healthy volunteers (Figure 1A). PBC patients in our study also demonstrated elevated serum bile acid levels at baseline compared to healthy volunteers (Figure 1B). Serum levels of IL-31 correlated positively with total bile acids ( $r=0.56$ ,  $p<0.0001$ , Figure 1C).<sup>[23]</sup>

## Seladelpar treatment decreased serum IL-31 levels in patients with PBC

At baseline, there were no significant differences in serum IL-31 levels between treatment groups ( $p=0.8962$ ). Mean changes from baseline to Month 3 for serum IL-31 levels among treatment groups are presented in Figure 2A. Seladelpar treatment resulted in substantial decreases in mean IL-31 levels from baseline to Month 3: seladelpar 5 mg (3.8 to 1.7 pg/mL,  $p=0.0002$ ), 10 mg (4.2 to 1.7 pg/mL,  $p=0.0003$ ) compared to placebo (4.3 to 3.9 pg/mL,  $p=0.28$ ). A robust dose-dependent percentage decrease in IL-31 levels was observed with the seladelpar treatments (Figure 2B): seladelpar 5 mg (-30%,  $p=0.0003$ ) and 10 mg (-52%,  $p<0.0001$ ) compared to placebo (+31%). Changes in IL-31 for individual patients are presented in Figure 2C. After 3-months of seladelpar treatment, 79% of patients in the 5 mg and 91% of patients in the 10 mg had decreased IL-31 levels compared to mixed responses in the placebo group (56% decrease and 44% increase). Seladelpar treatment lowered serum ALP as described previously (Placebo: -4.2%; seladelpar 5 mg: -35.9%; Seladelpar 10 mg: -44.3%, respectively)<sup>[8]</sup> and changes in IL-31 and changes in ALP were also correlated in the seladelpar treatment groups

(placebo  $r=0.11$ ,  $p=0.4179$ ; 5 mg  $r=0.56$ ,  $p<0.0001$ ; 10 mg  $r=0.40$ ,  $p=0.0028$ ). Seladelpar treatment did not result in changes in other type 2 cytokines, IL-4, IL-13, and IL-33, known to be involved in production of IL-31 in other diseases such as atopic dermatitis. (Supplementary Figure 1A-C, <http://links.lww.com/HEP/I187>).<sup>[27]</sup>

### **Correlation of itch intensity and serum IL-31 levels with seladelpar treatment**

At baseline, serum IL-31 levels were closely correlated with pruritus NRS ( $r=0.54$ ,  $p<0.0001$ , Figure 3A). A significant, dose-ordered association was maintained between changes in IL-31 and pruritus NRS in the seladelpar treatment groups and placebo group (Figure 3B): placebo ( $r=0.36$ ,  $p=0.0080$ ), seladelpar 5 mg ( $r=0.44$ ,  $p=0.0011$ ) and seladelpar 10 mg ( $r=0.54$ ,  $p<0.0001$ ). Patients having clinically meaningful improvement in pruritus (NRS $\geq 2$  decrease)<sup>[26]</sup> had greater dose-dependent reductions in serum IL-31 levels compared to those without pruritus improvement (NRS $< 2$  decrease) (Figure 3C). Baseline levels of IL-31 were 1.7- to 2.2-fold greater in those seladelpar patients with pruritus improvement (NRS $\geq 2$  decrease) than in placebo patients. In patients without improvement (NRS $< 2$  decrease), baseline IL-31 levels were 2.0- to 4.7-fold less than seladelpar patients who improved (Bottom of Figure 3C).

### **Treatment effects of seladelpar on serum IL-31 levels and pruritus in an open-label phase 2 study**

To confirm the findings in the ENHANCE study, we evaluated the effect of seladelpar on IL-31, bile acids and pruritus in a separate population enrolled in an open-label phase 2 study with PBC patients. Seladelpar treatment in this study had previously demonstrated improvements of pruritus, sleep disturbance and fatigue measured by visual analog scale (VAS, 0-100), 5-D itch and PBC-40 questionnaires at 1 year.<sup>[6]</sup> Similar to ENHANCE, we found substantial decreases in serum IL-31 levels with seladelpar treatment. A correlation of serum IL-31 and pruritus measured by VAS was also observed at baseline. An improvement of pruritus measured by VAS was correlated with a decrease in serum IL-31 levels with 1 year of seladelpar 10 mg treatment (Supplementary Figure 2, <http://links.lww.com/HEP/I187>).

### **Correlation between serum IL-31 levels and circulating bile acids**

The correlations of serum IL-31 with serum bile acids at baseline and with the change from baseline to Month 3 are shown in Table 2. There were no correlations between serum IL-31 levels and any of the unconjugated bile acids at baseline. Serum IL-31 levels were significantly and closely correlated with taurine- or glycine-conjugated bile acids ( $p < 0.0001$ ) at baseline: TCA ( $r = 0.64$ ), TCDCA ( $r = 0.54$ ), total bile acids ( $r = 0.54$ ), GCA ( $r = 0.52$ ), TDCA ( $r = 0.52$ ), TUDCA ( $r = 0.49$ ), GCDCA ( $r = 0.45$ ), total bile acids including unconjugated and conjugated UDCA ( $r = 0.43$ ), GUDCA ( $r = 0.40$ ), TLCA ( $r = 0.39$ ), GDCA ( $r = 0.31$ ) (Supplementary Figure 3, <http://links.lww.com/HEP/I187>). As for the observations at baseline, treatment for 3 months led to no significant correlations between changes in serum IL-31 levels and unconjugated bile acids. In marked contrast, strong and significant correlations between changes in serum IL-31 levels and changes in GCA ( $r = 0.67$ ,  $p < 0.0001$ ), total bile acids ( $r = 0.63$ ,  $p < 0.0001$ ) and TCA ( $r = 0.62$ ,  $p < 0.0001$ ) were seen with the seladelpar 10 mg treatment for 3 months. Seladelpar 10 mg treatment led to a significant decrease in total bile acids ( $p = 0.035$ ) which was attributed to the significant reductions in the major glycine-conjugated primary bile acids GCDCA ( $p = 0.033$ ) and GCA ( $p = 0.030$ ) (Supplementary Figure 4, <http://links.lww.com/HEP/I187>).

### **Effect of seladelpar treatment on serum autotaxin levels**

Autotaxin has been associated with pruritus in PBC.<sup>[9]</sup> Various antipruritic therapies including cholestyramine, rifampicin, nasobiliary drainage and ileal bile acid transporter (IBAT) inhibitors also reduce serum level of this enzyme.<sup>[9,28,29]</sup> In our study, we also found that baseline serum autotaxin levels were correlated with pruritus as well as serum IL-31 and total bile acids. However, seladelpar treatment did not result in decreases in autotaxin levels (Supplementary Figure 5, <http://links.lww.com/HEP/I187>).

## **DISCUSSION**

In the absence of effective and well tolerated treatment options, treatment for pruritus remains a high unmet medical need in patients with PBC. In this study we demonstrate that the improvement in pruritus from seladelpar treatment in patients with PBC is correlated with reductions in the known pruritogenic cytokine IL-31. We importantly demonstrate elevated baseline levels of IL-31 in patients, an association between reductions in IL-31 and seladelpar

treatment, as well as an association between pruritus improvement and reductions in IL-31. The reduction in IL-31 levels associated with improvements in bile acid levels. Seladelpar treatment did not, however, correlate with other related type 2 cytokines nor with autotaxin.

In previous studies, seladelpar treatment in these patients resulted in a robust improvement in pruritus and associated symptoms<sup>[6-8]</sup> but the mechanism by which this PPAR-delta agonist ameliorates pruritus is not well understood. IL-31 is a known critical mediator in pruritic skin diseases such as atopic dermatitis.<sup>[20]</sup> Two studies,<sup>[23,24]</sup> including ours, have reported elevation of serum IL-31 levels among PBC patients compared to healthy subjects. Of importance, FXR agonists, which are known to improve cholestatic markers (e.g. ALP) while causing or worsening pruritus in PBC patients,<sup>[30]</sup> dose-dependently increased serum IL-31 in clinical studies in NASH, PBC and PSC, and in mice with humanized livers.<sup>[23]</sup> Clearly cholestatic markers, IL-31 and pruritus can diverge with certain treatments.

In the current study, the majority of patients treated with seladelpar 10 mg demonstrated a remarkably uniform reduction in IL-31. We have observed close and significant correlations between IL-31 levels and severity of pruritus measured by NRS at baseline and changes at month 3 with seladelpar 10 mg treatment. These correlations were comparable to that of other pruritic diseases such as prurigo nodularis for which IL-31 has been therapeutically validated.<sup>[31-33]</sup> Pruritus NRS data were collected by patients using daily electronic diaries, which are the preferred method for accurately recording robust pruritus data, and this difference in methodology may explain why a previous report did not observe a correlation between pruritus and serum IL-31 in patients with PBC.<sup>[24]</sup>

Therapies to reduce bile acids such as bile acid sequestrants, nasobiliary drainage, inhibitors of IBAT, and the pan-PPAR agonist bezafibrate have demonstrated relief of pruritus in patients with PBC<sup>[34-36]</sup> thus demonstrating a connection between bile acids and pruritus. Receptor-mediated effects of bile acids via FXR and GPCRs are compelling in this regard, and these may represent components of itch signaling. Alternatively, bile acids can cause oxidative stress and mitochondrial injury in hepatocytes leading to inflammation. Other components of the hepatobiliary system are also impacted by bile acids, although the mechanisms of the impact of cholestatic levels of bile acids on the liver are also still subject to debate.<sup>[37]</sup>

At baseline, IL-31 in our study was clearly correlated with total bile acids and a variety of taurine- and glycine-conjugated bile acids but not with unconjugated bile acids. This correlation is maintained after the changes due to treatment with seladelpar in which both IL-31 and conjugated bile acids are decreased. This may suggest that conjugated bile acids, despite being considered less toxic, are more proximal to the regulation of IL-31 release.

We measured autotaxin levels in the patients with PBC in our study and found that they are correlated with pruritus, total bile acids and serum IL-31 levels at baseline. However, seladelpar did not lower the serum levels of autotaxin. This is concordant with the results of the BEZURSO (PBC) and FITCH (PBC and PSC) studies of bezafibrate where this pan-PPAR agonist reduced pruritus but did not reduce serum autotaxin levels.<sup>[36,38,39]</sup>

IL-31 has been reported to affect itch by directly acting on sensory neurons or adjacent keratinocytes, both of which possess high affinity IL-31 receptors.<sup>[18,40]</sup> *In vitro* and *in vivo* evidence indicates that IL-31 can lengthen and cause branching of such neurons.<sup>[17]</sup> Our finding that the type 2 cytokines IL-4, IL-13 and IL-33 are not affected by seladelpar suggests that the role of IL-31 in PBC itch is different than that seen in skin diseases. In cholestatic itch, IL-31 is putatively derived from the liver<sup>[23]</sup> and it may not respond to IL-4, IL-13 and IL-33 as it is known for T-helper type 2 cells in dermatological diseases. Furthermore, the observation that agonists of the bile acid receptor FXR both cause itch and increase IL-31 suggests a connection between cholestasis, the levels of bile acids, IL-31 and itch. Seladelpar improved cholestasis while lowering bile acids and IL-31 presenting a plausible explanation for its observed improvement of pruritus. This reinforces that IL-31 may serve as pruritogenic mediator in both skin and cholestatic diseases but may originate from different sources.

Whether IL-31, bile acids, and autotaxin act independently or coordinately by mutual potentiation is unknown. It seems clear that there are multiple likely components of pruritus signaling in cholestatic disease. Considering the well-established role of IL-31 and its receptor in itch, our findings suggest that IL-31 is also a component of the etiology of pruritus in patients with PBC and an addressable target in its treatment. Hopefully, IL-31 will be a guide for improved understanding of cholestatic liver diseases and facilitate therapeutic interventions in alleviating pruritus.

## REFERENCES

1. Lleo A, Wang GQ, Gershwin ME, Hirschfield GM. Primary biliary cholangitis. *Lancet* 2020;396:1915-1926.
2. Kowdley KV, Bowlus CL, Levy C, Mayo MJ, Pratt DS, Vuppalanchi R, et al. Application of the latest advances in evidence-based medicine in primary biliary cholangitis. *Am J Gastroenterol* 2023;118:232-242.
3. Mayo MJ, Carey E, Smith HT, Mospan AR, McLaughlin M, Thompson A, et al. Impact of pruritus on quality of life and current treatment patterns in patients with primary biliary cholangitis. *Dig Dis Sci* 2023;68:995-1005.
4. Lindor KD, Bowlus CL, Boyer J, Levy C, Mayo M. Primary biliary cholangitis: 2018 practice guidance from the American Association for the Study of Liver Diseases. *Hepatology* 2019;69:394-419.
5. Levy C, Manns M, Hirschfield G. New Treatment paradigms in primary biliary cholangitis. *Clin Gastroenterol Hepatol* 2023;21:2076-2087.
6. Kremer AE, Mayo MJ, Hirschfield G, Levy C, Bowlus CL, Jones DE, et al. Seladelpar improved measures of pruritus, sleep, and fatigue and decreased serum bile acids in patients with primary biliary cholangitis. *Liver Int* 2022;42:112-123.
7. Bowlus CL, Galambos MR, Aspinall RJ, Hirschfield GM, Jones DEJ, Dörffel Y, et al. A phase II, randomized, open-label, 52-week study of seladelpar in patients with primary biliary cholangitis. *J Hepatol* 2022;77:353-364.
8. Hirschfield GM, Shiffman ML, Gulamhusein A, Kowdley KV, Vierling JM, Levy C, et al. Seladelpar efficacy and safety at 3 months in patients with primary biliary cholangitis: ENHANCE, a phase 3, randomized, placebo-controlled Study. *Hepatology* 2023;78:397-415.
9. Kremer AE, Namer B, Bolier R, Fischer MJ, Oude Elferink RP, Beuers U. Pathogenesis and management of pruritus in PBC and PSC. *Dig Dis* 2015;33 Suppl 2:164-175.



10. Meixiong J, Vasavda C, Snyder SH, Dong X. MRGPRX4 is a G protein-coupled receptor activated by bile acids that may contribute to cholestatic pruritus. *Proc Natl Acad Sci U S A* 2019;116:10525-10530.
11. Yu H, Zhao T, Liu S, Wu Q, Johnson O, Wu Z, et al. MRGPRX4 is a bile acid receptor for human cholestatic itch. *Elife* 2019;8:e48431.
12. Hegade VS, Pechlivanis A, McDonald JAK, Rees D, Corrigan M, Hirschfield GM, et al. Autotaxin, bile acid profile and effect of ileal bile acid transporter inhibition in primary biliary cholangitis patients with pruritus. *Liver Int* 2019;39:967-975.
13. Vander Does A, Levy C, Yosipovitch G. Cholestatic itch: our current understanding of pathophysiology and treatments. *Am J Clin Dermatol* 2022;23:647-659.
14. Dillon SR, Sprecher C, Hammond A, Bilsborough J, Rosenfeld-Franklin M, Presnell SR, et al. Interleukin 31, a cytokine produced by activated T cells, induces dermatitis in mice. *Nat Immunol* 2004;5:752-760.
15. Roh YS, Choi J, Sutaria N, Belzberg M, Kwatra MM, Kwatra SG. IL-31 inhibition as a therapeutic approach for the management of chronic pruritic dermatoses. *Drugs* 2021;81:895-905.
16. Kabashima K, Irie H. Interleukin-31 as a clinical target for pruritus treatment. *Front Med (Lausanne)* 2021;8:638325.
17. Feld M, Garcia R, Buddenkotte J, Katayama S, Lewis K, Muirhead G, et al. The pruritus- and TH2-associated cytokine IL-31 promotes growth of sensory nerves. *J Allergy Clin Immunol* 2016;138:500-508.e24.
18. Nemmer JM, Kuchner M, Datsi A, Oláh P, Julia V, Raap U, et al. Interleukin-31 signaling bridges the gap between immune cells, the nervous system and epithelial tissues. *Front Med (Lausanne)* 2021;8:639097.
19. Furue M, Furue M. Interleukin-31 and pruritic skin. *J Clin Med* 2021;10:1906.

20. Kabashima K, Matsumura T, Komazaki H, Kawashima M; Nemolizumab-JP01 Study Group. Trial of nemolizumab and topical agents for atopic dermatitis with pruritus. *N Engl J Med* 2020;383:141-150.
21. Kabashima K, Matsumura T, Komazaki H, Kawashima M; Nemolizumab JP01 and JP02 Study Group. Nemolizumab plus topical agents in patients with atopic dermatitis (AD) and moderate-to-severe pruritus provide improvement in pruritus and signs of AD for up to 68 weeks: results from two phase III, long-term studies. *Br J Dermatol* 2022;186:642-651.
22. Kabashima K, Matsumura T, Hayakawa Y, Kawashima M; Nemolizumab-JP01 Study Group. Clinically meaningful improvements in cutaneous lesions and quality of life measures in patients with atopic dermatitis with greater pruritus reductions after treatment with 60 mg nemolizumab subcutaneously every 4 weeks: Subgroup analysis from a phase 3, randomized, controlled trial. *J Dermatolog Treat* 2023;34:2177096.
23. Xu J, Wang Y, Khoshdeli M, Peach M, Chuang JC, Lin J, et al. IL-31 levels correlate with pruritus in patients with cholestatic and metabolic liver diseases and is farnesoid X receptor responsive in NASH. *Hepatology* 2023;77:20-32.
24. Mu N, Lin F, Jiang Z, Liang Y, Yang Z. Implication of increased serum IL-31 for primary biliary cholangitis. *Immunol Invest* 2021;50:662-670.
25. Basile F, Santamaria A, Mannucci C, Rizzo L, Gangemi S, D'anna R, et al. Interleukin 31 is involved in intrahepatic cholestasis of pregnancy. *J Matern Fetal Neonatal Med* 2017;30:1124-1127.
26. Reich A, Riepe C, Anastasiadou Z, Mędrek K, Augustin M, Szepietowski JC, et al. Itch assessment with visual analogue scale and numerical rating scale: determination of minimal clinically important difference in chronic itch. *Acta Derm Venereol* 2016;96:978-980.
27. Yamamura K, Nakahara T. The dawn of a new era in atopic dermatitis treatment. *J Clin Med* 2022;11:6145.
28. Hegade VS, Kendrick SF, Dobbins RL, Miller SR, Thompson D, Richards D, et al. Effect of ileal bile acid transporter inhibitor GSK2330672 on pruritus in primary biliary cholangitis: a

double-blind, randomised, placebo-controlled, crossover, phase 2a study. *Lancet* 2017;389:1114-1123.

29. Thompson RJ, Arnell H, Artan R, Baumann U, Calvo PL, Czubkowski P, et al. Odevixibat treatment in progressive familial intrahepatic cholestasis: a randomised, placebo-controlled, phase 3 trial. *Lancet Gastroenterol Hepatol* 2022;7:830-842.
30. Nevens F, Andreone P, Mazzella G, Strasser SI, Bowlus C, Invernizzi P, et al. A Placebo-controlled trial of obeticholic acid in primary biliary cholangitis. *N Engl J Med* 2016;375:631-643.
31. Kwatra SG, Yosipovitch G, Legat FJ, Reich A, Paul C, Simon D, et al. Phase 3 Trial of Nemolizumab in Patients with Prurigo Nodularis. *N Engl J Med*. 2023;389:1579-1589.
32. Ständer S, Yosipovitch G, Legat FJ, Lacour JP, Paul C, Narbutt J, et al. Trial of Nemolizumab in Moderate-to-Severe Prurigo Nodularis. *N Engl J Med*. 2020;382:706-716.
33. Hashimoto T, Nattkemper LA, Kim HS, Kursewicz CD, Fowler E, Shah SM, et al. Itch intensity in prurigo nodularis is closely related to dermal interleukin-31, oncostatin M, IL-31 receptor alpha and oncostatin M receptor beta. *Exp Dermatol*. 2021;30:804-810.
34. Beuers U, Gerken G, Pusch T. Biliary drainage transiently relieves intractable pruritus in primary biliary cirrhosis. *Hepatology* 2006;44:280-281.
35. Hegade VS, Krawczyk M, Kremer AE, Kuczka J, Gaouar F, Kuiper EM, et al. The safety and efficacy of nasobiliary drainage in the treatment of refractory cholestatic pruritus: a multicentre European study. *Aliment Pharmacol Ther* 2016;43:294-302.
36. de Vries E, Bolier R, Goet J, Parés A, Verbeek J, de Vree M, et al. Fibrates for Itch (FITCH) in fibrosing cholangiopathies: a double-blind, randomized, placebo-controlled trial. *Gastroenterology* 2021;160:734-743.e6.
37. Cai SY, Boyer JL. The role of bile acids in cholestatic liver injury. *Ann Transl Med* 2021;9:737.

38. Corpechot C, Chazouillères O, Rousseau A, Le Gruyer A, Habersetzer F, Mathurin P, et al. A Placebo-controlled trial of bezafibrate in primary biliary cholangitis. *N Engl J Med* 2018;378:2171-2181.
39. Kremer AE, Le Cleac'h A, Lemoine S, Wolf K, De Chaisemartin L, Chollet-Martin S, et al. Antipruritic effect of bezafibrate and serum autotaxin measures in patients with primary biliary cholangitis. *Gut* 2019;68:1902-1903.
40. Bağcı IS, Ruzicka T. IL-31: A new key player in dermatology and beyond. *J Allergy Clin Immunol* 2018;141:858-866.

ACCEPTED

Figure 1. Serum IL-31 and total bile acid levels in healthy volunteers and patients with PBC.

(A) Patients with PBC had higher levels of serum IL-31 than healthy volunteers. (B) Patients with PBC had higher levels of total bile acids than healthy volunteers. (C) Serum IL-31 levels were significantly correlated with total bile acids in healthy volunteers and patients with PBC. Data are presented as means  $\pm$  SE.




Figure 2. Changes in serum IL-31 levels from baseline to Month 3. (A) Significant reduction in serum IL-31 levels by seladelpar treatment for 3 months. (B) Mean % change in serum IL-31 levels from baseline to Month 3. (C) Individual patient IL-31 levels at baseline and Month 3: majority of patients in the seladelpar 10 mg decreased IL-31 levels after 3 months of treatment. BL: Baseline, M3: Month 3. Data are presented as means  $\pm$  SE.




Figure 3. Serum IL-31 levels were correlated with pruritus NRS. (A) A close correlation between IL-31 and pruritus NRS was observed at baseline. (B) Correlation between changes in IL-31 and pruritus NRS was maintained. (C) Decrease of serum IL-31 levels was associated with improvement in pruritus. Data are presented as means  $\pm$  SE.




Table 1. Demographics and baseline characteristics of patients with primary biliary cholangitis.

Mean (SD)	Placebo (n=55)	Seladelpar 5 mg (n=53)	Seladelpar 10 mg (n=53)	Total (N=161)
Female, n (%)	54 (98%)	48 (91%)	50 (94%)	152 (94%)
Age, years	56 (7)	56 (9)	57 (10)	56 (9)
BMI, kg/m <sup>2</sup>	29 (6)	28 (5)	28 (7)	29 (6)
White, n (%)	49 (89%)	50 (94%)	46 (87%)	144 (90%)
Duration of PBC, years	8.9 (6.2)	9.3 (6.2)	8.9 (7.1)	9.0 (6.5)
AMA-positive, n (%)	49 (89%)	49 (92%)	47 (89%)	145 (90%)
UDCA received, n (%)	54 (98%)	50 (94%)	49 (92%)	153 (95%)
UDCA daily dose, mg/kg	15 (2)	15 (4)	14 (3)	15 (3)
Pruritus, NRS	2.7 (2.5)	2.8 (2.6)	2.5 (2.5)	2.7 (2.5)
□□□□□NRS□□ 4	6.1 (1.3)	6.3 (1.5)	6.0 (1.4)	6.1 (1.4)
□ NRS □ 4, n (%)	15 (28%)	15 (29%)	14 (26%)	44 (28%)
ALP (U/L)	282 (105)	281 (126)	263 (96)	275 (109)
ALT (U/L)	41 (20)	46 (24)	42 (20)	43 (21)
AST (U/L)	35 (14)	38 (18)	38 (14)	37 (15)
GGT (U/L)	200 (153)	202 (162)	208 (154)	203 (155)
Total Bilirubin (mg/dL)	0.67 (0.27)	0.70 (0.31)	0.65 (0.28)	0.67 (0.29)



Direct Bilirubin (mg/dL)	0.19 (0.12)	0.20 (0.15)	0.18 (0.12)	0.19 (0.13)
Albumin (g/dL)	4.2 (0.2)	4.1 (0.3)	4.1 (0.3)	4.1 (0.3)
ELF score	9.6 (1.0)	9.7 (0.9)	10.0 (1.0)	9.8 (1.0)
Cirrhosis, n (%)	5 (9%)	6 (11%)	6 (11%)	17 (11%)

---

ACCEPTED

Table 2. Correlations between serum IL-31 levels and bile acids at baseline and change from baseline at Month 3 in patients with primary biliary cholangitis.

Bile acids	Correlation with Serum IL-31: Baseline			Correlation with Serum IL-31: Change from Baseline at Month 3						
	Median (IQR)	All Patients (N=161)		Placebo (n=55)		Seladelpar 5 mg (n=53)		Seladelpar 10 mg (n=53)		
	(□M)	Coefficient (r)	p-value	Coefficient (r)	p-value	Coefficient (r)	p-value	Coefficient (r)	p-value	
Unconjugated	CA	0.04 (0.01, 0.13)	-0.01	0.89	0.00	0.98	0.01	0.96	0.03	0.82
	CDCA	0.14 (0.06, 0.34)	-0.05	0.54	-0.01	0.97	0.15	0.29	-0.01	0.95
	DCA	0.26 (0.13, 0.42)	-0.12	0.12	-0.03	0.81	-0.04	0.79	0.00	0.99
	LCA	0.05 (0.02, 0.10)	-0.11	0.16	-0.04	0.77	-0.08	0.60	0.10	0.46

Taurine- conjugated	TCA	0.18 (0.07, 0.56)	<b>p&lt;0</b> <b>.000</b> <b>1</b>	<b>0.46</b> <b>0.46</b> <b>4</b>	<b>0.00</b> <b>0.00</b> <b>4</b>	-0.20 0.1 6	<b>0.62</b> <b>0.62</b> <b>1</b>	<b>p&lt;0</b> <b>.000</b> <b>1</b>
	TCDC A	0.29 (0.12, 0.68)	<b>p&lt;0</b> <b>.000</b> <b>1</b>	<b>0.29</b> <b>0.29</b> <b>29</b>	<b>0.00</b> <b>0.00</b> <b>29</b>	0.01 0.9 3	<b>0.39</b> <b>0.39</b> <b>3</b>	<b>0.00</b> <b>0.00</b> <b>42</b>
	TDC A	0.09 (0.04, 0.22)	<b>p&lt;0</b> <b>.000</b> <b>1</b>	0.26 <b>0.26</b> <b>62</b>	0.0 <b>0.0</b> <b>62</b>	<b>0.31</b> <b>0.31</b> <b>27</b>	<b>0.29</b> <b>0.29</b> <b>27</b>	<b>0.03</b> <b>0.03</b> <b>5</b>
	TLCA	0.01 (0.01, 0.03)	<b>p&lt;0</b> <b>.000</b> <b>1</b>	0.01 <b>0.01</b> <b>5</b>	0.9 <b>0.9</b> <b>5</b>	0.01 0.9 3	0.25 <b>0.25</b> <b>5</b>	0.08 <b>0.08</b> <b>5</b>
Glycine- conjugated	GCA	0.96 (0.44, 2.47)	<b>p&lt;0</b> <b>.000</b> <b>1</b>	0.26 <b>0.26</b> <b>57</b>	0.0 <b>0.0</b> <b>57</b>	-0.03 0.8 2	<b>0.67</b> <b>0.67</b> <b>2</b>	<b>p&lt;0</b> <b>.000</b> <b>1</b>
	GCD CA	1.54 (0.82, 3.38)	<b>p&lt;0</b> <b>.000</b> <b>1</b>	0.10 <b>0.10</b> <b>6</b>	0.4 <b>0.4</b> <b>6</b>	0.17 0.2 3	<b>0.49</b> <b>0.49</b> <b>3</b>	<b>0.00</b> <b>0.00</b> <b>02</b>
	GDC A	0.64 (0.30, 1.16)	<b>p&lt;0</b> <b>.000</b> <b>1</b>	0.03 <b>0.03</b> <b>3</b>	0.8 <b>0.8</b> <b>3</b>	<b>0.28</b> <b>0.28</b> <b>45</b>	0.27 <b>0.27</b> <b>45</b>	0.05 <b>0.05</b> <b>6</b>

	GLC	0.08							
	A	(0.03, 0.17)	0.12	0.12	0.04	0.76	0.06	0.65	0.23 0.10
Total bile acids	TBA	5.18 (2.92, 10.66)		<b>p&lt;0</b>					<b>p&lt;0</b>
UDCA, Unconjugated	UDCA	1.99 (0.75, 5.06)	-0.07	0.39	0.02	0.87	0.06	0.66	<b>0.63</b> <b>.000</b>
UDCA, Taurine-conjugated	TUDCA	0.27 (0.13, 0.72)		<b>p&lt;0</b>					<b>p&lt;0</b>
UDCA, Glycine-conjugated	GUDCA	7.10 (3.82, 16.15)	<b>0.49</b>	<b>.000</b>	<b>0.44</b>	<b>0.00</b>	0.23	0.10	<b>0.38</b> <b>48</b>
Total bile acids + UDCA	TBA+UDCA	16.76 (9.71, 32.37)		<b>p&lt;0</b>					<b>p&lt;0</b>
	As		<b>0.43</b>	<b>.000</b>	<b>0.33</b>	<b>0.00</b>	0.14	0.34	<b>0.55</b> <b>.000</b>