



Jeffrey, M. R. (2023). *Elementary catastrophes underlying bifurcations of vector fields and PDEs*. (Nonlinearity). arXiv.org.
<https://doi.org/10.48550/arXiv.2310.14818>

Early version, also known as pre-print

License (if available):
CC BY

Link to published version (if available):
[10.48550/arXiv.2310.14818](https://doi.org/10.48550/arXiv.2310.14818)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is a pre-print version of the article (version of record). It first appeared online via ArXiv at <https://doi.org/10.48550/arXiv.2310.14818>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Elementary catastrophes underlying bifurcations of vector fields and PDEs

Mike R. Jeffrey*

April 24, 2023

Abstract

A practical method was proposed recently for finding local bifurcation points in an n -dimensional vector field F by seeking their ‘*underlying catastrophes*’. Here we apply the idea to a partial differential equation as an example of the role that catastrophes can play in reaction diffusion. What are these ‘underlying’ catastrophes? We then show they essentially define a restricted class of ‘solvable’ rather than ‘all classifiable’ singularities, by identifying degenerate zeros of a vector field F without taking into account its vectorial character. As a result they are defined by a minimal set of r analytic conditions that provide a practical means to solve for them, and a huge reduction from the calculations needed to classify a singularity, which we will also enumerate here. In this way, *underlying catastrophes* seem to allow us apply Thom’s *elementary catastrophes* in much broader contexts.

Contents

1	Introduction	2
2	Example: a reaction-diffusion catastrophe	5
3	Underlying catastrophes: the \mathcal{B}-\mathcal{G} conditions	11
4	\mathcal{B}-\mathcal{G} determinants and Boardman symbols	13
4.1	Thom-Boardman singularities	14
4.2	The case $\tau = 1, \dots, 1$, and the \mathcal{B} determinants	16
4.3	Proof of the above	17
4.4	Proof of theorem 4.6	20
4.5	The \mathcal{G} conditions for being ‘full’	22
5	Closing remarks	23
A	Counting minors	24

*Department of Engineering Mathematics, University of Bristol, Ada Lovelace Building, Bristol BS8 1TW, UK, email: mike.jeffrey@bristol.ac.uk

1 Introduction

Catastrophe theory was introduced by René Thom to explain how abrupt changes can arise out of smooth change. The theory was very influential in physics, particularly in optics [19, 29], and despite some overly speculative applications that attracted fair skepticism, its importance to science was never really in doubt, see for example the polemic [28] and its rebuttal [3].

These days catastrophes have been largely absorbed into the broader theory of bifurcations, but here we will argue that a wider application of the original concept is possible, using the idea of *underlying catastrophes* introduced in [12].

Not only do bifurcations become increasingly difficult to classify as we go to systems with more variables and more parameters, but to classify any bifurcation, its location must first be known. Here lies our problem. The known classification of bifurcations (e.g. [24, 2, 15, 10]) involve calculations whose implicitness, and sheer numerousness (as we will show here), make locating such points impossible beyond low order cases. Moreover it is unclear to what extent such classifications can be applied to spatiotemporal problems, like reaction-diffusion. Some authors have been drawn back to catastrophes as a simpler means to understand behaviour in dynamical systems and partial differential equations, hindered by the fact that catastrophes apply only to scalar potentials. Recent examples include the identification of butterfly catastrophes in reaction diffusion problems in [6, 13], a cusp catastrophe in a crowd density problem in [27], and attempts to apply catastrophe theory to specific contexts in liquid crystals [8] or variational PDEs [14]. The importance of nonlinear terms in reaction-diffusion has of course been clear right from the outset of the modern interest in pattern formation [7, 22, 25].

A more direct extension of catastrophes to such problems is made possible by a recent suggestion from [12], that *any* bifurcation actually has an elementary catastrophe *underlying* it. An *underlying catastrophe* was described in [12] as ‘a zero of a vector field encountering a singularity’. We will make this identification precise here. We also give an example of a reaction-diffusion problem where diffusion terms trigger an underlying catastrophe, separating a system’s steady state concentrations into different mixing regimes, the boundaries between which are clearly recognisable as a butterfly catastrophe. We do this without restricting attention to scalar catastrophes at any point.

We define an underlying catastrophe as follows.

Take a smooth vector field $\mathbf{F} : \mathbb{R}^n \times \mathbb{R}^r \rightarrow \mathbb{R}^n$ with components $\mathbf{F} =$

(f_1, f_2, \dots, f_n) . Write $\mathbf{F} = \mathbf{F}(\mathbf{x}, \boldsymbol{\alpha})$ for a variable $\mathbf{x} \in \mathbb{R}^n$ and parameter $\boldsymbol{\alpha} \in \mathbb{R}^r$, with gradient operator $\nabla = \frac{\partial}{\partial \mathbf{x}}$. A point where the vectors $\nabla f_1, \dots, \nabla f_n$, are linearly dependent creates a singularity, but we can typically assume that any $n-1$ of the gradient vectors $\nabla f_1, \dots, \nabla f_n$, are linearly independent. The Jacobian determinant $\mathcal{B}_1 = |\nabla(f_1, f_2, \dots, f_n)|$, then vanishes, and the degeneracy of the point can be characterized by a sequence of determinants $\mathcal{B}_i = |\nabla(\mathcal{B}_{i-1}, f_2, \dots, f_n)|$, letting $\mathcal{B}_0 = f_1$.

The vanishing of all \mathcal{B}_i for $i = 1, \dots, r$ will signal a catastrophe of codimension r . For primary forms

$$\mathbf{F} = \left(f(x_1, \boldsymbol{\alpha}) + \underline{\tau} \cdot \underline{x}, \lambda_2 x_2, \dots, \lambda_n x_n \right)$$

where $f(x_1, \boldsymbol{\alpha}) = x_1^{r+1} + \alpha_r x_1^{r-1} + \dots + \alpha_2 x_1 + \alpha_1$, (1)

with $\underline{\tau} \cdot \underline{x} = \tau_2 x_2 + \dots + \tau_n x_n$, the conditions $\mathcal{B}_i = 0$ reduce to just the one-dimensional conditions $\frac{\partial^r}{\partial x_1^r} f = 0$ that identify elementary catastrophes of the scalar field f .

More completely, the following was proposed in [12].

Definition 1.1. *A vector field $\mathbf{F} : \mathbb{R}^n \times \mathbb{R}^r \rightarrow \mathbb{R}^n$ exhibits an **underlying catastrophe** of codimension r if it has a point $(\mathbf{x}_*, \boldsymbol{\alpha}_*) \in \mathbb{R}^n \times \mathbb{R}^r$ at which $\mathbf{F} = 0$ and*

$$\mathcal{B}_1 = \dots = \mathcal{B}_r = 0. \tag{2}$$

*We say the catastrophe is **full** if the non-degeneracy conditions*

$$\mathcal{G}_{r, k_1 \dots k_{r-1}} \neq 0, \tag{3}$$

hold at $(\mathbf{x}_, \boldsymbol{\alpha}_*)$ for every $k_j \in \{1, \dots, n\}$, $j = 1, \dots, r-1$. Following Thom's elementary catastrophes we call these the *fold* ($r = 1$), *cusp* ($r = 2$), *swallowtail* ($r = 3$), etc.*

The functions $\mathcal{G}_{r, k_1 \dots k_{r-1}}$ are a set of extended determinants whose non-vanishing ensures non-degeneracy of the catastrophe and solvability of the conditions (2). They are more numerous than the \mathcal{B}_i s but it is straightforward to calculate them and just verify that they are nonzero. First, note that in defining the \mathcal{B}_i determinants,

$$\begin{aligned} \mathcal{B}_1 &= |\nabla(f_1, f_2, \dots, f_n)|, \\ \mathcal{B}_2 &= |\nabla(\mathcal{B}_1, f_2, \dots, f_n)|, \\ \mathcal{B}_3 &= |\nabla(\mathcal{B}_2, f_2, \dots, f_n)|, \dots \end{aligned} \tag{4}$$

we have chosen to replace the first component of \mathbf{F} each time, but one could choose to replace one of the other components. Let \mathcal{B}_{2,k_1} be the alternative to \mathcal{B}_2 with \mathcal{B}_1 instead placed in the k_1^{th} component of \mathbf{F} , and so iteratively let each $\mathcal{B}_{i,k_1\dots k_{i-1}}$ be an alternative to \mathcal{B}_i with $\mathcal{B}_{i-1,k_1\dots k_{i-2}}$ placed in the k_{i-1}^{th} component of \mathbf{F} , giving

$$\begin{aligned}\mathcal{B}_{2,k_1} &= \left| \frac{\partial(f_1, \dots, f_{k_1-1}, \mathcal{B}_1, f_{k_1+1}, \dots, f_n)}{\partial(x_1, \dots, x_n)} \right|, \\ \mathcal{B}_{3,k_1k_2} &= \left| \frac{\partial(f_1, \dots, f_{k_2-1}, \mathcal{B}_{2,k_1}, f_{k_2+1}, \dots, f_n)}{\partial(x_1, \dots, x_n)} \right|, \quad \dots\end{aligned}\quad (5)$$

and so on. Hence identifying $\mathcal{B}_{2,1} \equiv \mathcal{B}_2$, $\mathcal{B}_{3,11} \equiv \mathcal{B}_3$, etc. brings us back to the choices (4). The functions \mathcal{G}_{r_k} essentially establish an independence between these functions by evaluating extended determinants over $\mathbb{R}^n \times \mathbb{R}^r$ given by

$$\begin{aligned}\mathcal{G}_1 &= \left| \frac{\partial(f_1, \dots, f_n, \mathcal{B}_1)}{\partial(x_1, \dots, x_n, \alpha_1)} \right|, \\ \mathcal{G}_{2,k_1} &= \left| \frac{\partial(f_1, \dots, f_n, \mathcal{B}_1, \mathcal{B}_{2,k_1})}{\partial(x_1, \dots, x_n, \alpha_1, \alpha_2)} \right|, \\ \mathcal{G}_{3,k_1k_2} &= \left| \frac{\partial(f_1, \dots, f_n, \mathcal{B}_1, \mathcal{B}_{2,k_1}, \mathcal{B}_{3,k_1k_2})}{\partial(x_1, \dots, x_n, \alpha_1, \alpha_2, \alpha_3)} \right|, \quad \dots\end{aligned}\quad (6)$$

These provide the non-degeneracy conditions to be evaluated in (3). Again, although these $\mathcal{G}_{i,k}$ are numerous, one need only check that they are non-zero. For the primary forms (1) they extend the familiar non-degeneracy conditions for catastrophes of a scalar function f .

The key to underlying catastrophes is that the conditions (2) typically give a set of r algebraic equations which, along with $\mathbf{F} = 0$ and provided (3) hold, are solvable in n variables and r parameters. Were we to attempt to find a point in a system where all of the possible $\mathcal{B}_{i,k_1\dots k_{i-1}}$ vanished we would have roughly n^r conditions, requiring a system to have this many parameters to satisfy them in general. The complete classification of singularities and bifurcations contain many more cases, which in general are defined by more conditions still, the number of them growing superfactorially (as factorials of factorials, as we will show) in n and r . The underlying catastrophes reduce our attention, by satisfying (3), to cases with the minimal number of conditions $n + r$, and provides the simple determinant conditions (2) to find them.

We will first give a suggestion of the practical possibilities of these conditions with an example applying underlying catastrophes to a reaction-diffusion equation in section 2. We then give more formal statements and

general expressions for these ‘ $\mathcal{B}\mathcal{G}$ ’ determinants in section 3, recapping from [12]. To help understand how underlying catastrophes relate to established bifurcation theory, we will show that we can locate the conditions $\mathcal{B}_i = 0$ within the Thom-Boardman classification of singularities, treating the function \mathbf{F} as a general mapping rather than specifically a vector field. We outline the Thom-Boardman scheme in section 4, and show how the vast number of calculations involved (which we enumerate in appendix A), reduce to the $\mathcal{B}\mathcal{G}$ determinants of underlying catastrophes for corank 1 singularities. The proof of this result is instructive, so we explore it in some length in section 4.4, ending with some final remarks in section 5.

2 Example: a reaction-diffusion catastrophe

Let us write a reaction-diffusion equation for two species A and B on a spatiotemporal domain (t, x, y) . The usual introduction to Turing patterns starts from equations with a linear reaction term, some

$$\begin{aligned}\frac{\partial}{\partial t}A &= \nabla^2 d_1 A + \mu_1 A + \mu_2 B, \\ \frac{\partial}{\partial t}B &= \nabla^2 d_2 B + \nu_1 A + \nu_2 B,\end{aligned}\tag{7}$$

with reaction coefficients μ_i, ν_i , and diffusion constants d_i . Near a steady state one assumes that wave-like solutions permit us to replace the derivative ∇^2 by a factor $-k^2$.

At the end of his seminal paper [25], Turing noted that the linear reaction rates in (7) were valid for a system just leaving its homogeneous state, but more commonly one would expect to see a system passing from one pattern to another, and the study of such non-linear reaction-diffusion has continued intensively since, but let us revisit the problem using catastrophes. For an example take

$$\begin{aligned}\frac{\partial}{\partial t}A &= -k^2 d_1 A + \mu_1 A + \mu_2 B + \alpha_1 B^2 + \alpha_2 B^3, \\ \frac{\partial}{\partial t}B &= -k^2 d_2 B + \nu_1 A + \nu_2 B + \gamma_1 A^2 + \gamma_2 A^3,\end{aligned}\tag{8}$$

motivated by the form of various equations from Turing’s own planned follow-up work on phyllotaxis [7, 21], similar to the Swift-Hohenberg model [22], and various other models of biological potentials, chemical reactions, and crystal growth, such as [1, 6, 8, 11, 13, 14, 26] for example. We specifically choose nonlinear terms such that the righthand-side of (8) is simple enough to analyze by hand, but is not a gradient field (so that something

beyond standard catastrophe theory is required), and any nonlinear terms (including mixed terms like ‘ AB ’) may be added in general.

It will be convenient to make a linear transformation $(u, v) = (A + \frac{\gamma_1}{3\gamma_2}, B + \frac{\alpha_1}{3\alpha_2})$ to remove the quadratic term. Let $\mu_i = \nu_i = 0$ to simplify further (since we will still be left with a linear term from the diffusion), and let $\alpha_2 = \gamma_2 = -1$, then tidy up constants by defining $\alpha = \frac{1}{3}\alpha_1^2$, $\gamma = \frac{1}{3}\gamma_1^2$, $\beta = \frac{1}{3}\gamma_1 d_1 - \frac{2}{27}\alpha_1^3$, $\delta = \frac{1}{3}\alpha_1 d_2 - \frac{2}{27}\gamma_1^3$, $k_i = k^2 d_i$, leaving us with

$$\begin{aligned}\frac{\partial}{\partial t}u &= -k_1 u - \beta - \alpha v - v^3, \\ \frac{\partial}{\partial t}v &= -k_2 v - \delta - \gamma u - u^3,\end{aligned}\tag{9}$$

for some diffusion constants k_1, k_2 , and reaction constants $\alpha, \beta, \gamma, \delta$.

Our purpose now is to use this to illustrate the finding of underlying catastrophes. We will not explore the reaction diffusion problem thoroughly in any specific context, that is left to more exhaustive future work. For ease of discussion let us say that $u(t, x, y) < 0 < v(t, x, y)$ indicates v (i.e. B) is dominant over u (i.e. A), and vice versa, at some given (t, x, y) .

Let us get some basic intuition for this system’s dynamics, starting with its homogeneous steady states. This is purely for illustration so will not be complete.

To look at the homogeneous system, set the diffusion constants to $k_1 = k_2 = 0$. The righthand side is then divergence-free, equal to $-\nabla \times \phi$ with $\phi = \delta u + \beta v + \frac{1}{2}(\gamma u^2 + \alpha v^2) + \frac{1}{4}(u^4 + v^4)$. At $\alpha = \beta = \gamma = \delta = 0$ the righthand side is $(-v^3, -u^3)$, with a highly degenerate saddlepoint at the origin.

Perturbing any of the coefficients can cause this to degenerate into up to 9 steady states. Diffusion given by $k_1, k_2 > 0$, for instance, stabilizes the origin, creating an attractor where the species A and B are ‘balanced’ with concentrations $u = v = 0$. These are shown with and without diffusion in fig. 1.

For example, if we take $\beta = \delta = 0$ with $\alpha > 0$ and $\gamma > 0$, the concentration steady states are given by $(u_{ij}, v_{ij}) = (i\sqrt{-\gamma}, j\sqrt{-\alpha})$ with $i, j = -1, 0, +1$. The states $(\pm\sqrt{-\gamma}, 0)$ and $(0, \pm\sqrt{-\alpha})$ are centres, therefore surrounded by states of oscillatory concentrations, and these are stabilized to the steady state by small diffusion $k_1, k_2 > 0$. If α and γ have opposing signs, then only $(0, 0)$ is a centre, again stabilized by small diffusion. These are shown with small diffusion in fig. 2.

Clearly there are various other parameter regimes that must be considered, with different numbers and stabilities of the steady states, so for a more complete picture we turn to their catastrophes.




Figure 1: Flow in the (u, v) plane with $\alpha = \beta = \gamma = \delta = 0$. Left: zero diffusion $k_1 = k_2 = 0$. Right: with diffusion $k_1 = k_2 = 2$.




Figure 2: Flow in the (u, v) plane, $k_1 = k_2 = 0.2$, $\beta = 0$, $\delta = -1$, $\alpha = -3$, and left $\gamma = -3$, right $\gamma = 1$. Left: an initially balanced state (initial condition at the origin) would evolve to the state with v dominant. Right: an initially balanced state would remain close to balance, gaining a small u dominance.

First calculate the determinants \mathcal{B}_i from definition 1.1 to the righthand side of (9), i.e. taking

$$\mathbf{F} = -(k_1 u + \beta + \alpha v + v^3, k_2 v + \delta + \gamma u + u^3).$$

The calculations are straightforward, and the first four evaluate as

$$\mathcal{B}_1 = k_1 k_2 - ac, \quad (10)$$

$$\mathcal{B}_2 = 6(k_2 ua - vc^2), \quad (11)$$

$$\mathcal{B}_3 = 6(18k_2 uvc - k_2^2 a - c^3), \quad (12)$$

$$\mathcal{B}_4 = 72k_2(3uc^2 - 2k_2 vc - 9k_2 u^2 v), \quad (13)$$

$$\text{where } a = \alpha + 3v^2, \quad c = \gamma + 3u^2.$$

Where these vanish we will find sets of fold, cusp, swallowtail, and butterfly catastrophes, and one may solve to find their roots easily by computer. Or, more usefully, we can analyze them using the kind of parameterizations very beautifully expounded for the elementary catastrophes in [19]. We do this by applying successive \mathcal{B}_r conditions to express the catastrophe sets in the space of the coefficients $(\alpha, \beta, \gamma, \delta)$, taking the variables (u, v) as parameters.

First parameterize the family of steady states, where (9) vanishes, as

$$\begin{aligned}\beta(u, v) &= -k_1 u - \alpha v - v^3, \\ \delta(u, v) &= -k_2 v - \gamma u - u^3.\end{aligned}\tag{14}$$

One way to think of this is as a 4-dimensional manifold obtained by mapping from $(u, v; \alpha, \beta, \gamma, \delta)$ into $(\alpha, \beta, \gamma, \delta)$.

Pairs of steady states will collide to form folds where \mathcal{B}_1 vanishes, which we can parameterize as

$$\begin{aligned}0 &= \mathcal{B}_1 = k_1 k_2 - (\alpha + 3v^2)(\gamma + 3u^2) \\ \Rightarrow \quad \alpha(u, v) &= \frac{k_1 k_2}{\gamma + 3u^2} - 3v^2.\end{aligned}\tag{15}$$

These are a little easier to visualize, in the sense that they constitute a 3-dimensional manifold in the space of $(\alpha, \beta, \gamma, \delta)$, traced out by the functions (14) and (15) as (u, v) and γ vary.

Different branches of folds will collide at cusps, given by

$$\begin{aligned}0 &= \mathcal{B}_2 \propto \frac{k_1 k_2^2 u}{\gamma + 3u^2} - v(\gamma + 3u^2)^2 \\ \Rightarrow \quad \gamma(u, v) &= (k_1 k_2^2 u / v)^{1/3} - 3u^2,\end{aligned}\tag{16}$$

a 2-dimensional surface in the space of $(\alpha, \beta, \gamma, \delta)$ defined by the functions (14) to (16), parameterized by (u, v) .

These collide to form swallowtails. To find these it is useful to work in new variables $p = v/u$ and $q = uv$, and take p as a parameter. Dropping some multiplicative constants from our calculation of \mathcal{B}_3 , we obtain

$$\begin{aligned}0 &= \mathcal{B}_3 \propto 18k_2 uv (k_1 k_2^2 u / v)^{1/3} - \frac{k_1 k_2^3}{(k_1 k_2^2 u / v)^{1/3}} - k_1 k_2^2 u / v \\ \Rightarrow \quad q(p) &= \frac{1}{18} (k_1 k_2)^{1/3} (k_2^{1/3} p^{2/3} + k_1^{1/3} p^{-2/3}),\end{aligned}\tag{17}$$

where we let

$$u = (q/p)^{1/2}, \quad v = (pq)^{1/2}, \quad p = v/u, \quad q = uv.\tag{18}$$

The set $q(p)$ together with (14) to (16) gives us a curve of swallowtails in the space of $(\alpha, \beta, \gamma, \delta)$.

Finally these collide to form butterflies, given by

$$\begin{aligned} 0 &= \mathcal{B}_4 \propto 3u(\gamma + 3u^2)^2 - 2k_2v(\gamma + 3u^2) - 9k_2u^2v, \\ \Rightarrow 0 &= -5k_1^{2/3}k_2^{1/3}p^{-2/3} + 5k_1^{1/3}k_2^{2/3}p^{2/3} \\ \Rightarrow p &= (k_1/k_2)^{1/4}, \quad q = \frac{1}{9}(k_1k_2)^{1/2}. \end{aligned} \quad (19)$$

Hence butterflies are found at isolated points in the space of $(\alpha, \beta, \gamma, \delta)$ where

$$\begin{aligned} u &= \pm \frac{1}{3}(k_1k_2^3)^{1/8}, \quad \beta = \mp \frac{16}{27}(k_1^3k_2)^{3/8}, \quad \alpha = \frac{2}{3}(k_1^3k_2)^{1/4}, \\ v &= \pm \frac{1}{3}(k_1^3k_2)^{1/8}, \quad \delta = \mp \frac{16}{27}(k_1k_2^3)^{3/8}, \quad \gamma = \frac{2}{3}(k_1k_2^3)^{1/4}. \end{aligned} \quad (20)$$

In fig. 3 we plot the fold surface defined by (14)-(15) in the parameter space (β, δ) for some example values of (α, γ) .




Figure 3: Stability in the (β, δ) plane, $k_1 = k_2 = 1$. Left: $\alpha = \gamma = 1/5$. Right: $\alpha = \gamma = 2/3$. Numbers indicate how many steady states exist in each region, and letters indicate typical behaviour from an initially balanced concentration $u = v = 0$: 's' indicates the system evolves to a unique stable steady state with u or v in slight dominance either side of $\beta = \delta = 1$, 'ss' indicates the system will evolve to one of two stable steady states where either u or v is dominant, 'n' indicates there is a stable steady state that is not reachable from the origin so the concentrations diverge, and no letter indicates no stable steady states so the concentrations diverge.

In fig. 4 we plot in the folds in the diffusion plane (k_1, k_2) for some example values of $(\alpha, \beta, \gamma, \delta)$. The folds form the boundaries between regions that have different numbers of steady states as indicated, leading to different stable configurations for the system as described in the captions.

These are just cursory examples to illustrate the concept. By giving appropriate initial data and boundary conditions for the differential equations,




Figure 4: Stability in the (k_1, k_2) diffusion parameter plane, with labels similar to fig. 3. Left: $\alpha = \gamma = 1/5$, $\beta = \delta = 1$, the steady states lie at low concentrations. Right: $\alpha = \gamma = 2/3$, $\beta = \delta = -16/27$, the steady states lie at high concentrations.

we could go on to simulate the mixing of the two-species in the (x, y) plane at different times, or to study wave propagation through these concentrations. In [1], for example, underlying catastrophes of homogeneous steady states are used to determine the existence of heteroclinic connections that represent ‘wave-pinning’ patterns in cell polarity formation in plants. A number of other works have attempted to relate elementary catastrophes to the behaviours of partial differential equations. In [6], the signature shapes of a swallowtail are seen in bifurcation diagrams of an oxidation reaction, and in [20], butterfly sets are then shown in reaction diffusion by limiting to a scalar one species problem, and a similar treatment places cusps in a crowd density problem in [27]. A nice discussion of the problem of detecting high codimension bifurcation points in PDEs can be found in [14], before scalar catastrophe theory is used to develop numerical methods for the purpose. An attempt is made to relate catastrophes to the phases of liquid crystals in [8] by focussing on a scalar term like the free energy.

Typically in such studies it has been necessary to appeal to scalar functions to investigate the occurrence of catastrophes. The example above takes elements from these studies, and tries to show how catastrophes could be applied to them without reducing to scalar quantities. Hopefully this merely hints at the possible applications, and deeper study will require more detailed and specialised analysis with particular applications in mind.

Finally, but importantly, when deriving the catastrophes above we should have also calculated the non-degeneracy conditions \mathcal{G} from (25) at each stage. Though straightforward, these are more numerous than the \mathcal{B} determinants, and best done with a computer. The relevant conditions are \mathcal{G}_1 for the fold, $\mathcal{G}_{2,1}$ and $\mathcal{G}_{2,2}$ for the cusp, four determinants $\mathcal{G}_{3,ij}$ for the swallowtail and

eight determinants $\mathcal{G}_{4,ijk}$ for the butterfly, where i, j, k take values 1 and 2 (because the dimension of the system is 2).

We will not give all of these calculations, but we find they are indeed non-vanishing for nonzero k_1, k_2 . At the butterfly for example we find

$$\mathcal{G}_{4,ijk} = 103680(-1)^{i+k}(k_1k_2)^4(k_1/k_2)^{\frac{3}{4}(k+2j+3i-9)} \quad (21)$$

which is non-vanishing if and only if $k_1, k_2 \neq 0$, i.e. with diffusion.

The fact that these \mathcal{G} conditions are non-zero means the system is ‘full’ according to definition 3.1, meaning firstly that we can solve the conditions $\mathcal{B}_i = 0$, and moreover that these solutions are unique, i.e. we could not obtain different results by instead using the determinants from (24). Were any of these \mathcal{G} s vanishing, the associated catastrophe would not be ‘full’, and such cases require further case-specific study.

Not being full may imply that the point has some higher degeneracy, for example we indeed find that \mathcal{G}_1 vanishes at the set where the folds collide and become higher order catastrophes, i.e. the cusps, and the $\mathcal{G}_{2,1} = \mathcal{G}_{2,2} = 0$ at the set where the cusps are degenerate and become swallowtails, etc.

Not being full need always not imply an actual degeneracy of the system, however, as they may indicate the system has essentially become lower dimensional, so we just need to adjust the dimension of our calculations. If k_1 and k_2 vanish here then all of the \mathcal{G} conditions will vanish, and we can clearly see that the catastrophe calculations above all break down. What has actually happened in this case is that the righthand side of (9) is now divergence-free, and expressible as the derivative of a scalar potential, to which we could apply elementary catastrophe theory instead.

3 Underlying catastrophes: the \mathcal{B} - \mathcal{G} conditions

The \mathcal{B} - \mathcal{G} determinants from definition 1.1 in section 1 are straightforward to evaluate even for large n and r , so it is worth writing general expressions for any codimension.

Strictly speaking our interest is only local, so we should refer to *germs* rather than functions \mathbf{F} or vector fields, but we are not really adding at all to the rigorous theory of bifurcations or singularities here, only developing practical methods to locate them, so we will introduce such technicalities only as needed.

Recall that $\mathbf{F} : \mathbb{R}^n \times \mathbb{R}^r \rightarrow \mathbb{R}^n$ is taken to be a smooth vector field with components $\mathbf{F} = (f_1, f_2, \dots, f_n)$, and dependence written as $\mathbf{F} = \mathbf{F}(\mathbf{x}, \boldsymbol{\alpha})$ for a variable $\mathbf{x} \in \mathbb{R}^n$ and parameter $\boldsymbol{\alpha} \in \mathbb{R}^r$.

We can give more general formulae for the $\mathcal{B}\mathcal{G}$ determinants (4)-(6) as follows.

Definition 3.1. For $i = 1, 2, \dots$ we first define

$$\mathcal{B}_i = \left| \frac{\partial(\mathcal{B}_{i-1}, f_2, \dots, f_n)}{\partial(x_1, \dots, x_n)} \right|, \quad (22)$$

and for $i = 0$ define $\mathcal{B}_0 \equiv f_1$. To generalize this to allow placing \mathcal{B}_{i_1} in the k^{th} component of $\mathbf{F} = (f_1, \dots, f_n)$, denote the vector \mathbf{F} with its k^{th} row replaced by some scalar V by

$$(f_1, \dots, f_n) \setminus^k V = (f_1, \dots, f_{k-1}, V, f_{k+1}, \dots, f_n), \quad (23)$$

then let

$$\mathcal{B}_{i,K(i-1)} = \left| \frac{\partial(f_1, \dots, f_n) \setminus^{k_{i-1}} \mathcal{B}_{i-1,K(i-2)}}{\partial(x_1, \dots, x_n)} \right|, \quad (24)$$

where $K(i)$ denotes an i -length string $K(i) = k_1 \dots k_i$ of symbols $k_j \in \{1, \dots, n\}$, for $j = 1, \dots, i$. We define the case $i = 1$ as $\mathcal{B}_{1,K(0)} = \mathcal{B}_1$, and note that $\mathcal{B}_{i,1\dots 1} \equiv \mathcal{B}_i$. Finally let

$$\mathcal{G}_{r,K(r-1)} = \left| \frac{\partial(f_1, \dots, f_n, \mathcal{B}_1, \mathcal{B}_{2,K(1)}, \dots, \mathcal{B}_{r,K(r-1)})}{\partial(x_1, \dots, x_n, \alpha_1, \dots, \alpha_r)} \right|, \quad (25)$$

defining the case $r = 1$ as $\mathcal{G}_{1,K(0)} = \mathcal{G}_1 = \left| \frac{\partial(f_1, \dots, f_n, \mathcal{B}_1)}{\partial(x_1, \dots, x_n, \alpha_1)} \right|$.

A crucial assumption was stated before definition 1.1, namely that any set of $n - 1$ of the gradient vectors $\frac{\partial f_i}{\partial \mathbf{x}}$ are linearly independent. A useful way to express this property is with the following.

Definition 3.2. Let the **subrank** of a function $\mathbf{F} = (f_1, \dots, f_n)$ be the least number of dimensions spanned by any $n - 1$ of the gradient vectors $\frac{\partial f_1}{\partial \mathbf{x}}, \dots, \frac{\partial f_n}{\partial \mathbf{x}}$, or using the notation (23),

$$\text{subrank } \mathbf{F} = \min \left(\text{rank} \left[\left(\frac{\partial f_1}{\partial \mathbf{x}}, \dots, \frac{\partial f_n}{\partial \mathbf{x}} \right) \setminus^j \mathbf{0}, \quad j = 1, \dots, n \right] \right). \quad (26)$$

Definition 1.1 then applies to a vector field on \mathbb{R}^n for which

$$\text{subrank } \mathbf{F} = n - 1. \quad (27)$$

The property of being *full* in definition 1.1 guarantees that the conditions (22) are solvable to find $(\mathbf{x}_*, \boldsymbol{\alpha}_*) \in \mathbb{R}^n \times \mathbb{R}^r$, as we discuss in section 4.5.

If (27) is not satisfied then the determinants in (22) are not sufficient to locate a catastrophe, and one must instead solve all $\frac{1-n^r}{1-n}$ conditions $\mathcal{B}_{i,K(i-1)} = 0$, and there is no reason to expect these to be solveable in general unless a system has this many parameters.

Even if (27) is not satisfied, it is sometimes possible to trivially reduce the dimension n of the system until (27) *is* satisfied. We can illustrate this using the primary form of a catastrophe from (1). The vector field in (1) has subrank $n - 1$ provided $k_i \neq 0$ for all $i = 2, \dots, n$, such that at a singularity we have $\frac{\partial f_1}{\partial \mathbf{x}} = (0, k_2, \dots, k_n)$. If the subrank is less than $n - 1$, however, then we can remove redundant dimensions and apply definition 1.1 to the reduced system. For example if $k_i = 0$ for all $i > 2$ in (1), then subrank $\mathbf{F}(0) = 1$ so we cannot use definition 1.1, but if we reduce it to a planar system $\mathbf{F} = (f(x_1, \boldsymbol{\alpha}) + k_2 x_2, \lambda_2 x_2)$ then definition 1.1 applies.

4 \mathcal{B} - \mathcal{G} determinants and Boardman symbols

The idea set out in [12] essentially makes three propositions, namely that for a vector field $\mathbf{F} : \mathbb{R}^n \times \mathbb{R}^r \rightarrow \mathbb{R}^n$ in variables (x_1, \dots, x_n) and parameters $(\alpha_1, \dots, \alpha_r)$:

- (i) Any local bifurcation point of \mathbf{F} , namely where $\mathbf{F} = 0$ and its Jacobian has less than full rank, has an *underlying catastrophe*.
- (ii) An *underlying catastrophe* is a degenerate zero of \mathbf{F} that bifurcates under perturbation into a perturbation-dependent number of zeros. In particular, a corank 1 *underlying catastrophe* of codimension r would break up under perturbation into up to $r + 1$ zeroes of \mathbf{F} .
- (iii) For certain *primary forms* $(f(x_1, \alpha_1, \dots, \alpha_r), x_2, \dots, x_n)$ in terms of a scalar polynomial $f : \mathbb{R} \times \mathbb{R}^r \rightarrow \mathbb{R}$ (eq.(1) in section 1), the conditions defining an underlying catastrophe reduce to the familiar defining conditions of Thom's *elementary catastrophes*.

These properties are the motivation for the term ‘underlying catastrophes’, and we will show here the sense and conditions under which they are indeed true. The term ‘underlying’ comes from (i)-(ii), namely that we are identifying bifurcations only by looking for places where zeroes of \mathbf{F} encounter singularities, without any regard for the vectorial nature of \mathbf{F} , so in a sense we are looking only at the mapping ‘underlying \mathbf{F} ’ (ignoring that it maps to

a tangent space and this carries consequences for local stability). For cases of corank 1, giving us what Arnold classifies as series A_r with “one zero eigenvalue and $(r - 1)$ -fold degeneracy in the nonlinear terms” [2], we will show here that the singularity classification reduces precisely to the conditions $\mathcal{B}_i = 0$. The use of the term ‘catastrophe’ is then due to (iii), namely that these conditions then reduce further to the identifying conditions of the elementary catastrophes for the primary forms (1).

René Thom introduced a general classification of singularities in [23] for mappings $\mathbf{F} : \mathbb{R}^m \rightarrow \mathbb{R}^n$. They were defined through the transversality of certain submanifolds defined by kernels of \mathbf{F} and its derivatives. John Boardman provided a more explicit construction in [4], developed on by Mather and Morin [16, 18], which can be expressed as a sequence of extended Jacobian matrices containing minors of successive derivatives of \mathbf{F} , culminating in a convenient set of *Boardman symbols* for Thom’s singularities.

The Thom-Boardman singularity classification is less well known than perhaps it should be among mathematicians, and after half a century is still far less well utilised in applications than it could be. So let us begin with a slightly informal, methodological introduction to the Thom-Boardman procedure, before showing how the $\mathcal{B}\mathcal{G}$ conditions not only fit into the Thom-Boardman theory, but vastly simplify it for a significant class of singularities. We will restrict our attention to $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ but the extension for $\mathbf{F} : \mathbb{R}^m \rightarrow \mathbb{R}^n$ with $m \neq n$ is straightforward. Where possible we leave the more rigorous statements behind the theory to references.

4.1 Thom-Boardman singularities

The Thom-Boardman classification makes heavy use of *ranks* and *minors* of certain Jacobian matrices. The $k \times k$ minors of a $p \times q$ matrix M are the determinants of all $k \times k$ matrices taken from M by omitting any $p - k$ rows and $q - k$ rows. A $p \times q$ matrix M has rank s if there exists an $s \times s$ minor of M that is non-zero, but every larger minor *is* zero.

Let ∇ denote the derivative operator with respect to $\mathbf{x} = (x_1, \dots, x_n)$. We will be dealing with vectors and matrices of differing sizes below, so for some $\mathbf{F} = (f_1, \dots, f_n)$ and $\mathbf{G} = (g_1, \dots, g_m)$, for any n, m , let us be able to combine them as $(\mathbf{F}, \mathbf{G}) = (f_1, \dots, f_n, g_1, \dots, g_m)$, with Jacobians $\nabla(\mathbf{F}, \mathbf{G}) = (\nabla\mathbf{F}, \nabla\mathbf{G}) = (\nabla f_1, \dots, \nabla f_n, \nabla g_1, \dots, \nabla g_m)$. We refer to each component f_i or g_i as occupying a *row* of (\mathbf{F}, \mathbf{G}) , and similarly each ∇f_i or ∇g_i occupies a *row* of $\nabla(\mathbf{F}, \mathbf{G})$.

To define the *Boardman symbol* of a singularity we can define a sequence of extended Jacobians of \mathbf{F} .

Given $\mathbf{F} = (f_1, \dots, f_n) : \mathbb{R}^n \rightarrow \mathbb{R}^n$, first define the symbol $\Delta^1 \mathbf{F} = (f_1, \dots, f_n, \det \nabla \mathbf{F})$, and also for $i_1 > 1$ define $\Delta^{i_1} \mathbf{F} = \left(\mathbf{F}, m_1^{i_1}, m_2^{i_1}, \dots \right)$, where $m_j^{i_1}$, $j = 1, \dots, N_1$, are the $(n - i_1 + 1) \times (n - i_1 + 1)$ minors of $\nabla \mathbf{F}$.

Next define $\Delta^{i_2} \Delta^{i_1} \mathbf{F} = \left(\Delta^{i_1} \mathbf{F}, m_{2,1}^{i_2}, m_{2,2}^{i_2}, \dots \right)$, where $m_{2,j}^{i_2}$ are the $(n - i_2 + 1) \times (n - i_2 + 1)$ minors of $\nabla(\Delta^{i_1} \mathbf{F})$, and proceed iteratively, next defining $\Delta^{i_3} \Delta^{i_2} \Delta^{i_1} \mathbf{F} = \left(\Delta^{i_2} \Delta^{i_1} \mathbf{F}, m_{3,1}^{i_3}, m_{3,2}^{i_3}, \dots \right)$, where $m_{3,j}^{i_3}$ are the $(n - i_3 + 1) \times (n - i_3 + 1)$ minors of $\nabla(\Delta^{i_2} \Delta^{i_1} \mathbf{F})$, and so on, giving generally

$$\Delta^{i_j} \dots \Delta^{i_1} \mathbf{F} = \left(\Delta^{i_{j-1}} \dots \Delta^{i_1} \mathbf{F}, m_{j,1}^{i_j}, m_{j,2}^{i_j}, \dots \right), \quad (28)$$

where $m_{j,k}^{i_j}$ for $k = 1, \dots, N_j$, are the $(n - i_j + 1) \times (n - i_j + 1)$ minors of $\nabla(\Delta^{i_{j-1}} \dots \Delta^{i_1} \mathbf{F})$. We give a formula for the number N_j of minors in each stage of these calculations in appendix A.

Definition 4.1. *The Boardman symbol of \mathbf{F} at $\mathbf{x} = 0$ is the sequence $\tau = \tau_1, \dots, \tau_r$, such that each $\nabla \Delta^{\tau_{j-1}} \dots \Delta^{\tau_1} \mathbf{F}(0)$ has corank τ_j for $j = 1, \dots, r$, (including that $\nabla \mathbf{F}(0)$ has corank τ_1), and the symbol τ is taken to terminate at r such that $\tau_{r+1} = 0$.*

We call τ_j the j^{th} Boardman symbol.

Note that the symbols τ_1, \dots, τ_r , form a non-increasing sequence $\tau_1 \geq \tau_2 \geq \dots \geq \tau_r$. This means, for instance, that if the Jacobian of \mathbf{F} has corank one, then the Boardman symbol consists of a sequence of 1s only, and in this case we show, in the next section, that the minors $m_{j,k}^{i_j}$ are reducible to precisely the functions $\mathcal{B}_{i,K(i-1)}$ from definition 3.1.

The description above is usually made not in terms of functions $\Lambda^{\tau_r} \dots \Lambda^{\tau_1} \mathbf{F}$, or even their germs (loosely their ‘local representations’), but in terms of the ideals (loosely ‘linear combinations’) generated by the components of those germs, see e.g. [9, 17]. But the procedure is equivalent, and since the ranks of ideals are actually defined by referring back to the ranks of the functions $\nabla \Lambda^{i_r} \dots \Lambda^{i_1} \mathbf{F}$, the above seems both more direct and less unnecessarily technical, at least for the purposes of describing the calculations involved. A more technically complete presentation will be given in forthcoming work [5].

4.2 The case $\tau = 1, \dots, 1$, and the \mathcal{B} determinants

Let us consider more closely what these minors look like if $\nabla \mathbf{F}$ has corank one. In this case the Boardman symbols are just a sequence of symbols $\tau_1 = \dots = \tau_r = 1$, $\tau_{r+1} = 0$, for some $r \geq 1$. These are known as Σ^1 singularities or *Morin singularities* [17, 18].

We will show below that, if we assume a codimension r Morin singularity at some $\mathbf{x} = \mathbf{x}_0$, we can locate it by either:

- (a) solving to find where the requisite minors $m_{j,k}^{\tau_j}$ associated with the Boardman symbols $\tau_1 = \dots = \tau_r = 1$ all vanish at some $\mathbf{x} = \mathbf{x}_0$. The number of these minors grows super-factorially with r .
- (b) solving the conditions $\mathcal{B}_1 = \mathcal{B}_{2,k_1} = \dots = \mathcal{B}_{r,k_1 \dots k_{r-1}} = 0$ for all $k_1, \dots, k_{r-1} \in 1, \dots, n$, at $\mathbf{x} = \mathbf{x}_0$. There are $\sum_{m=0}^{r-1} n^m = \frac{1-n^r}{1-n}$ of these conditions.
- (c) restricting to vector fields with subrank $\mathbf{F} = n - 1$ (definition 3.2), and solving the conditions $\mathcal{B}_1 = \dots = \mathcal{B}_r = 0$ at $\mathbf{x} = \mathbf{x}_0$. There are only r of these conditions, and the \mathcal{G} conditions (definition 1.1) guarantee these have isolated roots.

In (a)-(b) the number of conditions can be expected to exceed the number of available parameters in a typical physical model, so they do not typically give a solveable system of equations. This in contrast to (c) which requires solving r equations in r parameters, so while the \mathcal{B} - \mathcal{G} conditions cannot detect all possible singularities, they restrict attention to those that can be solved for in closed form.

To show this we have the following series of results, which we state, and then give their proofs afterwards. Recall the definitions of the determinants \mathcal{B}_i and $\mathcal{B}_{i,K(i-1)}$ from (22)-(24). The statements “all $K(i-1)$ ” or “all $\mathcal{B}_{i,K(i-1)}$ ” are shorthand for writing that the string $K(r-1) = k_1 \dots k_{r-1}$ or function $\mathcal{B}_{i,k_1 \dots k_{i-1}}$ are taken for all $k_j = 1, \dots, n$ and for $j = 1, \dots, r-1$.

Below we will want the dependence on the variable $\mathbf{x} \in \mathbb{R}^n$ and parameter $\boldsymbol{\alpha} \in \mathbb{R}^r$ to be explicit, so write

$$\mathbf{F} = \mathbf{F}(\mathbf{x}; \boldsymbol{\alpha}), \quad \text{and} \quad \mathcal{B}_i = \mathcal{B}_i(\mathbf{x}; \boldsymbol{\alpha}).$$

The results below assume, for $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$, that

$$\mathbf{F}(0; 0) = 0 \quad \text{and} \quad \text{subrank } \mathbf{F}(0; 0) = n - 1. \quad (29)$$

First we will need to show that if one choice of \mathcal{B}_i vanishes, then all the permutations $\mathcal{B}_{i,K(i-1)}$ vanish. This is given by the following.

Lemma 4.1 (Vanishing of \mathcal{B}_i). *If $\mathcal{B}_1(0;0) = 0$ and $\mathcal{B}_{2,j}(0;0) = 0$ for some j , then $\nabla\mathcal{B}_{2,k}(0;0) = 0$ for all $k = 1, \dots, n$.*

Corollary 4.2. *If $\mathcal{B}_1(0;0) = \mathcal{B}_2(0;0) = \dots = \mathcal{B}_r(0;0) = 0$, then $\mathcal{B}_1(0;0) = \mathcal{B}_{2,K(1)}(0;0) = \dots = \mathcal{B}_{r,K(r-1)}(0;0) = 0$ for all $K(1), \dots, K(r-1)$, hence we can reduce each of the $\mathcal{B}_{i,K(i-1)}$ to one choice \mathcal{B}_i for $i = 1, \dots, r$.*

We will further need to show that the vanishing of \mathcal{B}_i implies vanishing of all of the minors involved in the i^{th} Boardman symbol. To show this we will make use of the following.

Lemma 4.3 (Vanishing minors). *If $\mathcal{B}_1(0;0) = 0$ and all $\mathcal{B}_{r,K(r-1)}(0;0) = 0$ for some $r > 1$, then any of the $n \times n$ minors of any $N \times n$ matrix formed from $N > n$ rows of the form $\nabla\mathcal{B}_{r-1,K(r-2)}(0;0)$ must vanish.*

Corollary 4.4. *If all $\mathcal{B}_1(0;0) = \dots = \mathcal{B}_{r,K(r-1)}(0;0) = 0$, then any of the $n \times n$ minors of any $N \times n$ matrix formed from $N > n$ rows*

$$\nabla F_j(0;0), \nabla\mathcal{B}_1(0;0), \dots, \nabla\mathcal{B}_{r-1,K(r-2)}(0;0) ,$$

must vanish.

Lastly we will need the converse of lemma 4.1, namely to say that $\mathcal{B}_i = 0$ if and only if all $\mathcal{B}_{i,K(i-1)} = 0$. We want to show this in the extended space $\mathbb{R}^n \times \mathbb{R}^r$.

Lemma 4.5 (Uniqueness of \mathcal{B}_i). *The condition $\mathcal{G}_{r,K(r-1)}(0;0) \neq 0$ for all $K(r-1) = k_1 \dots k_{r-1}$, with the assumption (29), implies that $\mathcal{B}_i(0;0) = 0$ if and only if all $\mathcal{B}_{i,K(i-1)}(0;0) = 0$ for $i = 1, \dots, r$.*

These allow us to prove the main result.

Theorem 4.6 (Underlying catastrophes and Thom-Boardman symbols). *An underlying catastrophe of codimension r , given by definition 1.1 with $\mathcal{B}_1(0;0) = \dots = \mathcal{B}_r(0;0) = 0$, is equivalent to a Thom-Boardman singularity with Boardman symbol $\tau = 1, \dots, 1$ (of length r).*

4.3 Proof of the above

Denote the linear subspace spanned by the rows of $\nabla\mathbf{F}(0;0)$ as $\mathcal{T} \subset \mathbb{R}^n$.

First take the trivial case $r = 0$, so there is no Boardman symbol (or the symbol is just 0), in particular the first Boardman symbol is trivially $\tau_1 = 0$.

So the rank of $\nabla\mathbf{F}(0;0)$ is n , i.e. $\mathcal{B}_1(0;0) = \det \nabla\mathbf{F}(0;0) \neq 0$ and there is no singularity at the origin, and hence the subspace \mathcal{T} is n -dimensional.

Then let us start with two simple and generally known results that we will use below.

Lemma 4.7 (Singularity). *If $\mathcal{B}_1(0;0) = 0$ and $\text{subrank}\mathbf{F}(0;0) = n - 1$, the linear subspace spanned by the rows of $\mathbf{F}(0;0)$ is an $(n - 1)$ -dimensional hyperplane $\mathcal{T} \subset \mathbb{R}^n$.*

Proof of lemma 4.7. If $\mathcal{B}_1(0;0) = \det \nabla\mathbf{F}(0;0) = 0$ and $\text{subrank}\mathbf{F}(0;0) = n - 1$, then $\nabla\mathbf{F}(0;0)$ has corank 1 (and hence defines a singularity with first Boardman symbol $\tau_1 = 1$), therefore the subspace \mathcal{T} is $(n - 1)$ -dimensional. \square

Lemma 4.8 (Equivalence of subspaces). *Let $M = \nabla\mathbf{F}$ and $\tilde{M}_j = \nabla(\mathbf{F} \setminus^j g) = (\nabla f_1, \dots, \nabla f_{j-1}, \nabla g, \nabla f_{j+1}, \dots, \nabla f_n)$, where $\mathbf{F} = (f_1, \dots, f_n)$ for smooth functions $f_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$. Let $M(0;0)$ and $\tilde{M}_j(0;0)$ have corank 1 and let $\text{subrank}\mathbf{F}(0;0) = n - 1$. Then the rows of $M(0;0)$ and $\tilde{M}(0;0)$ span the same $(n - 1)$ -dimensional subspace $\mathcal{T} \subset \mathbb{R}^n$.*

Proof of lemma 4.8. Since $M(0;0)$ and $\tilde{M}_j(0;0)$ both have corank 1, their rows span $(n - 1)$ -dimensional subspaces, say $\mathcal{T} \subset \mathbb{R}^n$ and $\tilde{\mathcal{T}}_j \subset \mathbb{R}^n$, respectively. By definition 3.2, $\text{subrank}\mathbf{F}(0;0) = n - 1$ means any $n - 1$ of the gradient vectors $\nabla f_1, \dots, \nabla f_n$, omitting ∇f_j for some $j \in \{1, \dots, n\}$, also span a subspace of exactly $(n - 1)$ dimensions, so let us call this \mathcal{T}'_j . Now the set \mathcal{T}' spanned by any $n - 1$ of the gradient vectors $\nabla f_1, \dots, \nabla f_n$, must be a subset both of the set \mathcal{T} spanned by $M(0;0)$, and the j^{th} set $\tilde{\mathcal{T}}_j$ spanned by $\tilde{M}_j(0;0)$, that is $\mathcal{T}' \subseteq \mathcal{T}$ and $\mathcal{T}' \subseteq \tilde{\mathcal{T}}_j$, but since each of these is $(n - 1)$ -dimensional this implies $\mathcal{T}' = \tilde{\mathcal{T}}_j = \mathcal{T}$ for every j . \square

We use the dimensionality of \mathcal{T} to prove the main lemmas. First is to show that the vanishing of \mathcal{B}_i implies the vanishing of all permutations $\mathcal{B}_{i,K(i-1)}$. We begin with $i = 1$.

Proof of lemma 4.1. Since $\mathcal{B}_{2,k} = \mathbf{F} \setminus^k \mathcal{B}_1$, then if $\mathcal{B}_{2,k}(0;0) = 0$ for some k this implies $\nabla\mathcal{B}_1(0;0)$ spans an $(n - 1)$ -dimensional linear subspace with $\{\nabla f_2, \dots, \nabla f_n\}$, and by lemma 4.8 this subspace must be \mathcal{T} , i.e. the same linear subspace spanned by the rows of $\nabla\mathbf{F}(0;0)$. Since lemma 4.1 also assumes $\mathcal{B}_1(0;0) = 0$, lemma 4.7 applies, hence $\nabla\mathcal{B}_{2,k}(0;0) = 0$ for all $k = 1, \dots, n$. \square

Then extend this to $i > 1$.

Proof of corollary 4.2. If $\mathcal{B}_1(0;0) = \mathcal{B}_2(0;0) = \dots = \mathcal{B}_r(0;0) = 0$, then applying lemma 4.1 at each order $i = 2, \dots, r$, implies that $\mathcal{B}_1(0;0) = \mathcal{B}_{2,K(1)} = \dots = \mathcal{B}_{r,K(r-1)}(0;0) = 0$ for every $K(1) \in [1, n]$, \dots , $K(r-1) \in [1, n]^{r-1}$. Hence at each order we can reduce $\mathcal{B}_{i,K(i-1)}$ to one choice \mathcal{B}_i for $i = 1, \dots, r$. \square

The next step is to relate this to the Boardman symbols by showing that if \mathcal{B}_i vanishes, then so do the numerous minors in the Thom-Boardman calculations. Again we show this first for $i = 1$.

Proof of lemma 4.3. If $\mathcal{B}_{r,K(r-1)}(0;0) = 0$ for all $K(r-1) \in [1, \dots, n]^{r-1}$, then since $\mathcal{B}_{r,k_1 \dots k_{r-2} k_{r-1}} = \det(\nabla \mathbf{F} \setminus^{k_{r-1}} \nabla \mathcal{B}_{r-1,k_1 \dots k_{r-2}})$, this implies by lemma 4.8 that we have $\nabla \mathcal{B}_{r-1,K(r-2)}(0;0) \in \mathcal{T}$ for all $K(r-2) = k_1 \dots k_{r-2} \in [1, n]^{r-2}$. Lemma 4.3 also assumes $\mathcal{B}_1(0;0) = 0$, so lemma 4.7 applies, so any $N \times n$ matrix formed from $N > n$ rows $\nabla \mathcal{B}_{r-1,K(r-2)}(0;0)$ has rank at most $n-1$, and any of its $n \times n$ minors vanish. \square

Then extend this to $i > 1$.

Proof of corollary 4.4. If all $\mathcal{B}_1(0;0) = \dots = \mathcal{B}_{r,K(r-1)}(0;0) = 0$, then all $\nabla \mathcal{B}_1(0;0) = \dots = \nabla \mathcal{B}_{r-1,K(r-2)}(0;0) \in \mathcal{T}$, and moreover lemma 4.7 applies, so any $N \times n$ matrix formed from $N > n$ rows

$$\nabla F_j(0;0), \nabla \mathcal{B}_1(0;0), \dots, \nabla \mathcal{B}_{r-1,K(r-2)}(0;0),$$

has rank at most $n-1$, and any of its $n \times n$ minors vanish. \square

It is slightly longer to prove the converse to lemma 4.1.

Proof of lemma 4.5. Denote the gradient derivative on $\mathbb{R}^n \times \mathbb{R}^r$ as $\square = (\nabla_{\mathbf{x}}, \nabla_{\alpha})$. Say $\mathbf{F}(0;0) = \mathcal{B}_1(0;0) = 0$ and $\det(\square \mathbf{F}(0;0), \square \mathcal{B}_1(0;0)) \neq 0$, so, by the inverse function theorem, the root $(0,0)$ is isolated.

Now say $\mathbf{F}(0;0) = \mathcal{B}_1(0;0) = \dots = \mathcal{B}_{r,1 \dots 1}(0;0) = 0$, then lemma 4.1 implies $\mathcal{B}_{i,K(i-1)}(0;0) = 0$ for all $K(i-1) = k_1 \dots k_{i-1}$. Assume $\mathcal{G}_{r,K(r-1)}(0;0) \neq 0$, i.e. $\det(\square \mathbf{F}(0;0), \square \mathcal{B}_1(0;0), \square \mathcal{B}_{2,k_1}(0;0), \dots, \square \mathcal{B}_{r,k_1 \dots k_{r-1}}(0;0)) \neq 0$ for all k_1, \dots, k_{r-1} , so, by the inverse function theorem, the root $(0,0)$ is isolated in (\mathbf{x}, α) space, and hence these roots must be the same or all k_1, \dots, k_{r-1} , hence $\mathcal{B}_i(0;0) = 0$ if and only if all $\mathcal{B}_{i,K(i-1)}(0;0) = 0$ for $i = 1, \dots, r$, local to the origin. \square

This brings us to the final proof of theorem 4.6, where we find precisely how the conditions $\mathcal{B}_i = 0$ appear (and are repeated many times over) in the procedure to obtain the Boardman symbols. To make this explicit we will proceed one codimension at a time through definition 4.1.

4.4 Proof of theorem 4.6

First take the trivial case $r = 0$, so there is no Boardman symbol (or the symbol is just 0), this means $\tau_1 = 0$, so the rank of $\nabla\mathbf{F}(0;0)$ is n , i.e. $\mathcal{B}_1(0;0) = \det \nabla\mathbf{F}(0;0) \neq 0$ and there is no singularity at the origin.

Now assume the first Boardman symbol is $\tau_1 = 1$. Then $\nabla\mathbf{F}(0;0)$ has corank 1 and defines a singularity, and $\mathcal{B}_1(0;0) = \det \nabla\mathbf{F}(0;0) = 0$. If the second Boardman symbol is $\tau_2 = 0$ so the complete symbol is just $\tau = 1$, and defines a *fold*, then the $(n+1) \times n$ matrix $\nabla\Delta^1\mathbf{F}(0;0) = (\nabla\mathbf{F}(0;0), \nabla\mathcal{B}_1(0;0))$ has rank n , so at least one of its $n \times n$ minors must be nonzero; we will look more closely at those minors in the next step.

Assume instead that $\tau_1 = \tau_2 = 1$. Then $\nabla\mathbf{F}(0;0)$ has corank 1, and moreover $\nabla\Delta^1\mathbf{F}(0;0)$ has corank 1, so all of the $n \times n$ minors of $\nabla\Delta^1\mathbf{F}(0;0)$ are zero. Those minors (with the exception of the minor $\nabla\mathbf{F}(0;0)$ which we already know vanishes) are precisely the functions we define as $\mathcal{B}_{2,k}$ for $k = 1, \dots, n$. Let

$$m_{2,k}^1 = \mathcal{B}_{2,k}(0;0) \quad \text{for } k = 1, \dots, n, \quad \text{and} \quad m_{2,n+1}^1 = \mathcal{B}_1(0;0). \quad (30)$$

Hence if $\tau_1 = \tau_2 = 1$ then $\mathcal{B}_1(0;0) = \mathcal{B}_{2,k}(0;0) = 0$ for all $k = 1, \dots, n$. Conversely, if $\mathcal{B}_1(0;0) = \mathcal{B}_{2,k}(0;0) = 0$, then by (22) we have $\mathcal{B}_{2,k}(0;0) = 0$ for all $k = 1, \dots, n$, implying $\tau_1 = \tau_2 = 1$.

Now if $\tau_3 = 0$ we are done and the singularity is a *cusp*, then the $2(n+1) \times n$ matrix $\nabla\Delta^1\Delta^1\mathbf{F} = \{\nabla\mathbf{F}, \nabla\mathcal{B}_1, \nabla\mathcal{B}_{2,1}, \dots, \nabla\mathcal{B}_{2,n}, \nabla\mathcal{B}_1\}$ must have rank n , implying that at least one of the $n \times n$ minors of $\nabla(\Delta^1\Delta^1\mathbf{F}(0;0))$ is nonzero; again we will look more closely at these in the next step.

Assume instead that $\tau_1 = \tau_2 = \tau_3 = 1$. Now $\nabla\mathbf{F}(0;0)$, $\nabla\Delta^1\mathbf{F}(0;0)$, and $\nabla\Delta^1\Delta^1\mathbf{F}(0;0)$ all have corank 1, so all of the $n \times n$ minors of $\nabla\Delta^1\Delta^1\mathbf{F}(0;0)$ are zero. Recalling

$$\nabla(\Delta^1\Delta^1\mathbf{F}) = (\nabla\mathbf{F}, \nabla\mathcal{B}_1, \nabla\mathcal{B}_{2,1}, \dots, \nabla\mathcal{B}_{2,n}, \nabla\mathcal{B}_1),$$

these minors number $\frac{(2(n+1))!}{n!(n+2)!}$ (the binomial coefficient for choosing n rows from $2(n+1)$). One of these is $\mathcal{B}_1 = \det \nabla\mathbf{F}$, another $2n$ of them are the functions $\mathcal{B}_{2,1}, \dots, \mathcal{B}_{2,n+1}$ repeated twice (because \mathcal{B}_1 is repeated twice in $\nabla\Delta^1\Delta^1\mathbf{F}$). Another n^2 are the determinants of Jacobian matrices in which

some k_2^{th} row of $\nabla \mathbf{F}$ is swapped for one row $\nabla m_{2,k_1}^1$, with $k_1, k_2 = 1, \dots, n$, and these are precisely the functions $\mathcal{B}_{3,k_1 k_2}$, for $k_1, k_2 = 1, \dots, n$. Let

$$m_{3,l(k_1,k_2)}^1 = \mathcal{B}_{3,k_1 k_2}(0;0) \quad \text{for } k_1, k_2 = 1, \dots, n, \\ \text{with } l(k_1, k_2) = k_1 + n(k_2 - 1), \quad (31a)$$

$$m_{3,j}^1 = \mathcal{B}_{2,j}(0;0) \quad \text{for } j = n^2 + 1, \dots, n^2 + n, \quad (31b)$$

$$m_{3,j}^1 = \mathcal{B}_{2,j}(0;0) \quad \text{for } j = n^2 + n + 1, \dots, n^2 + 2n, \quad (31c)$$

$$m_{3,j}^1 = \mathcal{B}_1(0;0) \quad \text{for } j = n^2 + 2n + 1, \dots, n^2 + 2n + 2, \quad (31d)$$

$$m_{3,j}^1 = 0 \quad \text{for } j = n^2 + n + 3, \dots, \frac{(2(n+1))!}{n!(n+2)!}. \quad (31e)$$

We have claimed in the last line that the remaining $m_{3,j}^1$ for $j = n^2 + n + 3, \dots, \frac{(2(n+1))!}{n!(n+2)!}$ vanish. These are the determinants of Jacobian matrices formed from $0 \leq d \leq n - 2$ rows of $\nabla \mathbf{F}$ and $2 \leq d' \leq n$ rows from $(\nabla \mathcal{B}_1, \nabla \mathcal{B}_{2,1}, \dots, \nabla \mathcal{B}_{2,n}, \nabla \mathcal{B}_1)$, but by lemma 4.3 this has rank $n - 1$, so all its $n \times n$ minors vanish. So $\mathcal{B}_{3,k_1 k_2}(0;0) = 0$ for all $k_1, k_2 = 1, \dots, n$ (as well as $\mathcal{B}_1(0;0) = \mathcal{B}_{2,k_1}(0;0) = 0$ for all $k_1 = 1, \dots, n$). Hence if $\tau_1 = \tau_2 = \tau_3 = 1$ then $\mathcal{B}_1(0;0) = \mathcal{B}_{2,k_1}(0;0) = \mathcal{B}_{3,k_1 k_2}(0;0) = 0$ for all $k_1, k_2 = 1, \dots, n$. Conversely, if $\mathcal{B}_1(0;0) = \mathcal{B}_2(0;0) = \mathcal{B}_3(0;0) = 0$, then by corollary 4.2, $\mathcal{B}_{2,k_1}(0;0) = \mathcal{B}_{3,k_1 k_2}(0;0) = 0$ for all $k_1, k_2 = 1, \dots, n$, implying $\tau_1 = \tau_2 = \tau_3 = 1$.

Now if $\tau_4 = 0$ the singularity is a *swallowtail*, and the $(2(n+1) + \chi) \times n$ matrix

$$\nabla \Delta^1 \Delta^1 \Delta^1 \mathbf{F} = \{ \nabla \mathbf{F}, \nabla \mathcal{B}_1, \nabla m_{2,1}^1, \dots, m_{2,n+1}^1, \nabla m_{3,1}^1, \dots, \nabla m_{3,\chi}^1 \}$$

where $\chi = \frac{(2(n+1))!}{n!(n+2)!}$, must have rank n since $\tau_3 = 0$, so at least one of its $n \times n$ minors must be nonzero; as usual, inspection of these is left to the next step.

And so on. At each successive symbol of length r , the vanishing of minors is equivalent to the vanishing of all of the quantities $\mathbf{F}(0;0)$, $\mathcal{B}_1(0;0)$, ... $\mathcal{B}_{r,k_1 \dots k_{r-1}}$. At the next order, assuming $\tau_1 = \tau_2 = \tau_3 = \tau_4 = 1$, for example, the minors include the functions $\mathcal{B}_{4,k_1 k_2 k_3}$, and we can define

$$m_{4,l(k_1,k_2,k_3)}^1 = \mathcal{B}_{4,k_1 k_2 k_3} \quad \text{for } k_1, k_2, k_3 = 1, \dots, n, \\ \text{where } l(k_1, k_2, k_3) = k_1 + n(k_2 - 1) + n^2(k_3 - 1), \quad (32)$$

while the remaining $m_{4,j}^1$ for $j = n^3 + 1, \dots, \frac{(2(n+1)+\chi)!}{n!((2(n+1)+\chi)-n)!}$, consist of the functions \mathcal{B}_1 , \mathcal{B}_{2,k_1} , $\mathcal{B}_{3,k_1 k_2}$, for $k_1, k_2 = 1, \dots, n$, as well as determinants of

Jacobian matrices formed from $0 \leq d \leq n-2$ rows of $\nabla \mathbf{F}$ and $2 \leq d' \leq n$ rows from $(\nabla \mathcal{B}_1, \nabla m_{2,1}^1, \dots, \nabla m_{2,n+1}^1, \nabla m_{3,1}^1, \dots, \nabla m_{3,\chi}^1)$, whose minors vanish by corollary 4.4. At each successive codimension, applying corollary 4.2 shows equivalence between the symbols $\tau_1 = \dots = \tau_r = 0$ and the conditions $\mathcal{B}_1(0;0) = \dots = \mathcal{B}_r(0;0) = 0$.

□

This does not mean that the ‘underlying catastrophes’ defined in [12] are equivalent to the Thom-Boardman singularities, indeed they are not. But they are consistent in that a codimension r underlying catastrophe (where $\mathbf{F}(0;0) = \mathcal{B}_1(0;0) = \dots = \mathcal{B}_r(0;0) = 0$) is a codimension r *Morin* singularity, i.e. the conditions are *sufficient* to define a zero of \mathbf{F} lying at such a singularity of \mathbf{F} viewed as a mapping.

Importantly in definition 1.1 we place a further restriction of ‘fullness’, which ultimately is not *necessary* to define a singularity in the sense of \mathbf{F} as a mapping, but ensures that \mathbf{F} is determined such that the conditions $\mathbf{F}(0;0) = \mathcal{B}_1(0;0) = \dots = \mathcal{B}_r(0;0) = 0$ are solveable. We complete this section by showing how this follows from the $\mathcal{G}_{r,K(r-1)}$ conditions.

4.5 The \mathcal{G} conditions for being ‘full’

The only place we have needed to make reference to the parameter dependence of \mathbf{F} above is in the proof of lemma 4.5. Taking $\mathbf{F} : \mathbb{R}^n \times \mathbb{R}^r \rightarrow \mathbb{R}^n$, denote again the gradient operator as $\square = (\nabla_{\mathbf{x}}, \nabla_{\alpha})$. The conditions $\mathcal{G}_{r,K(r-1)}(0;0) \neq 0$ simply state that the extended Jacobian

$$\mathcal{G}_r = \frac{\partial(\mathbf{F}, \mathcal{B}_1, \dots, \mathcal{B}_r)}{\partial(\mathbf{x}, \alpha_1, \dots, \alpha_r)} \quad (33)$$

has full rank $n + r$ at $(\mathbf{x}, \alpha) = (0,0)$, as do its variants $\mathcal{G}_{r,k_1 \dots k_{r-1}}$ for the alternative permutations of the $\mathcal{B}_{i,k_1 \dots k_{r-1}}$. At this point, the gradient vectors $\square F_1, \dots, \square F_n, \square \mathcal{B}_1, \dots, \square \mathcal{B}_r$, span an $(n+r)$ -dimensional linear subspace $\mathcal{T}_{\mathbf{x},\alpha} \subset \mathbb{R}^n \times \mathbb{R}^r$, which implies by the implicit function theorem that the problem $\mathbf{F} = \mathcal{B}_1 = \dots = \mathcal{B}_r = 0$ is solvable at isolated points $(\mathbf{x}, \alpha) = (\mathbf{x}_*, \alpha_{1*}, \dots, \alpha_{r*})$.

We can illustrate this using the primary form of a codimension r singu-

larity given in (1), namely

$$\mathbf{F} = \left(f(x_1, \boldsymbol{\alpha}) + \underline{\tau} \cdot \underline{x}, \lambda_2 x_2, \dots, \lambda_n x_n \right)$$

where
$$f(x_1) = x_1^{r+1} + \sum_{i=1}^r \alpha_i x_1^{i-1},$$

with $\underline{\tau} \cdot \underline{x} = \tau_2 x_2 + \dots + \tau_n x_n$, and where the τ_i and λ_i are non-zero constants.

Calculating \mathcal{B}_r from (22), and the function \mathcal{G}_r from (33), we have

$$\begin{aligned} \mathcal{B}_r &= (\lambda_2 \dots \lambda_n)^r f^{(r)}(x_1), \\ \mathcal{G}_r &= c_1 (\lambda_2 \dots \lambda_n)^{r+1} f^{(r+1)}(x_1), \end{aligned} \tag{34}$$

and calculating $\mathcal{B}_{r, k_1 \dots k_{r-1}}$ from (24) and $\mathcal{G}_{r, k_1 \dots k_{r-1}}$ from (25), we have

$$\begin{aligned} \mathcal{B}_{r, k_1 \dots k_{r-1}} &= c_2 \lambda_2^{p_2} \lambda_n^{p_n} \tau_2^{q_2} \dots \tau_n^{q_n} \mathcal{B}_r, \\ \mathcal{G}_{r, k_1 \dots k_{r-1}} &= c_3 \lambda_2^{s_2} \dots \lambda_n^{s_n} \tau_2^{t_2} \dots \tau_n^{t_n} \mathcal{G}_r. \end{aligned} \tag{35}$$

The actual values of the various numerical constants c_1, c_2, c_3 and $-r \leq p_i, q_i \leq 0$ and $-r - 1 \leq s_i, t_i \leq 0$ are not important, and depend quite complicatedly on the values of r and the k_i s. The point is that we need all λ_i and τ_i to be non-vanishing for the \mathcal{G} determinants to be non-vanishing, and thereby ensure that the \mathcal{B} determinants are non-trivial and have well-defined roots coinciding with those of the functions $f^{(r)}$. In a following paper [5] we intend to prove that the *underlying catastrophes* of definition 1.1 are in fact transformable, as mappings though not as vector fields, to the functions (1).

5 Closing remarks

The results above justify the methodology laid out in [12] and connect it with the theory of Thom-Boardman singularities. The hope is that these ideas make the application of Thom's ideas to vector fields and spatiotemporal systems much more practical, by providing new ways to find and characterize high codimension bifurcation points.

A number of examples were given in [12] to illustrate the importance of being *full*. A non-trivial example illustrating the importance of satisfying (27), even when (3) holds, is given by $\mathbf{F} = (x_1 + x_2^2, x_1^2 + \alpha_1 x_1 + \alpha_2 + x_2^2 + k x_1)$. If $k_1 = 0$ then $\text{subrank } \mathbf{F}(0; 0) = 0$ and we find $\mathcal{B}_1(0; 0) = \mathcal{B}_{21}(0; 0) = 0 \neq \mathcal{B}_{22}(0; 0)$, violating lemma 4.1, while the fullness conditions $\mathcal{G}_{2,1}(0; 0) \neq 0$

and $\mathcal{G}_{2,2}(0;0) \neq 0$ are still satisfied. If $k_1 \neq 0$ then (27) holds. One then finds that there is a fold at the origin as α_2 passes through zero, while the event that unfolds with α_1 is degenerate (not full), and there is no higher codimension catastrophe since $\mathcal{B}_{21}(0;0), \mathcal{B}_{22}(0;0) \neq 0$.

We have claimed that any bifurcation has an underlying catastrophe, and established the validity of the conjecture with a suitable definition that relates to zeros of vector field encountering certain corank 1 singularities. As already noted in [12], underlying catastrophes are not equivalence classes, and each one may represent several classes of bifurcations with different stability properties. The idea is that the concept of an underlying catastrophe is used to *find* a bifurcation point in a vector field, after which standard bifurcation analysis can be carried out to properly classify it, if required.

Here we have described more the relation of underlying catastrophes to singularities (rather than bifurcations), and this proves to be rather different, in that there appear to be singularities within Thom's classification that cannot be identified with any underlying catastrophe, because they require too many conditions to identify them. This perhaps rules out some singularities as unfindable by the solution of implicit conditions, but what singularities this applies to, and what this means for their role in applications, remains to be studied.

At present this analysis only applies to corank 1 cases. For these, the $\mathcal{B}\text{-}\mathcal{G}$ conditions reduce the number of minors needed to calculate the Boardman symbols, which increases factorially with the codimension r , to just $\mathcal{B}_1, \dots, \mathcal{B}_r$. For what other other singularities or bifurcations is this possible? It is not known yet whether the concept can be extended, but it seems obvious, for example, that corank 2 singularities should be related to *umbilic* catastrophes, though an extension of the $\mathcal{B}\text{-}\mathcal{G}$ conditions for these is not yet known.

Appendix

A Counting minors

The number of minors involved in calculating the Thom-Boardman symbols in section 4.1 grows 'superfactorially', that is, faster than factorially in the codimension r . To see this let us calculate the number, N_j , of new minors that must be calculated to find each successive j^{th} Boardman symbol, following exactly the procedure in section 4.1.

To find the first Boardman symbol τ_1 , one must calculate the $(n - i_1 + 1) \times (n - i_1 + 1)$ minors of $\nabla \mathbf{F} = (\nabla f_1, \dots, \nabla f_n)$, which we call $m_1^{i_1}, \dots, m_{N_1}^{i_1}$. Each minor is formed by choosing $(n - i_1 + 1)$ rows and columns from an $n \times n$ matrix, and so

$$N_1 = \binom{n}{n - i_1 + 1} \cdot \binom{n}{n - i_1 + 1}$$

where $\binom{n}{m}$ denotes the binomial coefficient. In finding the second Boardman symbol τ_2 , one then calculates the $(n - i_2 + 1) \times (n - i_2 + 1)$ minors of $\nabla \Delta^{i_1} \mathbf{F} = (\nabla f_1, \dots, \nabla f_n, m_1^{i_1}, \dots, m_{N_1}^{i_1})$, which we call $m_1^{i_2}, \dots, m_{N_2}^{i_2}$, where each minor is formed by choosing $(n - i_2 + 1)$ rows and columns from an $n \times (n + N_1)$ matrix, and so

$$N_2 = \binom{n}{n - i_2 + 1} \cdot \binom{n + N_1}{n - i_2 + 1}. \quad (36)$$

Continuing iteratively is easy to see that

$$N_3 = \binom{n}{n - i_3 + 1} \cdot \binom{n + N_1 + N_2}{n - i_3 + 1}$$

and so on.

That is, to calculate the Boardman symbol for a codimension r singularity we must evaluate \mathcal{N} minors, the number

$$\mathcal{N}(r) = \sum_{j=1}^r N_j \quad (37)$$

where

$$N_j = \binom{n}{n - i_j + 1} \cdot \binom{\sum_{k=0}^{j-1} N_k}{n - i_j + 1}, \quad N_0 = n. \quad (38)$$

and the brackets denote the binomial coefficient

$$\binom{n}{m} = \frac{n!}{m!(n-m)!}, \quad (39)$$

Table 1 shows $\mathcal{N}(r)$ for the first few codimensions r for corank 1 (when all $i_j = 1$), in systems of different dimension n , and the number of \mathcal{B} determinants this reduces to for an underlying catastrophe.

Thom-Boardman singularities

codim. $r = \rightarrow$ dim. $n = \downarrow$	1	2	3	4	5
1	2	4	8	16	32
2	3	6	21	231	26796
3	4	8	64	41728	12×10^{12}
4	5	10	220	94967015	3×10^{30}

underlying catastrophes

codim. $r = \rightarrow$ dim. $n = \downarrow$	1	2	3	4	5
1	2	3	4	5	6
2	3	4	5	6	7
3	4	5	6	7	8
4	5	6	7	8	9

Table 1: The number problem: these are the number of minors that are calculated to identify a singularity in the standard Thom-Boardman classification (top), versus the number of \mathcal{B} determinants needed to identify an underlying catastrophe (bottom), up to dimension $n = 4$ and codimension $r = 5$.

References

- [1] F. A. Al Saadi, A. R. Champneys, and M. R. Jeffrey. Wave-pinned patterns for cell polarity – a catastrophe theory explanation. *submitted*, 2022.
- [2] V. I. Arnold, V. S. Afrajmovich, Y. S. Il'yashenko, and L. P. Shil'nikov. *Dynamical Systems V: Bifurcation Theory and Catastrophe Theory*. Encyc. of Mathematical Sciences. Springer Verlag, 1994.
- [3] M. V. Berry. In support of catastrophe theory. *Nature*, 270:382–3, 1977.
- [4] J. M. Boardman. Singularities of differentiable maps. *Publications mathématiques de l'I.H.É.S.*, 33:21–57, 1967.
- [5] D. R. J. Chillingworth and M. R. Jeffrey. Finding and unfolding singularities of corank 1. *in draft*, 2023.
- [6] J. Cisternas, K. Rohe, and S. Wehner. Reaction–diffusion fronts and the butterfly set. *Chaos*, 30(113138):1–14, 2020.
- [7] J. H. P. Dawes. After 1952: The later development of Alan Turing's ideas on the mathematics of pattern formation. *Historia Mathematica*, 43(1):49–64, 2016.
- [8] L. Gibelli and S. Turzi. A catastrophe-theoretic approach to tricritical points with application to liquid crystals. *SIAM J. Appl. Math.*, 70(1):63–76, 2009.

- [9] C. G. Gibson. *Singular point of smooth mappings*, volume 25 of *Res. Notes in Math.* Pitman, 1979.
- [10] J. Guckenheimer and P. Holmes. *Nonlinear oscillations, dynamical systems, and bifurcations of vector fields.* Applied Mathematical Sciences 42. Springer, 2002.
- [11] M. P. Holl, A. J. Archer, and U. Thiele. Efficient calculation of phase coexistence and phase diagrams: application to a binary phase-field-crystal model. *J. Phys: Condens. Matter*, 33(115401):1–15, 2021.
- [12] M. R. Jeffrey. Catastrophe conditions for vector fields in \mathbb{R}^n . *J. Phys. A: Math. Theor. Special Issue on Clartons and the Asymptotics of Ideas: the Physics of Michael Berry*, 55(464006):1–25, 2022.
- [13] O. Y. Khachay and P. A. Nosov. On some numerical integration curves for PDE in neighborhood of “butterfly” catastrophe point. *Ural Math. Journal*, 2(2):127–40, 2016.
- [14] L. M. Kreusser, R. I. McLachlan, and C. Offen. Detection of high codimensional bifurcations in variational PDEs. *Nonlinearity*, 33:2335–63, 2020.
- [15] Y. A. Kuznetsov. *Elements of Applied Bifurcation Theory.* Springer, 3rd Ed., 2004.
- [16] J. N. Mather. On Thom-Boardman singularities. In M. M. Peixoto, editor, *Dynamical Systems*, pages 233–248. Academic Press, 1973.
- [17] D. Mond and J. J. Nuño Ballesteros. *Singularities of mappings.* Springer, 2020.
- [18] B. Morin. Calcul jacobien. *Ann. Sci. École Norm. Sup.*, 4(8):1–98, 1975.
- [19] T. Poston and I. N. Stewart. *Catastrophe theory and its applications.* Dover, 1996.
- [20] K. Rohe, J. Cisternas, and S. Wehner. Competing ternary surface reaction $CO + O_2 + H_2$ on Ir(111). *Proc. R. Soc. A*, 476(20190712):1–16, 2023.
- [21] P.T. Saunders, editor. *Collected Works of A.M. Turing: Morphogenesis.* North Holland, Amsterdam, 1992.
- [22] J. Swift and P. C. Hohenberg. Hydrodynamic fluctuations at the convective instability*. *Physical Review A*, 15(1):319–28, 1977.
- [23] R. Thom. Les singularités des application différentiables. *Ann. Inst. Fourier*, 6:43–87, 1955-6.
- [24] R. Thom. *Structural stability and morphogenesis.* Benjamin-Addison Wesley, translation by D.H.Fowler edition, 1975.
- [25] A. M. Turing. The chemical basis of morphogenesis. *Phil. Trans. Roy. Soc. B, Biological Sciences*, 237(641):37–72, 1952.

- [26] A. T. Winfree. Varieties of spiral wave behavior: An experimentalist's approach to the theory of excitable media. *Chaos*, 1(3):303–34, 1991.
- [27] Z. Xiaoping, S. Jiahui, and C. Yuan. Analysis of crowd jam in public buildings based on cusp-catastrophe theory. *Building and Environment*, 45:1755–61, 2010.
- [28] R. S. Zahler and H. J. Sussmann. Claims and accomplishments of applied catastrophe theory. *Nature*, 269:759–63, 1977.
- [29] E. C. Zeeman. *Catastrophe theory: selected papers 1972-1977*. Addison Wesley: Reading Massachusetts, 1977.