



# Advantages of robotic right colectomy over laparoscopic right colectomy beyond the learning curve: a systematic review and meta-analysis

Lelde Lauka, Francesco Brunetti, Nassiba Beghdadi, Margherita Notarnicola, Daniele Sommacale, Nicola de'Angelis

Department of Digestive and Hepatobiliary Surgery, Henri Mondor University Hospital, Créteil, France

*Contributions:* (I) Conception and design: All authors; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: L Lauka, N Beghdadi; (V) Data analysis and interpretation: L Lauka, N Beghdadi, N de'Angelis; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

*Correspondence to:* Nicola de'Angelis, MD, PhD. Department CARE, Unit of Digestive and Hepato-Pancreato-Biliary Surgery, Henri Mondor Hospital, AP-HP, University of Paris Est, UPEC, 51 Avenue du Maréchal de Lattre de Tassigny, 94010, Créteil, France.

Email: nicola.deangelis@aphp.fr.

**Background:** With the widespread application of robotic systems and the increasing number of studies comparing robotic right colectomy (RRC) and laparoscopic right colectomy (LRC), there is a need for an up-to-date systemic review and meta-analysis assessing the advantages of this technique.

**Methods:** The systemic review was performed in Medline, Scopus, Embase, Cochrane Oral Health Group Specialized Register and Google Scholar databases searching for studies comparing RRC and LRC, with no date restriction but limited to English and French literature. Two independent reviewers performed data extraction and qualitative synthesis. Random-effects models were used to summarize the risk ratio (RR) and mean differences (MD) with 95% confidence interval (CI).

**Results:** Twenty-six non-randomized clinical trials (NRCTs) and 1 RCT were included. Overall, 2,314 patients underwent RRC and 17,791 LRC. Operative time was significantly longer for RRC with a MD of 45.36 min (95% CI: 31.75–58.97;  $P < 0.00001$ ). Conversion rate was significantly lower in the RRC group with a RR of 0.47 (95% CI: 0.27–0.81;  $P = 0.007$ ,  $I^2 = 33\%$ ). Also, the number of harvested lymph node was significantly higher in the RRC group than the LRC group, with a MD of 2.03 (95% CI: 0.45–3.61;  $P = 0.01$ ,  $I^2 = 68\%$ ). Estimated blood loss favored RRC, with a MD of -8.68 (95% CI: -17.27 to -0.08;  $P = 0.05$ ,  $I^2 = 46\%$ ). There was no difference in the overall complication rate, mortality, anastomotic leakage, and time to first flatus. However, a significantly shorter hospital stay was associated with RRC, with a MD of -0.60 (95% CI: -1.01 to -0.19;  $P = 0.004$ ,  $I^2 = 64\%$ ). No quantitative analysis could be performed for oncological outcomes. RRC was associated with significantly higher costs (MD 3,185.50 USD; 95% CI: 720.98–5650.02;  $P = 0.01$ ,  $I^2 = 94\%$ ).

**Conclusions:** RRC is a safe procedure that may offer certain advantages over LRC as lower conversion rate, blood loss, hospital stay. However, this should be balanced out with increased operative time and higher costs.

**Keywords:** Laparoscopy; robotic surgery; right colectomy; mini-invasive surgery

Received: 10 February 2020. Accepted: 17 June 2020; Published: 20 October 2020.

doi: 10.21037/ales-20-36

View this article at: <http://dx.doi.org/10.21037/ales-20-36>

## Introduction

Over the last two decades robotic surgery has gained a solid place for colorectal procedures. The first tele-robotic-assisted laparoscopic colorectal surgeries were described in 2002 by Weber *et al.* (1) when he performed a right hemicolectomy and sigmoid colon resection for diverticulitis. Since 2007, studies comparing robotic and laparoscopic approaches have been reporting encouraging data on the safety and feasibility of robotic right colectomy (RRC). However, the application of robotic surgery is still associated with several drawbacks (mainly cost-related) in routine clinical practice. Recently, the introduction of the latest of robotic platforms, such as the da Vinci Xi robotic system, enhanced the interest towards this technique. Indeed, the new generation of surgical robot has been designed to overcome the previous limitations and it is now adapted for multi-quadrant surgeries (2,3). Since 2015, the publication rate on RRC has multiplied rapidly with 2 publications in 2005, 6 in 2010, 20 in 2015, 26 in 2018 and 36 in 2019. We found it essential to reevaluate the existing literature and analyze the reported outcomes by performing an up-to-date meta-analysis comparing RRC and LRC. More than 10 new studies have been published since the latest meta-analyses on the topic (4,5).

Therefore, the aim of the present article was to perform a systematic review and meta-analysis of randomized and non-randomized studies comparing the operative and postoperative outcomes of RRC *vs.* laparoscopic right colectomy (LRC). We present the following article in accordance with the PRISMA reporting checklist (available at <http://dx.doi.org/10.21037/ales-20-36>).

## Methods

A systematic review was performed following the Cochrane collaboration specific protocol (6) and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist (7). Studies comparing operative, postoperative, and oncological outcomes (for cancer patients) between laparoscopic and robotic right hemicolectomies were searched in the following databases without date restrictions: Medline (through PubMed), Scopus, Embase, Cochrane Oral Health Group Specialized Register and Google Scholar. A specific research equation was used for

each database using the following keywords and Mesh Terms: right colectomy, robotic, robotics, laparoscopy, laparoscopic. The most commonly used search equation was: ((right[All Fields] AND (“colectomy”[MeSH Terms] OR “colectomy”[All Fields])) AND (“robotics”[MeSH Terms] OR “robotics”[All Fields] OR “robotic”[All Fields])) AND (“laparoscopy”[MeSH Terms] OR “laparoscopy”[All Fields] OR “laparoscopic”[All Fields]).

Literature review was completed by an extensive search using the “related articles” function in PubMed. Moreover, the reference lists of the eligible records and of pertinent review articles, not included in this study, were double-checked in order to identify potential additional articles. Results were limited to human research, with review articles and case report being excluded. Articles were selected and reviewed if written in English or French. Literature search and selection were performed independently and blindly by two reviewers (LL and NB). Records were removed from the selection if both reviewers excluded the articles at the title/abstract screening levels. Disagreement was resolved with a discussion with a third reviewer (NdeA). Overall, the concordance rate between the two reviewers was 95%.

## Eligibility criteria

According to the PICOS schema, the following criteria were used for literature search and selection. P: population: adult patients who underwent surgical treatment for pathology of right colon (right colectomy); I: intervention: robotic (full or laparoscopy assisted) right colectomy with intracorporeal or extracorporeal anastomosis; C: comparisons: LRC with intracorporeal or extracorporeal anastomosis; O: outcomes: operative variables (operative time, conversion rate, estimated blood loss, number of retrieved lymph nodes), postoperative variables (morbidity, mortality, readmission, reoperation 30 days after surgery, time to first flatus, length of hospital stay), oncological outcomes (3 and 5 years overall survival and disease-free survival), costs (total, surgery related, hospital stay related); S: study design: randomized (RCT) and non-randomized controlled trials (NRCT).

Patients with both oncological and non-oncological pathologies were included. Studies with hybrid procedures involving laparoscopic and robotic procedural steps were not eligible.

### Data extraction

Two independent reviewers (LL and NB) performed the full-text analysis and data extraction by filling in an electronic database. Extracted data included first author name, year of publication, study design and type, number of included patients, study period, demographic variables (age, sex, BMI, ASA score, previous abdominal surgery), anastomosis type (intracorporeal or extracorporeal), operative and postoperative variables, oncological outcomes (see “outcomes” in eligibility criteria) and cost.

### Study quality assessment and risk of bias

Two independent reviewers (LL and NB) carried out the study risk of bias evaluation of the selected articles. RCTs were assessed using the Revised Cochrane risk-of-bias tool for randomized trials (RoB 2) (8). NRCTs were assessed using the Risk of Bias in Non-randomized Studies of Interventions (ROBINS-I) tool (9). Both tools were used as recommended in Cochrane Handbook for Systematic Reviews of Interventions 2nd edition (6).

### Statistical analysis

Data from the included studies were processed for the qualitative and quantitative analyses. For dichotomous outcome data, the risk ratio (RR) and 95% confidence interval (CI) were estimated using the Mantel-Haenszel method. For continuous data, the mean differences (MD) and 95% CI were estimated using inverse variance weighting. Whenever available, outcome measures [mean (SD) and median (interquartile range) or (range) values] were extracted for each surgical approach. If necessary and possible, outcome variables were calculated based on data available in the individual selected studies. In studies in which the mean or SD was not reported, these values were estimated from the median, range or interquartile range (6,10). Heterogeneity was assessed by the  $I^2$  statistic and values of 25%, 50%, and 75% were considered low, moderate, and high, respectively (11). The pooled estimates of the mean differences were calculated using random-effects models to take into account potential interstudy heterogeneity and to adopt a more conservative approach. The pooled effect was considered significant if  $P < 0.05$ . The meta-analysis was performed using RevMan software (version 5.3; Cochrane Collaboration).

## Results

### Literature search and selection

The initial search identified 129 articles that were screened for eligibility based on title and abstract evaluation. Ninety-two articles were rejected and the remaining 37 underwent full-text evaluation. No additional studies were identified through manual search, or cross-check of reference lists.

After full-text evaluation, 27 were selected for qualitative synthesis. Studies selection process and reasons for exclusion are shown in the PRISMA flow chart (Figure 1). Finally, only 25 studies were included in quantitative synthesis.

### Studies characteristics

The selected 27 studies were published between October 2007 and October 2019 with a total number of 20,478 patients included. They included 2,314 patients undergoing RRC and 17,791 patients undergoing LRC between 2002 and 2019. There was only 1 RCT, whereas the remaining 26 studies were NRCTs. All NRCTs had a retrospective design. General characteristics of included studies are reported in Table 1.


The mean age (SD) in the RRC group was 68.43 (6.95) years, with 52% of the patients being males. In the LRC group, the mean age was 68.47 (3.25) years with 51.27% of male patients.

There was no significant difference between groups regarding age (MD -0.24; 95% CI: -2.11, 1.63;  $P=0.80$ ), gender (RR 1.04; 95% CI: 0.97, 1.11;  $P=0.32$ ), previous abdominal surgeries (RR 0.80; 95% CI: 0.59, 1.08;  $P=0.15$ ), BMI (MD 0.21; 95% CI: -0.37, 0.79;  $P=0.48$ ), and ASA score (RR 1.11; 95% CI: 0.97, 1.26;  $P=0.12$ ).

Nine studies (12-20) included patients who underwent robotic and laparoscopic surgeries for right, left or rectal pathologies. Only data from subgroup analysis comparing RRC and LRC were extracted and used for this qualitative synthesis. Three studies (21-23) compared open, laparoscopic and robotic right colectomies. Subgroup analysis of laparoscopic and robotic approach was used for this synthesis.

One study by Trastulli *et al.* (24) described a subgroup of patients that received full RRC, whereas in the remaining studies laparoscopically assisted technique was performed (meaning the use of an assistant laparoscopic port).

Three studies (12,14,21), with the largest patient



**Figure 1** PRISMA Diagram. The flowchart shows the literature search and study selection process according to the PRISMA guidelines.

number (RRC n = 790, LRC n=15,899) were performed as a retrospective data collection from national registries. While there was no information about the type of anastomosis in each group, these studies were selected for further qualitative synthesis. Similarly, three monocentric studies (13,22,25) did not specify the anastomosis type but were included in the analysis. Five studies (26-30) included patients only with intracorporeal anastomosis and six studies (15,18,19,23,31,32) only with extracorporeal; in the rest of the studies both types of anastomosis were performed.

In the study of Trastulli *et al.* (24) patients in LRC group received both—intracorporeal and extracorporeal—anastomosis, only patients with intracorporeal anastomosis

(n=40) were included in the quantitative synthesis.

### Studies quality assessment

The assessment of the included RCT (33) classified the study as having “some concerns” according to RoB 2 tool (8). Three NRCTs were assessed as having moderate risk of bias (18,26,30). Twenty-one NRCTs were classified at serious risk of bias (13-16,19-25,27-29,31,32,34-38), all due to potential risks of confounding as none of these studies matched the laparoscopic group to robotic group before inclusion of the patients. Two NRCTs were classified at a critical risk of bias (12,17); in both studies it was due to critical bias of confounding that was assessed to be


**Table 1** General characteristics of included studies

Authors, year	Study period	Study design	No. of patients (total)	No. of patients (RRC)	No of patients (LRC)	Pathology
Gerbaud <i>et al.</i> , 2019	2013 to 2019	NRCT, Mono	101	42	59	Cancer, adenomas
Merola <i>et al.</i> , 2019	2012 to 2017	NRCT, Multi	188	94	94	Cancer
Yozgatli <i>et al.</i> , 2019	2015 to 2017	NRCT, Multi	96	35	61	Cancer
Blumberg, 2019	2013 to 2018	NRCT, Mono	122	21	101	Cancers, polyps
Park <i>et al.</i> , 2019	2010 to 2011	RCT, Mono	70	35	35	Cancer
Megevand <i>et al.</i> , 2019	2010 to 2015	NRCT, Mono	100	50	50	Cancer
Khorgami <i>et al.</i> , 2019	2012 to 2014	NRCT, Multi	7,685	442	7,243	All
Solaini <i>et al.</i> , 2019	2007 to 2017	NRCT, Multi	389	305	84	Cancer and benign lesions
Spinoglio <i>et al.</i> , 2018	2005 to 2015	NRCT, Mono	200	100	100	Cancer
Haskins <i>et al.</i> , 2018	2012 to 2014	NRCT, Multi	2,494	89	2,405	All
Nolan <i>et al.</i> , 2018	2011 to 2016	NRCT, Mono	106	10	96	All
Ngu <i>et al.</i> , 2018	2015 to 2017	NRCT, Mono	32	16	16	Cancer, sealed perforations
Widmar <i>et al.</i> , 2017	2012 to 2014	NRCT, Mono	282	119	163	Cancer
Lujan <i>et al.</i> , 2018	2009 to 2015	NRCT, Mono	224	89	135	All
Kang <i>et al.</i> , 2016	2007 to 2011	NRCT, Mono	63	20	43	Cancer
Dolejs <i>et al.</i> , 2017	2002 to 2004	NRCT, Multi	6,780	259	6,251	All
Widmar <i>et al.</i> , 2016	2009 to 2014	NRCT, Mono	276	69	207	All
de'Angelis <i>et al.</i> , 2016	2012 to 2015	NRCT, Mono	80	30	50	Cancer
Ferrara <i>et al.</i> , 2016	2008 to 2014	NRCT, Mono	28	13	15	Cancer
Guerrieri <i>et al.</i> , 2015	2013 to 2014	NRCT, Mono	29	18	11	Cancer
Trastulli <i>et al.</i> , 2015	2005 to 2014	NRCT, Multi	238	102	136	Cancer, adenomas
Davis <i>et al.</i> , 2014	2009 to 2011	NRCT, Multi	414	207	207	All
Casillas <i>et al.</i> , 2014	2005 to 2012	NRCT, Mono	162	52	110	Neoplasms
Lujan <i>et al.</i> , 2013	2008 to 2011	NRCT, Mono	47	22	25	All
Deutsch <i>et al.</i> , 2012	2004 to 2009	NRCT, Mono	65	18	47	All
deSouza <i>et al.</i> , 2010	2005 to 2009	NRCT, Mono	175	40	135	All
Rawlings <i>et al.</i> , 2007	2002 to 2005	NRCT, Mono	32	17	15	All

RRC, right robotic colectomy; LRC, laparoscopic right colectomy; NRCT, non-randomized control trial; Mono, monocentric; Multi, multicentric; RCT, randomized control trial.







**Figure 4** Forest plot of total costs. Mean difference (MD) and 95% CI were calculated using random-effects inverse variance weighting. Different size markers indicate study weight. RRC, robotic right colectomy;

of  $-0.16$  (95% CI:  $-0.61, 0.29$ ;  $P=0.49$ ). However, a high heterogeneity was found ( $I^2=95\%$ ).

Data about the length of hospital stay (LOS) was available and analyzed only from 22 studies. The meta-analysis showed a significantly shorter LOS in the RRC group. The MD was  $-0.60$  (95% CI:  $-1.01, -0.19$ ;  $P=0.004$ ,  $I^2=64\%$ ) (Figure 3D).

### Costs

Six studies compared the total cost per patient (19,23,26,27,31,32). de'Angelis *et al.* (32) reported the average costs per group and thus these data were not included in meta-analysis. Merola *et al.* (26) reported the mean cost in euros thus it was converted in American dollars (USD) according to actual conversion rate on February 2020. The USD value from two studies published more than 10 years ago (20,31) was converted to actual USD value as in February 2020. Total costs were significantly higher in the RRC group with a MD of  $3,185.50$  USD (95% CI:  $720.98, 5,650.02$ ;  $P=0.01$ ,  $I^2=94\%$ ) (Figure 4).

No subgroup analysis was performed analyzing surgical and hospitalization cost separately due to the lack of consistency in the calculations and descriptions of these costs in each study.

### Oncological outcomes

Three studies reported 3-year and/or 5-year OS and DFS for operated colon cancer patients (23,29,33). Meta-analyses on these outcomes were not performed due to insufficient statistical data provided in these studies. However, in all three studies oncological outcomes were comparable between the two surgical groups. Park *et al.* (33) reported a  $91.1\%$  (95% CI:  $78.8-100\%$ ) 5-year OS for all tumor stages in the RRC group and  $91.0\%$  (95% CI:  $81.3-100\%$ ) in the LRC group ( $P=0.678$ ). Five years DFS for all tumor stages were  $77.4\%$  (95% CI:  $60.6-92.1\%$ ) for RRC and  $83.6\%$  (95% CI:  $72.1-97.0\%$ ) for LRC ( $P=0.442$ ). In the study by Spinoglio *et al.* (29), the

5-year OS for RRC was  $77\%$  and for LRC  $73\%$  ( $P=0.64$ ). The 5-year DFS for RRC and LRC was  $85\%$  and  $83\%$  respectively ( $P=0.36$ ). No statistical difference was found in the study of Kang *et al.* (23) who reported a 5-year OS for RRC of  $73.1\%$  and for LRC of  $79.2\%$ , whereas the 5-year DFS was  $89.5\%$  and  $84\%$  respectively.

## Discussion

### Operative outcomes and learning curve

The present systemic review and meta-analysis provides the latest and the most comprehensive literature analysis on the comparison between robotic and LRC including 27 studies in the qualitative synthesis and 25 studies in the quantitative one. Overall, 13 studies (48%) (12,13,21,26-30,33,34,36-38) were published between 2018 and 2019, including the only RCT (33), highlighting the rising interest in the outcomes and potential benefits of RRC as compared to conventional laparoscopy.

In the present meta-analysis, we demonstrated significantly better operative outcomes for RRC regarding conversion rate and estimated blood loss, which confirm the safety of the robotic approach. Similar results have been reported in other meta-analyses published in last 5 years (4,5,40); however, this is the first meta-analysis to report a significantly higher number of harvested lymph node in the RRC group. The number of retrieved lymph node during an oncological colon resection has been widely discussed as one of the prognostic factors for oncological results. Recent studies (41,42) have reported that lymph node harvesting has a significant association with survival outcomes for both lymph node negative (stage I and II) and positive (stage III) colon cancers, confirming the lymph node harvesting as one of the quality criteria for oncological resection.

Operative time was significantly longer in RRC group as it has been shown in previous meta-analyses (4,5,40,43).



As suggested by Solaini *et al.* (4) the type of anastomosis (intracorporeal or extracorporeal) could be one of the factors impacting on the operative time. They performed a subgroup analysis on extracorporeal anastomosis and found no significant difference in the operative time between RRC and LRC. This finding suggests that future comparisons of RRC and LRC should be carried out only when the same type of anastomosis is performed, especially when the aim is to evaluate the impact of the surgical technique on operative time. This more consistent evaluation may potentially even demonstrate a benefit of the robotic technique when performing an intracorporeal anastomosis. Another important factor that affects operative time is surgeon's and surgical team's knowledge and experience with the robotic platform. The docking and targeting of the robotic system is most of the time included in the operative time count as these procedures are performed after the skin incision. Surgeon's control and familiarity with the robotic console as well as the scrub nurse's knowledge of the robotic instruments and their replacement also has a great impact on the overall operative time. Three studies (29,32,38) included in this systemic review took into account the possible impact of the learning curve on the operative time; in each of these studies the RRC was performed by a single surgeon. Spinoglio *et al.* (29) and Megevand *et al.* (38) reported a significant decrease in the operative time during the learning curve of the RRC with a mean operative time of 329 min in the earlier RRC procedures and 266 min in latest RRC series ( $P \leq 0.001$ ) (29). de'Angelis *et al.* (32) performed subgroup analyses categorizing RRC and LRC patients in 3 groups: cases 1–10, cases 11–20, cases 21+. They showed a trend toward a significant difference in the 11–20 case series, in favor of RRC with a mean operative time of 193.8 min (SD 24.24) in RRC group and 214.5 min (12.57) in LRC group ( $P=0.07$ ). It could be expected that once RRC will have become a routine procedure in many centers where surgeons and surgical teams have overpassed the learning curve, future studies will report substantially shorter operative time.

There are two studies published in 2017 that analyzed the learning curve for RRC. Parisi *et al.* (44) reported a series of 108 patients undergoing RRC with intracorporeal anastomosis. Based on a cumulative sum (CUSUM) analysis, 3 phases of learning curve were detected: cases 1–44: initial learning phase; cases 45–90: consolidation phase; cases 91–108: mastery phase. Mean operative time in phase 1 was 313.2 min (SD 72.8), in phase 2–285.8 min (62.1) and in phase 3–216.4 min (37.2); a significant difference was found

between both phase 1 and phase 2 and between phase 2 and phase 3 ( $P < 0.0001$ ). Similar results but with shorter learning curve were presented by Raimondi *et al.* (45). In a series of 23 RRC with intracorporeal anastomosis, the CUSUM analysis detected 2 phases of learning curve: cases 1–13: acquisition phase and cases 14–23: consolidation phase.

Mean operative time differed significantly: it was 281.5 min (SD 6.59) in the acquisition phase and 244.3 (8.959) in the consolidation phase ( $P=0.0026$ ).

### *Postoperative and oncological outcomes*

In the present meta-analysis, we did not find any differences regarding 30 days morbidity (total and Grade  $\geq 3$  complications), anastomotic leak rate, mortality, reoperation and readmission rates. Only meta-analysis by Ma *et al.* (5) has reported a decrease in overall complications in RRC group ( $P=0.05$ ). However, when the analysis was performed regarding the postoperative ileus, bleeding and anastomotic leakage rates, no differences were found between RRC and LRC. There exists an important inconsistency of definitions of postoperative complications as well as a wide range of non-surgical complications included in the total morbidity that may explain differences in study results. Although no decrease of morbidity was detected in our quantitative synthesis, we found a significantly shorter LOS in the RRC group as was previously described by Ma *et al.* (5). Two other meta-analyses have reported significantly shorter time to first flatus; our analysis did not show differences between the groups. This could be explained by different study selection for this outcome; we strictly included only studies where postoperative bowel activity was defined as "time to (first) flatus", studies with other descriptions, e.g., first bowel movement or return of bowel activity, were not considered.

To our knowledge, the present systemic review is the first attempt to perform a qualitative and quantitative synthesis comparing oncological outcomes (OS and DFS) between robotically and laparoscopically treated colon cancer patients. Three studies (23,29,33) provided data on 3-year and/or 5-year OS and DFS that were comparable between RRC and LRC groups. Although it was not possible to perform meta-analysis due to a lack of provided statistical variables, to date there exist no evidence of inferiority of RRC and it can be considered as oncologically safe approach as laparoscopy. Moreover, the finding of higher number of harvested lymph node during RRC may translate into improved oncological outcomes observable in

futures studies. Further well-designed studies with larger cohort of patients and sufficient follow-up are necessary to make concrete conclusions regarding this outcome.

### Costs

Similarly to other meta-analyses (4,40), the present quantitative synthesis showed a significant higher costs for RRC as compared with LRC, with a mean difference of more than 3,000 USD per patient case. Increased cost is one of the most important drawbacks for implementing the robotic systems in routine practice, therefore high-quality studies comparing expenses for robotic and laparoscopic approach are crucial. Unfortunately, the existing studies that were analyzed in the present systemic review (20,23,26,31,33), showed very high heterogeneity ( $I^2=95\%$ ) and high risk of bias. One of the most important reasons for heterogeneity is the different variables included in the cost analyses in each study (e.g., use of surgical and medical instruments, operating room time, cost of operating personnel, cost of anesthesia, length of hospital stay, costs of medicaments and/or consultation received after operation, laboratory and radiological exams etc.). It is clear that a cost of RRC can be a burden for its future implementation despite the increasing evidence of the safety and feasibility of this approach. However, upcoming years can make a dramatical change and decrease the cost as new robotic systems are appearing in the market.

### Limitations

The present meta-analysis is based on 24 NRCTs, which were retrospective, and only one RCT.

NRCTs are susceptible for selection, confounding and reporting bias and in general provide low level of evidence (46). Twenty-one of NRCTs were evaluated as having serious risk of bias, mainly due to confounding bias; only two studies (26,30) performed case matching between RRC and LRC before the analysis and one study (18) used propensity score matching.

In some of the outcome analysis, we observed high heterogeneity among the included studies. It can be explained by different sample size, study populations, insufficient or uncomplete data, and interstudy and/or intergroup differences in the surgical technique. Operative, postoperative outcomes and cost can be significantly affected by anastomosis type that varied among the studies or was not specified in large retrospective cohorts where data

were extracted from national databases (14,21).

### Future directions

The main attention is currently focusing on the comparison of different types of anastomosis during RRC. There are four ongoing studies about RRC that are registered in Clinicaltrials.gov medical library; three of them aim to compare outcomes of intracorporeal and extracorporeal anastomosis. Intracorporeal Versus Extracorporeal Anastomosis in Robotic Right Colectomy (INEXA) (ClinicalTrials.gov ID: NCT03130166) is a bicentric Danish RCT with a primary outcome measure: postoperative recovery. Robotic-assisted and Laparoscopic Right Colectomy Study (ANCOR) (ID: NCT03312569) is a multicentric prospective comparison study in USA of both anastomosis types that has set an incisional hernia rate as their first outcome measure.

Minimally Invasive Right Colectomy Anastomosis Study (MIRCAST) (ID: NCT03650517) is a large European observational, prospective, parallel cohorts study to compare both anastomosis type for 30 days postoperative morbidity. Finally, monocentric Danish prospective single-arm study with historical controls - Prospective Study of Extended Robotic Right Hemicolectomy With Complete Mesocolic Excision for Cancer (ID: NCT04190589)—aims to compare oncological quality of robotic versus open procedure.

### New robotic systems for abdominal surgery

Intuitive da Vinci ® robotic system has been dominating the market since its invention in 1994.

The latest generation—the da Vinci Xi—was introduced in 2014 and brought in several technical improvements compared to the pervious robotic platforms. The first single site RRC was described by Morelli *et al.* in 2013 (47). Da Vinci single site technology provides a possibility to perform single incision surgical procedures by adapted curved cannulas that triangulates instruments internally and separate arms externally, 5 lumen ports that can be introduced by 2.5 cm incision and semi rigid instrument. Authors of several case series on single site RRC (48-50) have stated that this approach is safe and feasible.

There is some evidence in the literature of surgical procedures performed by alternative robotic systems. Samalavicius *et al.* (51) has reported a right hemicolectomy performed with Senhance® Surgical System with Digital

Laparoscopy. This is a multi-port system with independent robotic arms that can be moved and docked separately to standard laparoscopic trocars. Surgeon works at a console using 3D-HD glasses and using handles that are based on laparoscopic instrument principles. Instruments used on the patients work similarly to the fundamentals of laparoscopy.

A more elaborate system names “Micro Hand S” has been developed in Central South University in collaboration with Tianjin University (China). It was produced to overcome the financial limitations that are often responsible for slow implementation of existing robotic system in developing countries. It resembles principles of da Vinci system with a surgeon’s console with 3D stereo image screen, two master manipulators, control panel and several pedals and a slave cart with three active slave manipulators (arms). Yi *et al.* (52) reported a detailed description on this technology and a first case series of abdominal procedures, among them— one successfully performed right colectomy. As stated by the authors, studies on larger patient series are crucial to confirm safety and feasibility of this system.

## Acknowledgments

*Funding:* None.

## Footnote

*Provenance and Peer Review:* This article was commissioned by the Guest Editors (Marco Milone and Ugo Elmore) for the series “Right Colectomy 2.0” published in *Annals of Laparoscopic and Endoscopic Surgery*. The article has undergone external peer review.

*Reporting Checklist:* The authors have completed the PRISMA reporting checklist. Available at <http://dx.doi.org/10.21037/ales-20-36>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/ales-20-36>). The series “Right Colectomy 2.0” was commissioned by the editorial office without any funding or sponsorship. The authors have no other conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related

to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Weber PA, Merola S, Wasielewski A, et al. Telerobotic-assisted laparoscopic right and sigmoid colectomies for benign disease. *Dis Colon Rectum* 2002;45:1689-94.
2. Protyniak B, Jorden J, Farmer R. Multi-quadrant robotic colorectal surgery: the da Vinci Xi vs Si comparison. *J Robot Surg* 2018;12:67-74.
3. Jimenez-Rodriguez RM, Quezada-Diaz F, Tchack M, et al. Use of the Xi robotic platform for total abdominal colectomy: a step forward in minimally invasive colorectal surgery. *Surg Endosc* 2019;33:966-71.
4. Solaini L, Bazzocchi F, Cavaliere D, et al. Robotic versus laparoscopic right colectomy: an updated systematic review and meta-analysis. *Surg Endosc* 2018;32:1104-10.
5. Ma S, Chen Y, Chen Y, et al. Short-term outcomes of robotic-assisted right colectomy compared with laparoscopic surgery: A systematic review and meta-analysis. *Asian J Surg* 2019;42:589-98.
6. Higgins JPT, Chandler J, Cumpston M, et al. *Cochrane Handbook for Systematic Reviews of Interventions*. 2nd edition. Chichester (UK): John Wiley & Sons, 2019.
7. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
8. Higgins JPT, Savovi J, Page MJ, et al. Revised Cochrane risk-of-bias tool for randomized trials (RoB 2.0) 2019. Available online: [https://www.unisa.edu.au/contentassets/72bf75606a2b4abcaf7f17404af374ad/rob2-0\\_indiv\\_main\\_guidance.pdf](https://www.unisa.edu.au/contentassets/72bf75606a2b4abcaf7f17404af374ad/rob2-0_indiv_main_guidance.pdf)
9. Sterne JA, Hernan MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:i4919.
10. Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample.

- BMC Med Res Methodol 2005;5:13.
11. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557-60.
  12. Khorngami Z, Li WT, Jackson TN, et al. The cost of robotics: an analysis of the added costs of robotic-assisted versus laparoscopic surgery using the National Inpatient Sample. *Surg Endosc* 2019;33:2217-21.
  13. Nolan HR, Smith BE, Honaker MD. Operative time and length of stay is similar between robotic assisted and laparoscopic colon and rectal resections. *J Robot Surg* 2018;12:659-64.
  14. Dolejs SC, Waters JA, Ceppa EP, et al. Laparoscopic versus robotic colectomy: a national surgical quality improvement project analysis. *Surg Endosc* 2017;31:2387-96.
  15. Ferrara F, Piagnerelli R, Scheiterle M, et al. Laparoscopy Versus Robotic Surgery for Colorectal Cancer: A Single-Center Initial Experience. *Surg Innov* 2016;23:374-80.
  16. Guerrieri M, Campagnacci R, Sperti P, et al. Totally robotic vs 3D laparoscopic colectomy: A single centers preliminary experience. *World J Gastroenterol* 2015;21:13152-9.
  17. Davis BR, Yoo AC, Moore M, et al. Robotic-assisted versus laparoscopic colectomy: cost and clinical outcomes. *JLS* 2014;18:211-24.
  18. Casillas MA, Leichtle SW, Wahl WL, et al. Improved perioperative and short-term outcomes of robotic versus conventional laparoscopic colorectal operations. *Am J Surg* 2014;208:33-40.
  19. Deutsch GB, Sathyanarayana SA, Gunabushanam V, et al. Robotic vs. laparoscopic colorectal surgery: an institutional experience. *Surg Endosc* 2012;26:956-63.
  20. Rawlings AL, Woodland JH, Vegunta RK, et al. Robotic versus laparoscopic colectomy. *Surg Endosc* 2007;21:1701-8.
  21. Haskins IN, Ju T, Skancke M, et al. Right Colon Resection for Colon Cancer: Does Surgical Approach Matter? *J Laparoendosc Adv Surg Tech A* 2018;28:1202-6.
  22. Widmar M, Keskin M, Strombom P et al. Lymph node yield in right colectomy for cancer: a comparison of open, laparoscopic and robotic approaches. *Colorectal Dis* 2017;19:888-94.
  23. Kang J, Park YA, Baik SH, et al. A Comparison of Open, Laparoscopic, and Robotic Surgery in the Treatment of Right-sided Colon Cancer. *Surg Laparosc Endosc Percutan Tech* 2016;26:497-502.
  24. Trastulli S, Coratti A, Guarino S, et al. Robotic right colectomy with intracorporeal anastomosis compared with laparoscopic right colectomy with extracorporeal and intracorporeal anastomosis: a retrospective multicentre study. *Surg Endosc* 2015;29:1512-21.
  25. Widmar M, Keskin M, Beltran P, et al. Incisional hernias after laparoscopic and robotic right colectomy. *Hernia* 2016;20:723-8.
  26. Merola G, Sciuto A, Pirozzi F, et al. Is robotic right colectomy economically sustainable? a multicentre retrospective comparative study and cost analysis. *Surg Endosc* 2020;34:4041-7.
  27. Blumberg D. Robotic colectomy with intracorporeal anastomosis is feasible with no operative conversions during the learning curve for an experienced laparoscopic surgeon developing a robotics program. *J Robot Surg* 2019;13:545-55.
  28. Solaini L, Cavaliere D, Pecchini F, et al. Robotic versus laparoscopic right colectomy with intracorporeal anastomosis: a multicenter comparative analysis on short-term outcomes. *Surg Endosc* 2019;33:1898-902.
  29. Spinoglio G, Bianchi PP, Marano A, et al. Robotic Versus Laparoscopic Right Colectomy with Complete Mesocolic Excision for the Treatment of Colon Cancer: Perioperative Outcomes and 5-Year Survival in a Consecutive Series of 202 Patients. *Ann Surg Oncol* 2018;25:3580-6.
  30. Ngu JC, Ng YY. Robotics confers an advantage in right hemicolectomy with intracorporeal anastomosis when matched against conventional laparoscopy. *J Robot Surg* 2018;12:647-53.
  31. deSouza AL, Prasad LM, Park JJ, et al. Robotic assistance in right hemicolectomy: is there a role? *Dis Colon Rectum* 2010;53:1000-6.
  32. de'Angelis N, Lizzi V, Azoulay D, et al. Robotic Versus Laparoscopic Right Colectomy for Colon Cancer: Analysis of the Initial Simultaneous Learning Curve of a Surgical Fellow. *J Laparoendosc Adv Surg Tech A* 2016;26:882-92.
  33. Park JS, Kang H, Park SY, et al. Long-term oncologic after robotic versus laparoscopic right colectomy: a prospective randomized study. *Surg Endosc* 2019;33:2975-81.
  34. Lujan HJ, Plasencia G, Rivera BX, et al. Advantages of Robotic Right Colectomy With Intracorporeal Anastomosis. *Surg Laparosc Endosc Percutan Tech* 2018;28:36-41.
  35. Lujan HJ, Maciel VH, Romero R, et al. Laparoscopic versus robotic right colectomy: a single surgeon's experience. *J Robot Surg* 2013;7:95-102.
  36. Gerbaud F, Valverde A, Danoussou D, et al. Experience With Transitioning From Laparoscopic to Robotic Right Colectomy. *JLS* 2019;23:e2019.00044.

37. Yozgatli TK, Aytac E, Ozben V, et al. Robotic Complete Mesocolic Excision Versus Conventional Laparoscopic Hemicolectomy for Right-Sided Colon Cancer. *J Laparoendosc Adv Surg Tech A* 2019;29:671-6.
38. Mégevand JL, Amboldi M, Lillo E, et al. Right colectomy: consecutive 100 patients treated with laparoscopic and robotic technique for malignancy. Cumulative experience in a single centre. *Updates Surg* 2019;71:151-6.
39. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205-13.
40. Rondelli F, Balzarotti R, Villa F, et al. Is robot-assisted laparoscopic right colectomy more effective than the conventional laparoscopic procedure? A meta-analysis of short-term outcomes. *Int J Surg* 2015;18:75-82.
41. Foo CC, Ku C, Wei R, et al. How does lymph node yield affect survival outcomes of stage I and II colon cancer? *World J Surg Oncol* 2020;18:22.
42. Lykke J, Rosenberg J, Jess P, et al. Lymph node yield and tumour subsite are associated with survival in stage I–III colon cancer: results from a national cohort study. *World J Surg Oncol* 2019;17:62.
43. Petrucciani N, Sirimarco D, Nigri GR, et al. Robotic right colectomy: A worthwhile procedure? Results of a meta-analysis of trials comparing robotic versus laparoscopic right colectomy. *J Minim Access Surg* 2015;11:22-8.
44. Parisi A, Scrucca L, Desiderio J, et al. Robotic right hemicolectomy: Analysis of 108 consecutive procedures and multidimensional assessment of the learning curve. *Surg Oncol* 2017;26:28-36.
45. Raimondi P, Marchegiani F, Cieri M, et al. Is right colectomy a complete learning procedure for a robotic surgical program? *J Robot Surg* 2018;12:147-55.
46. Song JW, Chung KC. Observational studies: cohort and case-control studies. *PLast Reconstr Surg* 2010;126:2234-42.
47. Morelli L, Guadagni S, Caprili G, et al. Robotic right colectomy using the Da Vinci Single-Site(R) platform: case report. *Int J Med Robot* 2013;9:258-61.
48. Chang TC, Chen YT, Yen MH, et al. Single-Incision Robotic Colectomy: Comparison of Short-Term Outcomes with Multiport Robotic Colectomy. *J Laparoendosc Adv Surg Tech A* 2020;30:183-7.
49. Spinoglio G, Lenti LM, Ravazzoni F, et al. Evaluation of technical feasibility and safety of Single-Site robotic right colectomy: three case reports. *Int J Med Robot* 2015;11:135-40.
50. Juo YY, Agarwal S, Luka S, et al. Single-Incision Robotic Colectomy (SIRC) case series: initial experience at a single center. *Surg Endosc* 2015;29:1976-81.
51. Samalavicius NE, Smolskas E, Deduchovas O, et al. Robotic right colectomy using the new Senhance(R) robotic platform: a three-trocar technique - a video vignette. *Colorectal Dis* 2019;21:1092-3.
52. Yi B, Wang G, Li J, et al. Domestically produced Chinese minimally invasive surgical robot system “Micro Hand S” is applied to clinical surgery preliminarily in China. *Surg Endosc* 2017;31:487-93.

doi: 10.21037/ales-20-36

**Cite this article as:** Lauka L, Brunetti F, Beghdadi N, Notarnicola M, Sommacale D, de’Angelis N. Advantages of robotic right colectomy over laparoscopic right colectomy beyond the learning curve: a systematic review and meta-analysis. *Ann Laparosc Endosc Surg* 2020;5:33.