

Impacts of ambient air quality on acute asthma hospital admissions during the COVID-19 pandemic in Oxford City, UK

Singh, Ajit; Morley, Gabriella L; Coignet, Cécile; Leach, Felix; Pope, Francis D; Neil Thomas, Graham; Stacey, Brian; Bush, Tony; Cole, Stuart; Economides, George; Anderson, Ruth; Abreu, Pedro; Bartington, Suzanne E

DOI:

[10.1136/bmjopen-2022-070704](https://doi.org/10.1136/bmjopen-2022-070704)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Singh, A, Morley, GL, Coignet, C, Leach, F, Pope, FD, Neil Thomas, G, Stacey, B, Bush, T, Cole, S, Economides, G, Anderson, R, Abreu, P & Bartington, SE 2024, 'Impacts of ambient air quality on acute asthma hospital admissions during the COVID-19 pandemic in Oxford City, UK: a time-series study', *BMJ open*, vol. 14, no. 1, e070704. <https://doi.org/10.1136/bmjopen-2022-070704>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.




When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

BMJ Open Impacts of ambient air quality on acute asthma hospital admissions during the COVID-19 pandemic in Oxford City, UK: a time-series study

Ajit Singh ,^{1,2} Gabriella L Morley,³ Cécile Coignet,⁴ Felix Leach ,⁵ Francis D Pope,¹ Graham Neil Thomas,² Brian Stacey,⁶ Tony Bush,^{5,7} Stuart Cole,⁸ George Economides,⁸ Ruth Anderson,⁸ Pedro Abreu,⁹ Suzanne E Bartington ²

To cite: Singh A, Morley GL, Coignet C, *et al*. Impacts of ambient air quality on acute asthma hospital admissions during the COVID-19 pandemic in Oxford City, UK: a time-series study. *BMJ Open* 2024;**14**:e070704. doi:10.1136/bmjopen-2022-070704

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-070704>).

AS and GLM are joint first authors.

Received 09 December 2022
Accepted 14 December 2023



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Suzanne E Bartington;
s.bartington@bham.ac.uk

ABSTRACT

Objectives The study aims to investigate the short-term associations between exposure to ambient air pollution (nitrogen dioxide (NO₂), particulate matter pollution—particles with diameter <2.5 µm (PM_{2.5}) and PM₁₀) and incidence of asthma hospital admissions among adults, in Oxford, UK.

Design Retrospective time-series study.

Setting Oxford City (postcode areas OX1–OX4), UK.

Participants Adult population living within the postcode areas OX1–OX4 in Oxford, UK from 1 January 2015 to 31 December 2021.

Primary and secondary outcome measures Hourly NO₂, PM_{2.5} and PM₁₀ concentrations and meteorological data for the period 1 January 2015 to 31 December 2020 were analysed and used as exposures. We used Poisson linear regression analysis to identify independent associations between air pollutant concentrations and asthma admissions rate among the adult study population, using both single (NO₂, PM_{2.5}, PM₁₀) and multipollutant (NO₂ and PM_{2.5}, NO₂ and PM₁₀) models, where they adjustment for temperature and relative humidity.

Results The overall 5-year average asthma admissions rate was 78 per 100 000 population during the study period. The annual average rate decreased to 46 per 100 000 population during 2020 (incidence rate ratio 0.58, 95% CI 0.42 to 0.81, p<0.001) compared to the prepandemic years (2015–2019). In single-pollutant analysis, we observed a significantly increased risk of asthma admission associated with each 1 µg/m³ increase in monthly concentrations of NO₂ 4% (95% CI 1.009% to 1.072%), PM_{2.5} 3% (95% CI 1.006% to 1.052%) and PM₁₀ 1.8% (95% CI 0.999% to 1.038%). However, in the multipollutant regression model, the effect of each individual pollutant was attenuated.

Conclusions Ambient NO₂ and PM_{2.5} air pollution exposure increased the risk of asthma admissions in this urban setting. Improvements in air quality during COVID-19 lockdown periods may have contributed to a substantially reduced acute asthma disease burden. Large-scale measures to improve air quality have potential to protect vulnerable people living with chronic asthma in urban areas.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This study investigated the association between ambient air pollution exposure and risk of admission to hospital for asthma.
- ⇒ The study includes an at-risk population of >800 000 person-years.
- ⇒ Associations were evaluated using single and multipollutant models.
- ⇒ Causation cannot be inferred from this observational study.
- ⇒ A single city study may limit wider generalisation of findings.

INTRODUCTION AND BACKGROUND

Asthma is a non-communicable chronic disease characterised by airway hyper-responsiveness, reversible airflow obstruction and inflammation.¹ As the most common respiratory condition worldwide, asthma affects more than 300 million people globally including 12% of the population in the UK² with 4.3 million UK adults requiring treatment for the condition.³ In 2019, asthma contributed to 368 years lived in disability (95% CI 238 to 546) and 410 per 100 000 Disability Adjusted Life Years (DALYs) (95% CI 280 to 588)⁴ among the UK population, with the greatest disease burden occurring in areas of high deprivation.³ Asthma can also be fatal and approximately 1200 people a year die from asthma in the UK.⁵ As there is no permanent cure, clinical management is directed towards preventing or reducing the severity of exacerbations and controlling symptoms including wheeze, cough and breathlessness.^{6 7} Primary prevention of asthma morbidity and mortality is therefore of major public health significance.

Asthma has multifactorial aetiology and is likely the result of complex gene and



environmental interactions.⁸ Environmental triggers may exacerbate symptoms so chronic asthma patients are advised to modify lifestyle and behaviours to reduce their exposure as a method for preventing exacerbations, alongside the use of inhaled or oral medication.⁶ In recent years, there has been increasing concern regarding the contribution of poor air quality to disease morbidity and mortality. Almost all of the world's population live in areas which do not meet 2021 WHO Global Air Quality Guidelines.^{9–11} In the UK, pollutants of major health concern include nitrogen dioxide (NO₂) which can cause airway inflammation, and fine particulate matter pollution—particles with diameter <2.5 μm (PM_{2.5}), which can enter the circulation, with long-term exposure to both pollutants linked to cardiorespiratory disease.^{12–15} People with existing chronic respiratory diseases such as asthma or chronic obstructive pulmonary disease are more vulnerable to PM_{2.5} particle and NO₂ exposure¹⁶ thereby exacerbating existing health inequalities. Long-term exposure to fine PM_{2.5} and NO₂ is recognised to be causally associated with increased risk of developing child and adult asthma.^{17,18} Mechanisms linking air pollution exposure to asthma include modification of inflammatory pathways, leading to airway remodelling and enhancement of sensitivity to environmental allergens.¹⁹ A recent meta-analysis of 22 studies undertaken in 12 countries showed a significant association between ambient air pollution exposure (NO₂, PM_{2.5}, CO and O₃) and risk of asthma exacerbations among adults and children. A 5-year time-series study (2009–2013) across primary care settings in London identified that an increase in short-term exposure to NO₂ and PM₁₀ was associated with a significant increase in respiratory consultations, inhaler prescriptions or both.²⁰

Improvements in ambient air quality have also been linked to reduced asthma morbidity. Implementation of emergency public health measures in response to the COVID-19 pandemic led to substantive reductions in industrial and transportation activities from spring 2020 and related changes in air pollution levels.^{21–23} Studies in major cities worldwide indicated lockdown periods were associated with a reduction in major air pollutant (NO₂, PM₁₀ and PM_{2.5}) concentrations.^{24–29} Studies in the UK revealed a similar pattern to global trends in terms of air quality changes,^{22,24,30} most notably NO₂ reductions in urban areas.²¹

The outbreak of COVID-19 led to concerns that this viral pandemic would increase the number of asthma exacerbations, resulting in an influx of emergency department visits and hospitalisations among asthma patients. However, a national analysis of anonymised patient records in Scotland and Wales identified substantial reductions in severe asthma exacerbations leading to hospital admission during COVID-19 lockdown periods, although mortality rates remained unchanged.³¹ A reduction in attendance to primary care for asthma exacerbations was also identified in a large-scale analysis of routine data in England.³² Similar reductions in asthma emergency attendances and admissions among children were

also identified across 15 countries worldwide,³³ suggested to be due to reduced exposure triggers and increased treatment adherence. While the trends described in air pollution are replicated in smaller cities and towns, the majority of existing research has been conducted in major cities^{24,30} with little opportunity to understand the relationship between changes in ambient air quality and asthma admissions among adults in smaller towns and cities in which 45% of the UK population live.³⁴

In this study, a time-series analysis was conducted to investigate the association between ambient air pollution exposure and risk of unplanned (emergency) admissions among adults with asthma living in Oxford, UK.

MATERIALS AND METHODS

This study is part of the Natural Environment Research Council (NERC) funded OxAria Study that aims to understand the impacts of COVID-19 on air and noise quality and associated health impacts in Oxford City (<https://oxaria.org.uk/>).

Study design

This study uses a retrospective time-series ecological design at the city level. The unit of analysis was a monthly count of the hospital admissions for adults, admitted with a primary diagnosis of acute asthma, during the study period of 1 January 2015–31 December 2020. Acute asthma is a serious condition involving progressive worsening of asthma symptoms, including breathlessness, wheeze, cough and chest tightness. Patients with severe or life-threatening acute asthma require immediate hospital treatment.

Setting

Oxford is a small historical city with an area of 46 km² and 68 m above sea level.³⁵ Oxford has a warm and temperate climate with mean annual temperature 10.3°C, mean annual cumulative rainfall 708 mm. The diverse population of this international city is approximately 152 000, of which 79.6% are adults.³⁴ The city has two universities with approximately 34 000 students enrolled. The city is a major centre for employment with a thriving economy, with an estimated 46 000 people usually coming to the city for work, on a daily basis, before the COVID-19 pandemic.³⁶ The population is served by Oxford University Hospitals National Health Service (NHS) Foundation Trust and comes under the jurisdiction of the Oxfordshire Clinical Commissioning Group (OCCG). A recent study by the OxAria team provides a detailed geographical, meteorological and environmental description of the study setting.²²

Study population

The study population is adults aged >18 years living within Oxford City postcode areas OX1–OX4 (see online supplemental figure S1).

Data collection

Health outcome data

Hospital Episode Statistics (HES) data were accessed through OCCG. The study sample was obtained from weekly aggregated HES admissions data to a single tertiary hospital (John Radcliffe Hospital, Oxford) where all patients admitted with a primary diagnosis of asthma, as the outcome of interest, were included. Cases were extracted based on the inclusion criteria of age (>18 years), residential postcode district (OX1–OX4) and Clinical Classifications identifier: ICD-10 (International Classification of Diseases 10th Revision code) coding for acute asthma (ICD-10, J45–J46).³⁷ Exclusion criteria were all cases of wheeze (ICD-10, R06.2), cough (ICD-10, R05) and influenza (ICD-10, J11). Missing values within the HES outcome dataset were minimal (<4%) over the total study duration. Here, missing values where present were replaced by interpolation based on the adjacent values. The mean monthly asthma hospital admissions rate was calculated and used as the unit of analysis.

Air pollution exposure and weather data

Hourly air pollutant concentrations (NO_2 , $\text{PM}_{2.5}$ and PM_{10}) for the period 1 January 2015 to 31 December 2020 were obtained at the Automatic Urban and Rural Network St Ebbe's urban background site (UKA00518),³⁸ located centrally in the OX1–OX4 postcode area (covers 3–5 miles radius). Monthly mean air pollution concentrations were calculated from hourly datasets and analysed for the study period.

Hourly meteorological data including relative humidity (RH) and air temperature (T) from 1 January 2015 to 31 December 2020 were obtained from Oxford Radcliffe Observatory³⁹—an urban background site, which was located in the OX2 postcode area. The proportion of missing hourly data was 16% (NO_2), 20% (PM_{10}) and 24% ($\text{PM}_{2.5}$) across the 6-year study duration. This was not replaced with imputed data given the potential for fluctuations over this time period which may result in an inaccuracy or bias.

COVID-19 lockdown classification

Data were obtained from 1 January 2015 to 31 December 2020. We consider two national lockdown periods in England in 2020: (1) lockdown 1 (23 March 2020–15 June 2020) and (2) lockdown 2 (5 November 2020–2 December 2020).^{22 23}

Statistical analysis

Mean annual air pollutant concentrations (NO_2 , $\text{PM}_{2.5}$, PM_{10}) for the study period (2015–2020) were calculated and compared with WHO guideline levels.¹¹ Correlation between air pollutant concentrations and meteorological variables (air temperature and RH) was explored using Pearson correlation matrices and coefficient.

The incidence rate of asthma admissions was determined for each year and the incidence rate ratio (IRR) compared with 2019 (baseline year), with a comparison

of 2020 admissions rate to a 5-year average by Poisson regression analysis. A Pearson correlation matrix was used to compare asthma admissions and air pollutant concentrations for 2020 and prepandemic years (2015–2019). To compare the trends between asthma admissions and air pollutant concentrations, values normalised by the mean were calculated and a time series was generated.

Poisson linear regression analysis was used to quantify independent associations between monthly air pollutant concentrations and incidence of asthma admissions among the adult study population, using single (NO_2 , $\text{PM}_{2.5}$, PM_{10}) and multipollutant (NO_2 and $\text{PM}_{2.5}$, NO_2 and PM_{10}) models, with adjustment for temperature and RH. All statistical analyses were conducted in STATA V.15 applying 95% CIs and two-tailed p values at <0.05 considered statistically significant.⁴⁰

Patient and public involvement

Public members were involved in the design and conduct of this study including public representation on the study steering committee.

RESULTS

Air quality

Overall mean annual average NO_2 and $\text{PM}_{2.5}$ concentrations for the total study duration exceeded the WHO Global Air Quality guideline average levels of $10 \mu\text{g}/\text{m}^3$ and $5 \mu\text{g}/\text{m}^3$, for all study years, respectively. A reduction in annual average NO_2 , PM_{10} and $\text{PM}_{2.5}$ concentrations (0.74, 0.42 and $0.97 \mu\text{g}/\text{m}^3$ per year, respectively) was observed from 2015 to 2019 which further reduced in 2020 (figure 1A). There was also evidence of seasonal peaks with higher pollutant levels in the winter months




Figure 1 (A) Time series (2015–2020) of monthly mean observed pollutant (NO_2 , PM_{10} and $\text{PM}_{2.5}$) concentrations at urban background site (St Ebbe's) in Oxford, where shaded lines represent the smooth fit at the 95% CI. (B) Box plots of the daily averaged air pollutant concentrations in 2020 vs 2015–2019. NO_2 , nitrogen dioxide; $\text{PM}_{2.5}$, particulate matter pollution—particles with diameter <2.5 μm .

for all three pollutants. The significant reductions in air pollutant concentrations in 2020 compared with 2015–2019 were attributed to COVID-19 national lockdown measures, primarily due to lower traffic emissions (figure 1B).²²

It is observed that the mean NO₂ concentration reduced by 26.7% to 10.7 µg/m³ in 2020 (95% CI 8.7 to 12.7 µg/m³), which was significantly (p<0.005) lower than the 5-year average (2015–2019) concentration of 14.6 µg/m³ (95% CI 12.6 to 16.7 µg/m³). Similarly, the mean PM₁₀ concentrations significantly (p<0.05) reduced by 18.6% in 2020 to 10.8 µg/m³ (95% CI 9.1 to 12.4 µg/m³) compared with 13.2 µg/m³ (95% CI: 11.2 to 15.2 µg/m³) from 2015 to 2019. Relatively, PM_{2.5} levels showed a greater reduction than the other two pollutants, with mean PM_{2.5} concentrations significantly (p<0.005) falling by 33.5% to 6.7 µg/m³ (95% CI 5.3 to 8.1 µg/m³) in 2020 from 10.1 µg/m³ in 2015–2019 (95% CI 8.4 to 11.8 µg/m³). However, a recent study using a machine learning for deweathering and detrending analysis has shown that these reductions were not only due to a reduction in emissions but also influenced by the weather.²²

PM_{2.5} and PM₁₀ levels were strongly positively correlated in both 2015–2019 (0.96) and 2020 (0.94) (online supplemental figure S2). A weakly positive correlation was also observed for NO₂ and PM_{2.5}, higher temperatures were associated with lower levels of air pollution, with a negative correlation between all temperature and all three pollutants although this was weaker in 2020. RH had a weakly positive association throughout the study period.

Asthma admissions

The overall incidence of asthma admissions for the total study period was 7.8 per month per 100 000 population. This varied by season, with the highest admissions rate during winter and spring, and the lowest in the summer months (online supplemental figure S3). We found that there were two notable peaks in admissions of adults with asthma in late 2016 and 2019 that coincided with peaks in PM concentrations (see figure 2). Overall, the asthma




Figure 2 Mean monthly rate of adult asthma admissions in Oxford asthma in 2020 vs a 5-year average. The shaded boxes represent the 95% CI.

Table 1 Annual incidence of adult hospital admissions for acute asthma by year (2015–2020), compared with 2019 as baseline

Year	Annual incidence rate*	IRR	95% CI	P value
2015	62.78	0.83	0.61 to 1.11	>0.05
2016	79.00	1.04	0.79 to 1.38	>0.05
2017	91.59	1.20	0.92 to 1.58	>0.05
2018	84.20	1.12	0.84 to 1.46	>0.05
2019	76.08	Baseline		
2020	45.70	0.60	0.43 to 0.82	<0.01

*Per 100 000 population.
IRR, incidence rate ratio.

admissions rate in 2020 was 42% lower than the 5-year average (IRR 0.58, 95% CI 0.42 to 0.81, p<0.001) (online supplemental table S1). There was no difference between the incidence of admissions in 2015–2018 compared with 2019, while there was a significantly reduced risk of admission in 2020 (IRR 0.6, p<0.01) compared with 2019 (table 1).

Further, figure 2 shows the monthly count of hospital admissions of adults with asthma in 2020 and compares with the monthly average pre-epidemic years (2015–2019). We found that the number of admissions for the whole first pandemic year after March 2020 was lower than the 5-year average. Although, there was a similar pattern in summer 2020 admissions compared with the 5-year average summer, where the admission rate remained lower. In contrast, a higher peak was noted in March 2020 compared with the previous 5 years. We also observe a peak in admissions during September, with admission rates nearly matching the 5-year average. There was a marked decline in admissions during the preceding months in 2020 which was not seen during the pre-pandemic years.

Understating the effect of air quality on asthma admissions in adults in response to COVID-19 measures in Oxford

Monthly mean normalised time series of air pollutant (NO₂, PM₁₀ and PM_{2.5}) concentrations and asthma admissions are presented in figure 3. Overall, variation in asthma admissions among adults mostly followed trends in air pollution over the 6-year study period, where a positive association between air pollutants and asthma admissions was observed in 2015–2019 with a relatively stronger correlation than in 2020 (figure 3 and online supplemental file). However, there was a negative correlation between air temperature and asthma admissions in 2020 and whereas a positive correlation was observed in pre-pandemic years (online supplemental table S2).

Further, to understand the association between air pollutants and asthma admissions among adults,




Figure 3 Normalised time series of monthly asthma admissions with air pollutants during 2015–2020. The shaded lines represent the smooth fit at the 95% CI. NO₂, nitrogen dioxide; PM_{2.5}, particulate matter pollution—particles with diameter <2.5 µm.

single-pollutant and multipollutant models were generated applying Poisson regression analysis (table 2). In the single-pollutant model, which was adjusted for temperature and RH during the study period, showed that the risk of asthma admission increased significantly with increasing concentrations of NO₂ and PM_{2.5}. This was a 4% increase in risk for every 1 µg/m³ increase in mean monthly NO₂ and an approximately 3% increase in risk for every 1 µg/m³ increase in mean monthly PM_{2.5}. For every 1 µg/m³ increase in mean monthly PM₁₀, there was also a 1.8% increase in the risk of asthma admission, however, this failed to reach significance despite narrow CIs.

Within the multipollutant model, the effect of each individual pollutant was non-significant. A weak correlation between pollutants makes it difficult for the model to separate the effects of correlated pollutants. Overall, in the multipollutants model, we identified a non-significant 2.6% increase in risk of asthma admission per month predicted for every 1 µg/m³ increase in NO₂ and a non-significant 1.7% increase in risk of admission for every 1 µg/m³ increase PM_{2.5}, which approximates the risk for PM₁₀.

DISCUSSION

This study examines for the first time the effects of ambient air quality on asthma-related hospital admissions in adults in Oxford city (UK), including impacts of the COVID-19 pandemic. A positive association was observed between risk of emergency asthma admission and air pollutant levels in Oxford, with hospital admission risk significantly increased even with modest increases in mean monthly NO₂ and PM_{2.5} concentrations in single-pollutant models. We also found a 42% reduction in the risk of acute asthma admission among adults living in Oxford in 2020 compared with the 5-year average, suggesting air quality improvements during COVID-19 lockdown periods may have contributed to a reduction in severe asthma exacerbations. However, this observed association is in the context of a general reduction in emergency admissions for all causes during the COVID-19 period.

The findings of this work, supported by recent studies, found that pollution levels—notably NO₂ concentrations—decreased significantly in 2020 compared with prepandemic years^{22 23 41 42} due to changes in anthropogenic activities. This pattern needs to consider in the context of already declining levels of air pollution in Oxford and exceptional weather in 2020, including a warmer and windier early part of the year, coinciding with the first national lockdown.^{22 41} Recent studies by the OxAria team highlighted a clear reduction in traffic volume of 69% and 38% during the first and second lockdowns, respectively, in Oxford, where NO₂ emissions from buses and cars reduced by 56% and 77%, respectively.^{22 23} While we did not attempt to explore contributions of specific emissions sources to changes in asthma admissions, the statistically significant effect of NO₂ and PM_{2.5} in the single-pollutant model was not replicated in the multipollutant model, which indicates that changes in emissions sources may be relevant for determining overall asthma exacerbation risk in this context.

The findings presented in our study regarding the consistently positive association between NO₂ and PM_{2.5} and asthma exacerbations rates are consistent with the existing evidence.^{43–52} The identified lockdown impact is also broadly consistent with a large-scale analysis of emergency admissions in Scotland and Wales which showed

Table 2 Incidence rate ratio (IRR) for adult asthma admissions of in response to changes in air pollutant concentrations using the Poisson model

Pollutants	Single-pollutant model			Multipollutant model		
	IRR	95% CI	P value	IRR	95% CI	P value
NO ₂	1.040	1.009 to 1.072	0.012	1.026	0.987 to 1.067	0.188
PM _{2.5}	1.029	1.006 to 1.052	0.013	1.017	0.988 to 1.045	0.254
PM ₁₀	1.018	0.999 to 1.038	0.07	NA	NA	NA
Temperature	0.98	0.950 to 1.020	0.340	0.99	0.960 to 1.030	0.620
Relative humidity	1.01	0.990 to 1.020	0.466	1.00	0.990 to 1.020	0.826

NA, not available; NO₂, nitrogen dioxide; PM_{2.5}, particulate matter pollution—particles with diameter <2.5 µm.



a 36% reduction in asthma-related hospitalisations for all ages population during the first 18 weeks of 2020,³¹ however, the fixed-effect meta-analysis model limited wider generalisability. Davies *et al* proposed multiple possible theories for their observations, including reductions in NO₂ and PM_{2.5} that were a national trend, however, unfortunately they were unable to substantiate this proposition as they did not collect concurrent air quality data. Further there exist differences in healthcare access and COVID-19 emergency public health regulation between the devolved nations and England. The first pandemic year, 2020, was also exceptional for the low rates of asthma admissions worldwide,⁵³ as previously reported in the paediatric population^{54–58} and reflected in our own local admissions dataset.

There are other hypotheses for the most likely contributing factors to the reduction in asthma admissions in 2020, including speculation that patients were more vigilant with their asthma prescriptions, and therefore, had improved compliance to preventative inhaled steroid medication.³¹ Given the fear about contracting COVID-19 when also diagnosed with a chronic condition⁵⁹ and the findings from the RECOVERY trial with adoption of steroids as a treatment for COVID-19,⁶⁰ it is plausible that patients improved compliance with their asthma plan, including compliance with any corticosteroid medication, during the pandemic and this contributed to a reduction in admissions during 2020.³¹ This explanation has not been widely tested, however, and it is unlikely that compliance continued throughout the entirety of 2020. An alternative theory is that many patients avoided healthcare services either due to fear that they may contract COVID-19 or a moral sense of duty to prevent overwhelming the NHS.^{59 61 62} There is weak evidence though for patients with asthma delaying presentation given there was no change in asthma mortality and no increase in acuity of admissions from the studies which looked at this metric.^{63 64} This suggests that delays in seeking healthcare assistance were not a reason for the lower admission rates observed here. Furthermore, asthma admission was chosen instead of attendance for the outcome in this study to reduce the chance that the change observed was due to a change in patient behaviour (delay in presentation). Nevertheless, it would have been beneficial for this study and any future studies to include a mechanism for measuring the extent to which a patient delayed presentation. Some studies have suggested that the low asthma admission rate may be due to low person-to-person transmission of other respiratory viruses, however, this is unlikely to completely explain the dramatic reduction in 2020 admissions observed in multiple countries as well as the UK.^{31 61 65 66}

Strengths and limitations

This study has some additional limitations including the ecological, observational design that precludes the establishment of causality. Due to the observational study design it is difficult to conclude that changes in

air pollution concentrations were wholly responsible for the reduction in asthma admissions during 2020. During the early pandemic phase several reports suggested that all-cause emergency admissions reduced, as the public responded to the ‘protect the NHS’ campaign or were too fearful of COVID-19 risks to present to hospital.⁶¹ However, acute asthma is a severe condition which requires urgent hospital management and therefore it is likely that most patients in this study accessed emergency care following an acute attack. Our findings are also consistent with those of Shah *et al*, who identified a statistically significant reduction in asthma patients who experienced an exacerbation during the lockdown period. The aggregated nature of the data collected on asthma admissions and the estimates of the population at risk means that caution must be exercised regarding any inferences at the individual level. Despite a defined population at risk, within a relatively small pollutant area, it is not known whether those admitted with asthma were exposed to the air pollutant concentrations measured. Further, asthma admissions data were aggregated to weekly time intervals due to the need for protection of patient anonymity; therefore, restricting our capability to undertake daily analysis. We were also unable to include further seasonal influences (eg, pollen exposure)⁶⁷ in the current study. Finally, we include data for ambient pollutant exposure only and recognise that adult patients with diagnosed asthma are likely to have spent a high duration of time indoors, particularly during lockdown periods. Indoor pollutants can have a significant impact on asthma conditions,⁶⁸ and it would be interesting to further investigate the potential impacts of indoor pollution on asthma admissions across the region. Further research is required to understand the respective changes in indoor and ambient pollutant concentrations during the COVID-19 pandemic and to better understand links to chronic disease outcomes among vulnerable patients. In particular, further studies should analyse admissions at the patient level to understand if pollutants have a particular impact on specific subgroups of the population with asthma which could be used to target future public health policies and guidance.

CONCLUSIONS

This study provides important insights into the effects of changes in ambient air quality on asthma exacerbations in adults during the COVID-19 pandemic in Oxford, UK. The COVID-19 national lockdown measures have led to pronounced changes in air pollutant (NO₂, PM₁₀ and PM_{2.5}) concentrations in Oxford. Our results show that adults with asthma had a significantly lower risk of unplanned hospital admission in 2020 as compared with the pre-pandemic years (2015–2019). While this study cannot attribute exact causation, our findings suggest that improvements to air quality could reduce asthma healthcare service burden. There is a need to identify which pollutants, sources and locations would be most

beneficial to target with future public health interventions to prevent severe asthma exacerbations requiring hospitalisation in adults.

Author affiliations

¹School of Geography Earth and Environmental Sciences, University of Birmingham, Birmingham, UK

²Institute of Applied Health Research, University of Birmingham, Birmingham, UK

³Institute of Microbiology and Infection, University of Birmingham, Birmingham, UK

⁴NHS Oxfordshire Clinical Commissioning Group, Oxford, UK

⁵Department of Engineering Science, University of Oxford, Oxford, UK

⁶Ricardo Energy and Environment, Didcot, UK

⁷Apertum, Oxfordshire, UK

⁸Oxfordshire County Council, Oxford, UK

⁹Oxford City Council, Oxford, UK

Twitter Ajit Singh @ajitsingh528 and Suzanne E Bartington @cleanair_doc

Acknowledgements We are grateful to staff at Oxfordshire Clinical Commissioning Group, Oxfordshire County Council and Oxford City Council for supporting this work.

Contributors SEB, FL, FDP, GNT, BS, TB, SC and GE contributed to funding acquisition for the wider OxAria study. SEB, GLM and GNT conceived and designed the current study. RA and CC acquired linked health data. TB and PA acquired air quality and meteorological data. GLM linked air quality and health data, conducted initial data analysis and drafted the initial report under the supervision of SEB, AS undertook data analysis, wrote the draft article and undertook data visualisation. All authors contributed to editing and critically revising the article and approved the final draft. SEB is guarantor of the work.

Funding This research was funded by the Natural Environment Research Council (grant no. NE/V010360/1). Its forerunning pilot project was funded by the National Institute for Health and Care Research (NIHR130095; NIHR Public Health Research).

Disclaimer The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Ethics approval Ethical approval for the OxAria study was provided by the University of Birmingham Research Ethics Committee (ERN_20-0831).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available in a public, open access repository. Air quality data are available from https://uk-air.defra.gov.uk/data/data_selector.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) licence, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Ajit Singh <http://orcid.org/0000-0003-0986-2064>

Felix Leach <http://orcid.org/0000-0001-6656-2389>

Suzanne E Bartington <http://orcid.org/0000-0002-8179-7618>

REFERENCES

- Holgate ST, Thomas M, O'Hehir RE. Chapter 7 - asthma. In: Holgate ST, Sheikh A, eds. *Middleton's allergy essentials*. Elsevier, 2017: 151–204.
- NICE. *What is the prevalence of asthma?* London, United Kingdom: National Institute for Health and Care Excellence, 2022. Available: <https://cks.nice.org.uk/topics/asthma/background-information/prevalence/>
- ALU. *Asthma UK, Asthma facts and statistics*. London, United Kingdom: Asthma + Lung UK, 2022. Available: <https://www.asthmaandlung.org.uk/media-centre/>
- IHME. *The global burden of disease study*. USA: Institute for Health Metrics and Evaluation, University of Washington, 2022. Available: <https://vizhub.healthdata.org/gbd-compare/>
- BLF. *Asthma statistics*. London, United Kingdom: British Lung Foundation, 2022. Available: <https://statistics.blf.org.uk/asthma>
- Hoffmann C, Maglakelidze M, von Schneidemesser E, et al. Asthma and COPD exacerbation in relation to outdoor air pollution in the metropolitan area of Berlin, Germany. *Respir Res* 2022;23:64.
- NICE. *Scenario: Newly-diagnosed asthma*. London, United Kingdom: National Institute for Health and Care Excellence, 2021. Available: <https://cks.nice.org.uk/topics/asthma/management/newly-diagnosed-asthma/>
- Subbarao P, Mandhane PJ, Sears MR. Asthma: epidemiology, etiology and risk factors. *Canadian Medical Association Journal* 2009;181:E181–90.
- WHO. *WHO's urban ambient air pollution database-update 2016*. Geneva: World Health Organization, 2016. Available: https://cdn.who.int/media/docs/default-source/air-quality-database/aq-d-2016/aap-database_summary_results_2016_v02.pdf?sfvrsn=384beb23_3
- WHO. *Billions of people still breathe unhealthy air: new WHO data*. Geneva, Switzerland: World Health Organization, 2022. Available: <https://www.who.int/news/item/04-04-2022-billions-of-people-still-breathe-unhealthy-air-new-who-data>
- WHO. *WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide*. Geneva, Switzerland: World Health Organization, 2021. Available: <https://apps.who.int/iris/handle/10665/345329>
- Madureira J, Brancher EA, Costa C, et al. Cardio-respiratory health effects of exposure to traffic-related air pollutants while exercising outdoors: a systematic review. *Environ Res* 2019;178:108647.
- PHE. *Health matters: air pollution*. London, United Kingdom: Public Health England, 2018. Available: <https://www.gov.uk/government/publications/health-matters-air-pollution/health-matters-air-pollution>
- PHE. *Review of interventions to improve outdoor air quality and public health*. London, United Kingdom: Public Health England, 2019. Available: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/938623/Review_of_interventions_to_improve_air_quality_March-2019-2018572.pdf
- ONS. *Road transport and air emissions*. London, United Kingdom: Office for National Statistics, 2019. Available: <https://www.ons.gov.uk/economy/environmentalaccounts/articles/roadtransportandairmissions/2019-09-16>
- Jiang X-Q, Mei X-D, Feng D. Air pollution and chronic airway diseases: what should people know and do? *J Thorac Dis* 2015;8:E31–40.
- Jacquemin B, Siroux V, Sanchez M, et al. Ambient air pollution and adult asthma incidence in six European cohorts (ESCAPE). *Environ Health Perspect* 2015;123:613–21.
- Khreis H, Kelly C, Tate J, et al. Exposure to traffic-related air pollution and risk of development of childhood asthma: a systematic review and meta-analysis. *Environment International* 2017;100:1–31.
- COMEAP. *Committee on the medical effects of air pollutants*. Chilton, United Kingdom: Centre for Radiation, Chemical and Environmental Hazards, 2022. Available: <https://www.gov.uk/government/groups/committee-on-the-medical-effects-of-air-pollutants-comeap>
- Ashworth M, Analitis A, Whitney D, et al. Spatio-temporal associations of air pollutant concentrations, GP respiratory consultations and respiratory Inhaler prescriptions: a 5-year study of primary care in the borough of Lambeth, South London. *Environ Health* 2021;20:54.
- AQEG. Report: estimation of changes in air pollution emissions, concentrations and exposure during the COVID-19 outbreak in the UK: rapid evidence review. In: *The Air Quality Expert Group, Department for Environment, Food and Rural Affairs (Defra)*. 2020. Available: https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2007010844_Estimation_of_Changes_in_Air_Pollution_During_COVID-19_outbreak_in_the_UK.pdf
- Singh A, Bartington SE, Song C, et al. Impacts of emergency health protection measures upon air quality, traffic and public health: evidence from Oxford, UK. *Environ Pollut* 2022;293:118584.

- 23 Singh A, Guo T, Bush T, *et al.* Impacts of COVID-19 lockdown on traffic flow, active travel and gaseous pollutant concentrations; implications for future emissions control measures in Oxford, UK. *Sustainability* 2022;14:16182.
- 24 Shi Z, Song C, Liu B, *et al.* Abrupt but smaller than expected changes in surface air quality attributable to COVID-19 lockdowns. *Sci Adv* 2021;7:eabd6696.
- 25 Mahato S, Pal S, Ghosh KG. Effect of lockdown amid COVID-19 pandemic on air quality of the Megacity Delhi, India. *Sci Total Environ* 2020;730:139086.
- 26 Kumar P, Hama S, Omidvarborna H, *et al.* Temporary reduction in fine particulate matter due to 'Anthropogenic emissions switch-Off' During COVID-19 Lockdown in Indian cities. *Sustain Cities Soc* 2020;62:102382.
- 27 Baldasano JM. COVID-19 lockdown effects on air quality by NO₂ in the cities of Barcelona and Madrid (Spain). *Sci Total Environ* 2020;741:140353.
- 28 Lian X, Huang J, Huang R, *et al.* Impact of city lockdown on the air quality of COVID-19-hit of Wuhan city. *Sci Total Environ* 2020;742:140556.
- 29 Liu F, Wang M, Zheng M. Effects of COVID-19 lockdown on global air quality and health. *Sci Total Environ* 2021;755:142533.
- 30 Jephcote C, Hansell AL, Adams K, *et al.* Changes in air quality during COVID-19 'lockdown' in the United Kingdom. *Environ Pollut* 2021;272:116011.
- 31 Davies GA, Alsallakh MA, Sivakumaran S, *et al.* Impact of COVID-19 lockdown on emergency asthma admissions and deaths: national interrupted time series analyses for Scotland and Wales. *Thorax* 2021;76:867–73.
- 32 Shah SA, Quint JK, Nwaru BI, *et al.* Impact of COVID-19 national lockdown on asthma exacerbations: interrupted time-series analysis of English primary care data. *Thorax* 2021;76:860–6.
- 33 Papadopoulos NG, Mathioudakis AG, Custovic A, *et al.* Childhood asthma outcomes during the COVID-19 pandemic: findings from the Pearl multi-national cohort. *Allergy* 2021;76:1765–75. 10.1111/all.14787 Available: <https://onlinelibrary.wiley.com/toc/13989995/76/6>
- 34 ONS. *Population estimates for the UK, England and Wales, Scotland and Northern Ireland: mid-2019, using April 2019 local authority district codes*. Fareham, England: Office for National Statistics, 2020. Available: <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/datasets/populationestimatesforukenglandandwalesscotlandandnorthernireland>
- 35 OP AM. *AM online projects*. Oxford, United Kingdom, 2022. Available: <https://en.climate-data.org/europe/united-kingdom/england/oxford-22/>
- 36 OCC. *Oxford Profile 2018 Key Facts Oxford City Council*. Oxford, United Kingdom, 2018. Available: https://www.oxford.gov.uk/downloads/file/5021/oxford_profile_2018
- 37 ICD. *International Statistical Classification of Diseases and Related Health Problems 10th Revision (ICD-10) Version for 2010*. International Classification of Diseases, United Nations System, 2010. Available: <https://icd.who.int/browse10/2010/en>
- 38 DEFRA. *UK air information*. Department for Environment Food & Rural Affairs, United Kingdom, 2022.
- 39 SoGE. *Daily data from the Radcliffe Observatory site in Oxford. School of Geography and Environment*. Oxford, United Kingdom: University of Oxford, 2022. Available: <https://www.geog.ox.ac.uk/research/climate/rms/daily-data.html>
- 40 STATA. *Stata 15: statistical software for data science*. 2022. Available: <https://www.stata.com/stata15/> [Accessed 10 Apr 2022].
- 41 Carslaw D. *Blog update on COVID-19 and changes in air pollution*. Ricardo, United Kingdom, Available: <https://ee.ricardo.com/news/blog-update-on-covid-19-and-changes-in-air-pollution>
- 42 Abreu P. *Air quality annual status report*. Oxford, United Kingdom: Oxford City Council, 2021. Available: https://www.oxford.gov.uk/downloads/file/7612/air_quality_annual_status_report_2020
- 43 Orellano P, Quaranta N, Reynoso J, *et al.* Effect of outdoor air pollution on asthma exacerbations in children and adults: systematic review and multilevel meta-analysis. *PLoS One* 2017;12:e0174050.
- 44 Guarneri M, Balmes JR. Outdoor air pollution and asthma. *The Lancet* 2014;383:1581–92.
- 45 Sunyer J, Spix C, Quénel P, *et al.* Urban air pollution and emergency admissions for asthma in four European cities: the APHEA project. *Thorax* 1997;52:760–5.
- 46 Muñoz X, Barreiro E, Bustamante V, *et al.* Diesel exhausts particles: their role in increasing the incidence of asthma: reviewing the evidence of a causal link. *Sci Total Environ* 2019;652:1129–38.
- 47 Wong KL, Wong WHS, Yau YS, *et al.* Asthma admission among children in Hong Kong during the first year of the COVID-19 pandemic. *Pediatr Pulmonol* 2022;57:3104–10.
- 48 Galán I, Tobías A, Banegas JR, *et al.* Short-term effects of air pollution on daily asthma emergency room admissions. *Eur Respir J* 2003;22:802–8.
- 49 Jacquemin B, Schikowski T, Carsin AE, *et al.* The role of air pollution in adult-onset asthma: a review of the current evidence. *Semin Respir Crit Care Med* 2012;33:606–19.
- 50 Tiotiu AI, Novakova P, Nedeva D, *et al.* Impact of air pollution on asthma outcomes. *Int J Environ Res Public Health* 2020;17:6212.
- 51 Kuo C-Y, Chan C-K, Wu C-Y, *et al.* The short-term effects of ambient air pollutants on childhood asthma hospitalization in Taiwan: a national study. *Int J Environ Res Public Health* 2019;16:203.
- 52 Lam HC-Y, Li AM, Chan EY-Y, *et al.* The short-term association between asthma Hospitalisations, ambient temperature, other meteorological factors and air Pollutants in Hong Kong: a time-series study. *Thorax* 2016;71:1097–109.
- 53 Onay ZR, Mavi D, Ayhan Y, *et al.* Did hospital admissions caused by respiratory infections and asthma decrease during the COVID-19 pandemic. *Medeni Med J* 2022;37:92–8.
- 54 Krivec U, Kofol Seliger A, Tursic J. COVID-19 lockdown dropped the rate of paediatric asthma admissions. *Arch Dis Child* 2020;105:809–10.
- 55 Williams TC, MacRae C, Swann OV, *et al.* Indirect effects of the COVID-19 pandemic on paediatric healthcare use and severe disease: a retrospective national cohort study. *Arch Dis Child* 2021;106:911–7.
- 56 Dann L, Fitzsimons J, Gorman KM, *et al.* Disappearing act: COVID-19 and Paediatric emergency department attendances. *Arch Dis Child* 2020;105:810–1.
- 57 Ashikkali L, Carroll W, Johnson C. The indirect impact of COVID-19 on child health. *Paediatr Child Health (Oxford)* 2020;30:430–7.
- 58 Vásquez-Hoyos P, Diaz-Rubio F, Monteverde-Fernandez N, *et al.* Reduced PICU respiratory admissions during COVID-19. *Arch Dis Child* 2021;106:808–11.
- 59 Lazzarini M, Barbi E, Apicella A, *et al.* Delayed access or provision of care in Italy resulting from fear of COVID-19. *Lancet Child Adolesc Health* 2020;4:e10–1.
- 60 The WHO Rapid Evidence Appraisal for COVID-19 Therapies (REACT) Working Group, Sterne JAC, Murthy S, *et al.* Association between administration of systemic corticosteroids and mortality among critically ill patients with COVID-19: a meta-analysis. *JAMA* 2020;324:1330.
- 61 Mulholland RH, Wood R, Stagg HR, *et al.* Impact of COVID-19 on accident and emergency Attendances and emergency and planned hospital admissions in Scotland: an interrupted time-series analysis. *J R Soc Med* 2020;113:444–53.
- 62 TFLH. *Over a third of people with lung conditions felt pressure to avoid or delay seeking treatment during lockdown London*. London, United Kingdom: Taskforce for Lung Health, 2020. Available: <https://www.blf.org.uk/taskforce/press-release/over-a-third-of-people-with-lung-conditions-felt-pressure-to-avoid-or-delay-seeking-treatment>
- 63 Chavasse R, Almario A, Christopher A, *et al.* The indirect impact of COVID-19 on children with asthma. *Arch Bronconeumol (Engl Ed)* 2020;56:768–9.
- 64 Sykes DL, Faruqi S, Holdsworth L, *et al.* Impact of COVID-19 on COPD and asthma admissions, and the pandemic from a patient's perspective. *ERJ Open Res* 2021;7:00822–2020.
- 65 Abe K, Miyawaki A, Nakamura M, *et al.* Trends in hospitalizations for asthma during the COVID-19 outbreak in Japan. *J Allergy Clin Immunol Pract* 2021;9:494–6.
- 66 Jones N. How Coronavirus lockdowns stopped flu in its tracks. *Nature* 2020. 10.1038/d41586-020-01538-8 [Epub ahead of print 21 May 2020].
- 67 Gonzalez-Barcala FJ, Aboal-Viñas J, Aira MJ, *et al.* Influence of pollen level on hospitalizations for asthma. *Arch Environ Occup Health* 2013;68:66–71.
- 68 Carazo Fernández L, Fernández Alvarez R, González-Barcala FJ, *et al.* Indoor air contaminants and their impact on respiratory pathologies. *Archivos de Bronconeumología (English Edition)* 2013;49:22–7.

Supplementary material:

Impacts of ambient air quality on acute asthma hospital admissions during the COVID-19 pandemic in Oxford City, UK; a time series study

Ajit Singh ^{a, b, †}, Gabriella L. Morley ^{c, †}, Cécile Coignet ^d, Felix Leach ^e, Francis D. Pope ^a, Neil Thomas ^b, Brian Stacey ^f, Tony Bush ^e, Stuart Cole ^g, George Economides ^g, Ruth Anderson ^g, Pedro Abreu ^h and Suzanne E. Bartington ^{b, *}

^a School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston Park Road, Birmingham, B15 2TT, UK; a.singh.2@bham.ac.uk ; f.pope@bham.ac.uk

^b Institute of Applied Health Research, University of Birmingham, Edgbaston Park Road, Birmingham, B15 2TT, UK; s.bartington@bham.ac.uk; g.n.thomas@bham.ac.uk

^c Institute of Microbiology and Infection, University of Birmingham, Edgbaston Park Road, Birmingham, B15 2TT, UK; GLM092@student.bham.ac.uk

^d NHS Oxfordshire Clinical Commissioning Group, Jubilee House, 5510 John Smith Drive, Oxford Business Park South, Oxford, OX4 2LH; cecile.coignet@nhs.net

^e Department of Engineering Science, University of Oxford, Parks Road, Oxford, OX1 3PJ, UK; felix.leach@eng.ox.ac.uk ; tony.bush@apertum.co.uk

^f Ricardo Energy & Environment, Gemini Building, Fermi Avenue, Harwell, Oxfordshire, OX11 0QR. Email: brian.stacey@ricardo.com

^g Oxfordshire County Council, County Hall, New Road, Oxford, OX1 1ND, UK; stuart.cole@oxfordshire.gov.uk; George.economides@oxfordshire.gov.uk; ruth.anderson@oxfordshire.gov.uk

^h Oxford City Council, Town Hall, St Aldate's, Oxford, OX1 1BX; pabreu@oxford.gov.uk

†Joint first authors

*Correspondence to: s.bartington@bham.ac.uk (S.E.B)



Figure S1 Map of Oxford postcode districts illustrating Oxford City in OX1, OX2, OX3 and OX4. (Source, https://en.wikipedia.org/wiki/File:OX_postcode_area_map.svg).




Figure S2 Correlation between daily mean air pollutant concentrations and meteorological variables during **a)** Covid-19 pandemic year (Corr₂₀₂₀) and **b)** pre-pandemic years (Corr₂₀₁₅₋₂₀₁₉) in Oxford. Here, RH and Temp denote relative humidity and temperature, respectively.




Figure S3 Time series of monthly adult asthma admissions, where the shaded lines represent the smooth fit at the 95% confidence interval.

Table S1 Incidence of adult hospital admissions for acute asthma in 2020 compared with pre-pandemic years (2015 – 2019).

Period	Incidence rate	IRR	95% CI	p value
Pre-pandemic (2015 – 2019) mean	78.29	Baseline		
2020	45.70	0.58	0.42 – 0.81	<0.001

*per 100,000 population

Table S2 Pearson correlation (r) matrix between air pollutant concentrations & meteorology and adult asthma admissions in 2020 versus 2015 – 2019.

Variables	NO ₂	PM _{2.5}	PM ₁₀	Temp	RH
Asthma admission 2015 – 2019	0.31	0.27	0.22	0.34	0.30
Asthma admission 2020	0.22	0.16	0.20	-0.38	0.14